Published in final edited form as:

Adv Enzymol Relat Areas Mol Biol. 2011 ; 77: 1–60.

STRUCTURE AND MECHANISM OF RND-TYPE MULTIDRUG EFFLUX PUMPS

HIROSHI NIKAIDO

Department of Molecular and Cell Biology, Barker Hall, University of California, Berkeley, California 94720-3202

I. Introduction

It has been known for a long time that Gram-negative bacteria are usually much more resistant to the actions of antibiotics and chemotherapeutic agents than the Gram-positive bacteria. The cells of the former group are surrounded by an extra membrane layer, the outer membrane (OM), and it was suspected that this additional membrane layer acted as a general barrier for the influx of agents. Indeed, the nonspecific diffusion channels of OM, the porins, limit the influx of small hydrophilic agents, because their channels are quite narrow (7 by 11 Å in the *Escherichia coli* OmpF porin) (1,2). In addition, the lipid bilayer domain of the OM is unusual in its extreme asymmetry by having the outer leaflet composed nearly exclusively of lipopolysaccharides containing only saturated fatty acid chains (3), and the very low fluidity of this outer leaflet decreases the spontaneous permeation rates of hydrophobic probe molecules nearly by two orders of magnitude, when compared with the conventional phospholipid bilayers containing many unsaturated fatty acid residues (4). Nevertheless, when the permeability coefficients of β -lactams across *E. coli* OM were measured, even the slowest-penetrating compounds were found to equilibrate across the outer membrane usually within a minute, i.e. in a time span that is much shorter than the generation time of *E. coli* (5), in part because the surface-to-volume ratio is very high in these small bacterial cells. It was thus clear that the OM barrier alone cannot generate significant levels of resistance, and another major process must work synergistically with this barrier (5). With β -lactams, the ubiquitous periplasmic β -lactamases can fulfill this role. However, with other classes of antibiotics, the nature of this second process remained obscure, until the constitutive multidrug efflux pumps were found to be widespread in Gram-negative bacteria and acted synergistically with the OM barrier (6,7).

Gram-negative bacteria usually contains multidrug efflux pumps belonging to several families, such as ABC (ATP-binding cassette), SMR (small multidrug resistance), MFS (major facilitator superfamily), MATE (multiple antibiotic and toxin extrusion), and RND (Resistance-Nodulation-Division) (8-10). Among these, most are pumps located in the cytoplasmic membrane, and must pump out drugs rapidly into the periplasm, because the drugs can penetrate back into cytosol frequently by spontaneous diffusion. Only the RND pumps (and a few exceptional pumps belonging to other families) exist in a tripartite form traversing both the OM and the inner membrane, in a manner first suggested by Wandersman and coworkers for a protein-secreting apparatus of Gram-negative bacteria (11). This complex involves, in addition to the RND pump protein located in the inner membrane, an outer membrane channel such as TolC of *E. coli* (belonging to OMF (outer membrane factor) family (12)), and a periplasmic adaptor protein (belonging to the MFP (membrane fusion protein) family (13)). As shown in Fig. 1, this construction allows the bacteria to pump out drug molecules directly into the external medium. This is a huge advantage for bacteria, because the drug in the medium has to cross the low permeability OM in order to re-enter the cells, in contrast to the drug molecules in the periplasm that can

penetrate easily the high permeability inner membrane. Thus the tripartite RND pumps are expected to produce drug resistance very effectively by working synergistically with the outer membrane barrier (6,7).

Wild-type strains of most Gram-negative bacteria are resistant to most lipophilic antibiotics (for *E. coli* they include penicillin G, oxacillin, cloxacillin, nafcillin, macrolides, novobiocin, linezolid, and fusidic acid), and this “intrinsic resistance” was often thought to be caused by the exclusion of drugs by the outer membrane barrier. Indeed breaching the outer membrane barrier does sensitize *E. coli* cells to the drugs just mentioned (14). However, the inactivation of the major and constitutively expressed RND pump AcrB also makes the bacteria nearly completely susceptible to these agents (the minimal inhibitory concentration [MIC] of a lipophilic penicillin, cloxacillin, goes down from 512 µg/ml in the wild type to only 2 µg/ml (15)) even in the presence of the intact outer membrane barrier. Thus the characteristic intrinsic resistance of gram-negative bacteria owes as much to the RND pumps as to the outer membrane barrier.

Since some tripartite pumps give resistance to drugs that cannot penetrate across the inner membrane, such as the dianionic compound carbenicillin, they were proposed to be able to capture drug molecules from the periplasm (6,7). Finally, in order to produce significant levels of drug resistance, the drugs that were pumped out into the periplasm by the simple pumps located in the inner membrane need to be captured and then extruded across the OM into the medium by tripartite RND pumps (16,17). For example, in *P. aeruginosa*, the expression of a simple tetracycline pump TetA produces only a very modest degree of resistance (MIC = 32 µg/ml) in the absence of the main tripartite pump MexB-MexA-OprM, but MIC is raised up to 512 µg/ml if the tripartite pump in addition is expressed at the normal, constitutive level. That the high resistance is not due to the pumping of tetracycline by the tripartite pump alone is seen from the fact that the tripartite pump, in the absence of TetA, produces an MIC of only 4 µg/ml (16). Thus these two types of pumps appear to act truly in a synergistic manner.

The first example of an RND pump that functions as a multidrug exporter was identified in 1993 (18); this was AcrB (called AcrE in that paper) in *E. coli*. The immediately upstream gene, *acrA*, was known to be involved in acriflavine resistance, but the mechanism was thought to be the decreased permeability of the envelope through the global chemical modification of its components. We showed that acriflavine penetrates rapidly across OM in both the wild-type and *acrAB*-inactivated strain, and yet it accumulates at a much higher level in the cells of the latter. We also noted the similarity of AcrB sequence to CzcA, a divalent cation efflux pump, and proposed that AcrB functions by actively pumping out acriflavine, not by strengthening the passive permeability barrier of OM. The extremely wide substrate specificity of this pump was indicated by the fact that inactivation of the *acrAB* genes made *E. coli* hypersusceptible to not only to several dyes, but also to detergents (such as SDS, Triton X-100, and bile salts) and to a wide range of antibiotics, including macrolides, β-lactams, tetracycline, chloramphenicol, fusidic acid, and novobiocin (but not aminoglycosides)(Fig. 2).

At about the same time, Poole and coworkers (19) found that *Pseudomonas aeruginosa* mutants capable of growth in the presence of an iron chelator, α,α'-dipyridyl, overproduced an outer membrane protein (OprM, called OprK in the paper), and that the *oprM* gene was a part of an operon with upstream genes *mexAB* that are homologous to *acrAB*. Indeed the inactivation of these genes made the organism hypersusceptible to chloramphenicol, tetracycline, ciprofloxacin, and streptomycin. Meanwhile our studies on clinical isolates of *P. aeruginosa* whose β-lactam resistance could not be explained fully by the combination of OM barrier and the periplasmic β-lactamase showed that unidentified multidrug efflux

pump(s) are overexpressed in these strains, leading to simultaneous resistance to a large number of agents (20,21). The most likely candidate for this pump was found to be the MexAB-OprM, the major constitutive RND multidrug efflux transporter of *P. aeruginosa*, through a collaborative with the Poole laboratory (22). MexAB-OprM extrudes not only fluoroquinolones, chloramphenicol, and tetracycline, but also novobiocin and a number of β -lactam antibiotics. These studies also gave the first indication that the pump functions as a tripartite complex, as *mexA*, *mexB*, and *oprM* genes comprised a single operon; in *E. coli*, the outer membrane channel TolC is coded elsewhere on the chromosome (23). Importantly, each of these three component proteins is essential for drug efflux, and the absence of even one component makes the entire complex totally nonfunctional (18,19,24). To this day, AcrB and MexB have been the most intensively studied RND drug transporters in bacteria, and have served as the prototype for biochemical and structural studies of such pumps.

Some of the observations made in this early period were important in formulating our concepts on the function of RND drug efflux transporters. 1. Some β -lactam compounds, such as carbenicillin and ceftriaxone, which contain multiple charged groups and therefore cannot diffuse across the inner membrane, as verified experimentally, are nevertheless very good substrates of MexABOprM (21) and AcrABTolC (15). This observation indicates that the RND pumps are capable of capturing substrates from the periplasm (6,7,21). The periplasmic capture also fits with the observation that the same pump can catalyze the efflux of substrates with diverse charges – uncharged, anionic, cationic, or zwitterionic (7). Export of these substrates from cytosol will generate different effects on membrane potential, which may be difficult to deal with. 2. Although the pumps often handle a very wide range of substrates, they all seem to have a significant lipophilic portion (7,15). This observation led to a hypothesis that many substrates become partially partitioned, from the periplasm, into the outer leaflet of the inner membrane (see Fig. 1), before being captured by the pump (7). This association with the membrane surface would involve a energetic cost in terms of decreased entropy, but a similar phenomenon is well known to occur with the interaction of amphiphilic anesthetics with lipid bilayers (25). An alternative interpretation of the requirement of lipophilic domain in the ligands, however, might be that hydrophobic interaction plays a predominant role in the binding of substrates at the binding pocket of the pump (see section IV).

Finally it should be mentioned that the tripartite architecture is not completely limited to RND pumps. In *E. coli*, an MFS efflux pump EmrB is known to form a tripartite structure together with the periplasmic adaptor EmrA and the outer membrane channel TolC (26), and to pump out weakly acidic or largely uncharged substrates, such as the proton uncoupler carbonylcyanide *m*-chlorophenylhydrazine (CCCP), nalidixic acid, and thiolactomycin (27). An ABC efflux pump, MacB, pumps out various macrolides with the periplasmic adaptor MacA, as well as the OM channel TolC (28). Although MacA has been crystallized (29), and an elegant proteoliposome reconstitution of MacB has been achieved (30), the totality of our knowledge on these non-RND systems cannot be compared with that on the RND systems, and we will concentrate on the RND systems in this review.

Recent reviews deal with various aspects of bacterial multidrug efflux: in addition to the comprehensive reviews cited earlier (9,10), there are reviews emphasizing different aspects, such as multidrug resistance (31,32), structure (33-35), and mechanisms (36). We will not discuss in this review the regulation of pump production (which was reviewed recently [37]) or the clinical microbiology of the pumps, as they are outside the scope of this review. We will also not discuss in detail the outer membrane channels, which will be dealt with by Dr. Ben Luisi in this volume, nor the periplasmic adaptor proteins which are reviewed recently by H. I. Zgurskaya and associates (38).

II. Biochemical and Genetic Studies

A. The Range of Substrates

Comparison of MIC values between the wild type and *ΔacrAB* strains (7,18,24) showed that AcrB of *E. coli* can handle a very wide range of compounds. These include cationic dyes such as acriflavine, crystal violet, ethidium bromide, and rhodamine 6G; antibiotics such as penicillins, cephalosporins, fluoroquinolones, macrolides, chloramphenicol, tetracyclines, novobiocin, fusidic acid, oxazolidinones, and rifampicin; detergents such as Triton X-100, sodium dodecylsulfate, and bile acids. The range extends even to simple solvents, such as pentane and cyclohexane (39,40)(Fig. 2). It was obvious that there is no structural similarity between most of these compounds. Furthermore, the charge states of the ligands are diverse, some nonionic, some anionic, some cationic, and others containing multiple ionizable groups (see Fig. 2). However, there was one common feature that the substrates were either relatively lipophilic or at least contained lipophilic domains (7). This led to the hypothesis that the substrates first interact with the membrane lipid bilayer (7), and become captured either from within the bilayer or from the bilayer-aqueous interface (7)(Fig. 1). However, alternative interpretations are possible, as mentioned earlier.

The capture of substrates from the periplasm or a location that is in rapid equilibrium with periplasm was already described in Introduction. In this connection, it is interesting that the general inhibitor of *P. aeruginosa* and *E. coli* RND pumps, MC-207,110 (phenylalanyl-arginyl- β -naphthylamide)(41), which apparently is a favored substrate of the pumps, affected the carbenicillin MIC only to a small extent (41). Thus the addition of this inhibitor decreased the MICs of drugs such as erythromycin and levofloxacin to almost exactly the same degree as the genetic inactivation of the MexB-MexA-OprM pump. In contrast the inhibitor produced a decrease of only fourfold in carbenicillin MIC, although the genetic inactivation decreased it 512-fold. A similar discrepancy, i.e. strong decrease of MIC by genetic inactivation vs. little decrease in the presence of the inhibitor, was also found for tetracycline and ethidium bromide (41). Since carbenicillin must be captured exclusively from periplasm, these results were sometimes interpreted to mean that this particular inhibitor acts more effectively for drugs that are predominantly captured in the cytosol. However, currently we are more inclined to assume that AcrB captures all drugs exclusively from the periplasm (see section IV). If so, these results are perhaps caused by the binding of various substrates to different subdomains of the large binding pocket of the transporter.

B. The Tripartite Assembly

As described in Introduction, the drug efflux function of AcrB requires the simultaneous presence of the periplasmic adaptor protein AcrA as well as the outer membrane channel TolC. Chemical cross-linking showed that the adaptor protein AcrA can be cross-linked to the RND transporter AcrB in intact cells (42), and more recently the whole trimeric complex was isolated from gently lysed *E. coli* cells without cross-linking (43). Interestingly the association of the component proteins was stabilized by the presence of substrates (43).

When the crystallographic structures of the transporters AcrB (44) and most recently MexB (45), the adaptor proteins MexA (46,47) and AcrA (48), and the outer membrane channel TolC (49) and OprM (50) became available (Fig. 3), it became possible to speculate on the details of interaction among these components. Space does not permit the author to discuss the details of these structures, but some of the prominent features will be listed. 1. AcrB (18) and also other RND pumps (51) were expected, from the amino acid sequences, to contain 12 transmembrane helices, with two extremely large periplasmic domains between helix 1 and 2 and between helix 7 and 8. Indeed one can estimate that the size of the periplasmic domain would almost surpass that of the transmembrane domain, a prediction borne out with

the x-ray crystallographic structures (44,45). What the amino acid sequence did not tell us was that these pumps existed as trimers, and that the tip of the large periplasmic domain, often called “TolC-docking domain,” had the dimension that would fit the internal end of the TolC channel (44). 2. The adaptor proteins had the very elongated shape as predicted from physicochemical studies (52). This shape was created by the very long α -helical hairpin structure on top of another elongated domain, the “lipoyl” domain that usually carries bound lipoic acid in various enzymes. Finally this structure was followed by a β -barrel domain. (The structure of the fourth domain at the opposite end of the molecule from the α -hairpin, called “membrane-proximal domain,” was solved recently (53) and will be discussed below). All these separate domains create a flexible structure, again as predicted by site-directed spin labeling studies (54). 3. The structure of TolC (49) was perhaps the most spectacular. It is a trimer containing the OM-traversing channel of a β -barrel, with each protomer contributing four transmembrane β -strands of the 12-stranded barrel. Even more interestingly, the channel in the β -barrel is continued by a long “helical tunnel,” approximately 130 Å long, that are made of long α -helical hairpins and that extends deep into the periplasm.

These structures obviously suggest that the tip of the periplasmic domain of AcrB, or the TolC-docking domain, contacts the end of the TolC periplasmic tunnel. This contact was also confirmed by introducing cysteine residues to the “top” of AcrB and the “bottom” of TolC channel, and observing the formation of disulfide linkage (55). Although this end of the TolC tunnel is closed in the first structure published, there are now structures in partially open states (56), and the function of TolC during the drug export process will be discussed in detail by B. Luisi in this volume.

Modeling efforts for the trimeric complex were also helped by studies that involved alteration of the structure of component proteins. For example, we found that the adaptor protein of another *E. coli* tripartite system, YhiU (also called MdtE), could not replace AcrA in the formation of tripartite system containing AcrB, and tested chimeric AcrA molecules containing different parts of YhiU (57). Within the 398 residue sequence of AcrA, residues 1-290 could be substituted by the YhiU sequence, but substitution of residues 290-357 produced a protein inactive in forming a functional tripartite complex with AcrB, suggesting that the C-terminal portion of the adaptor proteins was important in interaction with the RND pump proteins. Study of component interaction in vitro by the use of titration calorimetry gave also results (58) consistent with the chimeric AcrA data. Thus both the full length AcrA and its C-terminal half (residues 172-397) associated with AcrB, but the N-terminal half (residues 24-172) or the central fragment (residues 45-315) showed no evidence of interaction with AcrB. In this connection, it must be mentioned again that until very recently the published x-ray crystallographic structures of the adaptor proteins were incomplete, roughly corresponding to the central fragment just mentioned, thus lacking the most important C-terminal part that interacts with the RND transporter. In another approach, it was noted that AcrA can function with MexB and TolC, but the full spectrum of drug resistance found in MexAB-OprM system was not attained (59). Random mutagenesis of AcrA resulted in the isolation of mutants that works better with MexB, and some of them were located in the β -barrel domain.

As regards the interaction between AcrA and TolC, Koronakis and coworkers (60) were able to carry out a fine resolution analysis by cross-linking Cys residues of AcrA, introduced by site-directed mutagenesis, with amino groups on the surface of TolC, establishing how only one side of the α -helical coiled coil hairpin in AcrA interacts tightly with the intramolecular groove in the TolC helical bundle.

Hypothetical models of the tripartite assembly were proposed in the past. In one model (46), trimers of MexB and OprM are surrounded by either twelve or six MexA proteins. In another (61), nine AcrA protomers circle the trimers of AcrB and TolC. However there were some major problems in building these models. 1. The oligomeric state of the adaptor proteins is uncertain. Cross-linking studies showed that overexpressed AcrA existed as dimers and trimers in the cell (42), whereas AcrA lacking the N-terminal lipid modification behaved as monomers when isolated (52). Adaptor proteins have been crystallized as giant complexes containing more than ten protomers (46-48). These are obviously crystallographic artifacts that do not tell us anything about the physiological state of their self-association. 2. As mentioned above, the structure of the adaptor proteins used in modeling lacked the extreme N-terminus and a large C-terminal domain, the portion known to be essential for the interaction between the adaptor and the RND pump protein. 3. The stoichiometric ratio between the adaptor and the pump protein (or TolC/OprM) within the tripartite complex was not known.

A giant stride was recently made by the Koronakis group in this area (53). They have refined the data previously obtained for the MexA crystal (47) and succeeded in obtaining the structure of the hitherto missing domain containing the N-terminus and the large C-terminal segment. This is now known to form the fourth domain or the membrane-proximal domain of the adaptor, containing an α/β structure with a characteristic concave surface (Fig. 3). By using this MexA structure as a template, they built the complete structure of AcrA. Finally, by applying the method used earlier to study the interaction between TolC and AcrA (60), i.e. site-directed insertion of Cys followed by cross-linking, they were able to determine precisely where defined portions of AcrA fits into which surface of AcrB. Interestingly, in this cross-linking study in intact cells, AcrA seems to interact only with one contiguous stretches of AcrB within the periplasmic domain, thus suggesting a 1:1 stoichiometric ratio between these two proteins. Based on these observations, the “data-driven” model of AcrAB complex could be built, which is shown as Fig. 4.

Another significant advance in the study of component interaction was the in vitro examination of association between MexA and OprM (62). Such studies are difficult because both are membrane proteins (although some adaptor proteins appear to work in intact cells without the lipid modification at the N-terminus (52)), and the adaptor inserted into one membrane must interact with the OM channel inserted into another membrane (see Fig. 1). Reffay and coworkers (62) used an ingenious system of L₃ phase (sometimes called sponge phase) surfactants, where the distance between neighboring bilayers can be adjusted. In this system, the lateral diffusion rates of MexA become greatly diminished when it forms a complex with the OprM trimer, a much larger protein, in another bilayer. The MexA immobilization as a function of OprM concentration showed that the MexA monomer to OprM trimer stoichiometry was strongly dependent on pH (a finding reminiscent of an earlier study of AcrA conformation (54)), and ranged from 2 at pH 7.5 to 6 around pH 6. The former value is more or less consistent with the model of Symmons (53), but it is not yet clear how the second adaptor protein can be accommodated in the final tripartite complex. We note that these ratios are consistent with the abundance of MexA, MexB, and OprM protomers in the cell reported by the Nakae group (63), namely 2: 1: 1. However, in *E. coli* AcrA appears much more abundant, Tikhonova and Zgurskaya (43) reporting the abundance of 10: 1: 3 for AcrA, AcrB, and TolC protomers; this ratio, however, may reflect the fact that AcrA and TolC function also for other systems, AcrA for example for AcrD (64) and TolC not only for all other RND systems but also for some non-RND systems such as EmrAB and MacAB.

One observation that may be pertinent here is that some RND systems apparently contain a pair of adaptor proteins. For example, the TriABC-OpmH system of *P. aeruginosa*, which

pumps out triclosan, contains two adaptor protein genes (*triA* and *triB*) within the operon, and studies clearly showed that both genes are needed for triclosan resistance (65). The ZrpADBC system of *Serratia* sp. appears to be yet another example of such a system containing a pair of adaptor proteins (66). If we assume that the pump proteins in these systems function as trimers according to the functionally rotating mechanism (described in section III.C), each protomer of the pump is likely to contact a pair of the adaptor proteins, giving rise to the 2:1:1 ratio between the adaptor, pump, and OM channel proteins. However, an alternative arrangement of 1 : 1 : 1 stoichiometry is still possible, producing a situation that different adaptor-pump complexes would behave similarly to the different pump protomers in the MdtBC-type complexes (see section IV.E).

C. Kinetic Behavior of the AcrB Pump

In order to understand the behavior of the pump in intact cells, it is obviously essential to know its kinetic constants. Hints on the binding affinity of various substrates were obtained by using them as competitive inhibitors in the reconstitution assay of AcrB (67) (see below). In this assay, which depended on the export of fluorescent phospholipids by the purified AcrB protein, taurocholate, a conjugated bile salt, inhibited the reaction most strongly (presumably by competing as a substrate), the 50% inhibition occurring at 15 μ M. In contrast, antibiotics were less efficient inhibitors, and cloxacillin and erythromycin caused 50% inhibition only around 100 μ M. These results are important in showing that the conjugated bile salts are probably the preferred, and most likely the natural, substrates of the AcrB pump of *E. coli*, an organism that must live in an environment full of these membrane-disrupting detergents. In the same study (67), proton movement accompanying the pumping of substrates was also measured. Although a single concentration of substrates (0.2 mM) was used, again the conjugated bile salts were the most effective in increasing the proton flux. However, these assays do not measure substrate binding per se, and we do not know the true affinity of each substrate to the pump. (We should also mention that running similar assays in intact cells is difficult because both the substrate and the competitive inhibitor must diffuse across the outer membrane before binding to the pump.)

There was a recent attempt to measure the binding of substrates to purified AcrB protein by the use of fluorescence polarization (68). Because binding of fluorescent substrates to a large AcrB trimer is expected to slow the tumbling of these molecules, this method offers advantages. The apparent dissociation constants reported ranged from 5.5 μ M for rhodamine 6G to 74 μ M for ciprofloxacin. However, the method does not give reliable information on binding stoichiometry, and because the authors did not use, as controls, mutant AcrB proteins that are altered in substrate binding, it is unclear whether the binding occurred to the true binding sites, or to some extraneous pockets that bind lipophilic molecules in a non-specific manner.

The efflux of fluorescent dyes from cells after their energization was followed in several studies. Typically, cells are preloaded with dyes that become fluorescent only within the membrane or the cytosol. The preloading requires deenergization of the RND pump by proton conductor such as CCCP (carbonyl cyanide *m*-chlorophenylhydrazone), and re-energization of the cells is done by adding compounds such as glucose or formate. If everything goes perfectly, the time-dependent decrease in fluorescence intensity should follow the integrated form of the Michaelis-Menten equation, which should give us the kinetic constants K_m and V_{max} . In the first paper reporting the use of one of these probes (69), the results were not convincing because the efflux rates from *P. aeruginosa* strains either producing or not producing the MexB-MexA-OrpM efflux complex showed little difference, presumably because deenergization with cyanide could not be reversed rapidly. However, efflux assays were successfully performed since then as a qualitative assay, for example with NPN (*N*-phenyl-*n*-aphthylamine) and *E. coli* (41), and with DASPEI (2-[4-

dimethylamino]styryl-*N*-ethylpyridinium iodide) and *E. coli* (70). No effort to use the efflux curve for quantitative analysis of the transport process was reported. We have tried to do this quantitative analysis with a fluorescent dye, Nile Red using *E. coli* (J. Bohnert and H. Nikaido, unpublished results). However, the system is extremely complex, because CCCP used in the dye-loading period may still persist in the cells at least in the beginning, and the full membrane energization with carbon sources will take a few seconds, thus making the analysis of the early phase of the efflux curve quite difficult.

An early study that attempted to determine the kinetic constants of efflux was that of Narita et al. (63). Here the authors calculated the rate of ethidium efflux from the difference in the ethidium influx rates between efflux-incompetent and competent cells of *P. aeruginosa*, and reported a V_{max} of 100 nmol/s/(3×10^8 cells), said to correspond to a turnover number of 500 s^{-1} . However, the authors did not show how these numbers were arrived at. In fact, inspection of the data shows that the efflux rate plotted against intracellular ethidium concentration was essentially linear, and little sign of saturation can be observed. Thus these conclusions are not convincing.

Recently we had the first success with the estimation of kinetic constants of the AcrB pump (71). With attempts using intact cells, the difficulty is always with the estimation of the substrate concentration within the periplasm, where it is presumably captured by the pump. We solved this problem by using β -lactams as substrates. Since β -lactams are hydrolyzed by the periplasmic β -lactamase, and since we know the K_m and V_{max} of the enzyme, we can calculate the periplasmic concentration of β -lactams, C_p , from the Michaelis-Menten equation if we determine spectrophotometrically the β -lactam hydrolysis rate by intact cells. Once the periplasmic concentration C_p is known, the rate at which the β -lactams cross the outer membrane by simple diffusion through porin channels (V_{in}) can be calculated from the permeability coefficient of the OM, because it is proportional to the concentration difference of the β -lactam between the external medium and the periplasm. The efflux rate V_e is then the difference between the influx rate V_{in} and the hydrolysis rate V_h , or $V_{in} - V_h$. This approach was successfully applied initially to the influx, efflux, and hydrolysis of a cephalosporin, nitrocefin, and we found that the efflux occurs with the Michaelis-Menten saturation kinetics, with the apparent K_m of about 5 μM (Fig. 5A). There was little sign of positive cooperativity, a result that is interesting in view of the functional rotating mechanism of the pump discussed later in this review. A turnover number of about 10 s^{-1} was calculated. The low turnover number was as predicted in early studies, on the ground that the tripartite pumps may have to deal only with the small number of drug molecules that have trickled through the effective outer membrane barrier (7).

Interestingly, when the assay was extended to cephalosporins that were clinically useful, such as cephalothin, cefamandole, and cephaloridine, kinetics with strong cooperativity was observed (Fig. 5B). Although it is not clear why such kinetics were not observed with nitrocefin, we note that nitrocefin has a much lower K_m than these compounds. Interestingly, compounds showing strong cooperativity have also high V_{max} values, so that the ratio of V_{max} to $K_{0.5}$ (the substrate concentration giving a half-maximal rate; K_m in the case of non-cooperative kinetics) did not change much for the four compounds tested (range: $0.5\text{--}1.9 \times 10^{-2} \text{ cm}^3 \text{ mg}^{-1} \text{ s}^{-1}$) although the V_{max} for cephaloridine was 80 times higher than that for nitrocefin. The results on cephaloridine merit close attention for several reasons. 1. We have shown that cephaloridine cannot diffuse across the cytoplasmic membrane (15,21). Thus these results represent the kinetics of drug export initiated by the periplasmic capture of the drug. 2. The criterion for evaluation of the range of substrates for an efflux pump was usually the comparison of MICs (minimum inhibitory concentrations) of the drug in the wild type cells and in the pump gene deletion mutants. By this criterion, the presence or absence of AcrAB made little difference to cephaloridine MICs (15,70), and thus this compound was

thought to be either a nonsubstrate or at best a marginal substrate for the pump. However, the detailed kinetic analysis showed that cephaloridine may be pumped out very rapidly, with the turnover number approaching $1,000\text{ s}^{-1}$ (71), although the $K_{0.5}$ was very high (288 μM). Thus the MIC comparison can be a poor, and sometimes a misleading indicator of the pump function. In this case, the AcrAB function made little difference to the MIC values, because the bacteria are killed at β -lactam concentrations usually below 5 μM (73), and in this concentration range, the efflux is insignificant because of the high $K_{0.5}$ value and especially because of the strongly sigmoidal kinetics. The high turnover number for some substrates explain the observation that AcrAB can pump out simple solvents (39,40), as these compounds must traverse the OM bilayer rapidly because they are small and uncharged, and as they must be pumped out very rapidly by the efflux pump in order to produce resistance.

Interestingly, one cephalosporin, cefazolin, showed no sign of getting pumped out by the AcrAB pump (Fig. 5C). This compound has very hydrophilic substituents (a tetrazole and a thiadiazole) at 7- and 3-positions, respectively, and presumably the binding of the substrates to the AcrB binding site requires the presence of lipophilic moiety, preferably an aromatic ring, in at least one of these positions (see ref. 15).

Recently, binding of 24-, 25-, or 27-hydroxysterols to Niemann-Pick C1 protein (NPC1), a member of the RND family involved in cholesterol trafficking in vertebrate cells (74), was reported (75). The binding took place in spite of the presence of a high concentration (1 %) of the detergent Nonidet P40, and the half-saturation was reached with about 0.1 μM of 25-hydroxysterol. 7-, 19-, or 20-Hydroxysterol did not bind to this protein. In comparison with bacterial RND pumps, the NPC1 protein has an N-terminal extension consisting of an extramembranous domain of about 240 residues and an additional transmembrane segment. This extramembranous domain is secreted from cultured cells as a soluble dimer, which bound 25-hydroxysterol with a strong affinity ($K_D = 10\text{ nM}$) (76). Since this domain is uniquely present in NPC1, it is unclear whether these beautiful results would help us in our effort to understand the ligand binding process in efflux transporters (see section IV.B).

D. Genetic Studies of AcrB and Its Relatives

To identify domains responsible for pump functions, large segments of genes were replaced with sequences coming from another homolog, producing chimeric genes. For example, Elkins and Nikaido (77) used *E. coli* AcrB, which gives a strong resistance to ciprofloxacin, novobiocin, fusidic acid, erythromycin, and taurocholate, and AcrD, which generates no resistance to these agents except a modest one against taurocholate. Replacing both of the large periplasmic loops of AcrD with those from AcrB created a transporter pumping out all of the tested AcrB substrates efficiently. Replacing loops of AcrB with those from AcrD created a transporter that behaved like AcrD in terms of its substrate specificity. Finally, replacing the transmembrane regions of AcrD with corresponding sequences from AcrB did not alter the substrate specificity of the transporter. These results show that the substrate specificity is determined nearly entirely by the periplasmic domain (and that the critical binding of substrates presumably occurs here) and the transmembrane domain likely plays a role only in energy coupling. That the exchange of transmembrane domain does not alter the substrate specificity was also confirmed later by using *P. aeruginosa* MexB and MexY (78).

Zgurskaya and coworkers (79) used AcrB and *P. aeruginosa* MexB. The results of chimeric exchange suggested that the substrate specificity was determined largely by the second external loop (called PC2 in AcrB) of these transporters, containing residues 612 through 849, a conclusion consistent with the location of the substrate-binding pocket identified in the asymmetric trimer crystal of AcrB (see section IV.A). (This study was recently refined by the use of site-directed mutagenesis (80), which pinpointed the difference in macrolide

extrusion activity to residue 616, which is Gly in AcrB and Asn in MexB). They also showed that replacing the region coding for the transmembrane helices 8 through 12 of AcrB with the MexB sequence produced a somewhat MexB-like pattern for a few drugs (lower efflux of ethidium bromide and lincomycin, and higher efflux of cinnoxacin); currently it is difficult to explain these results in terms of structure, although the elongation of helix 8 into the periplasmic domain appears crucial in the conformational change that leads to drug extrusion (section III.C). Finally, this group showed that the N-terminal periplasmic loop (PN1 and PN2) and the first part of the C-terminal loop (PC1) essentially determine the interaction of the RND pumps with their cognate periplasmic adaptor proteins, a conclusion that is in complete agreement with the recent study based on the crystal structures (see Fig. 4).

Use of point mutations could in principle allow us to produce more detailed knowledge on the binding and export of substrates. In an approach pioneered by Lomovskaya's laboratory, random mutants of *P. aeruginosa* MexD that acquired the ability to extrude carbenicillin, which is not handled by the wild-type MexD, were isolated (81). All mutants mapped to the periplasmic domain, none to the transmembrane domain, further supporting the conclusion mentioned above. Among the residues identified, Phe608 (corresponding to Phe610 of AcrB) is a part of the substrate-binding pocket more recently identified (see below). Some others (Gln34, Gln89 [corresponding to Glu 89 of AcrB], Asn673[Thr676]) are on the walls of the large lateral cleft of the periplasmic domain, which will be discussed below (section IV.B) as a potential site of periplasmic entry for substrates. These residues are shown as space-filling models in the AcrB structure (Fig. 6).

Bohnert et al. (75) used an *E. coli* strain that lacked both AcrB and AcrF and instead overproduced another RND pump, YhiV. A spontaneous mutant obtained after repeated exposure to levofloxacin was shown to owe its increased resistance to a change of an aliphatic Val610 residue of the YhiV into an aromatic Phe residue. This mutant pump has an altered specificity. It produces a stronger resistance to relatively small, aromatic compounds such as levofloxacin, linezolid and tetracycline, but the resistance to a non-aromatic, bulky macrolides becomes weaker than in the parent protein. This residue, which corresponds to Val612 of AcrB, is seen to be a part of the substrate-binding pocket in the asymmetric AcrB trimer structure that will be discussed later (section IV.A).

Middlemiss and Poole (83) used a different approach and carried out an in vitro random mutagenesis of the *mexB* gene from *P. aeruginosa*. This is a more comprehensive approach, but is expected to generate more "noise." Indeed the group of mutants that decreased the level of resistance to most drugs included the presumed proton relay mutants in Asp407 and Asp408 of the transmembrane domain (see section III.B), or Gly220 mutant in the "peg" that is inserted into the periplasmic domain of the neighboring protomer. The mutants that are significantly altered in the substrate specificity, in contrast, often contained alterations in the periplasmic domain, as expected. Among these, alterations of Ala618 and Arg716 (corresponding to Ala618 and Arg717 of AcrB) occur on the opposing walls of the large lateral cleft. However, there were several mutants in the transmembrane domain, and their phenotypes remain to be explained.

Point mutations can also be introduced into the transmembrane domain by site-directed mutagenesis. In 1999, inspection of amino acid sequence of an RND-family toxic cation efflux pump CzcA of *Ralstonia* sp. and other RND pumps showed that there are several conserved charged residues in the transmembrane domain, including Asp402, Asp408, and Glu415 (84). Changing these residues into non-acidic residues abolished the cation pumping activity. This was followed up a few years later by the site-directed mutagenesis study of MexB (85). In MexB, the Asp402 and Asp408 of CzcA are replaced by a consecutive pair of

aspartate residues, Asp407 and Asp408 (corresponding to Asp407 and Asp408 in AcrB) on transmembrane segment (TMS) 4, and both of them were found to be essential for the pump function. This study also identified an essential Lys939 (corresponding to Lys940 of AcrB) located in TMS 10, and suggested that these three residues produce a proton-relay network. Finally, we have identified Thr978 of AcrB on TMS 11 as another, functionally essential component of this network (86)(Fig. 7). (This study also identified Arg971, located close to the cytosolic end of TMS11, as an essential residue (86), although it is far away from the network of other charged residues.) These results emphasize the role of the transmembrane domain in proton translocation, which presumably provide energy for drug pumping by the periplasmic domain.

Site-directed mutagenesis was also applied to residues assumed to be involved in the binding or movement of substrates. When we reported on the crystal structures of AcrB with ligands in the central cavity of the transmembrane domain (87), we were aware of the potential problem that the ligand might be binding to any hydrophobic pocket of the protein, which has nothing to do with the normal pathway of the exported ligand. Thus we changed residues that appeared to be involved in the binding through site-directed mutagenesis, and we obtained an apparently assuring result that conversion of Phe386 to alanine nearly totally abolished the resistance to ethidium, rhodamine 6G, and dequalinium (87). However, this result was obtained by using a very high copy number vector of the pUC series, and we now are aware that this approach may produce misleading results possibly by making the cell membrane leaky through a strong overproduction of intrinsic membrane proteins. Indeed, Phe386Ala mutation hardly affected susceptibility of *E. coli* to most drugs, when tested with expression by the vector pSPORT1 (88). Nevertheless, Hearn and coworkers (89), using a *Pseudomonas fluorescens* homolog of AcrB, EmhB, expressed from a medium copy number plasmid, found that decreased efflux of dequalinium was produced by the same change (Phe386Ala) as well as the change of Asn99 to alanine. Furthermore, changing Asp101, which is located at the “ceiling” of the central cavity, into alanine decreased the efflux of most drugs as well as of polycyclic hydrocarbons.

With AcrB mutant Asn101Ala, which will be described later in section III.A, the efflux activity was decreased by changing into alanine the residues Glu673, Phe664, or Phe666, all located in the right (looking into the center of molecule with the periplasmic domain up) wall or the bottom of the large periplasmic cleft, close to its entrance (89). With the Cu⁺ and Ag⁺ RND-type efflux pump CusA of *E. coli*, changing into leucine the methionine residues 573, 623, and 672, all close to the entrance of the periplasmic large cleft in the AcrB model, strongly reduced its activity (90). Together with the binding of substrates to this cleft observed in the Asn101Ala mutant AcrB (89), described below, these results favor the assumption that the periplasmic uptake of substrates by these pumps involves the transient binding to, and then entry from, the large periplasmic cleft.

Dastidar et al. (91) introduced cysteine residues to various positions of *Haemophilus influenzae* AcrB, chosen on the basis of their effects on substrate specificity in MexD (81) (described above), and measured, in intact cells, their accessibility to a relatively hydrophilic probe, fluorescein maleimide in the presence and absence of efflux substrates such as erythromycin, novobiocin, cloxacillin, ethidium bromide, Triton X-100, and the inhibitor Phe-Arg- β -naphthylamide. Most substrates, except ethidium, inhibited strongly the modification of cysteine at position 288, which corresponds to Gly290 of AcrB that is close to the bottom of the substrate-binding pocket in the asymmetric trimer structure (described in section IV.A). Interestingly, the modification of cysteine at position 601, corresponding to Phe617 of AcrB, one of the residues lining the substrate-binding pocket, is inhibited by Triton X-100, but not by other substrates. These data suggest the large size (and/or the

flexibility) of the binding pocket that would accommodate different ligands in different ways (see also section IV.B).

Murakami et al. (70) used cysteine scanning mutagenesis to examine the role of the narrow central “pore” of the trimer, which appears to be closed in the crystal structure of AcrB. Although alterations of the residues facing the center of the pore, including Asp101 just mentioned, were found to decrease the extrusion of drugs, the interpretation is complicated by the fact that the pore is made by a three-stranded coiled-coil structure composed of all three protomers of AcrB, so that inter-subunit disulfide bonds are formed in some cases. Now that the drug-binding site and the likely drug extrusion pathway were discovered within the AcrB protomer (section III.C), it seems unlikely that the interprotomeric central pore plays a direct role in substrate movement.

E. Membrane Vesicles and Proteoliposome Reconstitutions

In the biochemical studies of most bacterial transporters, the most useful approaches were often the use of membrane vesicles, either right-side-out or inverted. However, with RND family multidrug efflux pumps, this approach was not successful, although it was vigorously pursued in several laboratories including our own. The likely reasons for this failure include the hydrophobic nature of most substrates which allows a spontaneous equilibration of transported substrates across the membrane bilayer. Furthermore, “transport” by RND transporters may not change the transmembrane location of the substrates since the major pathway of substrate capture is from the periplasm. Another possibility may be that the periplasmic adaptor protein, which seems to be necessary to activate the transporter (see below), becomes stripped off during the preparation of the membranes.

The first successful functional reconstitution of an RND pump was achieved by Zgurskaya and Nikaido (67) in 1999. Cells containing *acrAB* genes on a high-copy-number pUC plasmid were grown without induction, and the membrane proteins were solubilized with 5% Triton X-100 overnight in the cold. The native AcrB protein contains four histidine residues among the eight residues at its C-terminus, and thus the protein could be purified by adsorption to a Cu^{2+} -chelate matrix followed by elution with imidazole, always in the presence of Triton X-100. Just before the reconstitution, AcrB was adsorbed again to the Cu^{2+} -chelate matrix, and Triton X-100 was exchanged with octyl- β -D-glucoside by elution with a buffer containing the latter detergent as well as imidazole. Since AcrB formed aggregates in octylglucoside within a few hours, it was necessary to carry out reconstitution into proteoliposomes immediately by the octylglucoside dilution method.

Innovative approaches were required for assays of the transport activity of AcrB, because most of the substrates of AcrB are quite hydrophobic (7) and are expected to diffuse spontaneously across the membrane bilayer, as mentioned above. Thus it was impossible to rely upon the quantitation of ligands accumulated within the intravesicular space. Inspired by the report that mammalian P-glycoprotein, which also transports hydrophobic compounds, has activity as a phospholipid flippase (92), we added 7-nitrobenz-2-oxa-1,3-diazol-4-yl (NBD)-labeled fluorescent phospholipid to the phospholipid mixture used in proteoliposome reconstitution. Phospholipids, however, are likely to become reinserted into the original bilayer even when they are expelled from it by the AcrB pump. In order to minimize this possibility, an excess of “acceptor” liposomes, which did not contain AcrB, were used to “trap” the fluorescent phospholipids extruded. Finally, the amount of the NBD-labeled phospholipids remaining in the AcrB-containing “donor” proteoliposomes was estimated by initially quenching the NBD fluorescence through the fluorescence energy transfer to rhodamine-labeled phospholipids added initially (together with NBD-labeled phospholipids) to the proteoliposome reconstitution mixture.

Phospholipid extrusion, as indicated by time-dependent increase in NBD fluorescence, required the addition of Mg^{2+} (Fig. 8), which is consistent with the observation that in intact cells, another RND-family multidrug efflux pump MexXY requires the presence of Mg^{2+} in the medium for its function (93). Remarkably, strong pumping also required the addition of lipid-free AcrA. This result was interpreted as the result of AcrA protein connecting the donor and acceptor vesicles (67). However, knowledge on the structure of AcrA (see Fig. 3 and Fig. 4) makes this hypothesis unlikely, because AcrA can connect to the second membrane only through TolC, which was not present in our system. Rather, AcrA could have activated the pumping activity of AcrB directly, in view of the fact that AcrD, a close homolog of AcrB, is activated in a reconstituted system by AcrA (64)(see below).

To show more directly that AcrB functions as a proton/drug antiporter, the system was energized by using the valinomycin-induced efflux of K^+ , which was converted into an interior-acidic proton gradient in the presence of KCl. By measuring the intravesicular pH with a fluorescent, membrane-impermeable pH indicator pyranine, we confirmed that proton efflux occurred accompanying the pH-gradient-driven operation of the pump in the presence of substrates (67) (Fig. 9). When the number of protons moved per vesicle is calculated, it appears that the AcrB pump is functioning extremely slowly, with a turnover rate of less than one per minute. We note that the assay was carried out without the addition of Mg^{2+} and AcrA, which were needed for the activation of AcrB according to the fluorescent phospholipid extrusion assay. Possibly an even more important factor could have been the location of the substrates, which were added to the extravesicular space that corresponded to cytosol as the acidified vesicle interior corresponded to the periplasm. Thus it seems likely that we were measuring the slow spontaneous influx of substrates into the intravesicular space, followed by their capture by the periplasmic domain of AcrB.

The affinity of substrates for AcrB was estimated in two ways. First, the substrates were added as potential competitors in the fluorescent phospholipid efflux assay. This showed that conjugated bile acids, such as taurocholate, were the most effective inhibitors of phospholipid efflux. Second, the potential substrates were added in the proton efflux assay, and again taurocholate was the most effective in producing proton efflux among the compounds tested (Fig. 9). These results fit the notion that the natural substrate for the AcrB pump are the bile salts, most of which exist in the conjugated form in the intestinal tract of higher animals.

Soon afterward, the successful reconstitution of another RND pump, CzcA protein of *Ralstonia*, was reported (83). CzcA is not a multidrug pump, but catalyzes the export of toxic divalent cations such as Zn^{2+} , Co^{2+} , and Cd^{2+} . The PCR-amplified gene was inserted into a commercial vector pASK-IBA3, which supplies a C-terminal, eight-residue tag. The protein was purified by affinity chromatography with a modified streptavidin matrix that binds the C-terminal tag (94). Solubilization and purification were performed in the presence of phospholipids (95). The purified CzcA was mixed with Triton X-100-treated liposomes, and the proteoliposomes were formed by removal of the detergent with Bio-Beads. A pH gradient was imposed across the membrane by diluting proteoliposomes containing 100 mM Tris-Cl at pH 5.0 into 100 mM Tris-Cl at pH 7.0. Since the substrates do not diffuse spontaneously across the membrane bilayer, the pumping activity can be assessed by traditional methods, and indeed vesicles were shown to accumulate divalent cations such as Zn^{2+} . Better results were obtained by acidifying the vesicle interior by the spontaneous efflux of NH_3 from proteoliposomes loaded with 0.5 M NH_4Cl . The data with Zn^{2+} were reported to show a sigmoidal kinetics with a Hill coefficient of 2, although only four concentrations of the substrate were used. These data, unfortunately, may not have much physiological relevance because the half-maximal rate was achieved at an absurdly high concentration of 6.6 mM, where micromolar (or even lower) values of $K_{0.5}$ was expected.

Furthermore, the experiment measured the movement of cations from outside (corresponding to cytosol) to intravesicular space (corresponding to periplasm), whereas the real pump is likely to export substrates from periplasm to the external medium. Indeed we now know that a similar cation pump, CusAB system captures ions such as Cu^+ from the periplasmic binding protein CusF before their efflux (96,97).

Reconstitution of another multidrug efflux RND pump AcrD was achieved in 2005 (64). AcrD was of special interest to us, as some of its substrates, aminoglycosides, are very hydrophilic and are not expected to cross membrane bilayers spontaneously. AcrD, in a form with a C-terminal hexahistidine tag, was expressed from a high copy number plasmid with isopropyl- β -D-thiogalactoside (IPTG) induction, and was purified in dodecylmaltoside. The AcrD preparations in dodecylmaltoside were used directly for proteoliposome reconstitution by octylglucoside dilution method, without the prior removal of dodecylmaltoside. Routine assays relied on the proton efflux determined with the intravesicular pH indicator pyranine, as described above. It was soon discovered that no acceleration of proton flux occurred, with aminoglycosides (70 μM) either outside or inside vesicles, unless AcrA (with the lipidation site removed) was added at the time of reconstitution, and thus to the interior of the vesicles. It was known that AcrA was required for the activity of AcrD in intact cells (77), but this was expected because AcrA (or one of its homologs) would be needed for the construction of a tripartite efflux machinery. In the proteoliposomes containing AcrD, a tripartite structure is not needed for efflux, and the result strongly suggested that AcrA and its homologs of the MFP family had another important function in directly activating the pumping function of the cognate RND efflux transporter. A similar situation was found more recently with the MFP MacA for the activation of an ABC efflux transporter MacB (30).

Results obtained with streptomycin are shown in Fig. 10. When streptomycin was added to the external medium, the drug is occupying a more alkaline compartment that is similar to the cytosol in intact cells. Under such a setup, we could not detect stimulation of proton efflux (curve 2); however, a strong stimulation was observed when streptomycin was added to the more acidic intravesicular space, corresponding to the periplasm (curve 3). These results are important as they were the first results showing that at least one drug was captured *exclusively* from the periplasm. Adding kanamycin, tobramycin, gentamicin, or amikacin to the interior of the vesicles similarly produced a strong increase in the efflux of protons (64). These aminoglycosides however seemed to produce some stimulation even when they were added to the outside medium, and we then concluded that these drugs may be captured both from the periplasm and from the cytosol. However, examination of the data (64) shows that the stimulation tended to be marginal or small when the drugs were added to the outside medium, and I now believe that data were overinterpreted at that time and that the capture occurs only from the periplasm with all aminoglycosides. We tested the intravesicular accumulation of [^3H]gentamicin by adding it to the external medium (64). Although the vesicles seemed to accumulate the drug slowly, the background accumulation was also significant, and the calculated turnover number of AcrD was in the range of 0.01 s^{-1} . This is insignificant in comparison of now determined turnover number of AcrB for cephalosporins, between $10\text{--}1,000 \text{ s}^{-1}$ (section II.C), and it fits with the notion that there is little evidence for the capture of aminoglycosides from inside the cytosol.

III. Structure of AcrB

A. Symmetric Structures of AcrB

Crystallography supplied essential pieces of knowledge on the mechanisms of RND pump complex. The crystallographic structures of the outer membrane channels and of the periplasmic adaptor proteins have been briefly described in section II.B above. Again, as

mentioned earlier, Murakami et al. solved the structure of AcrB as a homotrimer with a three-fold rotational symmetry in 2002 (44) (Fig. 3), the first crystallographic structure for a proton-motive-force-driven transporter solved.

We mentioned already that AcrB likely captures its substrates from the periplasm, and the prediction was made in the early days that drugs may first partition into the membrane-periplasm interface (Fig. 1). Indeed the AcrB trimer structure showed that there was a small opening (vestibule) between subunits at the bottom of the periplasmic domain, close to the external surface of the membrane bilayer (44), and this led to the top of the large central cavity in the transmembrane domain (Fig. 3). It was thus hypothesized that the drugs diffused through the vestibule into the central cavity, where it was captured. Indeed, co-crystallization of AcrB with various drugs (rhodamine 6G, ethidium, dequalinium, and ciprofloxacin) produced crystals with drug molecules within this central cavity (98). However, our conclusions were less than water-tight. First, the resolution was poor (3.5 to 3.8 Å), and thus the identification of the drug molecules could not be made with absolute confidence, although drugs with polyaromatic structures were used precisely for the ease of detection in the crystal structure. Second, the wall of the central cavity, where these drugs were thought to be bound, appeared quite open and totally unlike the traditional ligand-binding sites. In order to explain drug binding to such sites, we had to propose the existence of “composite” binding sites with the involvement of head groups of phospholipids that were assumed to be present in the central cavity (87). In order to confirm that the bound drugs were on their correct path to export, we applied site-directed mutagenesis to change Phe386, which seemed to be close to most substrates, into Ala, and found that the efflux activity was nearly abolished (87). However, as described already (section II.D), this result is likely to be an artifact resulting from the use of high copy number vectors. Thus our earlier conclusion that the drugs bind to the walls of the central cavity is not totally convincing. Nevertheless, the AcrB-YajC cocrystal obtained accidentally from *E. coli* grown in ampicillin-containing medium (99) was reported to contain two ampicillin molecules per AcrB protomer at this position near the ceiling of the central cavity, although the resolution was again poor (3.5 Å) and it is difficult to imagine that these ampicillin molecules survived the attack by the high levels of plasmid-coded β-lactamase during production and purification of the protein.

In one symmetric structure of AcrB, the revealed N-terminal structure was shown to narrow the cytosolic opening of the central cavity somewhat (100). However, it is unclear whether this has any bearing on the function of the pump, as the role of the central cavity in substrate binding is uncertain, as described above.

We obtained another set of drug cocrystallization data with a site-directed mutant of AcrB, Asn109Ala (88). In the simple hypothesis we had at that time for drug extrusion, the drug molecules are captured near the ceiling of the central cavity, enter the central pore bordered by three helices (roughly corresponding to Asp99 through Leu117) from the three protomers, and pass through this pore to reach the top of the periplasmic domain (see Fig. 3). However, the pore in the crystal structure appears closed. Since the side-chain of Asn109 appears to extend deeply into the center of the pore, we naively thought that changing it to alanine might weaken the interaction of helices from the three subunits, and “loosen up” the pore. The mutant AcrB functions almost normally as judged from the resistance levels to several antimicrobial agents. Interestingly, when the mutant protein was crystallized in the presence of several ligands (ethidium, rhodamine 6G, ciprofloxacin, nafcillin, and the inhibitor Phe-Arg-β-naphthylamide (41)), ligands were found to be bound not only to the ceiling of the central cavity (as seen earlier (98)) but also to the periplasmic domain, near the entrance of the large, deep, external cleft (Fig. 6). Again, there was the problem that the ligands could have become bound to any lipophilic pocket which may not be on the proper

pathway for export. Changing residues that appeared to be interacting with the ligands into alanine through site-directed mutagenesis produced a drastic decrease in resistance to most drugs in the case of Glu673, and more modest yet significant decreases with Phe666 and Phe664. Although we cannot exclude the possibility that they may inactivate the transporter by a mechanism unrelated to ligand binding, the more recent asymmetric AcrB structures (discussed in section III.C) are not inconsistent with the hypothesis that this is the portal for the periplasmic entry of substrates. The putative pathway for the drugs to the binding site is closed in the “access protomer” (see below), which resembles the protomers in the symmetric trimer in conformation; thus the drug molecules cannot go further, and may be forced to interact with the entrance of this pathway in the symmetric structure. Furthermore, there are random and site-directed mutation data that suggest the importance of residues within the cleft (section II.D)(see also Fig. 6).

Finally it should be noted that the symmetric AcrB crystals soaked in deoxycholate contained deoxycholate molecules within the external cleft of the periplasmic domain (101), in an essentially identical position to the various substrates mentioned above.

B. Attempts to Determine the Structure of AcrB during the Drug Export Cycle

Because of the uncertainties involved in the modeling of ligand electron density and in the possibility of non-specific interaction between the ligand and the protein surface, we tried a different approach. We crystallized AcrB mutants in which one of the transmembrane domain residues putatively involved in proton translocation, Asp407, Asp408, Lys940 (corresponding to Lys939 of MexB), or Thr978 (see section II.D above), was changed to Ala (86,102). In the crystal structure (44), the side-chains of these residues are very close, and appear to interact with each other either by salt bridge or H-bonding (Fig. 10). We assumed that during the normal function cycle of AcrB, a proton antiporter, one of these residues will be altered in its protonation state, most probably by the protonation of Asp, and this may disrupt the tight interactions among these residues and eventually cause conformational changes of the whole protein which would be coupled to ligand export. If so, our mutant proteins would mimic this transient, protonated intermediate state during the process of drug extrusion, and allow us to see this elusive structure that cannot be deduced from the ground state structures of the wild-type protein. When the crystal structures (without ligands) of these mutant proteins were examined (102), we found the following. 1. The tight interaction between Asp407, Asp408, Lys940, and Thr978 became indeed disrupted, and the Lys940 side-chain, which interacted strongly with both of the carboxyl groups of Asp407 and Asp408 in the wild type AcrB, went through a strong flipping motion and now faced away from these residues (Fig. 10). Remarkably, the alterations were quite similar regardless of which residue was changed to Ala (102). 2. An extensive conformational alterations were found in the transmembrane domain (102). Again, it was reassuring that the conformational alterations were similar in all four mutants, confirming that they were due to the disruption of the salt bridge/H-bonding network among these four residues, rather than to specific effects of individual mutations. Much of the changes involved the shortening or melting of helices, except that helix 8, one of the helices connecting the transmembrane domain to the periplasmic domain, became extended into the periplasmic domain (102). 3. When the structure of the periplasmic domain was examined, however, there were only very small changes (102), a disappointing result because we expected major conformational alterations in the periplasmic domain during substrate binding and extrusion (see above). The reason for this failure was suggested from the studies describing the asymmetric trimer structure of AcrB, described below.

C. Solution of the Asymmetric AcrB Trimer Structure

At about the same time as our study of the proton-relay network mutants, several papers appeared describing the asymmetric trimer structure of AcrB (103-105)(Fig. 11). In contrast to the symmetric structures seen before, in these structures each protomer, especially its periplasmic domain, takes a unique conformation (Fig. 11). All groups identified one protomer in which a pocket is opened up in the periplasmic domain, a likely substrate-binding site, and in the study by the Murakami group (103), this pocket was indeed found with a bound substrate molecule (minocycline or doxorubicin). The interpretation of electron density as the bound substrate was strengthened by using minocycline that was labeled with a heavy atom, bromine. This protomer is called the “binding” protomer by the Murakami group or T (for Tight) by others. Other protomers are called the “access” (or L for Loose) and “extrusion” (or O for Open) protomers. In the extrusion protomer, which shows the largest conformational alteration, the binding pocket becomes collapsed, and there appears to be an opening of the pathway from the binding pocket to the top (central funnel) of the periplasmic domain. Inspection of the structures of the transmembrane domains shows that the tight association of Asp407-Asp408-Lys940-Thr978 becomes disrupted in the extrusion protomer, likely as a consequence of the protonation of Asp407 and/or Asp408; indeed the arrangement of these side-chains here is nearly identical with that found in our mutant protein (102)(Fig. 10, green sticks). This is consistent with our hypothesis that the disruption of the network among the four residues mentioned alters the conformation of the periplasmic domain. However, the asymmetric structure also shows that extensive conformational changes in the periplasmic domain require complementary accommodation by the neighboring protomers (103-105). This was not possible in our construct, in which all protomers corresponded to the mutant protein.

As mentioned already, the extrusion protomer is the subunit where the conformation of the periplasmic domain is altered most strongly from the conformation seen in the symmetric trimer. This conformational alteration is presumably driven by the extension of helix 8 toward the periplasm, this change in turn is caused by the disruption of the salt bridge/H-bonding network among Asp407, Asp408, Lys940, and Thr978, as seen in the structures of both the wild-type protein in the asymmetric model (103-105) and the structure of the proton-relay network mutants (102).

The asymmetric structure was important because the structure of one single trimer suggests three stages in the drug extrusion process. Thus the protomers appear to go through a cyclic conformational change, from the open, “access” conformation (that would allow the access of substrates), through the ligand-bound, “binding” one, to finally the “extrusion” conformer, which propels the substrates out and whose transmembrane domain shows signs of disrupted network among proton-translocating residues (103-105)(as shown in Fig. 11B). Thus this mechanism is called the “functionally rotating mechanism”; it is reminiscent of the mechanism of F₁-ATPase, although there is nothing that corresponds to the physical rotation of its γ -subunit. The “power stroke” of this entire process occurs during the change of binding into extrusion conformation, which involves the largest movement of various sections in the periplasmic domain. This large conformational alteration is initiated apparently by the protonation of one or both of the Asp residues (Asp407 and Asp408) in the midst of the transmembrane domain, which leads to a significant elongation of TM helix 8 into the periplasmic domain and all the subsequent conformational changes in the periplasmic domain.

Recently, the reactivity of the carboxyl groups in AcrB was assessed by labeling with a hydrophobic reagent, dicyclohexylcarbodiimide, followed by CNBr cleavage of the protein and analysis of the fragments by mass spectrometry (106). It was found that Asp408 was especially reactive to carbodiimide, with a high pK_a of 7.4, implicating the protonation/

deprotonation of at least this Asp residue as a key step in the proton translocation pathway, and in eventually generating conformational alterations in AcrB that leads to drug extrusion.

IV. Mechanism of Drug Efflux

A. The Main Substrate-Binding Site

The possible route of substrate binding and extrusion is suggested by the structure of the asymmetric trimer (103-105). In the binding protomer, the periplasmic domain contains an expanded binding pocket containing several aromatic and hydrophobic residues (Phe136, 178, 610, 615, 617, and 628; Val139 and 612; Ile277 and 626; and Tyr327 (104)). This pocket was indeed found to contain drugs minocycline and doxorubicin (103), and its role is also supported by the previously described mutation data with the AcrB homolog YhiV (82) as well as the more recent data on AcrB (107). In the latter study, each of the Phe residues in the binding site was changed to Ala by site-directed mutagenesis. Among them, Phe610Ala mutant became strongly more susceptible to almost all drugs, and Phe178A and Phe615Ala had a somewhat less extensive effect. Changing Phe628 and Phe136, and especially Phe617, decreased the AcrB activity only to a limited range of substrates, perhaps understandable as these side-chains occupy a peripheral position in the binding pocket.

The significance of this binding site is also suggested by the observation that the pocket becomes collapsed in the extrusion protomer. Finally, its physiological significance is further supported by the comparison of its surface in AcrB and in the homology-modeled AcrD (36): although the surface is entirely hydrophobic in AcrB, in the aminoglycoside pump AcrD it contains many oxygen atoms that may function in the binding of the hydrophilic, basic substrates.

B. Pathways for Drugs: A Current Hypothesis

How do the substrates reach the binding pocket? All three groups who studied the asymmetric trimer structure agree that, in the binding protomer, there is an open pathway between the binding pocket and the external large cleft of the periplasmic domain of the binding protomer (102-104) (see Fig. 12). Although this cleft was earlier thought by some investigators to be the site of interaction between AcrA and AcrB, in the most up-to-date model of AcrA-AcrB interaction this cleft remains fully open (53) (Fig. 4). (The entry of the substrate from the cleft was advocated by Lomovskaya and Totrov (108), before the appearance of the asymmetric crystal structures.) We have seen that the possibility of substrate entry from the lateral cleft is also supported by the results of random mutagenesis data (for example ref. 81 and ref. 83 described in section II.D) and the results of site-directed mutagenesis (for example ref. 88 and ref. 89 in the same section)(see Fig. 6). Finally, cocrystallization studies showed the presence of bound substrates at the entrance of this large cleft (88), as described in section III.B (Fig. 6).

The Murakami group found in addition the second pathway that connects the binding site with the vestibule, the passageway between protomers, located close to the external surface of the inner membrane. A similar pathway between the binding site and the vestibule area could also be seen in the Darpin-containing crystal, that so far has shown the highest resolution (105). Thus there is a possibility that the substrate may go through the initial part of the vestibule, and then go into the binding pocket.

In the extrusion protomer, because of the conformational changes initiated most likely by the protonation of Asp408 in the transmembrane domain, the binding pocket becomes narrower, and the entrance pathway(s) become closed and at the same time, a new pathway from the binding pocket to the funnel-like structure at top becomes open (Fig. 12). This would favor the extrusion of drug molecules into the TolC channel.

Thus the basic principles of the drug pumping mechanism, based on the functionally rotating model, are now reasonably well understood. However, some details are still at the stage of speculation. For example, in order for the drug extrusion to occur with a reasonable efficiency, the drug molecule must remain in the binding pocket for some time. How would this be achieved, when both the drug entry pathway and the pocket are mostly surrounded by hydrophobic side-chains (Fig. 12)? Furthermore, how does the AcrB pump accommodate such a wide range of substrates, some of which are shown in Fig. 2? The answer to these questions is not immediately apparent from the publications describing asymmetric crystal structures (103-105), because the only drug molecules found in cocrystals are minocycline and doxorubicin, in spite of the trials with many dozens of drugs by the Murakami group. Because of this situation, we have tried a computer simulated docking approach (H. Nikaido, unpublished observation). With the binding protomer of minocycline-bound AcrB (103), the binding of about two dozens of known AcrB substrates was estimated. Minocycline was shown to bind to the apo-protein in a location and conformation found in the crystal structure, although this was not the complex predicted to be the most stable one. This result validates the use of the computational approach, at the same time suggesting that the energy scoring function of the program may not yet be perfect. Currently there is no way to include the movement of the backbone of this rather flexible protein in simulation. Thus the details of the predicted binding interactions of drugs give us only broad hints. Nevertheless, examination of the possible binding modes of many substrates was instructive, in a way somewhat similar to the usefulness of comparison of many homologous sequences. Many substrates (including minocycline, tetracycline, and carbenicillin) prefer to bind, at least in this modeling exercise, with their hydrophobic domain bound to the narrow central groove of the binding site, with their hydrophilic group(s) exposed either at the "top" and/or "bottom" minipockets within the site. These subdomains are much more hydrophilic than the central groove. This is very similar to the binding of minocycline (103) (for structure, see Fig. 2), which is bound to the groove mainly by using its hydrophobic middle of the molecule, and has its one polar domain (containing the acidic OH, an amino group, and an amide group) on top and its other polar group (an amino group) at the bottom. Murakami et al. (103) argue that the polar groups on top interact with Asn274 side-chain, and the amino group at bottom interacts with Gln176. However, none of these groups are in H-bonding distance, and are at best 4-5 Å away. Thus it seems best not to think of these interactions as being specific. Rather, the top and bottom of minocycline seem to fit into more hydrophilic, larger minipockets which may even be filled with water.

Interestingly, some ligands are predicted to bind nearly exclusively to the bottom pocket and its neighboring hydrophobic patch, completely avoiding the central groove where the bulk of minocycline binds. This group includes the inhibitors Phe-Arg- β -naphthylamide and NMP (naphthylmethylpiperazine) (109) as well as many of the binding modes of the fluoroquinolones and chloramphenicol. These observations may be pertinent to the finding that Phe-Arg- β -naphthylamide was very effective in preventing the efflux of levofloxacin and chloramphenicol, but not so effective for the efflux of carbenicillin or tetracycline by the AcrB homolog MexB (41).

In this light, we can imagine the movement of substrates as follows. Substrates are initially attracted to the hydrophobic surfaces of the pathways, either from the periplasm through the large external cleft of AcrB periplasmic domain, or from near the external surface of the plasma membrane through the vestibule. However, the protein-substrate complex is not in its most stable conformation here, because the hydrophilic part of the substrate is not stabilized optimally. The final stabilization is achieved at the binding site, because of the presence of two features here, one, interface(s) for hydrophobic interactions, and the other, rather large hydrophilic pockets next to it. The substrate range of AcrB is likely to be determined essentially by this structure of the binding site. The substrates bound here is then

ready for extrusion through the conformational change of the binding protomer into the extrusion protomer.

C. Biochemical Studies of the Conformational Alterations

Although the structure of the asymmetric AcrB trimer suggests strongly the mechanism of ligand extrusion coupled to proton influx, this mechanism still remains a hypothesis. We set out to prove this hypothesis by carrying out biochemical studies of the pump. The proposed cyclic change involves the opening and closing of the large external cleft in the periplasmic domain (Fig. 11). Thus if the opening of the cleft is prevented, we can predict that the cyclic conformational changes will be prevented and that the pump will cease to function. We introduced pairs of cysteine residues at various positions on the opposing walls of the cleft (110). Although single cysteine mutations produced little inhibition of transport, proteins containing double mutations (Asp566Cys and Thr678Cys; Phe666Cys and Thr678Cys; Phe666Cys and Gln830Cys) on the opposing walls of the cleft were strongly compromised in function. This is likely to be due to the forced closing of the cleft by disulfide bonds formed between these residues, but we cannot rule out the possibility that the trimeric assembly failed because of the premature formation of disulfide bonds between protomers. Indeed with cysteine pairs at other positions, there were data suggesting this interpretation. To avoid this problem, we expressed mutated AcrB in a host strain with a defective DsbA, a periplasmic disulfide oxidoreductase that plays a major role in the formation of disulfide bonds in this compartment (111,112). Indeed in this strain, the AcrB protein containing Phe666Cys and Thr678Cys (or Gln830Cys) largely retained its transport activity, presumably because disulfide bond formation did not occur in the absence of DsbA enzyme activity. We tried to see, by using these mutants, if the cross-linking and inactivation of AcrB can be observed in “real time” by using a fast-acting disulfide cross-linking agents based on the methanethiosulfonate groups (113). Indeed, addition of such cross-linkers to cells that are pumping out a fluorescent dye, ethidium, was shown to stop the function of the pump instantaneously (110), thus providing biochemical support for the rotating mechanism of drug transport.

A similar approach was also pursued by the group of K. M. Pos (114). They introduced pairs of cysteines at a wide range of positions, importantly including those that were expected to become closer in the access or the binding protomers, in addition to those in the extrusion protomer. The double cysteine mutant of Ser132 and Ala294, which are expected to become closer only in the access protomer, inactivated the pump as estimated by the decreased resistance to a number of drugs, and the pumping activity was increased to a detectable extent upon the addition of disulfide-reducing agent dithiothreitol to intact cells (114).

D. Use of Covalently Linked Trimers

The functional rotation mechanism of AcrB drug pumping process (103-105) predicts that if one of the three protomers is defective in the proton relay network, then the pumping action by the entire trimer should come to a halt. This hypothesis could not be tested by the disulfide cross-linking experiment described above, because all protomers had the same Cys mutations. We devised a way to test this hypothesis by creating a giant gene, in which three *acrB* sequences were connected together through a linker sequence (115). When the sequence of the cytosolic, horizontal helix (Met496 through Arg540) connecting the two halves of the transmembrane domain was used as the linker, the linked trimer was expressed well as a single giant protein, and produced drug resistance levels sometimes even higher than that produced by the monomeric *acrB* gene. When only one of the three component *acrB* sequences in the giant gene was changed to include mutations in the proton-relay network, then the entire trimer became inactive. When only one of the three *acrB* sequences was made to contain the Phe666Cys and Gln830Cys, then cross-linking of these two

cysteine residues by a fast-acting methanethiosulfonate cross-linker instantaneously inactivated the entire trimer, regardless of the position of the altered protomer in the giant sequence (Fig. 13). These results strongly support the notion that the trimer acts by a functionally rotating mechanism.

E. RND Systems Apparently Requiring Paired Transporters

The RND transporters we have discussed so far, AcrB, MexB, or AcrD, exist as homotrimers, and are now assumed to work by a functionally rotating mechanism. There are, however, RND transporter genes that occur in tandem, such as MdtBC, discovered by our group (116) as well as by Yamaguchi's group (117). Both groups showed that both of the RND transporters, MdtB and MdtC, had to be present for the full pumping activity. Interestingly, Baranova and Nikaido (116), who relied on the expression of these genes from the chromosome, found that MdtB or MdtC alone (in the presence of MFP MdtA) was completely inactive, whereas Nagakubo et al. (117), who expressed the genes from plasmids, reported that MdtC alone gives some marginal activity.

When the C-terminus of either the B or C protein was labeled with hexahistidine tag, metal affinity purification produces a complex containing both MdtB and C, confirming that these proteins produce a heteromultimeric complex (H. S. Kim and H. Nikaido, manuscript in preparation). The size of the complex showed that it was a trimer, and the average composition was B₂C. In order to establish the precise composition of the active trimer, the *mdtB* and *mdtC* genes were connected by linker sequences to produce covalently linked trimers of all possible combinations. The results showed that the highest activity was obtained only when the linked trimer contained two copies of MdtB and one copy of MdtC.

Since the heterotrimer contains two different protomers, the functionally rotating mechanism seems unlikely for this pump at least in its unmodified form. Furthermore, when the proton-translocation pathway residues such as Asp410 (corresponding to Asp407 of AcrB) were changed to Ala in one of the MdtB sequences, the complex lost its activity, although similar changes in the MdtC sequence (for example in Asp401, corresponding to Asp407 of AcrB) resulted only in a small change in activity (H. S. Kim and H. Nikaido, manuscript in preparation). Since we know from AcrB that protonation in this network of amino acids is needed to for the extrusion of substrate drugs (see above), we assume that MdtB performs this function by changing from the "binding" conformation to the "extrusion" conformation. We do not know at present what the role of MdtC is. It could simply play a role in mechanically stabilizing the trimeric complex, in facilitating the capture of substrates in the neighboring MdtB unit, or in actually capturing the substrates by itself. Regardless of the precise roles of protomers, it is clear that the operating principle here is somewhat different from that of the homotrimers. Interestingly, MdtC is a basic protein with the calculated pI of 8.4, in stark contrast to other RND transporters that are acidic (AcrB, 5.4; AcrD, 5.6; MdtB, 5.2). We also do not know if this heterotrimeric structure gives advantages, if any, in terms of pumping activity. Such RND pumps requiring a paired set of transporters are also found in related bacteria, such as SdeAB of *Serratia marcescens* (118), whose substrate range is limited as in MdtBC (bile acids and novobiocin). However, SmeJK of *Stenotrophomonas maltophilia* is reported to have a wider range of substrates (119).

V. AcrAB-TolC and Drug Resistance

We have so far outlined our current knowledge on the mechanism of AcrAB-TolC complex, beginning with the capture of substrates either from the periplasm or from the periplasm-inner membrane interface, followed by the conformational change in the AcrB periplasmic domain initiated by the flow of proton(s) through the AcrB transmembrane domain, finally resulting in the extrusion of drugs into the external medium through the long channel of

TolC. However, gram-negative RND-type efflux pumps have attracted attention because they are thought to play a large role in drug resistance, a major public health problem, which is becoming more and more acute because this type of multidrug efflux greatly diminishes the activity of most of the currently available antimicrobial agents (31). Thus this review will not be complete without discussing the effect of AcrAB-TolC on the drug resistance. Furthermore, the mechanisms and kinetics of the pump we have discussed so far cannot be confirmed unless they explain, quantitatively, the level of resistance finally achieved in the living bacterial cell.

The analysis is the simplest with β -lactams, as their target is in the periplasm, and as most of them are relatively hydrophilic so that we can neglect their diffusion across the inner membrane. We have attempted this analysis at the time when multidrug efflux pumps were not yet known (73). In that study we considered the influx across the outer membrane, which follows the Fick's first law of diffusion, and the drug degradation by the periplasmic β -lactamase, which follows the Michaelis-Menten kinetics. By solving these two simultaneous equations for the external drug concentration (C_o) that would just produce the periplasmic concentration that would kill the cell (C_{inh}), we could predict MIC values for β -lactams. Thus,

$$MIC = C_o = C_{inh} + \frac{V_{max}}{P \cdot A} \frac{C_{inh}}{(C_{inh} + K_m)} \quad (\text{Eqn } 1)$$

where V_{max} and K_m are the usual kinetic constants of the β -lactamase, and P the permeability coefficient of the β -lactam across the outer membrane. A is the area of the outer membrane per unit weight of bacteria.

This procedure was applied to four strains of *E. coli*, producing a widely different levels of the chromosomally coded AmpC β -lactamase (V_{max} of 0.013 to 16.6 nmol/mg/min with cephaloridine), and the predicted MIC values were very close to the observed values with all of the cephalosporins tested. However, with some penicillins, there was significant difference, which was most pronounced with carbenicillin, the observed MIC values usually 8 times higher than the predicted values. We can now understand these results at least qualitatively. With most cephalosporins, at very low periplasmic concentrations where the cells become killed (C_{inh} is usually less than 5 μM (73)), the efflux is insignificant especially because of its sigmoidal kinetics (71); thus exclusion of efflux from consideration produces only small errors. However, with carbenicillin, efflux is important as seen from the strongly decreased MIC values in *acrAB*-inactivated strain (15,72), whereas the chromosomal β -lactamase of *E. coli* works poorly with this compound (72); hence the prediction without consideration of efflux underestimates MIC values greatly.

We can also carry out a more quantitative analysis. The strain used in ref. 71 for the analysis of nitrocefin was a rather artificial construct, containing a mutant porin with a wider channel, to facilitate the analysis. Nevertheless, with this strain, when both the AcrAB efflux pump and the AmpC β -lactamase were deleted, the nitrocefin MIC dropped from 38 μM to 2.2 μM . We can calculate the theoretically expected MIC in the parent strain as follows.

When the periplasmic concentration reaches C_{inh} , the rate of efflux V_e is $V_{max}^e \cdot C_{inh} / (K_m^e + C_{inh})$, and the rate of hydrolysis is $V_{max}^h \cdot C_{inh} / (K_m^h + C_{inh})$, both from the Michaelis-Menten relationship, where the superscript "e" and "h" indicates constants related to efflux and hydrolysis, respectively. Then, since $V_{in} = P \cdot A \cdot (C_o - C_{inh})$, C_o or the MIC can be calculated as $C_{inh} + (V_e + V_h) / P / A$. As was noted earlier in the analysis without the consideration of efflux (73), the constant that is most difficult to estimate is C_{inh} . It has to be

smaller than 2.2 μM , the MIC of the $\Delta\text{acrAB } \Delta\text{ampC}$ mutant, but how much smaller? As there may be other efflux pumps that slowly pump out nitrocefin, we arbitrarily assume that C_{inh} is one-half of this concentration. Then the calculations give us the predicted MIC of 72 μM , or 36 $\mu\text{g/ml}$. Although this is twice the observed MIC of 18 $\mu\text{g/ml}$, the discrepancy is not too large, considering that MIC was determined under somewhat different conditions in comparison with the kinetic constants (for example temperature and medium).

An interesting exercise is to estimate the effect of outer membrane barrier on MIC values. We can see that the more permeable outer membrane would decrease the second term of the formula used above, by increasing P , and would make the MIC closer to C_{inh} . Thus the effect of efflux (or periplasmic hydrolysis) will become minimized. Conversely, in an organism with a more efficient outer membrane barrier, the term $(V_e + V_h)/P/A$ will become larger, and thus efflux (and hydrolysis) will contribute more effectively in raising the MIC values, the situation we find for example in *P. aeruginosa* (20-22).

When different strains were compared, it was evident that the MIC values did not have a linear relationship with the amount of efflux pumps produced. Here a relevant study is the rigorous analysis from the group of Frère (120) on the effect of β -lactamase level and of the target sensitivity to MIC. It was shown that if a high-affinity β -lactamase is overproduced, a 5,000-fold difference in the sensitivity of the target (penicillin-binding proteins) will produce only twofold difference in MIC. We can see in Eqn 1 that under certain conditions the $C_{inh}/(K_m + C_{inh})$ term approaches 1, so that the MIC will be affected solely by the balance between permeability and the enzyme concentration, i.e. by $(V_{max}/P/A)$, and not much by the sensitivity of the target, or C_{inh} . These conditions are attained, for example, when the K_m of the enzyme is very low, as with cefotaxime, in comparison with C_{inh} . Thus the presence of β -lactamase will “mask” the difference in target sensitivity. Of course, if the β -lactamase and efflux were totally absent, there will be a 5,000-fold difference in MICs. Although at the time of this work, the kinetic behavior of the AcrB pump was not known, the conclusions may be relevant, as the efflux follows saturation kinetics like the enzymatic hydrolysis.

It must be remembered, however, that the K_m of the AcrB pump is usually much higher than C_{inh} (see section II.C), and under such conditions the sensitivity of the target will continue to affect the magnitude of MIC. If we neglect the sigmoidal nature of the efflux kinetics (section II.C), we can calculate the situation with cephalothin as follows. Here the K_m (actually $K_{0.5}$) of efflux is about 100 μM (71), much larger than C_{inh} , which earlier we estimated to be around 1 μM for cephalosporins. Thus the $C_{inh}/(K_m + C_{inh})$ term of Eqn 1 may be approximated as C_{inh}/K_m , i.e. 0.01 in this case. The V_{max} for cephalothin efflux was about 1 nmol/s/mg cells (71). As P for cephalothin is around 10^{-5} cm/s (73) and A is 132 $\text{cm}^2/\text{mg cells}$, the $V_{max}/P/A$ term in Eqn 1 will be around 1,000 μM , and the second term of Eqn 1 will be close to 10 μM , yielding a predicted MIC of 11 μM (neglecting the periplasmic hydrolysis). If the target becomes twice more resistant, with the C_{inh} of 2 μM , the predicted MIC will be 22 μM , following the changes in target sensitivity in a nearly proportional manner. If the efflux rate is decreased by one-half, the predicted MIC will be 6 μM , an almost but not completely proportional change. In reality, the sigmoidal nature of the efflux kinetics is likely to produce significant deviations from proportionality in these situations.

Carrying out similar analysis for antibiotics that have targets in the cytoplasm is much more complicated, because we have to consider the additional step of fluxes across the cytoplasmic membrane. In addition, the efflux across the cytoplasmic membrane may be catalyzed by “singleton” pumps located in this membrane, rather than by AcrB, as shown recently by Tal and Schuldiner (121) for ethidium bromide (pumped out into the periplasm

by an MFS pump MdfA and an SMR pump EmrE). In this case, the drug efflux from cytosol to external medium would require the successive action by at least two pumps, the export into periplasm catalyzed by a “singleton” pump and then the export from periplasm to external medium catalyzed by the AcrB complex. The kinetic behavior of some of the singleton pumps has been elucidated in great detail, for example that of EmrE (122-124). However the kinetic constants of AcrB for non- β -lactam substrates are currently unknown, and building of mathematical models of transport of these compounds would be very difficult at present.

There are some interesting observations on the effect of the presence of AcrB on MICs of antibiotics with intracytoplasmic targets. Thus a mutation in the ribosomal protein L22 lowers the affinity of ribosomes to erythromycin (about fivefold increase in K_D), and makes the *E. coli* cells resistant to this drug, raising the growth-inhibitory concentration (at 50% growth) from 100 to 800 $\mu\text{g/ml}$ (125). In a strain with a deletion of the *tolC* gene (therefore in the virtual absence of efflux), this difference largely disappears, the 50% growth-inhibitory concentration increasing from 3.5 to only 5 $\mu\text{g/ml}$ by the introduction of the same L22 mutation (125). A similar observation was made in another laboratory (126), which found that the difference in the 50% growth-inhibitory concentrations between the wild type and an L22 mutant strain decreased from tenfold in the efflux-competent background to only 1.5-fold in the ΔacrAB background. These authors considered the possibility that erythromycin might increase the translation of AcrAB proteins, but could not find any evidence for such a phenomenon. Synergy between the efflux and the target mutations was also found in *Campylobacter* spp. (127). Thus these observations are the opposite of the masking of the target sensitivity by β -lactamase mentioned above.

An attempt has been made to explain this phenomenon of “resistance masking” in efflux-deficient strains by mathematical modeling (128,129). Since nothing is known about the kinetics of efflux, the efflux pump is assumed to operate at a substrate concentration far below the K_m , and that the efflux is proportional to intracellular drug concentration. Most importantly, because the increase in cell volume due to growth was assumed to be the only mechanism that decreases the drug concentration inside, the rate of net influx of drugs through a loosely defined “cell envelope” needed to be set extremely low, of the order of magnitude similar to the growth rate. This authors remains quite skeptical about the validity of such a modeling approach carried out in the almost total absence of experimental constraints, as nothing is known about the kinetics of OM diffusion of macrolides or about that of active efflux of these compounds. Possibly the exaggeration of the difference in target sensitivity by efflux could be brought about by the sigmoidal kinetics of the pump (71), but we need to know much more before we get into more realistic simulations.

VI. Coda

As a participant who was there at the discovery of *E. coli* AcrB as a multidrug efflux pump (18) 16 years ago, I am impressed and gratified by the amount of progress scientists have made in this area. At that time, the mere notion that a single transporter is capable of exporting such a wide variety of compounds (see Fig. 2) and that it may capture substrates in the periplasm was met with a great deal of skepticism, and the publication of our reports on the *P. aeruginosa* multidrug efflux process (20-21) was delayed for at least several months because the reviewers simply refused to believe in the existence of such a process.

In contrast, we now know the detailed 3D structures of all of the components of the tripartite pumping complex (Figs 3 and 4), and we understand the mechanism of drug extrusion at least in its basic outline. It seems likely that drugs are captured from the periplasm perhaps exclusively, because structures do not so far give any hint on how the drugs captured in the

cytosol may be brought out to the periplasmic binding site. We understand the pathway(s) through which drugs reach the binding site, and then how the protonation of one or two Asp residues in the transmembrane domain causes conformational changes that are transmitted to the periplasmic domain, to produce drug extrusion. We understand that the AcrB protomers within the trimeric complex act by a functionally rotating mechanism, so that all of them have to be functional for any drug molecule to be extruded, a situation that produces sigmoidal kinetics with some substrates. We are even approaching the stage where we can understand how various substrates bind to the various subdomains of the large binding sites (section IV.B).

However, there are still many questions that remain. I hope that I emphasized the need to correlate our biochemical knowledge of pump action to the final effect of the pump in raising MIC (section V), an area still in its infancy. Indeed, the kinetics of AcrB has been examined only for cephalosporins, and we totally lack comparable data for all other classes of antimicrobial compounds. We need to know more precisely how various drugs bind to the binding site. The computer docking studies (section IV.B) can give us only hints, and they remain to be proven by real biochemistry. We also do not have any idea on how AcrA stimulates the function of AcrB.

The functions of the adaptor (or MFP), AcrA and its homologs, have been reviewed (38). However, it should be mentioned that the MFP CusB of the Cus system, a toxic cation exporter of *E. coli*, is known to bind the substrate Cu^+ (96), and most excitingly, the direct transfer of metal ion from the periplasmic binding protein CusF to CusB was recently demonstrated (97). Thus it seems quite possible that in this system the substrate ion flows from the periplasmic binding protein to the periplasmic adaptor protein, and then finally to the periplasmic domain of an RND transporter, CusA.

There are also unresolved questions on the function of tripartite RND multidrug pumps in intact cells. For example, *P. aeruginosa* strains overexpressing a tripartite pump MexCD-OprJ were often found to have reduced function of other tripartite pumps MexAB-OprM and MexXY-OprM although the latter pumps are expressed at the normal level (130). Another curious observation is that TonB, a protein that functions in energizing the outer membrane siderophore receptors, is needed in *P. aeruginosa* for the function of the MexAB-OprM pump (131,132). As far as the authors are aware, there is no explanation of this effect at the molecular level, but this reminds us that a small protein YajC was accidentally co-crystallized with AcrB (99). Although the deletion of gene *yajC* had only a small effect on drug susceptibility, this observation certainly requires further follow-up study, especially because this co-crystal was obtained from cells expressing both YajC and AcrB at normal constitutive levels, without the lopsided, plasmid-driven overexpression of only AcrB.

Acknowledgments

The studies in the authors' laboratory are supported by a U.S. Public Health Service grant, AI 09644. I thank V. Koronakis for supplying the high resolution version of Fig. 4.

References

1. Nikaido H, Vaara M. Molecular basis of bacterial outer membrane permeability. *Microbiol. Rev.* 1985; 49:1–32. [PubMed: 2580220]
2. Nikaido H. Molecular basis of bacterial outer membrane permeability revisited. *Microbiol. Mol. Biol. Rev.* 2003; 67:593–656. [PubMed: 14665678]
3. Kamio K, Nikaido H. Outer membrane of *Salmonella typhimurium*: accessibility of phospholipid head groups to phospholipase C and cyanogen bromide-activated dextran. *Biochemistry.* 1976; 15:2561–2570. [PubMed: 820368]

4. Plésiat P, Nikaido H. Outer membranes of gram-negative bacteria are permeable to steroid probes. *Mol. Microbiol.* 1992; 6:1323–1333.
5. Nikaido H. Outer membrane barrier as a mechanism of antibiotic resistance. *Antimicrob. Agents Chemother.* 1989; 33:1831–1836. [PubMed: 2692513]
6. Nikaido H. Prevention of drug access to bacterial targets: permeability barriers and active efflux. *Science.* 1994; 264:382–388. [PubMed: 8153625]
7. Nikaido H. Multidrug efflux pumps of Gram-negative bacteria. *J. Bacteriol.* 1996; 178:5853–5859. [PubMed: 8830678]
8. Saier MH Jr, Paulsen IT, Sliwinski MK, Pao SS, Skurray RA, Nikaido H. Evolutionary origins of multidrug and drug-specific efflux pumps in bacteria. *FASEB J.* 1998; 12:265–274. [PubMed: 9506471]
9. Li X-Z, Nikaido H. Efflux-mediated drug resistance in bacteria. *Drugs.* 2004; 64:159–204. [PubMed: 14717618]
10. Li X-Z, Nikaido H. Efflux-mediated drug resistance in bacteria: An update. *Drugs.* 2009; 69:1555–1623. [PubMed: 19678712]
11. Wandersman C. Secretion across the bacterial outer membrane. *Trends Genet.* 1992; 8:317–322. [PubMed: 1365398]
12. Paulsen IT, Park JH, Choi PS, Saier MH Jr. A family of gram-negative bacterial outer membrane factors that function in the export of proteins, carbohydrates, drugs and heavy metals from gram-negative bacteria. *FEMS Microbiol. Lett.* 1997; 156:1–8. [PubMed: 9368353]
13. Dinh T, Paulsen IT, Saier MH Jr. A family of extracytoplasmic proteins that allow transport of large molecules across the outer membranes of gram-negative bacteria. *J. Bacteriol.* 1994; 176:3825–3831. [PubMed: 8021163]
14. Vaara M. Antibiotic-supersusceptible mutants of *Escherichia coli* and *Salmonella typhimurium*. *Antimicrob. Agents Chemother.* 1993; 37:2255–60. [PubMed: 8285603]
15. Nikaido H, Basina M, Nguyen V, Rosenberg EY. Multidrug efflux pump AcrAB of *Salmonella typhimurium* excretes only those β -lactam antibiotics containing lipophilic side chains. *J. Bacteriol.* 1998; 180:4686–92. [PubMed: 9721312]
16. Lee A, Mao MS, Warren MS, Misty A, Hoshino K, Okamura R, Ishida H, Lomovskaya O. Interplay between efflux pumps may provide either additive or multiplicative effects on drug resistance. *J. Bacteriol.* 2000; 182:3142–3150. [PubMed: 10809693]
17. Tal N, Schuldiner S. A coordinated network of transporters with overlapping specificities provides a robust survival strategy. *Proc. Natl. Acad. Sci. USA.* 2009 e-published.
18. Ma D, Cook DN, Alberti M, Pon NG, Nikaido H, Hearst JE. Molecular cloning and characterization of *acrA* and *acrE* genes of *Escherichia coli*. *J. Bacteriol.* 1993; 175:6299–6313. [PubMed: 8407802]
19. Poole K, Krebs K, McNally C, Neshat S. Multiple antibiotic resistance in *Pseudomonas aeruginosa*: Evidence for involvement of an efflux operon. *J. Bacteriol.* 1993; 175:7363–7372. [PubMed: 8226684]
20. Li XZ, Livermore DM, Nikaido H. Role of efflux pump(s) in intrinsic resistance of *Pseudomonas aeruginosa*: Resistance to tetracycline, chloramphenicol, and norfloxacin. *Antimicrob. Agents Chemother.* 1994; 38:1732–1741. [PubMed: 7986003]
21. Li XZ, Ma D, Livermore DM, Nikaido H. Role of efflux pump(s) in intrinsic resistance of *Pseudomonas aeruginosa*: active efflux as a contributing factor to beta-lactam resistance. *Antimicrob. Agents Chemother.* 1994; 38:1742–1752. [PubMed: 7986004]
22. Li XZ, Nikaido H, Poole K. Role of *mexA-mexB-oprM* in antibiotic efflux in *Pseudomonas aeruginosa*. *Antimicrob. Agents Chemother.* 1995; 39:1948–1953. [PubMed: 8540696]
23. Fralick JA. Evidence that TolC is required for functioning of the Mar/AcrAB efflux pump of *Escherichia coli*. *J. Bacteriol.* 1996; 178:5803–5805. [PubMed: 8824631]
24. Ma D, Cook DN, Alberti M, Pon NG, Nikaido H, Hearst JE. Genes *acrA* and *acrB* encode a stress-induced efflux system of *Escherichia coli*. *Mol. Microbiol.* 1995; 16:45–55. [PubMed: 7651136]
25. Smith IC, Auger M, Jarrell HC. Molecular details of anesthetic-lipid interaction. *Ann. N. Y. Acad. Sci.* 1991; 625:668–684. [PubMed: 2058915]

26. Lomovskaya O, Lewis K. Emr, an *Escherichia coli* locus for multidrug resistance. *Proc. Natl. Acad. Sci. USA*. 1992; 89:8938–8942. [PubMed: 1409590]
27. Furukawa H, Tsay JT, Jackowski S, Takamura Y, Rock CO. Thiolactomycin resistance in *Escherichia coli* is associated with the multidrug resistance efflux pump encoded by *emrAB*. *J. Bacteriol.* 1993; 175:3723–3729. [PubMed: 8509326]
28. Kobayashi N, Nishino K, Yamaguchi A. Novel macrolide-specific ABC-type efflux transporter in *Escherichia coli*. *J. Bacteriol.* 2001; 183:5639–5644. [PubMed: 11544226]
29. Yum S, Xu Y, Shunfu Piao S, Sim S, Kim H-M, Jo W-S, Kim K-J, Kweon H-S, Jeong M-H, Jeon H, Lee K, Ha N-C. Crystal structure of the periplasmic component of a tripartite macrolide-specific efflux pump. *J. Mol. Biol.* 2009; 387:1286–1297. [PubMed: 19254725]
30. Tikhonova EB, Devroy VK, Lau SY, Zgurskaya HI. Reconstitution of the *Escherichia coli* macrolide transporter: the periplasmic membrane fusion protein MacA stimulates the ATPase activity of MacB. *Mol. Microbiol.* 2007; 63:895–910. [PubMed: 17214741]
31. Nikaido H. Multidrug resistance in bacteria. *Annu. Rev. Biochem.* 2009; 79:119–146. [PubMed: 19231985]
32. Poole K. Efflux-mediated antimicrobial resistance. *J. Antimicrob. Chemother.* 2005; 56:20–51. [PubMed: 15914491]
33. Murakami S. Multidrug transporter, AcrB---the pumping mechanism. *Curr. Op. Struct. Biol.* 2008; 18:459–465.
34. Pos KM. Drug transport mechanism of the AcrB efflux pump. *Biochim. Biophys. Acta.* 2009; 1794:782–793. [PubMed: 19166984]
35. Higgins CF. Multiple molecular mechanisms for multidrug resistance transporters. *Nature*. 2007; 446:749–757. [PubMed: 17429392]
36. Nikaido H, Takatsuka Y. Mechanisms of RND multidrug efflux pumps. *Biochim. Biophys. Acta.* 2009; 1794:769–781. [PubMed: 19026770]
37. Nishino K, Nikaido E, Yamaguchi A. Regulation and physiological function of multidrug efflux pumps in *Escherichia coli* and *Salmonella*. *Biochim. Biophys. Acta.* 2009; 1794:834–843. [PubMed: 19230852]
38. Zgurskaya HI, Yamada Y, Tikhonova EB, Ge Q, Krishnamoorthy G. Structural and functional diversity of bacterial membrane fusion proteins. *Biochim. Biophys. Acta.* 2009; 1794:794–807. [PubMed: 19041958]
39. Nakajima H, Kobayashi K, Kobayashi M, Asako H, Aono R. Overexpression of the *robA* gene increases organic solvent tolerance and multiple antibiotic and heavy metal resistance in *Escherichia coli*. *Appl. Environ. Microbiol.* 1995; 61:2302–2307. [PubMed: 7793951]
40. White DG, Goldman JD, Demple B, Levy SB. Role of the *acrAB* locus in organic solvent tolerance mediated by expression of *marA*, *soxS*, or *robA* in *Escherichia coli*. *J. Bacteriol.* 1997; 179:6122–6126. [PubMed: 9324261]
41. Lomovskaya O, Warren MS, Lee A, Galazzo J, Fronko R, Lee M, Blais J, Cho D, Chamberland S, Renau T, Leger R, Hecker S, Watkins W, Hoshino K, Ishida H, Lee VJ. Identification and characterization of inhibitors of multidrug resistance efflux pumps in *Pseudomonas aeruginosa*: novel agents for combination therapy. *Antimicrob. Agents Chemother.* 2001; 45:105–116. [PubMed: 11120952]
42. Zgurskaya HI, Nikaido H. Cross-linked complex between oligomeric periplasmic lipoprotein AcrA and the inner-membrane-associated multidrug efflux pump AcrB from *Escherichia coli*. *J. Bacteriol.* 2000; 182:4264–4267. [PubMed: 10894736]
43. Tikhonova EB, Zgurskaya HI. AcrA, AcrB, and TolC of *Escherichia coli* form a stable intermembrane multidrug efflux complex. *J. Biol. Chem.* 2004; 279:32116–32124. [PubMed: 15155734]
44. Murakami S, Nakashima R, Yamashita E, Yamaguchi A. Crystal structure of bacterial multidrug efflux transporter AcrB. *Nature*. 2002; 419:587–93. 2002. [PubMed: 12374972]
45. Sennhauser G, Bukowska MA, Briand C, Grütter MG. Crystal structure of the multidrug exporter MexB from *Pseudomonas aeruginosa*. *J. Mol. Biol.* 2009; 389:134–145. [PubMed: 19361527]

46. Akama H, Matsuura T, Kashiwagi S, Yoneyama H, Narita S, Tsukihara T, Nakagawa A, Nakae T. Crystal structure of the membrane fusion protein, MexA, of the multidrug transporter in *Pseudomonas aeruginosa*. *J. Biol. Chem.* 2004; 279:25939–42. [PubMed: 15117957]
47. Higgins MK, Bokma E, Koronakis E, Hughes C, Koronakis V. Structure of the periplasmic component of a bacterial drug efflux pump. *Proc. Natl. Acad. Sci. USA.* 2004; 101:9994–9999. [PubMed: 15226509]
48. Mikolosko J, Bobyk K, Zgurskaya HI, Ghosh P. Conformational flexibility in the multidrug efflux system protein AcrA. *Structure.* 2006; 14:577–587. [PubMed: 16531241]
49. Koronakis V, Sharff A, Koronakis E, Luisi B, Hughes C. Crystal structure of the bacterial membrane protein TolC central to multidrug efflux and protein export. *Nature.* 2000; 405:914–9. [PubMed: 10879525]
50. Akama H, Kanemaki M, Yoshimura M, Tsukihara T, Kashiwagi T, Yoneyama H, Narita S, Nakagawa A, Nakae T. Crystal structure of the drug discharge outer membrane protein, OprM, of *Pseudomonas aeruginosa*: dual modes of membrane anchoring and occluded cavity end. *J. Biol. Chem.* 2004; 279:52816–52819. [PubMed: 15507433]
51. Tseng TT, Gratwick KS, Kollman J, Park D, Nies DH, Goffeau A, Saier MH Jr. The RND permease superfamily: an ancient, ubiquitous and diverse family that includes human disease and developmental proteins. *J. Mol. Microbiol. Biotechnol.* 1999; 1:107–125. [PubMed: 10941792]
52. Zgurskaya HI, Nikaido H. AcrA is a highly asymmetric protein capable of spanning the periplasm. *J. Mol. Biol.* 1999; 285:409–4. [PubMed: 9878415]
53. Symmons MF, Bokma E, Koronakis E, Hughes C, Koronakis V. The assembled structure of a complete tripartite bacterial multidrug efflux pump. *Proc. Natl. Acad. Sci. U.S.A.* 2009; 106:7173–7176. [PubMed: 19342493]
54. Ip H, Stratton K, Zgurskaya H, Liu J. pH-induced conformational changes of AcrA, the membrane fusion protein of *Escherichia coli* multidrug efflux system. *J. Biol. Chem.* 2003; 278:50474–50482. [PubMed: 14523004]
55. Tamura N, Murakami S, Oyama Y, Ishiguro M, Yamaguchi A. Direct interaction of multidrug efflux transporter AcrB and outer membrane channel TolC detected via site-directed disulfide cross-linking. *Biochemistry.* 2005; 44:11115–11121. [PubMed: 16101295]
56. Bavro VN, Pietras Z, Furnham N, Pérez-Cano L, Fernández-Recio J, Pei XY, Misra R, Luisi B. Assembly and channel opening in a bacterial drug efflux machine. *Mol. Cell.* 2008; 30:114–121. [PubMed: 18406332]
57. Elkins CA, Nikaido H. Chimeric analysis of AcrA function reveals the importance of its C-terminal domain in its interaction with the AcrB multidrug efflux pump. *J. Bacteriol.* 2003; 185:5349–56. [PubMed: 12949086]
58. Touzé T, Eswaran J, Bokma E, Koronakis E, Hughes C, Koronakis V. Interactions underlying assembly of the *Escherichia coli* AcrAB-TolC multidrug efflux system. *Mol. Microbiol.* 2004; 53:697–706. [PubMed: 15228545]
59. Krishnamoorthy G, Tikhonova EB, Zgurskaya HI. Fitting periplasmic membrane fusion proteins to inner membrane transporters: mutations that enable *Escherichia coli* AcrA to function with *Pseudomonas aeruginosa* MexB. *J. Bacteriol.* 2007; 190:691–698. [PubMed: 18024521]
60. Lobedanz S, Bokma E, Symmons MF, Koronakis E, Hughes C, Koronakis V. A periplasmic coiled-coil interface underlying TolC recruitment and the assembly of bacterial drug efflux pumps. *Proc. Natl. Acad. Sci. USA.* 2007; 104:4612–4617. [PubMed: 17360572]
61. Eswaran J, Koronakis E, Higgins MK, Hughes C, Koronakis V. Three's company: component structures bring a closer view of tripartite drug efflux pumps. *Curr. Opin. Struct. Biol.* 2004; 14:741–7. [PubMed: 15582398]
62. Reffay M, Gambin Y, Benabdelhak H, Phan G, Taulier N, Ducruix A, Hodges RS, Urbach W. Tracking membrane protein association in model membranes. *PLoS One.* 2009; 4:e5035. [PubMed: 19337368]
63. Narita S, Eda S, Yoshihara E, Nakae T. Linkage of the efflux-pump expression level with substrate extrusion rate in the MexAB-OprM efflux pump of *Pseudomonas aeruginosa*. *Biochem. Biophys. Res. Comm.* 2003; 308:922–926. [PubMed: 12927807]

64. Aires JR, Nikaido H. Aminoglycosides are captured from both periplasm and cytoplasm by the AcrD multidrug efflux transporter of *Escherichia coli*. J. Bacteriol. 2005; 187:1923–1929. [PubMed: 15743938]
65. Mima T, Joshi S, Gomez-Escalada M, Schweizer HP. Identification and characterization of TriABC-OpmH, a triclosan efflux pump of *Pseudomonas aeruginosa* requiring two membrane fusion proteins. J. Bacteriol. 2007; 189:7600–7609. [PubMed: 17720796]
66. Gristwood T, Fineran PC, Everson L, Salmond GP. PigZ, a TetR/AcrR family repressor, modulates secondary metabolism via the expression of a putative four-component resistance-nodulation-cell-division efflux pump. Mol. Microbiol. 2008; 69:418–435. [PubMed: 18485072]
67. Zgurskaya HI, Nikaido H. Bypassing the periplasm: reconstitution of the AcrAB multidrug efflux pump of *Escherichia coli*. Proc. Natl. Acad. Sci. USA. 1999; 96:7190–7195. [PubMed: 10377390]
68. Su CC, Yu EW. Ligand-transporter interaction in the AcrB multidrug efflux pump determined by fluorescence polarization assay. FEBS Lett. 2007; 581:4972–4976. [PubMed: 17910961]
69. Ocaktan A, Yoneyama H, Nakae T. Use of fluorescence probes to monitor function of the subunit proteins of the MexA-MexB-oprM drug extrusion machinery in *Pseudomonas aeruginosa*. J. Biol. Chem. 1997; 272:21964–21969. [PubMed: 9268332]
70. Murakami S, Tamura N, Saito A, Hirata T, Yamaguchi A. Extramembrane central pore of multidrug exporter AcrB in *Escherichia coli* plays an important role in drug transport. J. Biol. Chem. 2004; 279:3743–3748. [PubMed: 14576158]
71. Nagano K, Nikaido H. Kinetic behavior of the major multidrug efflux pump AcrB of *Escherichia coli*. Proc. Natl. Acad. Sci. 2009; 106:5854–5858. [PubMed: 19307562]
72. Mazzariol A, Cornaglia G, Nikaido H. Contributions of the AmpC β -lactamase and the AcrAB multidrug efflux system in intrinsic resistance of *Escherichia coli* K-12 to β -lactams. Antimicrob. Agents Chemother. 2000; 44:1387–1390. [PubMed: 10770787]
73. Nikaido H, Normark S. Sensitivity of *Escherichia coli* to various β -lactams is determined by the interplay of outer membrane permeability and degradation by periplasmic β -lactamases: a quantitative predictive treatment. Mol. Microbiol. 1987; 1:29–36. [PubMed: 3330755]
74. Chang TY, Chang CC, Ohgami N, Yamauchi Y. Cholesterol sensing, trafficking, and esterification. Annu. Rev. Cell Dev. Biol. 2006; 22:129–157. [PubMed: 16753029]
75. Infante RE, Abi-Mosleh L, Radhakrishnan A, Dale JD, Brown MS, Goldstein JL. Purified NPC1 protein. I. Binding of cholesterol and oxysterols to a 1278-amino acid membrane protein. J. Biol. Chem. 2008; 283:1052–1063. [PubMed: 17989073]
76. Infante RE, Radhakrishnan A, Abi-Mosleh L, Kinch LN, Wang ML, Grishin NV, Goldstein JL, Brown MS. Purified NPC1 protein: II. Localization of sterol binding to a 240-amino acid soluble luminal loop. J. Biol. Chem. 2008; 283:1064–1075. [PubMed: 17989072]
77. Elkins CA, Nikaido H. Substrate specificity of the RND-type multidrug efflux pumps AcrB and AcrD of *Escherichia coli* is determined predominantly by two large periplasmic loops. J. Bacteriol. 2002; 184:6490–6498. [PubMed: 12426336]
78. Eda S, Maseda H, Nakae T. An elegant means of self-protection in gram-negative bacteria by recognizing and extruding xenobiotics from the periplasmic space. J. Biol. Chem. 2003; 278:2085–2088. [PubMed: 12460990]
79. Tikhonova EB, Wang Q, Zgurskaya HI. Chimeric analysis of the multicomponent multidrug efflux transporters from Gram-negative bacteria. J. Bacteriol. 2002; 184:6499–6507. [PubMed: 12426337]
80. Wehmeier C, Schuster S, Fährnich E, Kern WV, Bohnert JA. Site-directed mutagenesis reveals amino acid residues in the *Escherichia coli* RND efflux pump AcrB that confer macrolide resistance. Antimicrob. Agents Chemother. 2009; 53:329–330. [PubMed: 18936189]
81. Mao W, Warren MS, Black DS, Satou T, Murata T, Nishino T, Gotoh N, Lomovskaya O. On the mechanism of substrate specificity by resistance nodulation division (RND)-type multidrug resistance pumps: the large periplasmic loops of MexD from *Pseudomonas aeruginosa* are involved in substrate recognition. Mol. Microbiol. 2002; 46:889–901. [PubMed: 12410844]
82. Bohnert JA, Schuster S, Fährnich E, Trittler R, Kern WV. Altered spectrum of multidrug resistance associated with a single point mutation in the *Escherichia coli* RND-type MDR efflux pump YhiV (MdtF). J. Antimicrob. Chemother. 2007; 59:1216–1222. [PubMed: 17062614]

83. Middlemiss JK, Poole K. Differential impact of MexB mutations on substrate selectivity of the MexAB-OprM multidrug efflux pump of *Pseudomonas aeruginosa*. J. Bacteriol. 2004; 186:1258–1269. [PubMed: 14973037]
84. Goldberg M, Pribyl T, Juhnke S, Nies DH. Energetics and topology of CzcA, a cation/proton antiporter of the resistance-nodulation-cell division protein family. J. Biol. Chem. 1999; 274:26065–26070. [PubMed: 10473554]
85. Guan L, Nakae T. Identification of essential charged residues in transmembrane segments of the multidrug transporter MexB of *Pseudomonas aeruginosa*. J. Bacteriol. 2001; 183:1734–1739. [PubMed: 11160105]
86. Takatsuka Y, Nikaido H. Threonine-978 in the transmembrane segment of the multidrug efflux pump AcrB of *Escherichia coli* is crucial for drug transport as a probable component of the proton relay network. J. Bacteriol. 2006; 188:7284–7289. [PubMed: 17015667]
87. Yu EW, Aires JR, Nikaido H. AcrB multidrug efflux pump of *Escherichia coli*: composite substrate-binding cavity of exceptional flexibility generates its extremely wide substrate specificity. J. Bacteriol. 2003; 185:5657–5664. [PubMed: 13129936]
88. Yu EW, Aires JR, McDermott G, Nikaido H. A periplasmic drug-binding site of the AcrB multidrug efflux pump: a crystallographic and site-directed mutagenesis study. J. Bacteriol. 2005; 187:6804–6815. [PubMed: 16166543]
89. Hearn EM, Gray MR, Foght JM. Mutations in the central cavity and periplasmic domain affect efflux activity of the resistance-nodulation-division pump EmhB from *Pseudomonas fluorescens* cLP6a. J. Bacteriol. 2006; 188:115–123. [PubMed: 16352827]
90. Franke S, Grass G, Rensing C, Nies DH. Molecular analysis of the copper-transporting efflux system CusCFBA of *Escherichia coli*. J. Bacteriol. 2003; 185:3804–3812. [PubMed: 12813074]
91. Dastidar V, Mao W, Lomovskaya O, Zgurskaya HI. Drug-induced conformational changes in multidrug efflux transporter AcrB from *Haemophilus influenzae*. J. Bacteriol. 2007; 189:5550–5558. [PubMed: 17526713]
92. Ruetz S, Gros P. Phosphatidylcholine translocase: a physiological role for the *mdr2* gene. Cell. 1994; 77:1071–1081. [PubMed: 7912658]
93. Mao W, Warren MS, Lee A, Mistry A, Lomovskaya O. MexXY-OprM efflux pump is required for antagonism of aminoglycosides by divalent cations in *Pseudomonas aeruginosa*. Antimicrob. Agents Chemother. 2001; 45:2001–2007. [PubMed: 11408215]
94. Schmidt TG, Skerra A. The Strep-tag system for one-step purification and high-affinity detection or capturing of proteins. Nature Protocols. 2007; 2:1528–35.
95. Hanada K, Yamato I, Anraku Y. Purification and reconstitution of *Escherichia coli* proline carrier using a site specifically cleavable fusion protein. J. Biol. Chem. 1988; 263:7181–7185. [PubMed: 3130379]
96. Bagai I, Liu W, Rensing C, Blackburn NJ, McEvoy MM. Substrate-linked conformational change in the periplasmic component of a Cu(I)/Ag(I) efflux system. J. Biol. Chem. 2007; 282:35695–35702. [PubMed: 17893146]
97. Bagai I, Liu W, Rensing C, Blackburn NJ, McEvoy MM. Direct metal transfer between periplasmic proteins identifies a bacterial copper chaperone. Biochemistry. 2008; 47:11408–11414. [PubMed: 18847219]
98. Yu EW, McDermott G, Zgurskaya HI, Nikaido H, Koshland DE Jr. Structural basis of multiple drug-binding capacity of the AcrB multidrug efflux pump. Science. 2003; 300:976–980. [PubMed: 12738864]
99. Törnroth-Horsefield S, Gourdon P, Horsefield R, Brive L, Yamamoto N, Mori H, Snijder A, Neutze R. Crystal structure of AcrB in complex with a single transmembrane subunit reveals another twist. Structure. 2007; 15:1663–1673. [PubMed: 18073115]
100. Das D, Xu QS, Lee JY, Ankoudinova I, Huang C, Lou Y, Degiovanni A, Kim R, Kim S-H. Crystal structure of the multidrug efflux transporter AcrB at 3.1 Å resolution reveals the N-terminal region with conserved amino acids. J. Struct. Biol. 2007; 158:494–502. [PubMed: 17275331]
101. Drew D, Klepsch MM, Newstead S, Flaig R, de Gier J-W, Iwata S, Beis K. The structure of the efflux pump AcrB in complex with bile acid. Mol. Memb. Biol. 2008; 25:67–682.

102. Su CC, Li M, Gu R, Takatsuka Y, McDermott G, Nikaido H, Yu EW. Conformation of the AcrB multidrug efflux pump in mutants of the putative proton relay pathway. *J. Bacteriol.* 2006; 188:7290–7296. [PubMed: 17015668]
103. Murakami S, Nakashima R, Yamashita E, Matsumoto T, Yamaguchi A. Crystal structures of a multidrug transporter reveal a functionally rotating mechanism. *Nature.* 2006; 443:173–179. [PubMed: 16915237]
104. Seeger MA, Schiefner A, Eicher T, Verrey F, Diederichs K, Pos KM. Structural asymmetry of AcrB trimer suggests a peristaltic pump mechanism. *Science.* 2006; 313:1295–1298. [PubMed: 16946072]
105. Sennhauser G, Amstutz P, Briand C, Storchenegger O, Grütter MG. Drug export pathway of multidrug exporter AcrB revealed by DARPin inhibitors. *PLoS Biol.* 2007; 5:e7. [PubMed: 17194213]
106. Seeger MA, von Ballmoos C, Verrey F, Pos KM. Crucial role of Asp408 in the proton translocation pathway of multidrug transporter AcrB: evidence from site-directed mutagenesis and carbodiimide labeling. *Biochemistry.* 2009; 48:5801–5812. [PubMed: 19425588]
107. Bohnert JA, Schuster S, Seeger MA, Fährnich E, Pos KM, Kern WV. Site-directed mutagenesis reveals putative substrate binding residues in the *Escherichia coli* RND efflux pump AcrB. *J. Bacteriol.* 2008; 190:8225–8229. [PubMed: 18849422]
108. Lomovskaya O, Totrov M. Vacuuming the periplasm. *J. Bacteriol.* 2005; 187:1879–1883. [PubMed: 15743933]
109. Bohnert JA, Kern WV. Selected arylpiperazines are capable of reversing multidrug resistance in *Escherichia coli* overexpressing RND efflux pumps. *Antimicrob. Agents Chemother.* 2005; 49:849–852. [PubMed: 15673787]
110. Takatsuka Y, Nikaido H. Site-directed disulfide cross-linking shows that cleft flexibility in the periplasmic domain is needed for the multidrug efflux pump AcrB of *Escherichia coli*. *J. Bacteriol.* 2007; 189:8677–8684. [PubMed: 17905989]
111. Bardwell JC, McGovern K, Beckwith J. Identification of a protein required for disulfide bond formation in vivo. *Cell.* 1991; 67:581–589. [PubMed: 1934062]
112. Kamitani S, Akiyama Y, Ito K. Identification and characterization of an *Escherichia coli* gene required for the formation of correctly folded alkaline phosphatase, a periplasmic enzyme. *EMBO J.* 1992; 11:57–62. [PubMed: 1740115]
113. Kenyon GL, Bruice TW. Novel sulfhydryl reagents. *Methods Enzymol.* 1977; 47:407–430. [PubMed: 927196]
114. Seeger MA, von Ballmoos C, Eicher T, Brandstatter L, Verrey F, Diederichs K, Pos KM. Engineered disulfide bonds support the functional rotation mechanism of multidrug efflux pump AcrB. *Nat. Struct. Mol. Biol.* 2008; 15:199–205. [PubMed: 18223659]
115. Takatsuka Y, Nikaido H. Covalently linked trimer of AcrB multidrug efflux pump provides support for the functional rotating mechanism. *J. Bacteriol.* 2009; 191:1729–1737. [PubMed: 19060146]
116. Baranova N, Nikaido H. The BaeSR two-component regulatory system activates transcription of the *yegMNOB* (*mdtABCD*) transporter gene cluster in *Escherichia coli* and increases its resistance to novobiocin and deoxycholate. *J. Bacteriol.* 2002; 184:4168–4176. [PubMed: 12107134]
117. Nagakubo S, Nishino K, Hirata T, Yamaguchi A. The putative response regulator BaeR stimulates multidrug resistance of *Escherichia coli* via a novel multidrug exporter system, MdtABC. *J. Bacteriol.* 2006; 184:4161–4167. [PubMed: 12107133]
118. Kumar A, Worobec EA. Cloning, sequencing, and characterization of the SdeAB multidrug efflux pump of *Serratia marcescens*. *Antimicrob. Agents Chemother.* 2005; 49:1495–1501. [PubMed: 15793131]
119. Crossman LC, Gould VC, Dow JM, Vernikos GS, Okazaki A, Sebahia M, Saunders D, Arrowsmith C, Carver T, Peters N, Adlem E, Kerhornou A, Lord A, Murphy L, Seeger K, Squares R, Rutter S, Quail MA, Rajandream M-A, Harris D, Churcher C, Bentley SD, Parkhill JD, Thomson NR, Avison MB. The complete genome, comparative and functional analysis of

- Stenotrophomonas maltophilia* reveals an organism heavily shielded by drug resistance determinants. *Genome Biol.* 2008; 9:R74. [PubMed: 18419807]
120. Lakaye B, Dubus A, Lepage S, Gros Lambert S, Frère J-M. When drug inactivation renders the target irrelevant to antibiotic resistance: a case story with β -lactams. *Mol. Microbiol.* 1999; 31:89–101. [PubMed: 9987113]
 121. Tal N, Schuldiner S. A coordinated network of transporters with overlapping specificities provides a robust survival strategy. *Proc. Natl. Acad. Sci. USA.* 2009; 106:9051–9056. [PubMed: 19451626]
 122. Adam Y, Tayer N, Rottem D, Schreiber G, Schuldiner S. The fast release of sticky protons: Kinetics of substrate binding and proton release in a multidrug transporter. *Proc. Natl. Acad. Sci. USA.* 2007; 104:17989–17994. [PubMed: 17984053]
 123. Rotem D, Steiner-Mordoch S, Schuldiner S. Identification of tyrosine residues critical for the function of an ion-coupled multidrug transporters. *J. Biol. Chem.* 2006; 281:18715–18722. [PubMed: 16672221]
 124. Schuldiner S. EmrE, a model for studying evolution and mechanism of ion-coupled transporters. *Biochim. Biophys. Acta.* 2009; 1794:748–762. [PubMed: 19167526]
 125. Lovmar M, Nilsson K, Lukk E, Vimberg V, Tenson T, Ehrenberg M. Erythromycin resistance by L4/L22 mutations and resistance masking by drug efflux pump deficiency. *EMBO J.* 2009; 28:736–744. [PubMed: 19197244]
 126. Moore SD, Sauer RT. Revisiting the mechanism of macrolide-antibiotic resistance mediated by ribosomal protein L22. *Proc. Natl. Acad. Sci. USA.* 2008; 105:18261–18266. [PubMed: 19015512]
 127. Cagliero C, Mouline C, Cloeckert A, Payot S. Synergy between efflux pump CmeABC and modifications in the ribosomal proteins L4 and L22 in conferring macrolide resistance in *Campylobacter jejuni* and *Campylobacter coli*. *Antimicrob. Agents Chemother.* 2006; 50:3893–3896. [PubMed: 16940070]
 128. Elf J, Nilsson K, Tenson T, Ehrenberg M. Bistable bacterial growth rate in response to antibiotics with low membrane permeability. *Phys. Rev. Lett.* 2006; 97:258104. [PubMed: 17280399]
 129. Fange D, Nilsson K, Tenson T, Ehrenberg M. Drug efflux pump deficiency and drug target resistance masking in growing bacteria. *Proc. Natl. Acad. Sci. USA.* 2009; 106:8215–8220. [PubMed: 19416855]
 130. Jeannot K, Elsen S, Köhler T, Attree I, van Delden C, Plésiat P. Resistance and virulence of *Pseudomonas aeruginosa* clinical strains overproducing the MexCD-OprJ efflux pump. *Antimicrob. Agents Chemother.* 2008; 52:2455–2462. [PubMed: 18474583]
 131. Zhao Q, Li XZ, Mistry A, Srikumar R, Zhang L, Lomovskaya O, Poole K. Influence of the TonB energy-coupling protein on efflux-mediated multidrug resistance in *Pseudomonas aeruginosa*. *Antimicrob. Agents Chemother.* 1998; 42:2225–2231. [PubMed: 9736539]
 132. Zhao Q, Poole K. Differential effects of mutations in *tonB1* on intrinsic multidrug resistance and iron acquisition in *Pseudomonas aeruginosa*. *J. Bacteriol.* 2002; 184:2045–2049. [PubMed: 11889114]
 133. Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC, Ferrin TE. UCSF Chimera—a visualization system for exploratory research and analysis. *J. Comput. Chem.* 2004; 25:1605–1612. [PubMed: 15264254]

Fig. 1.

An early schematic view of the tripartite pump complex. Note that amphiphilic drugs (empty and filled-in rectangles represent hydrophobic and hydrophilic parts of the molecule) are hypothesized to be captured from the periplasm-plasma membrane interface or possibly from the cytosol (or the cytosol-membrane interface). For the latter process, two possible pathways are envisaged (dashed arrows): either the substrate is flipped over to the outer surface of the membrane first and then follows the regular periplasmic capture pathway, or it follows a different capture pathway from the cytosol. For the AcrAB-TolC system, the efflux transporter corresponds to AcrB, the accessory protein to AcrA, and the OM channel to TolC. Modified from (7).

Fig. 2.
Examples of substrates for the AcrB pump.

Fig. 3.

Crystallographic structures of MexA, AcrB, and TolC. The structure of MexA (monomer), rather than that of AcrA, is shown here because the coordinates for the entire protein sequence have been published (53). The pictures were drawn by using Pymol (<http://www.pymol.org>), based on PDB files 1IWG (AcrB), 2V4D (MexA), and 2VDE (TolC). The approximate positions of membrane bilayers are shown in black horizontal lines. In the trimer structures of AcrB and TolC, each protomer is shown in cyan, mauve, and blue, and the MexA structure is colored in rainbow scheme from the blue of the N-terminus to the red of the C-terminus. In the periplasmic domain of AcrB, the boundaries of the “TolC-docking” and “Porter” subdomains are indicated. The large central cavity (red dotted lines) is connected to the periplasm through vestibules, the opening of one of which is shown as a red ellipse between the protomers colored in green and cyan. The apparently closed central pore that connects the central cavity to the top of TolC docking domain is indicated as red vertical lines. In MexA, the four domains are indicated. Although the AcrA and MexA are modified in the cell by the addition of a fatty acyl chain at the N-terminus plus the addition of glyceride (containing two additional fatty acyl chains), this is not shown in the figure. The β -barrel domain and the helical tunnel domain of TolC are indicated.

Fig. 4.

The model of AcrAB complex based on the recently refined AcrA structure and the cross-linking data. The periplasmic domain of an AcrB monomer is shown in a space-filling model, with each subdomain (PC for Periplasmic, C-terminal; PN for Periplasmic, N-terminal; DC for Docking, C-terminal; and DN, for Docking, N-terminal) colored differently. The AcrA is shown as a green ribbon model, and the domains [α -hairpin, lipoyl, β -barrel, and MP (membrane-proximal)] is identified in green letters. With permission from the authors (53).

Fig. 5. Kinetics of AcrAB-TolC-catalyzed efflux of nitrocefin (A), cephaloridine (B), and cefazolin (C). The rates of efflux from the periplasm (V_e) are plotted against the periplasmic drug concentration (C_p) for each drug. Nitrocefin efflux follows a classical Michaelis-Menten kinetics, but cephaloridine efflux shows a sigmoidal kinetics. For the cephaloridine and cefazolin the rates of β -lactamase-catalyzed hydrolysis (V_h) are also shown for comparison. From reference 71.

Fig. 6.

The view of the large cleft of AcrB periplasmic domain, seen from the outside. The opening of the cleft is approximately indicated by the large red ellipse. AcrB residues corresponding to those MexD residues affecting the substrate specificity (81), Phe610, Gln34, Glu89, and Thr676, are shown in space-filling models, as well as those that were shown to affect drug pumping by site-directed mutagenesis (88), Phe664, Phe666, and Glu673. The binding of drugs to the Asn109Ala mutant AcrB (88), to be discussed in section III.A, are shown in stick models (ciprofloxacin, yellow; nafcillin, green; ethidium, magenta; rhodamine 6G, salmon; and Phe-Arg- β -naphthylamide, orange).

Fig. 7.

Phospholipid extrusion assay of reconstituted AcrB. When a transmembrane pH gradient was generated by diluting the vesicles (internal pH, 7.0) into a buffer of pH 6.0, donor vesicles not containing any AcrB (bottom trace) do not show much change in fluorescence. Nor do the AcrB-containing vesicles in the absence of pH gradient (middle trace). However, when AcrB-containing vesicles are exposed to pH gradient, there is the expected gradual increase in NBD fluorescence in the presence of Mg^{2+} . Note that the rate of fluorescence increase is strongly accelerated if AcrA (15 $\mu\text{g/ml}$) is also added. From (67).

Fig. 8.

Proton flux assay of reconstituted AcrB. AcrB-containing vesicles were made in the presence of 1 mM pyranine, a fluorescent pH indicator, and the extravesicular pyranine was removed by gel filtration. The interior-acidic pH gradient was generated by diluting the vesicles containing 0.1 M KCl into 0.1 M NaCl and then by adding valinomycin (arrow). The acidic internal pH was maintained for several minutes in the absence of drug (curve 1). When erythromycin (curve 2), chloramphenicol (curve 3), cloxacillin (curve 4), glycocholate (curve 5), or taurocholate (curve 6) were present at 0.2 mM, the internal pH returned to its initial value more rapidly, presumably due to the proton efflux accompanying the influx of drugs by the AcrB antiporter. From (67).

Fig. 9.

Proton flux assay of reconstituted AcrD with streptomycin. AcrD-containing proteoliposomes were made in KCl, and an interior-acidic pH gradient was generated by diluting them into NaCl and by adding valinomycin at 30 s. The insets show the experimental set-up. The gray object denotes AcrD, with its periplasmic domain shown as ellipses. The black rectangle denotes AcrA, and the substrate (streptomycin, 70 μ M) is shown as "S". Curve 1, AcrD proteoliposomes containing Mg^{2+} and AcrA inside, but with no streptomycin. Curve 2, the same proteoliposome as in Curve 1, to which streptomycin was added to the external medium. Curve 3, AcrD-proteoliposomes made on the same day that contained Mg^{2+} , AcrA, and streptomycin in the intravesicular space. From (64).

Fig. 10.

Putative salt-bridge/H-bonding network in the transmembrane domain of the wild type AcrB protomer (in yellow stick model) and the Asp407Ala mutant (in green stick model), based on PDB files 1IWG and 2HQC. The view is along the line perpendicular to the membrane plane, from the periplasmic side. Note the prominent flipping of the Lys940 side-chain, which is also seen with other mutants of the network (not shown). From (102).

Fig. 11.

Periplasmic domain of the asymmetric crystal structure of AcrB (A) and the proposed functionally rotating mechanism (B), from the studies of Murakami et al (103), drawn by using Pymol from PDB file 2DRD. The view is from the top (periplasm), and the transmembrane domains were removed for clarity. The three protomers each with a distinctive conformation are shown in different colors. In part A, the binding protomer (green) contains the bound ligand minocycline (in stick model). When the protonation of the Asp407, Asp408, or both changes the alignment of proton relay residues shown in Fig. 10, the conformational alteration is transmitted to the periplasmic domain, which is changed into the Extrusion protomer that extrudes the drug molecule outward. After reprotonation of Asp residue(s) in the transmembrane domain, the initial conformation of the Access protomer is restored. Thus each protomer was proposed to go through cyclic changes of conformation (part B, where each of the protomer retains its initial color scheme) in a functionally rotating manner (103-105). Note that the large external cleft of the periplasmic domain becomes closed in the Extrusion protomer.

Fig. 12.

Cut-out view of the binding protomer with the bound minocycline in a green stick representation (A) and the extruding protomer (B), both from PDB file 2DRD, drawn by using the UCSF Chimera package (133). The wide passageway from the external cleft (dashed green arrow) seen A becomes closed in B. The passageway in A was also enlarged so that the extremely hydrophobic surface in this area can be seen more easily. Modified from (36) with permission from Elsevier.

Fig. 13.

Use of the covalently linked AcrB trimer showing that inactivation of one protomer is sufficient in inactivating the entire trimer. Covalently linked trimers were constructed as shown in the scheme on the left, with only one protomer containing Phe666Cys and Gln830Cys mutations. These residues are located on the opposite walls of the cleft, and the linked AcrB trimer remains active as seen by the slow net entry of the dye ethidium bromide (because of efflux), when it is expressed in *dsbA* mutant strain, in which the disulfide bond formation is inefficient in the periplasmic space (curve 1, part A). Real-time inhibition of pump function through closure of the periplasmic cleft is then achieved by the addition of a fast-acting methanethiosulfonate SH crosslinker MTS-2-MTS (curve 2). However, the pump stopped functioning when the two Cys residues were cross-linked by the addition of a cross-linking methanethiosulfonate agent MTS-2-MTS (curve 3).