Published in final edited form as:

Biochem Biophys Res Commun. 2011 April 1; 407(1): 79–85. doi:10.1016/j.bbrc.2011.02.113.

AUGMENTED OXYGEN-MEDIATED TRANSCRIPTIONAL ACTIVATION OF CYTOCHROME P450 (CYP)1A EXPRESSION AND INCREASED SUSCEPTIBILITIES TO HYPEROXIC LUNG INJURY IN TRANSGENIC MICE CARRYING THE HUMAN CYP1A1 OR MOUSE 1A2 PROMOTER *IN VIVO*[¶]

WEIWU JIANG^{1,*}, XANTHI I. COUROUCL^{1,*}, LIHUA WANG¹, ROBERTO BARRIOS², and BHAGAVATULA MOORTHY^{1,†}

¹ DEPARTMENT OF PEDIATRICS, BAYLOR COLLEGE OF MEDICINE, HOUSTON, TX

² WEILL CORNELL MEDICAL COLLEGE, HOUSTON, TX

Abstract

Supplemental oxygen administration is frequently administered to preterm and term infants having pulmonary insufficiency. However, hyperoxia contributes to the development of bronchopulmonary dysplasia (BPD) in premature infants. Cytochrome P450 (CYP)A enzymes have been implicated in hyperoxic lung injury. In this study, we tested the hypothesis that hyperoxia induces CYP1A1 and 1A2 enzymes by transcriptional activation of the corresponding promoters *in vivo*, and transgenic mice expressing the human CYP1A1 or the mouse 1A2 promoter would be more susceptible to hyperoxic lung injury than wild type (WT) mice. Adult WT (CD-1) (12 week-old) mice, transgenic mice carrying a 10 kb human CYP1A1 promoter and the luciferase (luc) reporter gene (CYP1A1-luc), or mice expressing the mouse CYP1A2 promoter (CYP1A2-luc) were maintained in room air or exposed to hyperoxia for 24–72 h. Hyperoxia exposure of CYP1A1-luc mice for 24 and 48 h resulted in 2.5- and 1.25-fold increases, respectively in signal intensities, compared to room air controls. By 72 h, the induction had declined to control levels. CYP1A2-luc mice also showed enhanced luc expression after 24–48 h, albeit to a lesser extent than those expressing the CYP1A1 promoter. Also, these mice showed decreased levels of endogenous CYP1A1 and 1A2 expression after prolonged hyperoxia, and were also more susceptible to lung injury than similarly exposed WT mice, with CYP1A2-luc mice showing the greatest injury. Our results support the hypothesis that hyperoxia induces CYP1A enzymes by transcriptional activation of its corresponding promoters, and that decreased endogenous expression of these enzymes contribute to the increased susceptibilities to hyperoxic lung injury in the transgenic animals. In summary, this is the first report providing direct evidence of hyperoxia-mediated induction of CYP1A1 and CYP1A2 expression *in vivo* by mechanisms entailing transcriptional activation of the corresponding promoters, a phenomenon that has implications for hyperoxic lung injury, as well as other pathologies caused by oxidative stress.

[¶]This work was supported in part by National Institute of Health grants RO1HL070921, R01HL087174, and RO1ES01969 to BM.

© 2011 Elsevier Inc. All rights reserved.

[†]Corresponding Author: Bhagavatula Moorthy, Ph.D., Associate Professor of Pediatrics, Baylor College of Medicine, 6621 Fannin, F.C. 530.01, Houston, TX 77030, bmoorthy@bcm.edu, Tel: (832) 824-3266, FAX: (832) 825-3204.

*The first two authors contributed equally to this work.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Keywords

Cytochrome P4501A1; CYP1A2; hyperoxia; promoter; luciferase reporter; *in vivo*; mice; gene regulation; transgenic mice; lung injury

Introduction

Supplemental oxygen is frequently used in the treatment of pulmonary insufficiency in pre-term and term neonates [1,2]. Considerable evidence links oxygen exposure to the development of neonatal diseases such as bronchopulmonary dysplasia (BPD), a major cause of morbidity and mortality in premature infants [1–6]. Hyperoxia causes lung injury in animal models [2,7,8], and reactive oxygen species (ROS) such as superoxide anion, hydroxyl radical, and hydrogen peroxide are most likely to play important roles in mediating lung injury [9–13].

The cytochrome P450 (CYP) system of enzymes is a superfamily of heme-containing proteins that is involved in the metabolism of numerous exogenous and endogenous compounds [14]. The CYP1A family comprises of two proteins, i.e. CYP1A1 and 1A2, which are classically induced by planar aromatic hydrocarbons, such as benzo[*a*]pyrene found in cigarette smoke, or 3-methylcholanthrene [15]. Previous work in rodent models has shown that hyperoxia induces CYP1A enzymes [9, 12, 13, 16–18], but the mechanisms involved are not clearly understood. In the classic model of CYP1A1 induction, the ligand binds to the arylhydrocarbon receptor (AHR), a cytosolic protein, and causes translocation of the AHR-ligand complex into the nucleus. There, it forms a heterodimer with the arylhydrocarbon nuclear translocator (ARNT), and this heterodimer activates a battery of genes that are under the control of the *Ah* gene locus [15, 19, 20]. CYP1A1 induction involves the interaction of the AHR-ligand-ARNT complex with multiple Ah-responsive elements (AHREs) within the *CYP1A1* gene promoter region to activate transcription [15, 20–23]. The mechanisms involved in the hyperoxic induction of CYP1A1 are not fully understood, but an AHR-dependent mechanism has been proposed [16–20,24].

Whereas induction of CYP has been implicated in hyperoxic lung injury [25,26], several groups, including us have shown that CYP1A1 may play a protective role. Pretreatment of rats [27,28] or mice [29] with inducers of CYP1A enzymes attenuates hyperoxic lung injury. Additionally, we showed that pre-treating rats with a CYP1A-specific inhibitor, 1-aminobenzotriazole (ABT), followed by subsequent exposure to 95% O₂ severely potentiates hyperoxic lung injury [12]. These observations support the hypothesis that CYP1A1 and CYP1A2 may play protective roles against oxygen-mediated lung injury.

The mechanisms of induction of CYP1A1 by hyperoxia are not well understood. Therefore, in this study we tested the hypothesis that hyperoxia will induce transcriptional activation of human CYP1A1 or mouse CYP1A2 *in vivo* in transgenic mice carrying the human CYP1A1 or mouse CYP1A2 promoters, and that these mice display altered susceptibilities to hyperoxic lung injury.

Materials and Methods

Chemicals

Sucrose, NADP⁺, ethoxyresorufin, glucose 6-phosphate, and glucose 6-phosphate dehydrogenase were purchased from Sigma-Aldrich (St. Louis, MO). Polyvinylidene difluoride membranes and buffer components for electrophoresis and Western blotting were

obtained from Bio-Rad Laboratories (Hercules, CA). The primary monoclonal antibody to CYP1A1 was a gift from Dr. P.E. Thomas (Rutgers University, Piscataway, NJ).

Animal studies and treatment protocol

Transgenic mice expressing the human 13.2 Kb CYP1A1 promoter to drive luciferase expression (CYP1A1-luc) or mice expressing the 8.4 kb mouse CYP1A2 promoter (CYP1A2-luc) on CD-1 background were obtained from Xenogen Corporation (Alameda, CA) [27,28]. Twelve week-old male wild type (WT) (CD-1), CYP1A1-luc, or CYP1A2-luc mice were maintained in room air or exposed to hyperoxia (> 95% O₂) for 24, 48, or 72 h, and luciferase expression was measured by bioluminescent imaging. After the imaging, the mice were sacrificed, and lung injury was assessed in 3 animals, and endogenous expression of CYP1A1 and 1A2 was determined in 4 different mice from each group.

Bioluminescent imaging

The substrate luciferin was administered into the peritoneal cavity at a dose of 150 mg/kg (20 mg/ml) approximately 5 min prior to imaging. Mice were anesthetized with isoflurane/oxygen and placed on the imaging station. Ventral and dorsal images were collected for 1 second using the IVIS imaging system 100 (Xenogen Corporation) [30], and luciferase expression was determined *in vivo* by bioluminescent imaging (Xenogen Corporation (Alameda, CA) [30–32]. Photons emitted from the liver region were quantified using Series Livingimage software [30–32].

Enzyme assays

Ethoxyresorufin-O-deethylase (EROD) (CYP1A1) and methoxyresorufin O-deethylase (MROD) activities in the liver and lung microsomes were assayed essentially as described previously [9,17,18].

Electrophoresis and Western Blotting

Liver or lung microsomes (~ 20 µg) were subjected to SDS-polyacrylamide gel electrophoresis in 10% acrylamide gel followed by Western blotting to detect CYP1A1 and 1A2 proteins, as reported in the recent articles from our laboratory [9,17,18].

Real-time reverse transcriptase polymerase chain reaction (RT-PCR)

Total RNA (500 ng) from normoxic and hyperoxic mice was subjected to one step real time quantitative TaqMan RT-PCR. Gene-specific primers in the presence of TaqMan reverse transcription reagents and RT reaction mix (Applied Biosystems) were used to reverse transcribe RNA, and TaqMan Universal PCR Master Mix and Assays-on-Demand Gene Expression probes (Applied Biosystems) were used for PCR amplification, as reported earlier [18,31,32]. The relative mRNA levels for P4501A1 were normalized to their 18S content, and relative expression levels of CYP1A1 and 1A2 genes were calculated as reported earlier [18,31,32].

Lung injury

Lung injury was assessed by histology, as described previously [17,28].

Statistical analyses

All statistical analyses were conducted using two-way ANOVA, followed by modified t-tests.

Results

Effect of hyperoxia on CYP1A1/1A2 promoter activation (luciferase reporter expression)

The major goal of this investigation was to determine the mechanisms of induction of CYP1A1 and 1A2 by hyperoxia *in vivo*. As shown in figure 1, hyperoxia exposure for 24–48 h resulted in marked induction of CYP1A1 (Figure 1A,B)- and CYP1A2 (Figure 1C,D)-promoter driven luciferase expression *in vivo* in the transgenic mice, compared to those maintained in room air. Luciferase expression declined after 72 h of hyperoxia (Figure 1A–D). Quantitative imaging analyses showed that hyperoxia for 24 and 48 h induced CYP1A1-driven luciferase activities by 2.5- and 1.25-fold, respectively over the corresponding room air controls (Figure 1B). By 72 h, the luciferase expression was significantly lower than room air controls. In CYP1A2-luc mice, hyperoxia induced luciferase activities by 1.3- and 0.76-fold at the 24 and 48 h time points, respectively, over the corresponding room controls (Figure 1D). Similar to the responses noted in CYP1A1-luc mice, hyperoxia for 72 h significantly attenuated CYP1A2-driven luciferase activities, compared to room air controls (Figure 1D).

Effect of MC on endogenous CYP1A1/1A2 enzyme activities and apoprotein contents

In order to determine if CYP1A1-luc or CYP1A2-luc mice will display altered endogenous expression of CYP1A1 and/or 1A2, we determined the activities of EROD and MROD, reflecting catalytic functionalities of CYP1A1 and 1A2, respectively (Figure 2). We also determined the CYP1A1 and 1A2 apoprotein contents. As shown in figure 2A, hyperoxia elicited a 45% induction of hepatic EROD activities in the WT (CD-1) mice at the 24 h time point. EROD activities declined by 48 h, and were significantly lower than room air controls after 72 h of hyperoxia (Figure 2A). In CYP1A1-luc mice, room air controls displayed significantly higher expression of EROD than WT mice (Figure 2A). Hyperoxia exposure resulted in suppression of EROD activities by 45, 45, and 60% after for 24, 48, or 72 h, respectively (Figure 2A). CYP1A2-luc mice also showed similar trends, with hyperoxia diminishing the EROD activities at 24, 48, and 72 h, albeit to a lesser extent than that observed in the CYP1A1-luc mice (Figure 2A).

Hepatic MROD, which reflects CYP1A2 activity, was also induced by hyperoxia by about 36% at 24 h. The activity declined by 48 h, and was significantly lesser than room air controls by 72 h (Figure 2B). MROD activities were higher in the air-breathing CYP1A1-luc and CYP1A2-luc mice, compared to the WT controls (Figure 2B). Hyperoxia exposure suppressed the MROD activities in the CYP1A1-luc, as well as CYP1A2-luc mice (Figure 2B). Western blot analyses showed expression patterns of CYP1A1 and 1A2 (Figure 2C) that correlated with enzyme activity data.

In pulmonary tissues, hyperoxia elicited a 0.6- and 0.4-fold induction of EROD (CYP1A1) activity at the 24 h and 48 time points, respectively over room air controls in the WT mice (Figure 2D), but the activities were suppressed (40%) after 72 h of hyperoxia (Figure 2D). In CYP1A1-luc mice, hyperoxia for 24 h induced EROD activities by 28%, but after 48–72 h, the activities were significantly suppressed (43–46%) compared to room air controls (Figure 2D). CYP1A2-luc mice showed the highest EROD activities in room air (Figure 2D), but hyperoxia for 48 and 72 h diminished CYP1A1 activities by 68 and 88%, respectively, compared to room air controls (Figure 2D). Western blot analyses showed that the CYP1A1 apoprotein contents correlated with enzyme activities (Figure 2E).

Effect of hyperoxia on CYP1A1 and 1A2 mRNA expression

In WT mice, hyperoxia caused a 3.5- and 2-fold induction of hepatic CYP1A1 mRNA levels at the 1 and 8 days, respectively, versus controls (Figure 3A). Induction declined by 72 h. In

the CYP1A1-luc mice, the extent of induction of endogenous CYP1A1 mRNA after 24 h of hyperoxia was lesser than that observed in the WT mice, and by 48 h, the CYP1A1 levels declined to control (Figure 3A). In the CYP1A2-luc mice, the degree of induction by hyperoxia was modest, being 1-, 0.6-fold, respectively at the 24 and 48 h time points, respectively (Figure 3A).

In contrast to CYP1A1, where hyperoxia for 24 h differentially induced this gene in WT and transgenic mice, hepatic CYP1A2 mRNA expression was augmented in the hyperoxic WT, CYP1A1-luc, and CYP1A2 mice to similar levels (Figure 3B). In each of the genotypes, i.e. WT, CYP1A1-luc, and CYP1A2-luc mice, hyperoxia for 72 h caused a marked decline in CYP1A2 mRNA expression, compared to the corresponding air-breathing controls (Figure 3B).

Similar to our observations in liver, hyperoxia caused a much greater induction in the levels of pulmonary CYP1A1 mRNA in the WT mice than the CYP1A1-luc or CYP1A2-luc mice (Figure 3C). In WT mice, hyperoxia-exposed samples showed an induction of about 2-fold compared to controls at the 24 h time point. The induction declined by 48 h of hyperoxia, and was even more suppressed by 72 h (Figure 3C). In CYP1A1-luc mice, hyperoxia caused a 1-fold increase at the 24 h time point over room air controls. Induction declined after 48 and 72 h time points of hyperoxia (Figure 3C). In CYP1A2-luc mice, hyperoxia elicited induction of CYP1A1 mRNA expression at 24 h and significantly suppressed it by 48 h (Figure 3C).

Effect of hyperoxia on lung injury

Hyperoxia for 24 or 48 h did not increase pulmonary edema in the WT mice as shown by lung weight/body weight ratios, or by histology (not shown). After 72 h of hyperoxia WT mice showed increased cellularity, infiltration of inflammatory cells, and intra-alveolar edema and (Figure 4). CYP1A1-luc mice were much more susceptible to hyperoxic lung injury than similarly exposed WT mice (Figure 4). CYP1A1-luc showed injury even at 48 h (not shown), and by 72 h these mice displayed perivascular inflammation, intra-alveolar edema, and widening of the adventitia of the pulmonary artery (Figure 4). CYP1A2-luc mice showed the greatest lung injury after 72 h, with widespread intra-alveolar edema and proteinaceous material filling the alveoli (Figure 4).

Discussion

In this study, we tested the hypothesis that hyperoxia induces human CYP1A1 and 1A2 enzymes by transcriptional activation of the corresponding promoters *in vivo*, and transgenic mice expressing the human CYP1A1 or 1A2 promoters would be more susceptible to hyperoxic lung injury than WT mice.

Our results showing marked induction of luciferase expression in transgenic mice expressing CYP1A1-luc or 1A2-luc (Figure 1A–1D), supporting the hypothesis that hyperoxia elicited CYP1A induction by transcriptional activation of the corresponding promoters. The observation that hyperoxia elicited greater degree induction of luciferase expression in the CYP1A1-luc mice compared to that of CYP1A2-luc mice was similar to our previous findings on the induction of CYP1A1 and 1A2 by 3-methylcholanthrene [30,31]. In fact, we show for the first time that hyperoxia induces CYP1A genes by transcriptional activation of the CYP1A promoters *in vivo*. Thus, these transgenic models can be very useful in determining the mechanisms of induction of CYP1A *in vivo*.

The finding that hyperoxia caused marked increases of hepatic EROD (Figure 2A) and MROD activities (Figure 2B) in the WT mice indicated induction of CYP1A1 and 1A2

expression, as EROD and MROD activities are known to reflect CYP1A1 and 1A2 activities, respectively [9,17,18]. These results were consistent with our previous findings showing induction of CYP1A1 enzymes in rats and mice by hyperoxia *in vivo* [9, 17, 18].

The fact that hyperoxia induced the endogenous CYP1A1 and 1A2 activities in the transgenic mice suggests that introduction of the promoters into the mice did not affect the phenomenon of induction, and that the molecular properties governing gene expression of the CYP1A-luc mice and the endogenous CYP1A genes are similar [31–33].

That hyperoxia also caused induction of CYP1A1 and 1A2 apoproteins in liver (Figure 2A–B), and CYP1A1 alone in the lungs of WT and transgenic mice was in agreement with our enzyme activity data. The results in Figure 3A showing a robust induction of hepatic CYP1A1 mRNA in the WT mice was consistent with the hypothesis that hyperoxia induced CYP1A1 activity and protein expression by inducing its message [18,24]. The fact that hyperoxia caused much lesser induction of CYP1A1 mRNA in the livers of CYP1A1-luc and CYP1A2-luc mice was probably due to decreased accessibility of the ligand-AHR-ARNT complex and other co-activators to the endogenous CYP1A1 enhancer elements, presumably through alteration in chromatin remodeling as a result of incorporation of the CYP1A1 promoter transgene.

The fact that hyperoxia for 24 h elicited induction of hepatic CYP1A2 expression in the CYP1A1-luc similar to that in WT mice (Figure 3B) supported the hypothesis that introduction of human CYP1A1 promoter into the genome of mice did not alter the susceptibility of CYP1A2 gene to hyperoxic induction. On the other hand the diminution of CYP1A2 mRNA induction in the CYP1A2-luc mice may have been due to decreased accessibility of the ligand-AHR-ARNT complex to CYP1A2 enhancer(s) that are responsible for induction of the CYP1A2 mRNA [31]. These results also suggest that mechanisms governing transcriptional activation of exogenous CYP1A promoters by hyperoxia are similar to those of the endogenous counterparts.

The increased susceptibility of CYP1A1-luc and 1A2-luc to hyperoxic lung injury was probably due to decreased endogenous expression of CYP1A enzymes in these mice by prolonged hyperoxia. Our previous results showed that CYP1A enzymes play beneficial roles against hyperoxic lung injury by detoxifying ROS such as F₂-isoprostanes [35]. F₂-isoprostanes have been shown to be elevated in rats exposed to hyperoxia [36], and these products are formed by reactive oxygenase species mediated peroxidation of lipid hydroperoxides, resulting in protein and DNA oxidation, eventually leading to lung injury and cell death [10]. In fact, levels of F₂-isoprostanes are elevated in premature infants suffering from BPD [36], suggesting that these compounds may have an etiologic role in lung injury.

Thus, further studies on the molecular regulation of CYP1A genes by hyperoxia could lead to novel strategies for the prevention and/or treatment of BPD in premature infants and ARDS in adults.

Acknowledgments

We thank Dr. Paul Thomas (Rutgers University) for providing us monoclonal antibodies against CYP1A1/1A2.

Abbreviations

BPD bronchopulmonary dysplasia (BPD)

CYP	cytochrome P450
AHR	Ah receptor
EROD	ethoxyresorufin O-demethylase
MROD	methoxyresorufin O-demethylase
luc	luciferase
ANOVA	analyses of variance
AHREs	Ah response elements
ARNT	Ah receptor nuclear translocator
RT-PCR	reverse transcriptase-Polymerase chain reaction
WT	wild type

References

1. Northway WH Jr, Rosan RC. Radiographic features of pulmonary oxygen toxicity in the newborn: Bronchopulmonary dysplasia. *Radiology*. 1968; 91:49–58. [PubMed: 4871379]
2. Fisher AB. Oxygen therapy. Side effects and toxicity. *Am Rev Respir Dis*. 1980; 122:61–69. [PubMed: 7458051]
3. Bland, RD.; Coalson, JJ. Chronic lung disease in early infancy. M. Dekker; New York: 2000.
4. Baraldi E, Filippone M. Chronic lung disease after premature birth. *N Engl J Med*. 2007; 357:1946–1955. [PubMed: 17989387]
5. Fanaroff AA, Stoll BJ, Wright LL, Carlo WA, Ehrenkranz RA, Stark AR, Bauer CR, Donovan EF, Korones SB, Laptook AR, Lemons JA, Oh W, Papile LA, Shankaran S, Stevenson DK, Tyson JE, Poole WK. Trends in neonatal morbidity and mortality for very low birthweight infants. *Am J Obstet Gynecol*. 2007; 196:147.e141–148. [PubMed: 17306659]
6. Short EJ, Klein NK, Lewis BA, Fulton S, Eisengart S, Kercsmar C, Baley J, Singer LT. Cognitive and academic consequences of bronchopulmonary dysplasia and very low birth weight: 8-year-old outcomes. *Pediatrics*. 2003; 112:e359. [PubMed: 14595077]
7. Freeman BA, Crapo JD. Hyperoxia increases oxygen radical production in rat lungs and lung mitochondria. *J Biol Chem*. 1981; 256:10986–10992. [PubMed: 7287745]
8. Budinger GR, Mutlu GM, Urich D, Soberanes S, Buccellato LJ, Hawkins K, Chiarella SE, Radigan KA, Eisenbart J, Agrawal H, Berkelhamer S, Hekimi S, Zhang J, Perlman H, Schumacker PT, Jain M, Chandel NS. Epithelial Cell Death is an Important Contributor to Oxidant-Mediated Acute Lung Injury. *Am J Respir Crit Care Med*. 2010
9. Moorthy B, Nguyen UT, Gupta S, Stewart KD, Welty SE, Smith CV. Induction and decline of hepatic cytochromes P4501A1 and 1A2 in rats exposed to hyperoxia are not paralleled by changes in glutathione S-transferase-alpha. *Toxicol Lett*. 1997; 90:67–75. [PubMed: 9020404]
10. Morel Y, Barouki R. Down-regulation of cytochrome P450 1A1 gene promoter by oxidative stress. Critical contribution of nuclear factor 1. *J Biol Chem*. 1998; 273:26969–26976. [PubMed: 9756946]
11. Yang F, Coalson JJ, Bobb HH, Carter JD, Banu J, Ghio AJ. Resistance of hypotransferrinemic mice to hyperoxia-induced lung injury. *Am J Physiol*. 1999; 277:L1214–1223. [PubMed: 10600893]
12. Moorthy B, Parker KM, Smith CV, Bend JR, Welty SE. Potentiation of oxygen-induced lung injury in rats by the mechanism-based cytochrome P-450 inhibitor, 1-aminobenzotriazole. *J Pharmacol Exp Ther*. 2000; 292:553–560. [PubMed: 10640292]
13. Moorthy, B. The CYP1A subfamily. In: Ioannides, E., editor. *Cytochromes P450. Role in Drug Metabolism and Toxicity of Drugs and other Xenobiotics*. RSC Publishing; Cambridge, UK: 2008. p. 97-135.

14. Guengerich FP. Cytochrome P450: what have we learned and what are the future issues? *Drug Metab Rev.* 2004; 36:159–197. [PubMed: 15237850]
15. Whitlock JP Jr, Okino ST, Dong L, Ko HP, Clarke-Katzenberg R, Ma Q, Li H. Cytochromes P450 5: induction of cytochrome P4501A1: a model for analyzing mammalian gene transcription. *Faseb J.* 1996; 10:809–818. [PubMed: 8666157]
16. Okamoto T, Mitsuhashi M, Fujita I, Sindhu RK, Kikkawa Y. Induction of cytochrome P450 1A1 and 1A2 by hyperoxia. *Biochem Biophys Res Commun.* 1993; 197:878–885. [PubMed: 8267628]
17. Couroucli XI, Welty SE, Geske RS, Moorthy B. Regulation of pulmonary and hepatic cytochrome P4501A expression in the rat by hyperoxia: implications for hyperoxic lung injury. *Mol Pharmacol.* 2002; 61:507–515. [PubMed: 11854430]
18. Jiang W, Welty SE, Couroucli XI, Barrios R, Kondraganti SR, Muthiah K, Yu L, Avery SE, Moorthy B. Disruption of the Ah receptor gene alters the susceptibility of mice to oxygen-mediated regulation of pulmonary and hepatic cytochromes P4501A expression and exacerbates hyperoxic lung injury. *J Pharmacol Exp Ther.* 2004; 310:512–519. [PubMed: 15123765]
19. Nebert DW, Roe AL, Dieter MZ, Solis WA, Yang Y, Dalton TP. Role of the aromatic hydrocarbon receptor and [Ah] gene battery in the oxidative stress response, cell cycle control, and apoptosis. *Biochem Pharmacol.* 2000; 59:65–85. [PubMed: 10605936]
20. Whitlock JP Jr. Induction of cytochrome P4501A1. *Annu Rev Pharmacol Toxicol.* 1999; 39:103–125. [PubMed: 10331078]
21. Nukaya M, Moran S, Bradfield CA. The role of the dioxin-responsive element cluster between the Cyp1a1 and Cyp1a2 loci in aryl hydrocarbon receptor biology. *Proc Natl Acad Sci U S A.* 2009; 106:4923–4928. [PubMed: 19261855]
22. Chiaro CR, Patel RD, Marcus CB, Perdew GH. Evidence for an aryl hydrocarbon receptor-mediated cytochrome p450 autoregulatory pathway. *Mol Pharmacol.* 2007; 72:1369–1379. [PubMed: 17720764]
23. Sato W, Suzuki H, Sasaki T, Kumagai T, Sakaguchi S, Mizugaki M, Miyairi S, Yamazoe Y, Nagata K. Construction of a system that simultaneously evaluates CYP1A1 and CYP1A2 induction in a stable human-derived cell line using a dual reporter plasmid. *Drug Metab Pharmacokinet.* 2010; 25:180–189. [PubMed: 20460824]
24. Bhakta KY, Jiang W, Couroucli XI, Fazili IS, Muthiah K, Moorthy B. Regulation of cytochrome P4501A1 expression by hyperoxia in human lung cell lines: Implications for hyperoxic lung injury. *Toxicol Appl Pharmacol.* 2008; 233:169–178. [PubMed: 18824009]
25. Hazinski TA, France M, Kennedy KA, Hansen TN. Cimetidine reduces hyperoxic lung injury in lambs. *J Appl Physiol.* 1989; 67:2586–2592. [PubMed: 2606866]
26. Hazinski TA, Noisin E, Hamon I, DeMatteo A. Sheep lung cytochrome P4501A1 (CYP1A1): cDNA cloning and transcriptional regulation by oxygen tension. *J Clin Invest.* 1995; 96:2083–2089. [PubMed: 7560103]
27. Mansour H, Brun-Pascaud M, Marquetty C, Gougerot-Pocidal MA, Hakim J, Pocidal JJ. Protection of rat from oxygen toxicity by inducers of cytochrome P-450 system. *Am Rev Respir Dis.* 1988a; 137:688–694. [PubMed: 2830813]
28. Sinha A, Muthiah K, Jiang W, Couroucli X, Barrios R, Moorthy B. Attenuation of hyperoxic lung injury by the CYP1A inducer beta-naphthoflavone. *Toxicol Sci.* 2005; 87:204–212. [PubMed: 15958656]
29. Mansour H, Levacher M, Azoulay-Dupuis E, Moreau J, Marquetty C, Gougerot-Pocidal MA. Genetic differences in response to pulmonary cytochrome P-450 inducers and oxygen toxicity. *J Appl Physiol.* 1988b; 64:1376–1381. [PubMed: 3378972]
30. Zhang W, Moorthy B, Chen M, Muthiah K, Coffee R, Purchio AF, West DB. A Cyp1a2-luciferase transgenic CD-1 mouse model: responses to aryl hydrocarbons similar to the humanized AhR mice. *Toxicol Sci.* 2004; 82:297–307. [PubMed: 15329440]
31. Jiang W, Wang L, Zhang W, Coffee R, Fazili IS, Moorthy B. Persistent induction of cytochrome P450 (CYP)1A enzymes by 3-methylcholanthrene in vivo in mice is mediated by sustained transcriptional activation of the corresponding promoters. *Biochem Biophys Res Commun.* 2009; 390:1419–1424. [PubMed: 19900403]

32. Jiang W, Wang L, Kondraganti SR, Fazili IS, Couroucli XI, Felix EA, Moorthy B. Disruption of the gene for CYP1A2, which is expressed primarily in liver, leads to differential regulation of hepatic and pulmonary mouse CYP1A1 expression and augmented human CYP1A1 transcriptional activation in response to 3-methylcholanthrene in vivo. *J Pharmacol Exp Ther*. 2010; 335:369–379. [PubMed: 20732958]
33. Operana TN, Nguyen N, Chen S, Beaton D, Tukey RH. Human CYP1A1^{GFP} expression in transgenic mice serves as a biomarker for environmental toxicant exposure. *Toxicol Sci*. 2007; 95:98–107. [PubMed: 17065433]
34. Fessel JP, Porter NA, Moore KP, Sheller JR, Roberts LJ 2nd. Discovery of lipid peroxidation products formed in vivo with a substituted tetrahydrofuran ring (isofurans) that are favored by increased oxygen tension. *Proc Natl Acad Sci U S A*. 2002; 99:16713–16718. [PubMed: 12482927]
35. Moorthy B, Wang L, Couroucli XI, Khan FM, Jiang W. Mice lacking the genes for cytochrome P450 (CYP)1A1 or 1A2 display increased levels of lipid peroxidation products and enhanced susceptibility to hyperoxic lung injury in vivo. *Toxicologist*. 2008; 97:772.
36. Cotton RB, Hazinski TA, Morrow JD, Roberts LJ, Zeldin DC, Lindstrom DP, Lappalainen U, Law AB, Steele S. Cimetidine does not prevent lung injury in newborn premature infants. *Pediatr Res*. 2006; 59:795–800. [PubMed: 16641218]

Figure 1.

Bioluminescent imaging of CYP1A1-luc (A, B) or CYP1A2-luc (C,D) mice following hyperoxia exposures. CYP1A1-luc (A) or 1A2-luc mice (C) were maintained in room air or exposed to hyperoxia (> 95% O₂) for 24–72 h, and luciferase expression was analyzed by bioluminescent imaging in real time at the indicated time points. Quantitation of bioluminescent imaging data of CYP1A1-luc (B) or 1A2-luc (D) mice was conducted using IVIS imaging software. Values represent mean ± S.E. (n = 5). *, Statistically significant differences between room air and hyperoxic mice at P < 0.05, as determined by two-way ANOVA.

Figure 2.

Effect of hyperoxia on endogenous hepatic EROD (CYP1A1) (A), MROD (B), activities and apoprotein contents (C), and pulmonary EROD (D) and apoprotein expression (E). Adult male WT (CD-1), CYP1A1-luc or CYP1A2-luc mice were maintained in room air or exposed to hyperoxia at the indicated time points, and hepatic EROD (A), MROD (B) activities were determined in the liver microsomes. Data represent mean $\pm$ SE of fold induction versus room air controls from at least 4 individual animals. C, Representative Western blot showing effect of hyperoxia on endogenous hepatic CYP1A1 and 1A2 (C) expression in mice. PC, positive control showing induction of CYP1A1/1A2 by 3-methylcholanthrene. D and E represent pulmonary EROD and apoprotein expression, respectively, in microsomes of WT, CYP1A1-luc, and CYP1A2-luc mice.

Figure 3.

Effect of hyperoxia on endogenous hepatic CYP1A1 (A), CYP1A2 (B), and pulmonary mRNA (C) in mice. Adult male WT (CD-1), CYP1A1-luc or CYP1A2-luc mice were maintained in room air or exposed to hyperoxia for 24–72 h, and hepatic CYP1A1 (A), CYP1A2 (B), and pulmonary CYP1A1 (C) mRNA levels were determined by real time RT-PCR at the indicated time points. Data represents mean $\pm$ S.E. of fold induction versus controls from at least four individual animals. *, Statistically significant differences between room air and hyperoxic mice at $P < 0.05$, as determined by two-way ANOVA.

Figure 4.

Effect of hyperoxia on lung injury in WT, CYP1A1-Luc, and CYP1A2-mice. WT, CYP1A1-luc, and CYP1A2-luc mice were exposed to hyperoxia as described under Materials and Methods, and lung tissues were processed for histological analyses. The figure shows representative light micrographs of lung sections of mice maintained in room air or exposed to hyperoxia for 72 h. Air-breathing mice (upper panels) showed normal lung architecture. Hyperoxia for 72 h showed increased lung injury in WT mice. Single arrow shows infiltration of inflammatory cells, and double arrow shows areas of intra-alveolar edema. CYP1A1-luc mice showed even more injury after 72 h of hyperoxia, with lungs showing perivascular and intra-alveolar edema and marked widening of the adventitia of the pulmonary artery. CYP1A2-luc mice showed the greatest susceptibility to lung injury, as evidenced by intra-alveolar edema and proteinaceous materials filling the alveoli.