Published in final edited form as:

Pharmacol Biochem Behav. 2011 September ; 99(3): 408–413. doi:10.1016/j.pbb.2011.05.021.

Chronic cocaine self-administration attenuates the anxiogenic-like and stress potentiating effects of the benzodiazepine inverse agonist, FG 7142

R. Parrish Waters and Ronald E. See

Department of Neurosciences, Medical University of South Carolina, Charleston SC 29425, USA

Abstract

Stress is a well-known risk factor in relapse to drug abuse. Several forms of stress in animals have been used with varied degrees of success to elicit reinstatement of drug-seeking after chronic drug self-administration. Here, we tested the ability of the benzodiazepine (BZ) inverse agonist, FG 7142, to elicit anxiety-like behavior and potentiate stress responses in rats as measured by standard behavioral and hormonal indices and for its ability to affect reinstatement of cocaine-seeking in rats with a prior history of cocaine self-administration. FG 7142 elicited anxiety-like behavior on the elevated plus maze (EPM) in cocaine-naïve rats, and cocaine-naïve rats injected with FG 7142 exhibited increased plasma corticosterone levels following EPM exposure. However, in animals with a history of cocaine self-administration, FG 7142 failed to affect elevated plus maze performance and did not affect plasma corticosterone response to the EPM. Furthermore, FG 7142 failed to reinstate cocaine-seeking, nor did it alter conditioned cue-induced reinstatement. These data indicate that the anxiety-related and stress potentiating qualities of BZ inverse agonism are attenuated in cocaine-experienced animals and do not lead to reinstatement of cocaine-seeking.

Keywords

benzodiazepine; cocaine; FG 7142; reinstatement; relapse; self-administration

1. Introduction

Relapse to cocaine-seeking or cocaine-taking is a foremost obstacle in the treatment of cocaine addiction, and a major instigator of relapse in humans and animal models is exposure to a stressor (Shaham et al., 2000). Abstinent human cocaine addicts often report an increase in craving for cocaine following stressful stimuli (Sinha et al., 2006), and cocaine-seeking behavior can be elicited in animals extinguished from cocaine self-administration using anxiogenic-like stressors, such as intermittent footshock stress (Erb et al., 1996). Notably, the ability of stress to induce reinstatement does not generalize to all stressors, in that predator odor, restraint stress, and stress-predictive cues (e.g., a tone previously associated with footshock) all fail to elicit the reinstatement of drug-seeking (Shaham et al., 2000).

© 2011 Elsevier Inc. All rights reserved.

Corresponding author: Dr. Ronald E. See, Department of Neurosciences, Medical University of South Carolina, 173 Ashley Avenue, BSB 416B, Charleston SC 29425, 843-792-2487, seere@musc.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Although using footshock to reinstate cocaine-seeking is useful in understanding the clinical phenomenon of stressor induced relapse (Epstein et al., 2006), engaging stress-related systems by other means may offer some advantages in developing models of relapse. Of particular note are pharmacological agents that act on central neural mechanisms that mediate anxiety-like behavior and other stress-related responses and also elicit the reinstatement of extinguished drug-seeking (See and Waters, 2011). For example, central administration of corticotropin releasing factor (CRF), an anxiogenic peptide closely linked to mounting a stress response (Sutton et al., 1982), can reinstate cocaine-seeking by stimulating CRF receptors in multiple brain areas (Erb et al., 2001; Erb and Stewart, 1999; Shalev et al., 2010). Furthermore, the anxiogenic noradrenergic α -2 receptor antagonist, yohimbine, reinstates drug-seeking for methamphetamine (Shepard et al., 2004), cocaine (Feltenstein and See, 2006), alcohol (Le et al., 2005), and heroin (Banna et al., 2010). The reinstating properties of yohimbine are presumed to rely on presynaptic α -2 antagonism and subsequent increased norepinephrine release, which is associated with a stress response (Korf et al., 1973). However, some evidence suggests that yohimbine-induced reinstatement may involve additional mechanisms, including CRF release (Brown et al., 2009).

Expanding the repertoire of neuroactive pharmacological agents that can elicit the reinstatement of drug-seeking may elucidate additional mechanisms involved in relapse. Agonist action at the benzodiazepine (BZ) binding site of the GABA_A receptor complex has been well characterized to decrease the magnitude of stress responses and the anxiety associated with the application of stress (Marin et al., 1997). While not a primary means of treatment in addiction, BZ agonists also provide some relief from the anxiety associated with cocaine withdrawal (Paine et al., 2002), a contributing cause of relapse in human cocaine addicts (Goeders, 2002). Furthermore, BZ agonists also have a demonstrated efficacy in reducing conditioned cue-induced reinstatement of cocaine-seeking (Goeders et al., 2009). The interactions of BZ agonists with stress responses and the reinstatement of cocaine-seeking demonstrate a potential role for BZ receptors in stress-related reinstatement of drug-seeking, and present a target to investigate the interactions between a pharmacologically-induced stress response and the reinstatement of drug-seeking.

The BZ inverse agonist, FG 7142, is an established anxiogenic compound that elicits panic attacks in humans (Dorow et al., 1983) and increases anxiety-like behavior in rodents (File et al., 1985; Pellow et al., 1987). Furthermore, FG 7142 increases stress associated physiological markers, such as plasma corticosterone levels (Pellow and File, 1985) and central noradrenergic activity (Ida et al., 1991; Mason et al., 1998). Here, we assessed the anxiogenic-like and stress potentiating properties of FG 7142 in cocaine-naïve and cocaine-experienced animals using the elevated plus maze and measurement of plasma corticosterone. We also determined the ability of FG 7142 to reinstate extinguished cocaine-seeking, as well as to alter the reinstatement of cocaine-seeking in the presence of conditioned cues after repeated FG 7142 administration.

2. Methods

2.1. Subjects

Male, Sprague-Dawley rats (initial weight 275–350 g; Charles River, Wilmington, MA, USA) were individually housed in a temperature- and humidity- controlled vivarium on a 12:12 reverse light-dark cycle (lights off 6 AM – 6 PM). All experimental procedures occurred during the dark cycle. Animals received water and standard rat chow (Harlan, Indianapolis, IN, USA) *ad libitum* for the duration of each experiment, except for the first 4 days of cocaine self-administration, during which animals were maintained on 20 g of standard rat chow per day to facilitate the acquisition of lever responding (Bongiovanni and See, 2008). Housing and care of the rats were carried out in accordance with the “Guide for

the Care and Use of Laboratory Rats” (Institute of Laboratory Animal Resources on Life Sciences, National Research Council).

2.2. Surgery

Rats were anesthetized using a mixture of ketamine hydrochloride and xylazine (66 and 1.3 mg/kg, respectively, intramuscular [IM]), followed by equithesin (0.5 ml/kg with a solution of 8.0 mg/ml pentobarbital sodium, 34.0 mg/ml chloral hydrate, and 42.6 mg/ml magnesium sulfate heptahydrate dissolved in a 44% propylene glycol, 10% ethanol solution, intraperitoneal [IP]). Ketorolac (Sigma, St. Louis, MO, USA) was administered as a preoperative analgesic (2 mg/kg, IP). Chronic indwelling catheters were implanted into the right jugular vein using previously described methods (Feltenstein et al., 2007). Catheter patency was maintained by flushing with 0.1 ml of 70 U/ml heparinized saline immediately prior to self-administration sessions, and a 0.1 ml antibiotic solution of cefazolin (100 mg/ml; dissolved in saline) and 0.1 ml of 100 U/ml heparinized saline regimen following self-administration sessions. Polyethylene caps were used to cover the tips of catheters when the rats were not connected to infusion pumps. To verify catheter patency, rats occasionally received a 0.1 ml infusion of methohexital sodium (10 mg/ml; dissolved in 0.9% physiological saline), a short acting barbiturate that produces a rapid loss of muscle tone when administered intravenously.

2.3. Drugs

Cocaine hydrochloride (National Institute on Drug Abuse, Research Park Triangle, NC, USA) was dissolved in 0.9% physiological saline at 4 mg/ml. The beta-carboline, FG 7142 (N-methyl- β -carboline-3-carboxamide; Sigma, Sydney, Australia), was suspended at concentrations of 1, 5, 10, 20, and 30 mg/ml in saline (0.9% w/vol) using 3 drops of Tween 80 per 1 ml saline. This suspension, or the vehicle (saline plus Tween 80) was administered at a volume of 1 ml/kg IP immediately prior to testing. Yohimbine hydrochloride was placed into solution (1.25 mg/ml; Sigma-Aldrich, St. Louis, MO, USA) using sterile distilled water and administered IP 30 min prior to testing.

2.4. Cocaine Self-administration

Rats lever pressed for cocaine in self-administration chambers (30×20×20 cm) linked to a computerized data collection program (MED-PC, Med Associates, Inc., St. Albans, VT, USA). Each chamber was equipped with two retractable levers, with white stimulus lights above each lever, a tone generator, and a house light on the wall opposite the levers. This chamber was contained within a sound-attenuating cubicle equipped with a ventilation fan. Rats self-administered cocaine during daily 2-h sessions according to an FR-1 schedule of reinforcement. At the start of each session, the animal's catheter was connected to a liquid swivel (Instech, Plymouth Meeting, PA, USA) via polyethylene 20 tubing that was encased in steel spring leashes (Plastics One Inc., Roanoke, VA, USA). The house light signaled the initiation of the session and remained illuminated throughout the entire session. Lever presses on the active lever resulted in a 2-s activation of the infusion pump (0.2 mg/50 μ l infusion; approximately 0.6 mg/kg/infusion) and a 5-s presentation of a stimulus complex, consisting of activation of the white stimulus light above the active lever and activation of the tone generator (78 dB, 4.5 kHz). After each infusion, responses on the active lever were recorded, but had no programmed consequences during a 20-s time-out period. During the sessions, responses on the inactive lever were also recorded, but had no programmed consequences. Daily cocaine self-administration continued until each rat had obtained self-administration criteria of 10 sessions with at least 10 infusions per session.

2.5. Elevated Plus Maze (EPM)

Elevated plus maze (EPM) testing occurred in a room dimly lit with red light (4 lumen LED bulbs $\times$ 3) during the animals' dark phase. Animals with a history of cocaine self-administration as described above were tested on the first day of abstinence. A group of cocaine-naïve animals were also tested for comparison. Cocaine-naïve animals were catheterized to allow blood sampling, and transported/handled daily to match the handling history of cocaine-experienced animals. FG 7142 or vehicle was administered via IP injection prior to placement on the EPM. We used a custom built EPM consisting of a platform constructed with opaque black plastic. The maze contained two open arms (50.80×11.11 cm) with clear plastic ledges (0.64 cm tall) on both sides and the end, and two closed arms (50.80×11.11 cm) with opaque black walls (40.64 cm tall) on both sides and the end. Animals were allowed 10 min to explore the apparatus, and the first 5 min of each trial was scored. Behavior was recorded using Ethovision 7.0 software (Noldus, Asheville, NC). We analyzed time spent in the open arms as an inverse measure of anxiety (Pellow et al., 1985). Total locomotion was estimated by counting closed arm entries to assess differences in general activity (Hogg, 1996).

2.6. Corticosterone Radioimmunoassay

Blood samples (100 μ l) were obtained from jugular catheters immediately prior to injection of FG 7142 or vehicle, and again 10 min later (following EPM exposure); maximal corticosterone responses to environmental stressors (e.g., restraint stress) are typically observed in this time frame (Sakakura et al., 1976). Whole blood samples were centrifuged for 20 min at 10,000 G and 4°C. Plasma was stored at -80°C until analysis for corticosterone (ng/ml) using a commercially available radioimmunoassay kit (MP Biomedical, Solon, OH) and a gamma counter.

2.7. Extinction and Reinstatement

Following chronic cocaine self-administration, subjects underwent a series of daily 2-h extinction sessions, whereby responses on either the active or inactive lever were recorded, but resulted in no programmed consequences (i.e., no infusions and no conditioned stimulus presentations). Animals continued under extinction conditions for at least seven days, and until they reached the criteria of 2 consecutive days with <15 active lever presses per session. Reinstatement trials began once an animal met the extinction criteria. Animals were run in three separate cohorts: FG 7142 low dose response testing (0, 1, 5, and 10 mg/kg), FG 7142 high dose response testing (0, 10, 20, and 30 mg/kg), and repeated vehicle and FG 7142 (10 mg/kg) followed by cue-induced reinstatement tests.

FG 7142 and Reinstatement—This experiment examined the dose response effect of FG 7142 on the reinstatement of cocaine-seeking. Following extinction, rats underwent five reinstatement tests, with intervening extinction sessions (criteria of 2 consecutive days of <15 active lever presses/2-h session). Reinstatement tests consisted of a 2-h session in the self-administration chamber, during which lever presses were recorded, but resulted in no programmed consequences. Immediately prior to entering the chambers, the first group of rats received one of four doses of FG 7142 (0, 1, 5, 10 mg/kg) in a counterbalanced fashion. Following these trials, animals underwent reinstatement testing with yohimbine (1.25 mg/kg), injected 30 min prior to testing. Another group of rats received one of four doses of FG 7142 (0, 10, 20, 30 mg/kg) in a counterbalanced fashion. Our laboratory has previously utilized multiple counterbalanced reinstatement trials of this type (Feltenstein et al., 2009; Feltenstein and See, 2006; Reichel and See, 2010).

FG 7142 and Cue-induced Reinstatement—This experiment assessed the effect of repeated pretreatment with FG 7142 on conditioned cue-induced reinstatement. Once extinction criteria were met, animals first received a vehicle injection 2 h after each extinction session for three days. The following day, animals underwent a cue-induced reinstatement test, during which active lever presses resulted in the presentation of drug-associated cues (light and tone), but no cocaine infusions. The subjects then received an FG 7142 (10 mg/kg) injection 2 h after each extinction session for three days, followed by a second cue-induced reinstatement test.

2.8. Statistical analysis

Planned comparisons (two-tailed t-tests) were used to compare EPM performance in animals treated with vehicle or FG 7142. Pre- and post-manipulation plasma corticosterone levels were compared using repeated measures analysis of variance (ANOVA) and subsequent post hoc comparisons (Bonferroni posttest). Mixed factor ANOVA was used to compare active and inactive lever presses during self-administration and extinction. One-way ANOVA and post-hoc comparisons (Bonferroni posttest) were used to compare responding during reinstatement and extinction. The alpha was set at $p < 0.05$ and all data are expressed as the mean $\pm$ S.E.M.

3. Results

3.1. Elevated Plus Maze

In cocaine-naïve animals, FG 7142 (10 mg/kg) produced an anxiety-like profile on the EPM (Fig. 1), with significantly less time spent in the open arms ($t_{16} = 2.33$, $p = 0.034$) when compared with vehicle injection. This effect was not observed in rats with prior cocaine self-administration experience as cocaine-experienced subjects treated with FG 7142 or vehicle spent an equivalent amount of time exploring the open arms of the EPM. FG 7142 did not affect total locomotor behavior in either group, as determined by closed arm entries (data not shown).

3.2. Plasma Corticosterone

ANOVA revealed that FG 7142 influenced the corticosterone response of cocaine-naïve rats to EPM exposure ($F_{1,12} = 17.80$, $p = 0.003$). Bonferroni post-hoc tests ($p < 0.05$) indicated that animals injected with FG 7142 exhibited a significant increase in plasma corticosterone following EPM exposure, and that these levels were significantly higher than animals injected with vehicle alone and exposed to the EPM (Fig. 2). In contrast, FG 7142 did not influence the corticosterone response of cocaine-experienced animals to the EPM. No animals exhibited a significant increase in corticosterone to the EPM alone (vehicle treated + EPM exposure).

3.3. Cocaine Self-administration and Extinction

All animals readily acquired and maintained cocaine self-administration criteria. Lever pressing did not significantly differ between experimental cohorts. Daily cocaine intake during self-administration was 16.6 ± 0.37 mg/kg per 2-h session (average of the last 3 days). All animals met extinction criteria following self-administration, with an average of 12.0 ± 0.74 days of extinction (summarized in Fig. 3 and Fig. 4). Inactive lever responding was uniformly low in all animals and did not differ between groups during or after self-administration.

3.4. Reinstatement of Cocaine-seeking

FG 7142 and Reinstatement—FG 7142 failed to reinstate cocaine-seeking at any dose (Fig. 3A and 3B); however, statistical analysis indicated a significant reduction in lever pressing at the 10 and 20 mg/kg doses ($F_{4,39}=4.03, p=0.009$; Bonferroni $p<0.05$ compared to extinction). Following testing with FG 7142 in the first cohort (Fig. 3A), yohimbine (1.25 mg/kg) significantly reinstated cocaine-seeking in the same animals ($F_{4,123}=5.47, p=0.0004$; Bonferroni $p<0.05$ compared to extinction).

FG 7142 and Cue-induced Reinstatement—Presentation of cocaine associated cues significantly reinstated active lever pressing ($F_{8,35}=5.35, p=0.0004$; Bonferroni $p<0.05$ for both cue-induced reinstatement tests compared to extinction). Prior repeated treatment of animals with FG 7142 (10 mg/kg) on the three days preceding the second cue-induced reinstatement test did not influence lever pressing in response to cue presentation (Fig. 4).

4. Discussion

The current study confirmed the anxiogenic-like and stress-potentiating qualities of FG 7142 in cocaine-naïve animals. We found these effects of FG 7142 to be attenuated in animals with a history of cocaine self-administration. Furthermore, FG 7142 failed to reinstate extinguished cocaine-seeking across a wide dose range or alter subsequent cue-induced reinstatement after repeated treatment. Our results suggest that chronic cocaine self-administration decreased the efficacy of BZ inverse agonism to induce anxiety and potentiate responses to stress. This is in contrast to other anxiogenic-like and stress-inducing environments or compounds, which show potentiation after chronic cocaine experience (Alves et al., 2008; Erb et al., 2003; Magalhaes et al., 2002; Yang et al., 1992).

Anxiety measures in rodent models (Landgraf and Wigger, 2002) allow for the assessment of an individual subject's stress coping strategy by placing them into a novel environment with stress-inducing qualities (i.e., the open arms of the EPM) and recording their behavioral reactions to this stress. The EPM is an established measure of anxiety (behavior induced by the stressful qualities of the open arms), and has been routinely used to assess the anxiogenic-like effects of pharmacological stressors, including FG 7142 (Pellow et al., 1987). In the current study, FG 7142 increased anxiety-like behavior on the EPM in cocaine-naïve animals, and elicited an increase in plasma corticosterone levels following EPM exposure, an effect not exhibited by animals treated with vehicle. These results suggest that FG 7142 elicits an anxiety-like state, and also potentiates an animal's stress response. FG 7142 failed to exhibit these effects in cocaine-experienced animals, as treatment with FG 7142 or vehicle resulted in similar levels of open arm exploration in the EPM and equivalent plasma corticosterone responses to EPM exposure. This finding was unexpected, as chronic cocaine exposure has been reported to increase the behavioral and physiological responses of rats to environmental stressors used to assess anxiety, including the forced swim test (Magalhaes et al., 2002) and the defensive withdrawal test (Yang et al., 1992), as well as to pharmacological stressors, such as CRF (Erb et al., 2003).

Cocaine-experienced animals treated with vehicle or FG 7142 exhibited non-significantly elevated plasma corticosterone levels following exposure to the EPM. This observation could reflect a slight increase in the EPM stress response in these animals as compared to cocaine-naïve animals, although unapparent in the animals' behavior on the EPM. Moderate cocaine experience, such as the regimen used herein, has previously been shown to lack an effect on anxiety-like behavior in the EPM (Mantsch et al., 2008). These minimal effects of cocaine experience on EPM performance likely result from the lower stress-inducing qualities of this test when compared to tests with more potent stress inducing qualities (e.g., forced swim test) that elicit potentiated responses from cocaine-experienced animals

(Magalhaes et al., 2002). At any rate, contrary to its effects in cocaine-naïve animals, FG 7142 did not influence anxiety-like behavior on the EPM, and failed to potentiate the stress response (i.e., further increase plasma corticosterone) of cocaine-experienced animals to the EPM. These results suggest that the anxiogenic-like and stress potentiating qualities of FG 7142 are diminished following chronic cocaine self-administration.

Reinstatement of cocaine-seeking has been reported after multiple anxiogenic-like stressors, including acute intermittent footshock stress (Erb et al., 1996; Korte and De Boer, 2003), food deprivation stress (Carroll, 1985), and yohimbine administration (Feltenstein and See, 2006; Pellow et al., 1985), although this effect does not generalize to all anxiogenic-like stressors (Shaham et al., 2000). While the anxiogenic-like and stress-inducing qualities of FG 7142 have been previously established (Evans and Lowry, 2007; File et al., 1985; Pellow and File, 1985), we found that FG 7142 failed to reinstate cocaine-seeking. In fact, two of the higher doses of FG 7142 (10 and 20 mg/kg; Fig 3B) decreased active lever responding. While this reduction could be interpreted as freezing behavior (Conti et al., 1990) or a sedative effect (Chuck et al., 2006), this possibility is unlikely as the magnitude of change was quite small due to the very low baseline responding, and was not seen for inactive lever presses.

In contrast to FG 7142, which did not reinstate cocaine seeking, these same animals exhibited yohimbine induced reinstatement, similar to that previously reported (Anker and Carroll, 2010; Feltenstein and See, 2006). Repeated pretreatment with FG 7142 had no effect on subsequent cue-induced reinstatement, also in contrast to what has been observed with yohimbine administration (Kupferschmidt et al., 2009). Some previous studies have directly compared the behavioral and physiological effects of yohimbine and FG 7142, with some notable similarities and differences between the two drugs. Both of these compounds elicit anxiety-like behaviors, including decreased open arm time on the EPM (Atack et al., 2005; Pellow et al., 1985), decreased time engaged in social interactions (Guy and Gardner, 1985), and decreased exploration of a novel environment (Mason et al., 1998). Furthermore, both FG 7142 and yohimbine activate similar brain regions associated with stress responses as indicated by the induction of fos expression, including the bed nucleus of the stria terminalis (BNST), frontal cortex (Funk et al., 2006; Singewald et al., 2003), and basolateral amygdala (Hale et al., 2010). All of these brain structures have been implicated in the reinstatement of cocaine-seeking (Capriles et al., 2003; Erb and Stewart, 1999; McLaughlin and See, 2003), including yohimbine induced reinstatement (Funk et al., 2006; Nair et al., 2011). While similarities exist, some previously noted differences between yohimbine and FG 7142 might contribute to their divergent effects on reinstatement behavior. For example, yohimbine, but not FG 7142, increased CRF mRNA in the BNST (Funk et al., 2006), a phenomenon closely associated with stress-induced reinstatement (Erb et al., 2000). Also, while yohimbine potently and reliably elicits norepinephrine release in the frontal cortex, an effect believed to be important in the reinstatement of cocaine-seeking, reports on the ability of FG 7142 to affect this measure are inconsistent (Dazzi et al., 2002; Mason et al., 1998). The modest (compared to yohimbine) effects of FG 7142 on these specific substrates may help explain its inability to reinstate cocaine-seeking.

The failure of FG 7142 to reinstate cocaine-seeking may also relate to its unique pharmacological profile. In contrast to yohimbine, which increases norepinephrine levels through presynaptic α -2 blockade (Charney et al., 1983) and also affects other monoamines (Dzung Le et al., 2009; Millan et al., 2000), FG 7142 works via inverse agonism of the BZ receptor (Dorow et al., 1983). The lack of an FG 7142 effect on EPM behavior and the corticosterone response associated with exposure to this apparatus in cocaine-experienced animals, as well as the lack of an effect on the reinstatement of cocaine-seeking, may be due to a down-regulation of central BZ receptors associated with chronic cocaine intake, as

previously reported (Goeders, 1991; McAllister et al., 1987). In spite of this finding, BZ agonists successfully decrease self-administration behavior (Goeders et al., 1989), as well as cue-induced reinstatement (Goeders et al., 2009) in cocaine-experienced animals. Thus, it appears that inverse agonist action at the BZ receptor may be selectively or preferentially altered by chronic cocaine intake.

In conclusion, we have demonstrated that the anxiogenic-like qualities of FG 7142 are attenuated in animals with a history of cocaine self-administration. Furthermore, administration of FG 7142 does not influence the corticosterone response of cocaine-experienced animals to EPM exposure. This adaptive response and the unique pharmacological properties of FG 7142 may account for the lack of an effect of FG 7142 on the reinstatement of cocaine-seeking. Future studies designed to directly assess BZ receptor levels and function in specific brain regions involved in addiction and stress responses in cocaine-naïve and cocaine-experienced animals may further reveal the role of this receptor in chronic cocaine-induced neuroplasticity and treatment response.

Research Highlights

- Relapse to cocaine abuse can be elicited by stress.
- Pharmacological stressors can serve as probes in understanding this phenomenon.
- FG 7142 potentiates the stress response of rats to the EPM.
- These effects are absent in cocaine trained rats.
- FG 7142 does not influence extinguished cocaine seeking behavior.

Acknowledgments

This research was supported by NIDA grant DA021690 (RES) and NIH grant C06 RR015455.

References

- Alves CJ, Magalhaes A, Summavielle T, Melo P, De Sousa L, Tavares MA, et al. Hormonal, neurochemical, and behavioral response to a forced swim test in adolescent rats throughout cocaine withdrawal. *Ann NY Acad Sci.* 2008; 1139:366–373. [PubMed: 18991883]
- Anker JJ, Carroll ME. Reinstatement of cocaine seeking induced by drugs, cues, and stress in adolescent and adult rats. *Psychopharmacology.* 2010; 208:211–222. [PubMed: 19953228]
- Atack JR, Hutson PH, Collinson N, Marshall G, Bentley G, Moyes C, et al. Anxiogenic properties of an inverse agonist selective for alpha3 subunit-containing GABA A receptors. *Br J Pharmacol.* 2005; 144:357–366. [PubMed: 15655523]
- Banna KM, Back SE, Do P, See RE. Yohimbine stress potentiates conditioned cue-induced reinstatement of heroin-seeking in rats. *Behav Brain Res.* 2010; 208:144–148. [PubMed: 19931573]
- Bongiovanni M, See RE. A comparison of the effects of different operant training experiences and dietary restriction on the reinstatement of cocaine-seeking in rats. *Pharmacol Biochem Behav.* 2008; 89:227–233. [PubMed: 18230406]
- Brown ZJ, Tribe E, D'Souza NA, Erb S. Interaction between noradrenaline and corticotrophin-releasing factor in the reinstatement of cocaine seeking in the rat. *Psychopharmacology.* 2009; 203:121–130. [PubMed: 18985323]
- Capriles N, Rodaros D, Sorge RE, Stewart J. A role for the prefrontal cortex in stress- and cocaine-induced reinstatement of cocaine seeking in rats. *Psychopharmacology.* 2003; 168:66–74. [PubMed: 12442201]

- Carroll ME. The role of food deprivation in the maintenance and reinstatement of cocaine-seeking behavior in rats. *Drug Alcohol Depend.* 1985; 16:95–109. [PubMed: 4075974]
- Charney D, Heninger G, Redmond D Jr. Yohimbine induced anxiety and increased noradrenergic function in humans: effects of diazepam and clonidine. *Life Sci.* 1983; 33:19–29. [PubMed: 6865647]
- Chuck TL, McLaughlin PJ, Arizzi-LaFrance MN, Salamone JD, Correa M. Comparison between multiple behavioral effects of peripheral ethanol administration in rats: sedation, ataxia, and bradykinesia. *Life Sci.* 2006; 79:154–161. [PubMed: 16487981]
- Conti L, Maciver C, Ferkany J, Abreu M. Footshock-induced freezing behavior in rats as a model for assessing anxiolytics. *Psychopharmacology.* 1990; 102:492–497. [PubMed: 1982903]
- Dazzi L, Ladu S, Spiga F, Vacca G, Rivano A, Pira L, et al. Chronic treatment with imipramine or mirtazapine antagonizes stress- and FG7142-induced increase in cortical norepinephrine output in freely moving rats. *Synapse.* 2002; 43:70–77. [PubMed: 11746735]
- Dorow R, Horowski R, Paschelke G, Amin M. Severe anxiety induced by FG 7142, a beta-carboline ligand for benzodiazepine receptors. *Lancet.* 1983; 2:98–99. [PubMed: 6134976]
- Dzung Le A, Funk D, Harding S, Juzysch W, Fletcher PJ. The role of noradrenaline and 5-hydroxytryptamine in yohimbine-induced increases in alcohol-seeking in rats. *Psychopharmacology.* 2009; 204:477–488. [PubMed: 19229522]
- Epstein D, Preston K, Stewart J, Shaham Y. Toward a model of drug relapse: an assessment of the validity of the reinstatement procedure. *Psychopharmacology.* 2006; 189:1–16. [PubMed: 17019567]
- Erb S, Funk D, Le AD. Prior, repeated exposure to cocaine potentiates locomotor responsivity to central injections of corticotropin-releasing factor (CRF) in rats. *Psychopharmacology.* 2003; 170:383–389. [PubMed: 12955298]
- Erb S, Hitchcott PK, Rajabi H, Mueller D, Shaham Y, Stewart J. Alpha-2 adrenergic receptor agonists block stress-induced reinstatement of cocaine seeking. *Neuropsychopharmacology.* 2000; 23:138–150. [PubMed: 10882840]
- Erb S, Salmaso N, Rodaros D, Stewart J. A role for the CRF-containing pathway from central nucleus of the amygdala to bed nucleus of the stria terminalis in the stress-induced reinstatement of cocaine seeking in rats. *Psychopharmacology.* 2001; 158:360–365. [PubMed: 11797056]
- Erb S, Shaham Y, Stewart J. Stress reinstates cocaine-seeking behavior after prolonged extinction and a drug-free period. *Psychopharmacology.* 1996; 128:408–412. [PubMed: 8986011]
- Erb S, Stewart J. A role for the bed nucleus of the stria terminalis, but not the amygdala, in the effects of corticotropin-releasing factor on stress-induced reinstatement of cocaine seeking. *J Neurosci.* 1999; 19:RC35. [PubMed: 10516337]
- Evans AK, Lowry CA. Pharmacology of the beta-carboline FG-7142, a partial inverse agonist at the benzodiazepine allosteric site of the GABA A receptor: neurochemical, neurophysiological, and behavioral effects. *CNS Drug Rev.* 2007; 13:475–501. [PubMed: 18078430]
- Feltenstein MW, Altar CA, See RE. Aripiprazole blocks reinstatement of cocaine seeking in an animal model of relapse. *Biol Psychiatry.* 2007; 61:582–590. [PubMed: 16806092]
- Feltenstein MW, Byrd EA, Henderson AR, See RE. Attenuation of cocaine-seeking by progesterone treatment in female rats. *Psychoneuroendocrinology.* 2009; 34:343–352. [PubMed: 18977603]
- Feltenstein MW, See RE. Potentiation of cue-induced reinstatement of cocaine-seeking in rats by the anxiogenic drug yohimbine. *Behav Brain Res.* 2006; 174:1–8. [PubMed: 16920204]
- File SE, Pellow S, Braestrup C. Effects of the [beta]-carboline, FG 7142, in the social interaction test of anxiety and the holeboard: correlations between behaviour and plasma concentrations. *Pharmacol Biochem Behav.* 1985; 22:941–944. [PubMed: 3839582]
- Funk D, Li Z, Le AD. Effects of environmental and pharmacological stressors on c-fos and corticotropin-releasing factor mRNA in rat brain: relationship to the reinstatement of alcohol seeking. *Neuroscience.* 2006; 138:235–243. [PubMed: 16359808]
- Goeders N. Cocaine differentially affects benzodiazepine receptors in discrete regions of the rat brain: persistence and potential mechanisms mediating these effects. *J Pharmacol Exp Ther.* 1991; 259
- Goeders NE. Stress and cocaine addiction. *J Pharmacol Exp Ther.* 2002; 301:785–789. [PubMed: 12023504]

- Goeders NE, Clampitt DM, Keller C, Sharma M, Guerin GF. Alprazolam and oxazepam block the cue-induced reinstatement of extinguished cocaine seeking in rats. *Psychopharmacology*. 2009; 201:581–588. [PubMed: 18797849]
- Goeders NE, McNulty MA, Mirkis S, McAllister KH. Chlordiazepoxide alters intravenous cocaine self-administration in rats. *Pharmacol Biochem Behav*. 1989; 33:859–866. [PubMed: 2559416]
- Guy AP, Gardner CR. Pharmacological characterisation of a modified social interaction model of anxiety in the rat. *Neuropsychobiology*. 1985; 13:194–200. [PubMed: 2864655]
- Hale MW, Johnson PL, Westerman AM, Abrams JK, Shekhar A, Lowry CA. Multiple anxiogenic drugs recruit a parvalbumin-containing subpopulation of GABAergic interneurons in the basolateral amygdala. *Prog Neuropsychopharmacol Biol Psychiatry*. 2010; 34:1285–1293. [PubMed: 20647026]
- Hogg S. A review of the validity and variability of the elevated plus-maze as an animal model of anxiety. *Pharmacol Biochem Behav*. 1996; 54:21–30. [PubMed: 8728535]
- Ida Y, Elsworth JD, Roth RH. Anxiogenic [beta]-carboline FG 7142 produces activation of noradrenergic neurons in specific brain regions of rats. *Pharmacol Biochem Behav*. 1991; 39:791–793. [PubMed: 1784607]
- Korf J, Aghajanian GK, Roth RH. Increased turnover of norepinephrine in the rat cerebral cortex during stress: role of the locus coeruleus. *Neuropharmacology*. 1973; 12:933–938. [PubMed: 4750561]
- Korte SM, De Boer SF. A robust animal model of state anxiety: fear-potentiated behaviour in the elevated plus-maze. *Eur J Pharmacol*. 2003; 463:163–175. [PubMed: 12600708]
- Kupferschmidt D, Tribe E, Erb S. Effects of repeated yohimbine on the extinction and reinstatement of cocaine seeking. *Pharmacol Biochem Behav*. 2009; 91:473–480. [PubMed: 18824018]
- Landgraf R, Wigger A. High vs low anxiety-related behavior rats: an animal model of extremes in trait anxiety. *Behav Genet*. 2002; 32:301–314. [PubMed: 12405513]
- Le AD, Harding S, Juzysch W, Funk D, Shaham Y. Role of alpha-2 adrenoceptors in stress-induced reinstatement of alcohol seeking and alcohol self-administration in rats. *Psychopharmacology*. 2005; 179:366–373. [PubMed: 15551068]
- Magalhaes A, Tavares M, Sousa L. Postnatal cocaine exposure: effects on behavior of rats in forced swim test. *Ann NY Acad Sci*. 2002; 965:529–534. [PubMed: 12105128]
- Mantsch JR, Baker DA, Francis DM, Katz ES, Hoks MA, Serge JP. Stressor- and corticotropin releasing factor-induced reinstatement and active stress-related behavioral responses are augmented following long-access cocaine self-administration by rats. *Psychopharmacology*. 2008; 195:591–603. [PubMed: 17899015]
- Marin RH, Martijena ID, Arce A. Effect of diazepam and a beta-carboline on open-field and T-maze behaviors in 2-day-old chicks. *Pharmacol Biochem Behav*. 1997; 58:915–921. [PubMed: 9408196]
- Mason K, Heal DJ, Stanford SC. The anxiogenic agents, yohimbine and FG 7142, disrupt the noradrenergic response to novelty. *Pharmacol Biochem Behav*. 1998; 60:321–327. [PubMed: 9632213]
- McAllister, K.; Goeders, N.; Dworkin, S.; Boisse, N.; Samoriski, G.; Guarino, J., et al. Chronic cocaine modifies brain benzodiazepine receptor densities. In: Harris, LS., editor. *Proceedings of the Problems of Drug Dependence*. Rockville, MD USA: 1987. p. 101-108.
- McLaughlin J, See RE. Selective inactivation of the dorsomedial prefrontal cortex and the basolateral amygdala attenuates conditioned-cued reinstatement of extinguished cocaine-seeking behavior in rats. *Psychopharmacology*. 2003; 168:57–65. [PubMed: 12845418]
- Millan MJ, Newman-Tancredi A, Audinot V, Cussac D, Lejeune F, Nicolas JP, et al. Agonist and antagonist actions of yohimbine as compared to fluparoxan at alpha(2)-adrenergic receptors (AR)s, serotonin (5-HT)(1A), 5-HT(1B), 5-HT(1D) and dopamine D(2) and D(3) receptors. Significance for the modulation of frontocortical monoaminergic transmission and depressive states. *Synapse*. 2000; 35:79–95. [PubMed: 10611634]
- Nair SG, Navarre BM, Cifani C, Pickens CL, Bossert JM, Shaham Y. Role of dorsal medial prefrontal cortex dopamine D1-family receptors in relapse to high-fat food seeking induced by the anxiogenic drug yohimbine. *Neuropsychopharmacology*. 2011; 36:497–510. [PubMed: 20962767]

- Paine TA, Jackman SL, Olmstead MC. Cocaine-induced anxiety: alleviation by diazepam, but not buspirone, dimenhydrinate or diphenhydramine. *Behav Pharmacol.* 2002; 13
- Pellow S, Chopin P, File SF, Briley M. Validation of open: closed arm entries in an elevated plus-maze as a measure of anxiety in the rat. *J Neurosci Methods.* 1985; 14:149–167. [PubMed: 2864480]
- Pellow S, File SE. The effects of putative anxiogenic compounds (FG 7142, CGS 8216 and Ro 15-1788) on the rat corticosterone response. *Phys Behav.* 1985; 35:587–590.
- Pellow S, Johnston A, File S. Selective agonists and antagonists for 5-hydroxytryptamine receptor subtypes, and interactions with yohimbine and FG 7142 using the elevated plus-maze test in the rat. *J Pharm Pharmacol.* 1987; 39:917. [PubMed: 2892916]
- Reichel C, See R. Modafinil effects on reinstatement of methamphetamine seeking in a rat model of relapse. *Psychopharmacology.* 2010; 210:337–346. [PubMed: 20352413]
- Sakakura M, Saito Y, Takebe K, Yamashita I, Ishii K. Time course of hypothalamic CRF activity after the administration of two different stresses. *Endocrinol Jpn.* 1976; 23:413–416. [PubMed: 1087604]
- See RE, Waters RP. Pharmacologically-induced stress: a cross-species probe for translational research in drug addiction and relapse. *Am J Transl Res.* 2011; 3:81–89. [PubMed: 21139808]
- Shaham Y, Erb S, Stewart J. Stress-induced relapse to heroin and cocaine seeking in rats: a review. *Brain Res Brain Res Rev.* 2000; 33:13–33. [PubMed: 10967352]
- Shalev U, Erb S, Shaham Y. Role of CRF and other neuropeptides in stress-induced reinstatement of drug seeking. *Brain Res.* 2010; 1314:15–28. [PubMed: 19631614]
- Shepard JD, Bossert JM, Liu SY, Shaham Y. The anxiogenic drug yohimbine reinstates methamphetamine seeking in a rat model of drug relapse. *Biol Psychiatry.* 2004; 55:1082–1089. [PubMed: 15158427]
- Singewald N, Salchner P, Sharp T. Induction of c-Fos expression in specific areas of the fear circuitry in rat forebrain by anxiogenic drugs. *Biol Psychiatry.* 2003; 53:275–283. [PubMed: 12586446]
- Sinha R, Garcia M, Paliwal P, Kreek M, Rounsaville B. Stress-induced cocaine craving and hypothalamic-pituitary-adrenal responses are predictive of cocaine relapse outcomes. *Arch Gen Psychiatry.* 2006; 2006:8.
- Sutton RE, Koob GF, Le Moal M, Rivier J, Vale W. Corticotropin releasing factor produces behavioural activation in rats. *Nature.* 1982; 297:331–333. [PubMed: 6978997]
- Yang X-M, Gorman A, Dunn AJ, Goeders N. Anxiogenic effects of acute and chronic cocaine administration: neurochemical and behavioral studies. *Pharmacol Biochem Behav.* 1992; 41:643–650. [PubMed: 1584846]

Fig. 1. Effects of acute FG 7142 on EPM performance. FG 7142 increased the anxiety-like behavior in cocaine-naïve animals (vehicle $n=8$; FG 7142 $n=10$), but not in cocaine-experienced animals (vehicle $n=10$; FG 7142 $n=11$). Cocaine-naïve animals injected with FG 7142 explored the open arms of the maze significantly less than animals injected with vehicle ($*p<0.05$), an effect that was absent in cocaine-experienced animals.

Fig. 2.

Effects of acute FG 7142 and EPM exposure on plasma corticosterone levels. FG 7142 administration and EPM exposure elicited a significant increase ($*p<0.05$) in plasma corticosterone in cocaine-naïve animals (vehicle $n=8$; FG 7142 $n=6$), an effect that was not observed in cocaine-experienced animals (vehicle $n=5$; FG 7142 $n=6$).

Fig. 3.

Acute FG 7142 effects after chronic cocaine self-administration and extinction. FG 7142 failed to increase lever responding across a low (Fig. 3A; $n=26$) or high (Fig. 3B; $n=8$) dose range, with some doses of FG 7142 (Fig. 3B) leading to decreased responding ($*p<0.05$). In contrast, acute yohimbine (1.25 mg/kg) significantly reinstated cocaine-seeking ($*p<0.05$).

Fig. 4.
The effects of pretreatment with FG 7142 on subsequent cue-induced reinstatement.
Presentation of conditioned drug-paired cues reinstated cocaine-seeking ($*p < 0.05$, $n = 4$).
Pretreatment with FG 7142 did not influence this behavior.