Published in final edited form as:

Lancet Infect Dis. 2004 June ; 4(6): 323–324. doi:10.1016/S1473-3099(04)01038-2.

Global warming and malaria: a call for accuracy

Paul Reiter, Christopher J Thomas, Peter M Atkinson, Simon I Hay, Sarah E Randolph, David J Rogers, G Dennis Shanks, Robert W Snow, and Andrew Spielman

PR is professor of medical entomology, Institut Pasteur, Paris, France; CJT is senior lecturer in spatial ecology and Wolfson Institute fellow in health and environment, School of Biological and Biomedical Sciences, University of Durham, UK; PA is professor of geography, University of Southampton, UK; SIH is a Wellcome Trust research fellow, SER is professor of parasite ecology, and DJR is professor of ecology, Department of Zoology, University of Oxford, UK; GDS is at the US Army Centre for Health Promotion and Preventive Medicine, MD, USA; RWS is professor of tropical public health, University of Oxford; and AS is professor of tropical public health, immunology, and infectious diseases, Harvard School of Public Health, Boston, MA, USA. SIH and RWS are also at the KEMRI Wellcome Trust collaborative programme, Kenya.

For more than a decade, malaria has held a prominent place in speculations on the impacts of global climate change. Mathematical models that “predict” increases in the geographic distribution of malaria vectors and the prevalence of the disease have received wide publicity. Efforts to put the issue into perspective^{1–5} are rarely quoted and have had little influence on the political debate. The model proposed by Frank C Tanser and colleagues⁶ in *The Lancet* and the accompanying Commentary by Simon Hales and Alistair Woodward⁷ are typically misleading examples.

The relation between climate and malaria transmission is complex and varies according to location,² yet Tanser et al base their projections on thresholds derived from a mere 15 African locations. Slight adjustments of values assigned to such thresholds and rules can influence spatial predictions strongly.⁸ The authors invest considerable effort in assessing the sensitivity of their model to climate change scenarios but do not report the internal sensitivities to thresholds and rules. The predictive skill of their model is low (63% sensitivity, 95% CI 61–65%) but they consider projections acceptable if prevalence is projected “to within a month” (presumably ± 1 month?), thereby biasing their model towards success. A model covering an entire year in a parasite-positive site would always be correct, although in such areas it would be relatively insensitive to climate. By contrast, sites in which transmission is seasonal would provide a more reliable test of accuracy, but estimation is more difficult because climate sensitivity is greater. Furthermore, because parasite clearance in communities is not instantaneous,⁹ spot samples of parasitaemia on survey dates are not a suitable indicator of the duration of the transmission season. Lastly, “person/months” are unsuitable as a measure of transmission: an extension of season from 1 to 4 months will have more impact than from 10 to 12 months. According to their model, an extension of transmission from 11 to 12 months results in 10^6 more person/months in a population of 10^6 people, whereas an extension from 1 to 5 months gives the same increase in a population of 250 000.

What Tanser and colleagues have modelled is merely the duration of the transmission season, which they interpret as “heightened transmission” and increased incidence. A greater failing is their reliance on “parasite-ratio studies”. The relations between transmission

season and parasite prevalence, and parasite prevalence and clinical disease, are unclear but unlikely to be linear. Moreover, they use 1995 data for human populations, although these are projected to double by 2030. In addition, the proportion living in urban areas—with a specific climate¹⁰ and orders of magnitude less malaria transmission^{11,12}—is projected to rise from 37% to 53%.¹³ For all these reasons, we do not accept the model as a “baseline against which interventions can be planned”.

It is regrettable that many involved in this debate ignore the rich heritage of literature on the subject. For example, in 1937, in his classic textbook,¹⁴ L W Hackett stated: “Everything about malaria is so moulded and altered by local conditions that it becomes a thousand different diseases and epidemiological puzzles. Like chess, it is played with a few pieces, but is capable of an infinite variety of situations”. A pressing question in Hackett’s time was the changing distribution of the disease in Europe. On the role of climate, he wrote: “Certainly, climate lays down the broad lines of malaria distribution ... Nevertheless, although this is a very simple and plausible explanation ... even the early malariologists felt that there was something unsatisfactory about it ... malaria has not so much receded as it has contracted, oftentimes toward the north ... Thus in Germany it is the northern coast which is still malarious, the south is free ... There is, therefore, no climatic reason why (malaria) should have abandoned south Germany or the French Riviera”.

We quote Hackett because we feel that the classic components of science—unbiased observation and systematic experimentation—cannot be sidestepped with models that omit many of his chess pieces. Yet Hales and Woodward⁷ begin by stating: “The present geographical distribution of malaria is explained by a combination of environmental factors (especially climate) and social factors (such as disease-control measures)”. In our opinion, “even the early malariologists” would surely disagree: much of the decline of malaria in Europe took place without control measures during a period when the climate was warming.

The text by Hales and Woodward that follows displays a lack of knowledge. Thus, “Most people at risk of malaria live in areas of stable transmission ... ” is simply wrong. It is true that in many parts of the world malaria is termed “stable” because transmission remains relatively constant from year to year, the disease is endemic, the collective immunity is high, and epidemics are uncommon. However, in many other regions, the disease is endemic but “unstable” because annual transmission varies considerably, and the potential for epidemics is great. Climatic factors, particularly rainfall, are sometimes, but by no means always, relevant.¹⁵

Again, “On the fringes of endemic zones, where transmission is limited by rainfall ... there are strong seasonal patterns, and occasional major epidemics” is also wrong. In many regions, far from any “fringes”, malaria is endemic, stable, but highly seasonal. For example, in semi-arid regions of Mali, transmission is restricted to the rainy season, from July to September. The same 3 months constituted the transmission season for *Plasmodium falciparum* in Italy before it was eliminated.¹⁶ Paradoxically, in parts of the Sudan, rainfall is restricted to a month at most, but malaria is transmitted throughout the year. Female *Anopheles gambiae* survive drought and heat by resting in dwellings and other sheltered places.¹⁷ Blood feeding and transmission continue, but the mosquitoes do not develop eggs until the rains return. This phenomenon, termed gonotrophic dissociation, is remarkably similar to the winter survival strategy of *Anopheles atroparvus*, the principal vector of malaria in Holland until the mid 20th century.¹⁶

By contrast, malaria is unstable in many regions that normally have abundant rainfall, and epidemics occur during periods of drought. An illustrative example is the catastrophic 1934–35 epidemic in Ceylon (now Sri Lanka), estimated to have killed 100 000 people.¹⁸ Worst

hit was the south-western quadrant of the country, where average annual rainfall is greater than 250 cm, and malaria was endemic, but unstable and relatively infrequent. The dominant vector, *Anopheles culicifacies*, breeds along the banks of rivers and tends to be scarce in normal years. In the years 1928–33 there was abundant rainfall, river flow was high, *A. culicifacies* was rare, and the human population was exceptionally malaria-free. However, after failure of two successive monsoons, the drying rivers produced colossal numbers of *A. culicifacies*, and the resulting epidemic was exacerbated by the low collective immunity. In the drier parts of the island, where *A. culicifacies* was dominant but transmission was more stable, immunity protected the population from the worst ravages of the disease.

Hales and Woodward state that “the underlying problem” of the future “extension of seasonality” of malaria is “pollution of the atmosphere”, and call for rich countries to “recognise their obligations to the poorest by substantially reducing fossil-fuel consumption”. We understand public anxiety about climate change, but are concerned that many of these much-publicised predictions are ill informed and misleading. We urge those involved to pay closer attention to the complexities of this challenging subject.

References

1. Reiter P. From Shakespeare to Defoe: malaria in England in the little ice age. *Emerg Infect Dis.* 2000; 6:1–11. [PubMed: 10653562]
2. Reiter P. Climate change and mosquito-borne disease. *Environ Health Perspect.* 2001; 109(suppl 1): 141–61. [PubMed: 11250812]
3. Hay SI, Cox J, Rogers DJ, et al. Climate change and the resurgence of malaria in the East African highlands. *Nature.* 2002; 415:905–09. [PubMed: 11859368]
4. Shanks GD, Hay SI, Stern DI, Biomndo K, Snow RW. Meteorologic influences on *Plasmodium falciparum* malaria in the highland tea estates of Kericho, Western Kenya. *Emerg Infect Dis.* 2002; 8:1404–08. [PubMed: 12498655]
5. Rogers DJ, Randolph SE. The global spread of malaria in a future, warmer world. *Science.* 2000; 289:1763–66. [PubMed: 10976072]
6. Tanser FC, Sharp B, le Sueur D. Potential effect of climate change on malaria transmission in Africa. *Lancet.* 2003; 362:1792–98. [PubMed: 14654317]
7. Hales S, Woodward A. Climate change will increase demands on malaria control in Africa. *Lancet.* 2003; 362:1775. [PubMed: 14654310]
8. Thomas CJ, Davies G, Dunn CE. Mixed picture for changes in stable malaria distribution with future climate in Africa. *Trends Parasitol.* 2004; 20:216–20. [PubMed: 15105021]
9. Smith T, Charlwood JD, Kihonda J, et al. Absence of seasonal variation in malaria parasitaemia in an area of intense seasonal transmission. *Acta Trop.* 1993; 54:55–72. [PubMed: 8103627]
10. Arnfield A. Two decades of urban climate research: a review of turbulence, exchanges of energy and water, and the urban heat island. *Int J Climatol.* 2003; 23:1–26.
11. Snow R, Trape J, Marsh K. The past, present and future of childhood malaria mortality in Africa. *Trends Parasitol.* 2001; 17:593–97. [PubMed: 11756044]
12. Robert V, Macintyre K, Keating J, et al. Malaria transmission in urban sub-Saharan Africa. *Am J Trop Med Hyg.* 2003; 68:169–76. [PubMed: 12641407]
13. United Nations. World urbanization prospects: the 2001 revision. Data tables and highlights. United Nations; New York: 2002.
14. Hackett, LW. Malaria in Europe, an ecological study. Oxford University Press; London: 1937.
15. Gilles, HM.; Warrell, DA., editors. Bruce-Chwatt's essential malariology. Edward Arnold; London: 1993.
16. Bruce-Chwatt, LJ.; de Zulueta, J. The rise and fall of malaria in Europe, a historico-epidemiological study. Oxford University; Oxford: 1980.
17. Omer SM, Cloudsley-Thompson JL. Survival of female *Anopheles gambiae* Giles through a 9-month dry season in Sudan. *Bull World Health Organ.* 1970; 42:319–30. [PubMed: 5310144]

18. Dunn, C. Malaria in Ceylon: an enquiry into its causes. Bailliere, Tindall and Cox; London: 1937.