Published in final edited form as:

Pharmacol Biochem Behav. 2010 October ; 96(4): 521–525. doi:10.1016/j.pbb.2010.07.015.

LOCAL CEREBRAL GLUCOSE UTILIZATION IN RATS EXPOSED TO AN ENRICHED ENVIRONMENT: A COMPARISON TO IMPOVERISHMENT

A.K. Läck, K.E. Gill, and L.J. Porrino

Department of Physiology and Pharmacology, Wake Forest University School of Medicine
Winston-Salem, NC 27157

Abstract

Environmental enrichment and environmental impoverishment have been shown to differentially alter brain function. Here, we investigate the effects of enrichment versus impoverishment on cerebral use of glucose in rodents. Rats were housed from postnatal day 28 to day 58 in either a socially and environmentally enriched environment or an impoverished environment devoid of other rats or environmental stimuli. Locomotor activity was measured at the end of the enrichment/impoverishment period. Following the duration of the exposure to these environments, cerebral metabolic rate of glucose utilization was determined using quantitative 2-[¹⁴C] deoxyglucose autoradiography in 37 brain regions in the cerebral cortex, forebrain, brain stem and thalamus. There were no differences in locomotor activity between the conditions. The nucleus accumbens core and shell had significantly higher rates of glucose utilization in enriched compared to impoverished animals. These data suggest that environment has a significant effect on the brain function which may help to explain the beneficial and protective effects of enrichment against drug abuse and addiction.

Keywords

Environmental enrichment; Nucleus accumbens; Cerebral glucose utilization; Rat

Introduction

Recent research has explored the effects of environmental enrichment on neurochemistry and behavior, from changes in the dopaminergic markers (Zakharova et al., 2009; Zhu et al., 2005; Zhu et al., 2004), locomotor behavior, (Bowling et al., 1993; Del Arco and Mora, 2008; Del Arco et al., 2007; Hoffmann et al., 2009; Neugebauer et al., 2004; Van Waas and Soffie, 1996) and learning (reviewed in (Van Waas and Soffie, 1996)), to sensitivities to the effects of drugs (Bardo et al., 2001; Howes et al., 2000; McCool and Chappell, 2009; Solinas et al., 2008). Generally, environmental enrichment is described as “a combination of complex inanimate and social stimulation” (Rosenzweig et al., 1978)

© 2010 Elsevier Inc. All rights reserved.

Corresponding Author: Linda J. Porrino, Ph.D. Dept. Physiology & Pharmacology Wake Forest University School Medicine Medical Center Blvd. Winston-Salem, NC 27157 USA lporrino@wfubmc.edu Phone: 336-716-8506 Fax: 336-716-8501.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

In the case of rodents, enrichment involves rearing animals with peers in large cages with a variety of novel objects and handling the animals daily. Animals reared in an impoverished environment are housed in solitary conditions without regular handling or novel stimuli. It is clear that rearing conditions, either impoverished/isolated or enriched/socially housed, differentially modify aspects of neuronal function and associated behavior. These modifications have implications for an unlimited number of conditions, including learning and memory, behavior, psychological disorders, and drug abuse.

Enriched mice and rats display improved learning and memory skills in a variety of tests including the T-maze and water maze (reviewed in (van Praag et al., 2000)). These animals also show decreased locomotor levels (Bowling et al., 1993; Del Arco and Mora, 2008; Del Arco et al., 2007; Hoffmann et al., 2009; Neugebauer et al., 2004; Van Waas and Soffie, 1996) which may be explained by the antidepressant effects of enrichment (Brenes et al., 2008), or a decreased response to novelty. As with decreased locomotor behavior, enriched rats that are prenatally exposed to cocaine spend less time following their partners in a social interaction test, while isolated, prenatally cocaine-exposed rats exhibit more play solicitations (Neugebauer et al., 2004). Behavioral differences may be attributed to known alterations in synaptic plasticity and dopaminergic markers associated with rearing rats in enriched or impoverished environments.

Dopamine is heavily implicated in drug abuse and addiction, thus, differences in sensitivity to the effects of drugs and alcohol and associated neural and behavioral changes have been one major focus of environmental enrichment studies. Isolation/impoverishment has been shown to lead to higher levels of self administration of ethanol (McCool and Chappell, 2009), amphetamine (Bardo et al., 2001), and cocaine (Howes et al., 2000), while enrichment has been shown to decrease self-administration of amphetamine (Bardo et al., 2001; Green et al., 2002) and reduce the effects of other drugs of abuse. These effects may be associated with a reduction in cell surface levels of dopamine transporter (Zhu et al., 2005) and reduced DA uptake (Zhu et al., 2004) in the mPFC of rats reared in enriched environments compared to those reared in impoverished environments. Additionally, it has been shown that non-contingent cocaine administration raises levels of tyrosine hydroxylase, the rate-limiting enzyme in the synthesis of dopamine, in isolated/impoverished rats, but not enriched rats, indicating that the impoverished rats may have heightened reward sensitivity and vulnerability to drug abuse (Zakharova et al., 2009).

One approach to the analysis of the consequences of environmental enrichment is the use of metabolic mapping with the quantitative autoradiographic 2- $[^{14}\text{C}]$ deoxyglucose method (2-DG) for the measurement of rates of local cerebral glucose utilization (Sokoloff et al., 1977). Because of the close coupling between energy metabolism and functional activity in the central nervous system, this method provides a means to identify those brain regions in which functional activity is altered by pharmacological or physiological stimuli or other manipulations (Porrino et al., 1984). Unlike electrophysiology or the measurement of changes in specific protein or mRNA levels, the 2DG method can evaluate the effects throughout the entire central nervous system, thus providing a broad visualization of the distribution of alterations in functional activity at all brain levels. Therefore, the purpose of the present study was to map differences in the distribution of rates of cerebral metabolic activity in rats reared in enriched or impoverished environments, in order to determine whether social and environmental housing conditions during development alter patterns of brain activation in adulthood.

Methods

Animals

All animal procedures were performed in accordance with protocols approved by Wake Forest University School of Medicine Animal Care and Use Committee and were consistent with the NIH *Guide for the Care and Use of Laboratory Animals*. Male Sprague-Dawley rats (n = 22; 28 days old) were obtained from Harlan Industries (Indianapolis, IN). All animals were maintained in a temperature- and humidity-controlled vivarium with a 12-h light-dark cycle (lights on at 07:00). Food and water were available ad libitum.

Environmental Conditions

Rats were assigned randomly to either an enriched environment group or an impoverished environment group. Enriched rats (n = 10) were housed in standard polycarbonate cages (25.9 × 47.6 × 20.9 cm) in groups of two. The enriched environment contained 4 novel objects that were rotated weekly and the rats were handled daily, for at least 5 minutes. The impoverished rats (n = 12) were singly housed in standard polycarbonate cages (25.9 × 47.6 × 20.9 cm). Impoverished rats had no objects in their cages and were handled only briefly during cage changes. Rats were housed in these conditions for a period of 30 days at which time they were tested for locomotor activity followed by surgery and cerebral glucose utilization measurement.

Locomotor Activity

At the end of the enrichment period animals were tested for locomotor activity. Locomotor activity was measured in open-field Plexiglas® test chambers (42 × 42 × 30 cm) by electronic counters that detected interruptions of 8 independent infrared photocell beams (Omnitech, Columbus, OH). Photocell counts were recorded and stored at 10-min intervals for a total of 30 minutes per session. The following measures were calculated: horizontal activity, or the number of total beam interruptions; forward locomotor, or ambulatory activity; vertical activity, or rearing and stereotypy.

Surgical Procedures

Surgical procedure for the 2[¹⁴C]DG studies followed that described by (Crane and Porrino, 1989). Briefly, rats were lightly anesthetized with a mixture of halothane, nitrous oxide and oxygen. Polyethylene catheters were then inserted into the femoral vein and artery. Catheters ran subcutaneously and exited at the nape of the neck. A minimum of 36 h was allowed prior to initiation of the 2[¹⁴C]DG procedure to ensure full recovery from anesthesia.

2[¹⁴C] Deoxyglucose method

Local cerebral glucose utilization was measured in the home cages of the animals according to the method of Sokoloff et al. (1977). Rats were initially given an intravenous pulse of 2-[¹⁴C]deoxyglucose (DG) (specific activity 50–55 mCi/mmol; New England Nuclear, Boston, MA) at a dose of 75 µCi/kg. Timed arterial blood samples were then drawn over the next 45 min. Samples were centrifuged and plasma concentrations were measured for 2[¹⁴C]DG and glucose. At the end of the 45-min sampling procedure, animals were sacrificed using 50mg sodium pentobarbital. Brains were rapidly removed and frozen in isopentane at –45°C, then stored at –80°C until sectioning. Coronal sections (20 µm) were cut in a cryostat maintained at –22°C. Sections were picked up on glass coverslips, dried on a hotplate (60°C), and then exposed to film (Kodak MRM film, Kodak, Rochester, New York) along with calibrated [¹⁴C] standards (Amersham, Arlington Heights, IL) for 12–15 days. Autoradiograms were then analyzed by quantitative densitometry with a computerized

image processing system (MCID Imaging Research, Cambridge, England). Optical density measurements for each brain structure were made in a minimum of five sections. Local cerebral glucose utilization was calculated using the operational equation defined by Sokoloff et al. (1977). Rates of glucose utilization were determined in 38 brain structures selected on the basis of previous reports (Whitlow et al., 2002); according to the rat brain atlas (Paxinos and Watson, 1997).

Statistical Analysis

Behavioral data for the treatment groups were analyzed by means of Student's t-test for independent samples. Rates of glucose utilization were analyzed in specific brain region groups (mesocorticolimbic, nigrostriatal etc.) using Condition (impoverished vs. enriched) $\times$ Region ANOVAs with the specific brain areas within the group as a repeated measures variable and treatment as the between subject variable. The Greenhouse-Geisser correction was used if assumptions of sphericity were not met. ANOVAs were conducted for the following brain region groups: Mesocorticolimbic (11 areas), Nigrostriatal and related areas (8 areas), Other limbic regions (8 areas), Sensory/motor cortex and thalamus (8 areas), Other areas (2 areas). Where appropriate, post-hoc t-tests were performed on specific regions between condition groups with alpha adjusted for number of tests performed. Data are presented as mean $\pm$ SEM μ mol/100 g/min.

Results

Locomotor Activity

The effects of rearing in impoverished and enriched environments on locomotor behavior are shown in Figure 1. Animals housed in groups of two in an enriched environment produced no significant difference in total distance, horizontal activity or stereotypy when compared to animals in an impoverished environment.

Cerebral Glucose Utilization

The rates of local cerebral glucose utilization in enriched and impoverished animals are displayed in Table 1. Cerebral metabolic rates were globally higher in environmentally enriched animals compared to impoverished animals. This pattern was observed in 33 out of the 37 individual brain regions that were measured. Analysis of brain region groups revealed a main effect of region in the mesocorticolimbic group ($F(3.742, 67.36) = 135.5, p < 0.001$), the nigrostriatal group ($F(3.227, 58.09) = 231.5, p < 0.001$), the limbic group ($F(3.540, 60.18) = 229.4, p < 0.001$), the sensory/motor group ($F(3.983, 67.72) = 170.2, p < 0.001$) and the "other" group ($F(1, 16) = 29.19, p < 0.001$). Subsequently there was a significant interaction of Condition $\times$ Region in the mesocorticolimbic group ($F(3.742, 67.36) = 2.663, p = 0.043$). Based on evident differences between 2-DG means in the condition groups, two post-hoc t-tests were performed on the nucleus accumbens core and shell. Both the nucleus accumbens core ($p = 0.01$) and shell ($p = 0.02$) showed a significant difference between impoverished and enriched conditions. Significant interactions were not found in any other brain regions.

Discussion

The results of the present study demonstrate that pair housing and environmental enrichment produce alterations in functional activity, as measured by higher rates of glucose consumption, in the nucleus accumbens core and shell. In addition, while not significant, this pattern was observed in the majority of discrete brain regions that were measured.

The nucleus accumbens is a brain region that is heavily innervated by dopamine neurons and highly implicated in reward processing and the reinforcing properties of drugs of abuse. Our finding that impoverishment and enrichment differentially affect glucose utilization is consistent with previous reports of alterations in this region and dopaminergic differences due to enrichment or impoverishment. Greater functional activity may reflect increases in dendritic spine density in the nucleus accumbens that have been reported with environmental enrichment (Comery et al., 1996; Kolb et al., 2003). There is also evidence that basal levels of dopamine are reduced in the nucleus accumbens of mice reared in isolation (Eells et al., 2006).

Alterations in glucose utilization in the nucleus accumbens may be associated with the protective effects of enrichment against drug abuse and addiction (for review see (Stairs and Bardo, 2009)). Isolation/impoverishment leads rodents to self-administer greater amounts of ethanol (Deehan et al., 2007; McCool and Chappell, 2009), while enrichment has been shown to reduce the motivational properties of ethanol (de Carvalho et al., 2010). Rodents that have been reared in an enriched environment are less likely to self-administer low doses of amphetamine (Bardo et al., 2001; Green et al., 2002) or cocaine (Howes et al., 2000). While they are more sensitive to the acute locomotor stimulating effects of amphetamine, enriched rats are resistant to sensitization after repeated injections (Bowling and Bardo, 1994; Bowling et al., 1993). Amphetamine has also been shown to increase levels of glutamate in the nucleus accumbens to a much greater extent in enriched animals than in impoverished animals (Rahman and Bardo, 2008). In addition to reducing reward sensitivity in drug-naïve animals, environmental enrichment decreases the negative effects of prenatal and juvenile cocaine exposure (Neugebauer et al., 2004) and prevents the reinstatement of a previously established cocaine-induced conditioned place preference (Solinas et al., 2008). These results are supported by data that show that rats reared in isolation show greater reward sensitivity, even to sucrose, than socially housed, enriched rats (Brenes et al., 2008).

The 2-deoxyglucose method is most often used to measure changes in task related functional activity as reflected by changes in rates of local cerebral metabolism. However, it can also be used to assess baseline differences in brain glucose metabolism associated with different states or phenotypes. Task-related changes generally occur over a relatively short time frame usually in the range of seconds and minutes, whereas basal rates of glucose utilization generally have longer time lines (minutes, hours, days and weeks) and reflect intrinsic ongoing brain activity. There are numerous processes in the brain that require energy including the restoration and maintenance of chemical gradients, protein synthesis and trafficking, transcription, neurotransmitter synthesis and processing, intracellular signaling, axonal transport, etc. (Ames, 2000). The differences in rates of glucose utilization observed here between rats raised in impoverished and enriched conditions undoubtedly are differences in intrinsic brain activity, and are not directly related to a task or to a response to external stimuli. More likely, the differences reflect the accumulation of an animal's metabolic history and may be amplified in its interactions with the external world.

The data presented here show no differences in locomotor activity between the impoverished and enriched animals. This is in contrast to previous studies which suggest that environmentally enriched animals have decreased locomotor activity compared to animals housed in an impoverished environment (Brenes et al., 2008, 2009; Elliott and Grunberg, 2005; Green et al., 2003; Smith et al., 2009; Van Waas and Soffie, 1996; Zaias et al., 2008). The discrepancy between these results may be due to differences in the number of animals in socially housed groups. For each of the previously mentioned studies, enriched animals were housed in groups of three to twelve per cage, whereas animals in the current study were housed only in pairs. Therefore, the lack of locomotor differences observed here may be due to more limited social interaction in our enriched groups. If this paradigm had

been extended with additional social stimuli, it is plausible that overt behavioral changes would have appeared..

Another possible explanation for the discrepancy between this and other reports is the age at testing. Other evidence suggests that the benefits of enrichment and social housing are more robust in adolescent animals (Zakharova et al., 2009) although this is not a consistent finding (Bowling and Bardo, 1994). Since the animals in the present study were adult at the time of testing, this may also have contributed to the lack of locomotor differences between the enriched and impoverished groups. Despite the lack of behavioral differences, there were apparent differences in brain function in reward-related areas. Thus, these data suggest that brain changes due to enrichment occur before there are any overt behavioral changes. Furthermore, it is apparent that even small changes in environment are sufficient to produce alterations in brain function, even in the absence of significant changes in behavior.

In addition to the significant difference in glucose metabolism observed in the nucleus accumbens, there was also a trend for higher glucose utilization in the cingulate cortex, dorsolateral caudate, basolateral amygdala, and the CA3 region of the hippocampus of enriched animals. These trends are consistent with literature on other changes associated with enrichment. For example enrichment has been shown to benefit certain neurological disorders such as Huntington's disease (van Dellen et al., 2000), which is known to degenerate the caudate, an area associated with cognition and motor function (Douaud et al., 2006; Hedreen and Folstein, 1995). Environmental enrichment also decreases stress related neuronal activity in the amygdala (Abraham and Kovacs, 2000). Lastly, environmental enrichment has been shown to improve hippocampal dependent memory (Duffy et al., 2001), and lead to increases in cerebral glucose utilization in the nucleus accumbens of Tryon rats, a strain bred for their ability to learn. (Gonzalez-Lima et al., 1994). While our measures of metabolism in these regions were not statistically significant they are measurements of basal levels of metabolism and are therefore resting state. On this note, the trends we observed in these regions may have been significantly different if metabolism had been measured during a task.

In summary, the present findings confirm that environmental enrichment affects glucose metabolism in the nucleus accumbens, a brain region known to be an integral part of drug addiction. In addition, the paradigm of enrichment used in this study was not sufficient to produce locomotor effects in these animals. These results may reflect mechanisms underlying the reported behavioral benefits of environmental enrichment.

Acknowledgments

The authors would like to thank Brooke Legg, Donald Floyd and Mack Miller for their contributions. This project was funded by the Center for Neurobiological Investigation of Drug Abuse grant DA06634 from the National Institute of Drug Abuse. AKL was supported by RJR-Leon Golberg fellowship in Pharmacology and Toxicology.

References

- Abraham IM, Kovacs KJ. Postnatal handling alters the activation of stress-related neuronal circuitries. *Eur J Neurosci.* 2000; 12:3003–3014. [PubMed: 10971641]
- Bardo MT, Klebaur JE, Valone JM, Deaton C. Environmental enrichment decreases intravenous self-administration of amphetamine in female and male rats. *Psychopharmacology (Berl).* 2001; 155:278–284. [PubMed: 11432690]
- Bowling SL, Bardo MT. Locomotor and rewarding effects of amphetamine in enriched, social, and isolate reared rats. *Pharmacol Biochem Behav.* 1994; 48:459–464. [PubMed: 8090815]

- Bowling SL, Rowlett JK, Bardo MT. The effect of environmental enrichment on amphetamine-stimulated locomotor activity, dopamine synthesis and dopamine release. *Neuropharmacology*. 1993; 32:885–893. [PubMed: 8232791]
- Brenes JC, Padilla M, Fornaguera J. A detailed analysis of open-field habituation and behavioral and neurochemical antidepressant-like effects in postweaning enriched rats. *Behav Brain Res*. 2008
- Brenes JC, Padilla M, Fornaguera J. A detailed analysis of open-field habituation and behavioral and neurochemical antidepressant-like effects in postweaning enriched rats. *Behav Brain Res*. 2009; 197:125–137. [PubMed: 18786573]
- Comery TA, Stamoudis CX, Irwin SA, Greenough WT. Increased density of multiple-head dendritic spines on medium-sized spiny neurons of the striatum in rats reared in a complex environment. *Neurobiol Learn Mem*. 1996; 66:93–96. [PubMed: 8946401]
- Crane AM, Porrino LJ. Adaptation of the quantitative 2-[14C]deoxyglucose method for use in freely moving rats. *Brain Res*. 1989; 499:87–92. [PubMed: 2804673]
- de Carvalho CR, Pandolfo P, Pamplona FA, Takahashi RN. Environmental enrichment reduces the impact of novelty and motivational properties of ethanol in spontaneously hypertensive rats. *Behav Brain Res*. 2010; 208:231–236. [PubMed: 19962407]
- Deehan GA Jr, Cain ME, Kiefer SW. Differential rearing conditions alter operant responding for ethanol in outbred rats. *Alcohol Clin Exp Res*. 2007; 31:1692–1698. [PubMed: 17651466]
- Del Arco A, Mora F. Prefrontal cortex-nucleus accumbens interaction: in vivo modulation by dopamine and glutamate in the prefrontal cortex. *Pharmacol Biochem Behav*. 2008; 90:226–235. [PubMed: 18508116]
- Del Arco A, Segovia G, Garrido P, de Blas M, Mora F. Stress, prefrontal cortex and environmental enrichment: studies on dopamine and acetylcholine release and working memory performance in rats. *Behav Brain Res*. 2007; 176:267–273. [PubMed: 17097747]
- Douaud G, Gaura V, Ribeiro MJ, Lethimonnier F, Maroy R, Verny C, Krystkowiak P, Damier P, Bachoud-Levi AC, Hantraye P, Remy P. Distribution of grey matter atrophy in Huntington's disease patients: a combined ROI-based and voxel-based morphometric study. *Neuroimage*. 2006; 32:1562–1575. [PubMed: 16875847]
- Duffy SN, Craddock KJ, Abel T, Nguyen PV. Environmental enrichment modifies the PKA-dependence of hippocampal LTP and improves hippocampus-dependent memory. *Learn Mem*. 2001; 8:26–34. [PubMed: 11160761]
- Eells JB, Misler JA, Nikodem VM. Early postnatal isolation reduces dopamine levels, elevates dopamine turnover and specifically disrupts prepulse inhibition in *Nurr1*-null heterozygous mice. *Neuroscience*. 2006; 140:1117–1126. [PubMed: 16690213]
- Elliott BM, Grunberg NE. Effects of social and physical enrichment on open field activity differ in male and female Sprague-Dawley rats. *Behav Brain Res*. 2005; 165:187–196. [PubMed: 16112757]
- Gonzalez-Lima F, Ferchmin PA, Eterovic VA, Gonzalez-Lima EM. Metabolic activation of the brain of young rats after exposure to environmental complexity. *Dev Psychobiol*. 1994; 27:343–351. [PubMed: 8001724]
- Green TA, Cain ME, Thompson M, Bardo MT. Environmental enrichment decreases nicotine-induced hyperactivity in rats. *Psychopharmacology (Berl)*. 2003; 170:235–241. [PubMed: 12845407]
- Green TA, Gehrke BJ, Bardo MT. Environmental enrichment decreases intravenous amphetamine self-administration in rats: dose-response functions for fixed- and progressive-ratio schedules. *Psychopharmacology (Berl)*. 2002; 162:373–378. [PubMed: 12172690]
- Hedreen JC, Folstein SE. Early loss of neostriatal striosome neurons in Huntington's disease. *J Neuropathol Exp Neurol*. 1995; 54:105–120. [PubMed: 7815073]
- Hoffmann LC, Schutte SR, Koch M, Schwabe K. Effect of "enriched environment" during development on adult rat behavior and response to the dopamine receptor agonist apomorphine. *Neuroscience*. 2009; 158:1589–1598. [PubMed: 19110038]
- Howes SR, Dalley JW, Morrison CH, Robbins TW, Everitt BJ. Leftward shift in the acquisition of cocaine self-administration in isolation-reared rats: relationship to extracellular levels of dopamine, serotonin and glutamate in the nucleus accumbens and amygdala-striatal FOS expression. *Psychopharmacology (Berl)*. 2000; 151:55–63. [PubMed: 10958117]

- Kolb B, Gorny G, Soderpalm AH, Robinson TE. Environmental complexity has different effects on the structure of neurons in the prefrontal cortex versus the parietal cortex or nucleus accumbens. *Synapse*. 2003; 48:149–153. [PubMed: 12645040]
- McCool BA, Chappell AM. Early social isolation in male Long-Evans rats alters both appetitive and consummatory behaviors expressed during operant ethanol self-administration. *Alcohol Clin Exp Res*. 2009; 33:273–282. [PubMed: 19032581]
- Neugebauer NM, Cunningham ST, Zhu J, Bryant RI, Middleton LS, Dwoskin LP. Effects of environmental enrichment on behavior and dopamine transporter function in medial prefrontal cortex in adult rats prenatally treated with cocaine. *Brain Res Dev Brain Res*. 2004; 153:213–223.
- Paxinos, G.; Watson, C. The rat brain in stereotaxic coordinates. Academic Press; London: 1997.
- Porrino LJ, Lucignani G, Dow-Edwards D, Sokoloff L. Correlation of dose-dependent effects of acute amphetamine administration on behavior and local cerebral metabolism in rats. *Brain Res*. 1984; 307:311–320. [PubMed: 6540614]
- Rahman S, Bardo MT. Environmental enrichment increases amphetamine-induced glutamate neurotransmission in the nucleus accumbens: a neurochemical study. *Brain Res*. 2008; 1197:40–46. [PubMed: 18242591]
- Rosenzweig MR, Bennett EL, Hebert M, Morimoto H. Social grouping cannot account for cerebral effects of enriched environments. *Brain Res*. 1978; 153:563–576. [PubMed: 698794]
- Smith MA, Iordanou JC, Cohen MB, Cole KT, Gergans SR, Lyle MA, Schmidt KT. Effects of environmental enrichment on sensitivity to cocaine in female rats: importance of control rates of behavior. *Behav Pharmacol*. 2009; 20:312–321. [PubMed: 19584714]
- Sokoloff L, Reivich M, Kennedy C, Des, Rosiers MH, Patlak CS, Pettigrew KD, Sakurada O, Shinohara M. The [14C]deoxyglucose method for the measurement of local cerebral glucose utilization: theory, procedure, and normal values in the conscious and anesthetized albino rat. *J Neurochem*. 1977; 28:897–916. [PubMed: 864466]
- Solinas M, Chauvet C, Thiriet N, El Rawas R, Jaber M. Reversal of cocaine addiction by environmental enrichment. *Proc Natl Acad Sci U S A*. 2008; 105:17145–17150. [PubMed: 18955698]
- Stairs DJ, Bardo MT. Neurobehavioral effects of environmental enrichment and drug abuse vulnerability. *Pharmacol Biochem Behav*. 2009; 92:377–382. [PubMed: 19463254]
- van Dellen A, Blakemore C, Deacon R, York D, Hannan AJ. Delaying the onset of Huntingtons's in mice. *Nature*. 2000; 404:721–722. [PubMed: 10783874]
- van Praag H, Kempermann G, Gage FH. Neural consequences of environmental enrichment. *Nat Rev Neurosci*. 2000; 1:191–198. [PubMed: 11257907]
- Van Waas M, Soffie M. Differential environmental modulations on locomotor activity, exploration and spatial behaviour in young and old rats. *Physiol Behav*. 1996; 59:265–271. [PubMed: 8838604]
- Whitlow CT, Freedland CS, Porrino LJ. Metabolic mapping of the time-dependent effects of delta 9-tetrahydrocannabinol administration in the rat. *Psychopharmacology (Berl)*. 2002; 161:129–136. [PubMed: 11981592]
- Zaias J, Queeney TJ, Kelley JB, Zakharova ES, Izenwasser S. Social and physical environmental enrichment differentially affect growth and activity of preadolescent and adolescent male rats. *J Am Assoc Lab Anim Sci*. 2008; 47:30–34. [PubMed: 18351719]
- Zakharova E, Miller J, Unterwald E, Wade D, Izenwasser S. Social and physical environment alter cocaine conditioned place preference and dopaminergic markers in adolescent male rats. *Neuroscience*. 2009; 163:890–897. [PubMed: 19580849]
- Zhu J, Apparsundaram S, Bardo MT, Dwoskin LP. Environmental enrichment decreases cell surface expression of the dopamine transporter in rat medial prefrontal cortex. *J Neurochem*. 2005; 93:1434–1443. [PubMed: 15935059]
- Zhu J, Green T, Bardo MT, Dwoskin LP. Environmental enrichment enhances sensitization to GBR 12935-induced activity and decreases dopamine transporter function in the medial prefrontal cortex. *Behav Brain Res*. 2004; 148:107–117. [PubMed: 14684252]

Research Highlights

- Environmental enrichment and social housing during adolescence produced increased functional brain activity in adulthood when compared to rats reared in an impoverished environment.
- Differences in functional activity, as reflected by rates of glucose utilization, were greatest in the nucleus accumbens shell and core with rates higher in those rats reared in an enriched environment.
- These differences in brain activity occurred in the absence of differences in spontaneous locomotor behavior.

Fig 1. Effect of environment on spontaneous locomotor activity in rats. Data shown are means $\pm$ S.E.M. for cumulative distance or photocell beam breaks during a 30min test period. Data were analyzed by Student's t-tests.

TABLE 1

Local glucose utilization in the brains of adult male rats in impoverished or enriched environment

Region	Impove rished (n=12)	En riched (n= 10)	% Diff
Mesocorticolimbic system			
Cingulate cortex, area 3	91.1 ± 3.2	95.5 ± 3.9	4. 9%
Infralimbic cortex	71.3 ± 2.6	73.9 ± 2.7	3. 7%
Agranular insular cortex	82.1 ± 2.7	88.2 ± 3.8	7. 4%
Anterior cingulate cortex	99.9 ± 3.4	109.3 ± 3.6	9. 4%
Anterior nucleus accumbens	86.2 ± 2.7	90.0 ± 3.3	4. 4%
Nucleus accumbens, core	75.8 ± 2.4	84.3 ± 1.9*	11 .1%
Nucleus accumbens, shell	73.1 ± 3.2	85.4 ± 3.6*	16 .9%
Basolateral amygdala	77.9 ± 3.0	86.2 ± 3.5	10 .7%
Central nucleus, amygdala	41.1 ± 1.4	43.1 ± 1.3	4. 7%
Median forebrain bundle	63.2 ± 1.5	65.4 ± 1.6	3. 6%
Ventral tegmental area	74.8 ± 3.8	71.5 ± 2.9	-4. 5%
Nigrostriatal system and related areas			
Dorsolateral caudate	108.5 ± 4.1	120.2 ± 4.9	10 .7%
Dorsomedial caudate	92.9 ± 4.1	100.6 ± 3.9	8. 3%
Ventral caudate	97.6 ± 3.3	103.4 ± 3.2	6. 0%
Globus pallidus	61.1 ± 2.8	63.3 ± 2.9	3. 6%
Medial habenula	70.8 ± 3.1	71.8 ± 2.6	1. 5%
Lateral habenula	96.7 ± 3.5	104.5 ± 3.6	8. 1%
Substantia nigra, pars compacta	72.0± 2.6	74.1 ± 1.7	3. 0%
Substantia nigra, pars reticulata	58.7 ± 2.8	63.0 ± 2.7	8. 3%
Other limbic areas			
Medial thalamus	107.2 ± 4.5	118.5 ± 5.0	10 .5%
Lateral thalamus	92.8 ± 3.0	96.7 ± 3.8	4. 2%

Region	Impove rished (n=12)	En riched (n= 10)	% Diff
Hippocampus CA1	64.3 ± 2.7	69.8 ± 2.6	8. 4%
Hippocampus CA3	69.7 ± 3.0	76.7 ± 2.4	10 .0%
Hippocampus Dentate gyrus	54.5 ± 2.2	58.4 ± 2.3	7. 2%
Ventromedial hypothalamus	50.0 ± 2.0	49.7 ± 2.0	−0. 7%
Dorsal raphe	72.6 ± 3.3	72.9 ± 5.0	0. 3%
Median raphe	96.9 ± 3.8	95.6 ± 4.3	−1. 4%
Sensory/motor cortex and thalamus			
Motor cortex	96.5 ± 3.6	105.4 ± 4.0	9. 2%
Auditory cortex	127.7 ± 4.2	133.1 ± 4.1	4. 2%
Visual cortex	97.8 ± 3.3	106.7 ± 4.6	9. 1%
Corpus collosum	34.1 ± 1.8	37.3 ± 1.9	9. 4%
Medial geniculate	107.1 ± 3.8	109.1 ± 4.5	1. 8%
Lateral geniculate	87.4 ± 3.2	91.6 ± 3.2	4. 8%
Superior colliculus	89.6 ± 3.8	98.6 ± 3.1	10 .1%
Inferior colliculus	114.3 ± 5.3	128.0 ± 6.7	12 .0%
Other areas			
Locus Coeruleus	64.5 ± 3.0	62.9 ± 2.3	−2. 6%
Cerebellum	73.4 ± 3.5	76.1 ± 4.1	3. 7%

Values are means ± SEM of the number of animals in parentheses, expressed as $\mu\text{mol}/100\text{g}/\text{min}$.

*
P < 0.05 Difference between impoverished and enriched animals, Post-hoc Student's t test.