Published in final edited form as:

Neurosci Biobehav Rev. 2011 November ; 35(10): 2058–2063. doi:10.1016/j.neubiorev.2010.11.010.

Functional connectivity and cholinergic modulation in auditory cortex

Raju Metherate*

Department of Neurobiology and Behavior, Center for Hearing Research, University of California, Irvine

Abstract

Although it is known that primary auditory cortex (A1) contributes to the processing and perception of sound, its precise functions and the underlying mechanisms are not well understood. Recent studies point to a remarkably broad spectral range of largely subthreshold inputs to individual neurons in A1—seemingly encompassing, in some cases, the entire audible spectrum—as evidence for potential, and potentially unique, cortical functions. We have proposed a general mechanism for spectral integration by which information converges on neurons in A1 via a combination of thalamocortical pathways and intracortical long-distance, “horizontal,” pathways. Here, this proposal is briefly reviewed and updated with results from multiple laboratories. Since spectral integration in A1 is dynamically regulated, we also show how one regulatory mechanism—modulation by the neurotransmitter acetylcholine (ACh)—could act within the hypothesized framework to alter integration in single neurons. The results of these studies promote a cellular understanding of information processing in A1.

Keywords for indexing

thalamocortical; intracortical; acetylcholine; nicotinic; muscarinic; frequency receptive field; spectral integration; intracellular; synapse

1. Introduction

Spectral integration is an important function of A1, as indicated by physiological responses of neurons to spectrally complex stimuli (Mendelson and Cynader, 1985; Phillips et al., 1985; Rauschecker and Tian, 2000; Tian and Rauschecker, 2004; Whitfield and Evans, 1965) and behavioral deficits following cortical lesions in animals and humans (Harrington et al., 2001; Kelly and Whitfield, 1971; Ohl et al., 1999; Phillips, 1998; Phillips and Farmer, 1990; Wetzel et al., 1998). Spectral integration by single neurons is represented most simply by frequency receptive fields, which typically are constructed using pure tone stimuli and extracellular recordings of action potentials. Receptive fields thus delineated are “classical,” suprathreshold (derived from action potentials) receptive fields that do not reflect the presence of subthreshold inputs. Classical frequency receptive fields are of similar breadth throughout the lemniscal (primary) auditory system (Calford et al., 1983), a finding that seems to suggest A1 does not contribute unique functions to spectral processing. However,

*Address for correspondence: Raju Metherate, Ph.D., University of California, Irvine, 2205 McGaugh Hall, Irvine, CA 92697-4550, (949) 824-6141, fax: (949) 824-2447, raju.metherate@uci.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

when subthreshold (also known as subliminal, surround, nonclassical) receptive fields are considered, spectral integration in A1 neurons may be considerably more extensive than in subcortical neurons (see next section). Although the functions of subthreshold inputs are not yet fully appreciated, we have proposed a simple framework of functional connectivity by which spectral integration, including the contribution of subthreshold inputs, may take place in A1 (Kaur et al., 2004; Kaur et al., 2005; Metherate et al., 2005). This framework is intended to promote an understanding of the mechanisms and functions of spectral integration in A1, and is reviewed and updated below.

Spectral integration in A1 is dynamic, varying, for example, with behavioral state (e.g., sleep vs. waking) (Edeline et al., 2001) or with attention to spectral cues in a behavioral task (Fritz et al., 2003; Weinberger, 2004). Rapid changes in spectral processing are likely due to physiological mechanisms such as neuromodulation (Edeline, 2003), which also may contribute to lasting changes in frequency representations subsequent to behavioral training (Kilgard and Merzenich, 1998; Recanzone et al., 1993). Thus, spectral integration is dynamically regulated, at times rapidly and perhaps continuously. In the second part of this review I will note some mechanisms by which one neuromodulatory agent, the neurotransmitter ACh, can regulate the proposed framework for spectral integration in A1.

2. Breadth of subthreshold inputs to neurons in A1

Studies of classical frequency receptive fields indicate that there is little change in receptive field breadth as one ascends through the lemniscal auditory pathway (e.g., from the central nucleus of the inferior colliculus to the ventral division of the medial geniculate body, MGv, to A1) (Calford et al., 1983), suggesting no change in spectral integration. However, measures of classical receptive fields are based on single- or multi-unit activity (i.e., tone-evoked action potential discharge) and do not take into account the presence or absence of subthreshold responses to stimuli outside the frequency receptive field. It is now well known that classical receptive fields underestimate the breadth of synaptic inputs to cortical neurons. For instance, blockade of GABAergic inhibition results in an expansion of the frequency receptive field (Foeller et al., 2001; Muller and Scheich, 1988; Wang et al., 2000; Wang et al., 2002), indicating the presence of normally-subthreshold excitatory postsynaptic potentials (EPSPs) (but see Kurt et al., 2006). More direct evidence is provided by *in vivo* intracellular recordings, which reveal extensive subthreshold receptive fields (Kaur et al., 2004; Ojima and Murakami, 2002; Tan et al., 2004; Wehr and Zador, 2003). These studies also demonstrate that throughout subthreshold and suprathreshold receptive fields, synaptic responses reflect overlapping (“balanced”) inputs from both excitatory and inhibitory neurons that are driven by afferent input. Disruption by GABA receptor blockade of the excitatory-inhibitory balance leads to expansion of classical receptive fields (above). Intracellular studies are complemented by other studies that measure extracellular tone-evoked local field potentials (LFPs, which reflect synchronous synaptic potentials in a local group of neurons), in some cases simultaneously with multiunit spike recordings. These studies show LFP receptive fields that are similar in breadth to intracellular synaptic receptive fields, and in some cases several octaves broader than classical receptive fields at the same recording site (Eggermont, 1996; Galvan et al., 2001; Kaur et al., 2004; Norena and Eggermont, 2002). At higher intensities, the bandwidths of LFP receptive fields often exceed the experimenters’ measurement capacity; e.g., a maximum bandwidth of eight octaves in one recent study (Happel et al., 2010). The breadth of intracellular and LFP receptive fields suggests that some neurons in A1 integrate over much, if not all, of the audible spectrum (Happel et al., 2010; Metherate et al., 2005; Schulze and Langner, 1999).

3. Contributions of thalamocortical and intracortical circuits to spectral integration in A1

To examine mechanisms of spectral integration, we focused on how information about characteristic frequency (CF) and spectrally-distant nonCF stimuli—that is, the “center” and one “edge” of a frequency receptive field—converges on single neurons in A1. (CF is the frequency eliciting the lowest threshold response; experimentally, we define nonCF as ~3 octaves from CF so as to maximize differences in underlying circuitry.) The relevant circuits—thalamocortical and long-range “horizontal” intracortical pathways—are illustrated schematically in Fig. 1, which also illustrates our hypothesis. The contribution of the thalamocortical pathway is relatively straightforward and involves relaying information mostly about CF and near-CF stimuli from the MGv. Physiological and anatomical studies show that thalamocortical projections link MGv and A1 neurons with similar CFs (Budinger et al., 2000; Imig and Morel, 1984; Winer et al., 1999). Thalamocortical arbors labeled by injections of tracer into small portions of MGv cover regions of A1 containing neurons with similar CFs (Velenovsky et al., 2003), and paired recordings of neurons in MGv and A1 show that cells with correlated discharge exhibit CFs within one-third of an octave (Miller et al., 2001). These data imply that thalamocortical axons project to restricted portions of A1 and do not diverge throughout A1. Thus, direct thalamocortical projections are not responsible for the broad spectral integration observed in physiological studies. Rather, it seems that thalamocortical inputs mediate cortical responses to CF and near-CF stimuli. [Note, however, that this scheme does not incorporate projections from auditory nonlemniscal thalamic regions outside of MGv. Nonlemniscal neurons often have broad receptive fields and project diffusely to superficial layers of temporal cortex; their poorly understood functions may include modulation of cortical excitability (e.g., see discussion in Weinberger, 2004), and also could contribute to breadth of tuning in A1].

According to the framework in Fig. 1, neurons in A1 respond to nonCF stimuli via intracortical horizontal pathways. This scheme is supported by recent findings from several laboratories. One approach has been to deliver into A1 the GABA-A receptor agonist muscimol to inhibit locally generated (cortical), but not afferent (thalamocortical) activity, and then determine the effect on cortical responses to CF and nonCF stimuli. In one study (Kaur et al., 2004), muscimol only partly reduced the initial (first ~10 ms) LFP response to CF stimuli, but fully suppressed longer-latency response components, consistent with inhibition of cortical neurons but not thalamocortical inputs. In contrast, muscimol in some cases fully suppressed all response to nonCF stimuli, suggesting the critical involvement of intracortical pathways. Although a more recent study has questioned the sole use of muscimol to silence cortical activity, and instead suggested simultaneous activation of GABA-A receptors and blockade of GABA-B receptors (Liu et al., 2007), a third study found no difference between the two approaches (Happel et al., 2010). The latter study confirmed that cortical silencing eliminates responses to nonCF, but not CF, stimuli (Happel et al., 2010). Thus, it appears that thalamocortical inputs generate (or, more correctly, initiate) responses to CF stimuli, whereas responses to nonCF stimuli involve intracortical projections from neurons with spectrally-distant CFs (Fig. 1).

Recent studies also show that thalamocortical inputs trigger a rapid and local intracortical amplification of the response to CF stimuli (Happel et al., 2010; Liu et al., 2007), which explains why cortical silencing reduces (partly) the response to CF stimuli. In addition, stimuli within ~1 octave (“near-CF” stimuli) trigger cortical responses that similarly rely on local intracortical amplification of inputs (Happel et al., 2010; Liu et al., 2007).

Although a complete dissociation between the pathways relaying information about CF and spectrally distant stimuli may be unlikely, one prediction of the model (Fig. 1) is that since

thalamocortical vs. intracortical projections terminate in different cortical layers, the neural activity elicited by CF vs. nonCF stimuli similarly will have different laminar profiles. Consistent with this prediction, current-source density (CSD) profiles elicited by CF and nonCF stimuli are different, with CF stimuli eliciting major current sinks in lower layer 3 and upper layer 4 (the location of thalamocortical terminals) and nonCF stimuli eliciting current sinks in multiple layers (consistent with intracortical projection patterns) (Kaur et al., 2005). Of course, the model in Fig. 1 is overly simplistic and does not incorporate different cell types, multiple thalamic and cortico-cortical projections, and a host of other features of the auditory forebrain. Rather, the model is intended to illustrate the general concept as a starting point for understanding connectivity in A1, and to provide a framework for understanding functional modulation and plasticity. The latter issue is reviewed next.

4. Cholinergic modulation of thalamocortical and intracortical connectivity

The final section of this review will touch upon dynamic regulation of processing in A1 via regulation of functional connectivity. Regulatory mechanisms are undoubtedly diverse, but most attention is focused on a subset of neurotransmitters that exert neuromodulatory actions (Edeline, 2003). This section will focus on ACh, a neurotransmitter that plays important roles in arousal, attention, and sensory learning (Hasselmo, 1999; Sarter et al., 2001; Weinberger, 2004). Cholinergic actions at both major subtypes of ACh receptor—nicotinic (nAChR) and muscarinic (mAChR) receptors—may regulate spectral integration in A1. In particular, cholinergic modulation of thalamocortical and intracortical transmission may underlie regulation of spectral processing as reflected in changes to frequency receptive fields.

Many studies of cholinergic modulation *in vivo* have shown that activation of mAChRs enhance cortical responses to sensory inputs (Chen and Yan, 2007; McKenna et al., 1989; Metherate et al., 1988; Sillito and Kemp, 1983; Zhang et al., 2005). A number of cellular effects, not all of which are directly excitatory, contribute to muscarinic modulation of sensory responses. Muscarinic stimulation can increase postsynaptic membrane resistance, and, thereby, responses to afferent inputs, due to decreased conductance of several K^+ channels (Halliwell and Adams, 1982; Krnjevic et al., 1971; Madison et al., 1987; McCormick and Prince, 1986). Accordingly, activation of cholinergic synapses in auditory cortex increases postsynaptic excitability via a slow EPSP associated with increased membrane resistance and decreased afterhyperpolarization potentials (AHPs) (Cox et al., 1994; Metherate et al., 1992). Stimulation of cortically-projecting neurons from the nucleus basalis enhances afferent responses in A1 evoked by thalamic (Metherate and Ashe, 1993; Metherate et al., 1992) or acoustic stimulation (Chen and Yan, 2007; Edeline et al., 1994; Zhang et al., 2005), in part by enhancing tone-evoked EPSPs and simultaneously reducing tone-evoked inhibition (Froemke et al., 2007; Metherate and Ashe, 1993; Metherate et al., 1992). The use of the auditory thalamocortical slice preparation (Cruikshank et al., 2002), which preserves the thalamocortical pathway from MGv to A1 along with (unspecified) long-range intracortical pathways, showed that the cholinergic agonist carbachol suppressed intracortical EPSPs while having lesser or no effects on thalamic-evoked EPSPs (Hsieh et al., 2000). Since all of the above effects are blocked by muscarinic receptor antagonists such as atropine, they demonstrate the functional consequences of muscarinic cellular actions for auditory processing in A1.

ACh also acts at nAChRs, which are widely distributed in the auditory system (Morley and Happe, 2000). Studies to date support two main functions of nAChRs in sensory cortex: presynaptic regulation of thalamocortical transmission and postsynaptic excitation of GABAergic interneurons (for review, see Metherate, 2004). Both functions relate to hypothesized mechanisms of spectral integration (Fig. 2). The distribution of nAChRs in

cortex varies among species, but generally relates to the widely-held hypothesis that nAChRs regulate thalamocortical transmission in general (Clarke, 2004). Studies in cat and rat show a dense concentration of nAChRs in layers 3–4 where thalamocortical inputs terminate, and also layers 1 and 5/6 (Clarke et al., 1984; Clarke et al., 1985; Lavine et al., 1997; London et al., 1985; Parkinson et al., 1988; Prusky et al., 1987; Sahin et al., 1992). In the mouse, nAChRs are less prominent in the middle layers (Marks et al., 1992; Rogers et al., 1998; Ross et al., 2000; Zoli et al., 1998)). In addition to cortical locations of nAChRs, in several species (mouse, rat, primate, human) nAChRs are found in the subcortical white matter region containing the auditory thalamocortical pathway, as evidenced by positron emission tomography (PET) and radioligand studies (Chattopadhyay et al., 2005; Ding et al., 2004; Easwaramoorthy et al., 2007). Thus, thalamocortical transmission in cat and rat may be regulated by nAChRs located at or near thalamocortical terminals; this mechanism is supported by functional studies (Clarke, 2004; Gil et al., 1997; Lambe et al., 2003). In rodents and primates, nAChRs associated with thalamocortical axons suggest an additional mechanism. Using the auditory thalamocortical slice, we recently showed that activation of nAChRs in the thalamocortical pathway increases axon excitability and regulates thalamocortical relay (Kawai et al., 2007), providing the first functional evidence that ACh—or any neurotransmitter—can act on the myelinated portion of thalamocortical axons. Figure 2 shows a hypothetical arrangement of axonal and presynaptic nAChRs that individually or together may regulate cortical responses to thalamic inputs.

Other studies have demonstrated rapid nicotinic excitation of postsynaptic neurons, and in sensory cortex, as in other cortical areas, such actions are most prominent in GABAergic interneurons (Alkondon et al., 2000; Christophe et al., 2002; Frazier et al., 1998; Frazier et al., 2003; Gullledge et al., 2007; Ji and Dani, 2000; Ji et al., 2001; Jones and Yakel, 1997; Porter et al., 1999). In cortical layer 1, which has the highest density among cortical layers of cholinergic axons and synapses (Mechawar et al., 2000; Mechawar et al., 2002), nicotinic agonists excite nearly all GABAergic interneurons (Christophe et al., 2002; Gullledge et al., 2007). Notably, activation of nAChRs in layer 1 produces IPSPs in layer 2/3 interneurons but not pyramidal cells, suggesting that nicotinic excitation of layer 1 disinhibits deeper layers. Some GABAergic interneurons in cortical layers 2–5 also can be excited by nicotinic agonists (Gullledge et al., 2007; Porter et al., 1999; Xiang et al., 1998). Interneurons excited by nAChR activation include low-threshold spiking, regular-spiking, late-spiking and irregular spiking neurons, but do not include fast-spiking neurons. Thus, nicotinic excitation of interneurons can shape processing of afferent information by inhibiting some neurons and disinhibiting others.

It is tempting to speculate on the effect of endogenous ACh on A1 function when it is acting simultaneously at multiple nicotinic and muscarinic receptors and receptor subtypes. Although several studies have examined the effects of ACh released endogenously or exogenously in A1, the effects observed have been largely attributed to mAChRs (Ashe et al., 1989; Chen and Yan, 2007; Froemke et al., 2007; McKenna et al., 1988; Metherate et al., 1990). It is hard to reconcile such largely-muscarinic effects of ACh with the high density of nAChRs in A1 and robust effects of selective nicotinic drugs (above) (Gioanni et al., 1999; Kawai et al., 2007; Lavine et al., 1997; Liang et al., 2006). Possible explanations for overlooked nicotinic effects include rapid desensitization of nAChRs or ineffective agonist concentrations at the relevant receptor locations (Metherate, 2004; Role and Berg, 1996). Nonetheless, the studies reviewed above using selective agonists and antagonists suggest that activation of mAChRs would tend to increase postsynaptic excitability while decreasing intracortical transmission via presynaptic receptors, whereas, in contrast, activation of nAChRs may enhance thalamocortical transmission. The role of nAChRs in regulating transmission along intracortical pathways is unknown, but may involve excitation of GABAergic interneurons. Thus, the combined muscarinic and nicotinic actions of ACh

could enhance thalamocortical transmission while suppressing intracortical excitatory transmission. Given the hypothesized contribution of thalamocortical and intracortical inputs to frequency receptive fields (Fig. 1), the integrated actions of ACh could produce (Fig. 2): suppression of responses to nonCF stimuli (mediated by intracortical horizontal inputs), facilitation of responses to CF and near-CF stimuli (mediated by thalamocortical inputs), and enhancement of responsiveness to remaining inputs (due to enhanced postsynaptic excitability). The net effect could be to reduce receptive field breadth, lower the threshold to CF stimuli, and enhance responses to stimuli within the “sharpened” receptive field (Fig. 2, right). Note how the increased receptive field selectivity differs from that produced by another neuromodulatory agent, norepinephrine, which can sharpen receptive fields via generally suppressive effects that are greater at nonCF than at CF (Edeline, 2003). Future studies will test the predictions in Fig. 2, but as these examples suggest, understanding the cellular actions of ACh can promote a deeper understanding of the mechanisms regulating auditory processing in A1.

Acknowledgments

The author’s research cited here was supported by the National Institutes of Health (NIDCD DC02967 and NIDA DA12929) and the National Science Foundation (IBN 9510904).

References

- Alkondon M, Pereira EF, Almeida LE, Randall WR, Albuquerque EX. Nicotine at concentrations found in cigarette smokers activates and desensitizes nicotinic acetylcholine receptors in CA1 interneurons of rat hippocampus. *Neuropharmacology*. 2000; 39:2726–2739. [PubMed: 11044743]
- Ashe JH, McKenna TM, Weinberger NM. Cholinergic modulation of frequency receptive fields in auditory cortex: II. Frequency-specific effects of anticholinesterases provide evidence for a modulatory action of endogenous ACh. *Synapse*. 1989; 4:44–54. [PubMed: 2772838]
- Budinger E, Heil P, Scheich H. Functional organization of auditory cortex in the Mongolian gerbil (*Meriones unguiculatus*). IV. Connections with anatomically characterized subcortical structures. *Eur J Neurosci*. 2000; 12:2452–2474. [PubMed: 10947822]
- Calford MB, Webster WR, Semple MM. Measurement of frequency selectivity of single neurons in the central auditory pathway. *Hearing Res*. 1983; 11:395–401.
- Chattopadhyay S, Xue B, Collins D, Pichika R, Bagnera R, Leslie FM, Christian BT, Shi B, Narayanan TK, Potkin SG, Mukherjee J. Synthesis and evaluation of nicotine alpha4beta2 receptor radioligand, 5-(3'-18F-fluoropropyl)-3-(2-(S)-pyrrolidinylmethoxy)pyridine, in rodents and PET in nonhuman primate. *J Nucl Med*. 2005; 46:130–140. [PubMed: 15632043]
- Chen G, Yan J. Cholinergic modulation incorporated with a tone presentation induces frequency-specific threshold decreases in the auditory cortex of the mouse. *Eur J Neurosci*. 2007; 25:1793–1803. [PubMed: 17432966]
- Christophe E, Roebuck A, Staiger JF, Lavery DJ, Charpak S, Audinat E. Two types of nicotinic receptors mediate an excitation of neocortical layer I interneurons. *J Neurophysiol*. 2002; 88:1318–1327. [PubMed: 12205153]
- Clarke PB. Nicotinic modulation of thalamocortical neurotransmission. *Prog Brain Res*. 2004; 145:253–260. [PubMed: 14650920]
- Clarke PB, Pert CB, Pert A. Autoradiographic distribution of nicotine receptors in rat brain. *Brain Res*. 1984; 323:390–395. [PubMed: 6098346]
- Clarke PB, Schwartz RD, Paul SM, Pert CB, Pert A. Nicotinic binding in rat brain: autoradiographic comparison of [3H]acetylcholine, [3H]nicotine, and [125I]-alpha-bungarotoxin. *J Neurosci*. 1985; 5:1307–1315. [PubMed: 3998824]
- Cox CL, Metherate R, Ashe JH. Modulation of cellular excitability in neocortex: muscarinic receptor and second messenger-mediated actions of acetylcholine. *Synapse*. 1994; 16:123–136. [PubMed: 7910986]

- Cruikshank SJ, Rose HJ, Metherate R. Auditory thalamocortical synaptic transmission, in vitro. *J Neurophysiol.* 2002; 87:361–384. [PubMed: 11784756]
- Ding YS, Fowler JS, Logan J, Wang GJ, Telang F, Garza V, Biegon A, Pareto D, Rooney W, Shea C, Alexoff D, Volkow ND, Vocci F. 6-[18F]Fluoro-A-85380, a new PET tracer for the nicotinic acetylcholine receptor: studies in the human brain and in vivo demonstration of specific binding in white matter. *Synapse.* 2004; 53:184–189. [PubMed: 15236351]
- Easwaramoorthy B, Pichika R, Collins D, Potkin SG, Leslie FM, Mukherjee J. Effect of acetylcholinesterase inhibitors on the binding of nicotinic $\alpha 4\beta 2$ receptor PET radiotracer, (18)F-nifene: A measure of acetylcholine competition. *Synapse.* 2007; 61:29–36. [PubMed: 17068780]
- Edeline J-M, Hars B, Maho C, Hennevin E. Transient and prolonged facilitation of tone-evoked responses induced by basal forebrain stimulations in the rat auditory cortex. *Exp Brain Res.* 1994; 97:373–386. [PubMed: 8187850]
- Edeline JM. The thalamo-cortical auditory receptive fields: regulation by the states of vigilance, learning and the neuromodulatory systems. *Exp Brain Res.* 2003; 153:554–572. [PubMed: 14517594]
- Edeline JM, Dutrieux G, Manunta Y, Hennevin E. Diversity of receptive field changes in auditory cortex during natural sleep. *Eur J Neurosci.* 2001; 14:1865–1880. [PubMed: 11860482]
- Eggermont JJ. How homogeneous is cat primary auditory cortex? Evidence from simultaneous single-unit recordings. *Auditory Neurosci.* 1996; 2:79–96.
- Foeller E, Vater M, Kossel M. Laminar analysis of inhibition in the gerbil primary auditory cortex. *J Assoc Res Otolaryngol.* 2001; 2:279–296. [PubMed: 11669400]
- Frazier CJ, Rollins YD, Breese CR, Leonard S, Freedman R, Dunwiddie TV. Acetylcholine activates an α -bungarotoxin-sensitive nicotinic current in rat hippocampal interneurons, but not pyramidal cells. *J Neurosci.* 1998; 18:1187–1195. [PubMed: 9454829]
- Frazier CJ, Strowbridge BW, Papke RL. Nicotinic receptors on local circuit neurons in dentate gyrus: a potential role in regulation of granule cell excitability. *J Neurophysiol.* 2003; 89:3018–3028. [PubMed: 12611982]
- Fritz J, Shamma S, Elhilali M, Klein D. Rapid task-related plasticity of spectrotemporal receptive fields in primary auditory cortex. *Nat Neurosci.* 2003; 6:1216–1223. [PubMed: 14583754]
- Froemke RC, Merzenich MM, Schreiner CE. A synaptic memory trace for cortical receptive field plasticity. *Nature.* 2007; 450:425–429. [PubMed: 18004384]
- Galvan VV, Chen J, Weinberger NM. Long-term frequency tuning of local field potentials in the auditory cortex of the waking guinea pig. *J Assoc Res Otolaryngol.* 2001; 2:199–215. [PubMed: 11669394]
- Gil Z, Connors BW, Amitai Y. Differential regulation of neocortical synapses by neuromodulators and activity. *Neuron.* 1997; 19:679–686. [PubMed: 9331357]
- Gioanni Y, Rougeot C, Clarke PB, Lepouse C, Thierry AM, Vidal C. Nicotinic receptors in the rat prefrontal cortex: increase in glutamate release and facilitation of mediodorsal thalamo-cortical transmission. *Eur J Neurosci.* 1999; 11:18–30. [PubMed: 9987008]
- Gulledge AT, Park SB, Kawaguchi Y, Stuart GJ. Heterogeneity of phasic cholinergic signaling in neocortical neurons. *J Neurophysiol.* 2007; 97:2215–2229. [PubMed: 17122323]
- Halliwel JV, Adams PR. Voltage-clamp analysis of muscarinic excitation in hippocampal neurons. *Brain Res.* 1982; 250:71–92. [PubMed: 6128061]
- Happel MF, Jeschke M, Ohl FW. Spectral integration in primary auditory cortex attributable to temporally precise convergence of thalamocortical and intracortical input. *J Neurosci.* 2010; 30:11114–11127. [PubMed: 20720119]
- Harrington IA, Heffner RS, Heffner HE. An investigation of sensory deficits underlying the aphasia-like behavior of macaques with auditory cortex lesions. *Neuroreport.* 2001; 12:1217–1221. [PubMed: 11338194]
- Hasselmo ME. Neuromodulation: acetylcholine and memory consolidation. *Trends Cogn Sci.* 1999; 3:351–359. [PubMed: 10461198]

- Hsieh CY, Cruikshank SJ, Metherate R. Differential modulation of auditory thalamocortical and intracortical synaptic transmission by cholinergic agonist. *Brain Res.* 2000; 880:51–64. [PubMed: 11032989]
- Imig TJ, Morel A. Topographic and cytoarchitectonic organization of thalamic neurons related to their targets in low-, middle-, and high-frequency representations in cat auditory cortex. *J Comp Neurol.* 1984; 227:511–539. [PubMed: 6470221]
- Ji D, Dani JA. Inhibition and disinhibition of pyramidal neurons by activation of nicotinic receptors on hippocampal interneurons. *J Neurophysiol.* 2000; 83:2682–2690. [PubMed: 10805668]
- Ji D, Lape R, Dani JA. Timing and location of nicotinic activity enhances or depresses hippocampal synaptic plasticity. *Neuron.* 2001; 31:131–141. [PubMed: 11498056]
- Jones S, Yakel JL. Functional nicotinic ACh receptors on interneurons in the rat hippocampus. *J Physiol.* 1997; 504 (Pt 3):603–610. [PubMed: 9401968]
- Kaur S, Lazar R, Metherate R. Intracortical pathways determine breadth of subthreshold frequency receptive fields in primary auditory cortex. *J Neurophysiol.* 2004; 91:2551–2567. [PubMed: 14749307]
- Kaur S, Rose HJ, Lazar R, Liang K, Metherate R. Spectral integration in primary auditory cortex: laminar processing of afferent input, in vivo and in vitro. *Neuroscience.* 2005; 134:1033–1045. [PubMed: 15979241]
- Kawai H, Lazar R, Metherate R. Nicotinic control of axon excitability regulates thalamocortical transmission. *Nature Neurosci.* 2007; 10:1168–1175. [PubMed: 17704774]
- Kelly JB, Whitfield IC. Effects of auditory cortical lesions on discriminations of rising and falling frequency-modulated tones. *J Neurophysiol.* 1971; 34:802–816. [PubMed: 5097158]
- Kilgard MP, Merzenich MM. Cortical map reorganization enabled by nucleus basalis activity. *Science.* 1998; 279:1714–1718. [PubMed: 9497289]
- Krnjevic K, Pumain R, Renaud L. The mechanism of excitation by acetylcholine in the cerebral cortex. *J Physiol (Lond).* 1971; 215:247–268. [PubMed: 5579661]
- Kurt S, Crook JM, Ohl FW, Scheich H, Schulze H. Differential effects of iontophoretic in vivo application of the GABA(A)-antagonists bicuculline and gabazine in sensory cortex. *Hear Res.* 2006; 212:224–235. [PubMed: 16442250]
- Lambe EK, Picciotto MR, Aghajanian GK. Nicotine induces glutamate release from thalamocortical terminals in prefrontal cortex. *Neuropsychopharmacol.* 2003; 28:216–225.
- Lavine N, Reuben M, Clarke PBS. A population of nicotine receptors is associated with thalamocortical afferents in the adult rat: laminar and areal analysis. *J Comp Neurol.* 1997; 380:175–190. [PubMed: 9100131]
- Liang K, Poytress BS, Chen Y, Leslie FM, Weinberger NM, Metherate R. Neonatal nicotine exposure impairs nicotinic enhancement of central auditory processing and auditory learning in adult rats. *Eur J Neurosci.* 2006; 24:857–866. [PubMed: 16848798]
- Liu BH, Wu GK, Arbuckle R, Tao HW, Zhang LI. Defining cortical frequency tuning with recurrent excitatory circuitry. *Nat Neurosci.* 2007; 10:1594–1600. [PubMed: 17994013]
- London ED, Waller SB, Wamsley JK. Autoradiographic localization of [3H]nicotine binding sites in the rat brain. *Neurosci Lett.* 1985; 53:179–184. [PubMed: 3982705]
- Madison DV, Lancaster B, Nicoll RA. Voltage clamp analysis of cholinergic action in the hippocampus. *J Neurosci.* 1987; 7:733–741. [PubMed: 3559710]
- Marks MJ, Pauly JR, Gross SD, Deneris ES, Hermans-Borgmeyer I, Heinemann SF, Collins AC. Nicotine binding and nicotinic receptor subunit RNA after chronic nicotine treatment. *J Neurosci.* 1992; 12:2765–2784. [PubMed: 1613557]
- McCormick DA, Prince DA. Mechanism of action of acetylcholine in the guinea-pig cerebral cortex in vitro. *J Physiol (Lond).* 1986; 375:169–194. [PubMed: 2879035]
- McKenna TM, Ashe JH, Hui GK, Weinberger NM. Muscarinic agonists modulate spontaneous and evoked unit discharge in auditory cortex of cat. *Synapse.* 1988; 2:54–68. [PubMed: 3420531]
- McKenna TM, Ashe JH, Weinberger NM. Cholinergic modulation of frequency receptive fields in auditory cortex: I. Frequency-specific effects of muscarinic agonists. *Synapse.* 1989; 4:30–43. [PubMed: 2672402]

- Mechawar N, Cozzari C, Descarries L. Cholinergic innervation in adult rat cerebral cortex: a quantitative immunocytochemical description. *J Comp Neurol.* 2000; 428:305–318. [PubMed: 11064369]
- Mechawar N, Watkins KC, Descarries L. Ultrastructural features of the acetylcholine innervation in the developing parietal cortex of rat. *J Comp Neurol.* 2002; 443:250–258. [PubMed: 11807835]
- Mendelson JR, Cynader MS. Sensitivity of cat primary auditory cortex (AI) neurons to the direction and rate of frequency modulation. *Brain Res.* 1985; 327:331–335. [PubMed: 3986511]
- Metherate R. Nicotinic acetylcholine receptors in sensory cortex. *Learn Mem.* 2004; 11:50–59. [PubMed: 14747517]
- Metherate R, Ashe JH. Ionic flux contributions to neocortical slow waves and nucleus basalis-mediated activation: whole-cell recordings in vivo. *J Neurosci.* 1993; 13:5312–5323. [PubMed: 8254377]
- Metherate R, Ashe JH, Weinberger NM. Acetylcholine modifies neuronal acoustic rate-level functions in guinea pig auditory cortex by an action at muscarinic receptors. *Synapse.* 1990; 6:364–368. [PubMed: 2287993]
- Metherate R, Cox CL, Ashe JH. Cellular bases of neocortical activation: modulation of neural oscillations by the nucleus basalis and endogenous acetylcholine. *J Neurosci.* 1992; 12:4701–4711. [PubMed: 1361197]
- Metherate R, Kaur S, Kawai H, Lazar R, Liang K, Rose HJ. Spectral integration in auditory cortex: Mechanisms and modulation. *Hear Res.* 2005; 206:146–158. [PubMed: 16081005]
- Metherate R, Tremblay N, Dykes RW. The effects of acetylcholine on response properties of cat somatosensory cortical neurons. *J Neurophysiol.* 1988; 59:1231–1251. [PubMed: 2897434]
- Miller LM, Escabi MA, Schreiner CE. Feature selectivity and interneuronal cooperation in the thalamocortical system. *J Neurosci.* 2001; 21:8136–8144. [PubMed: 11588186]
- Morley BJ, Happe HK. Cholinergic receptors: dual roles in transduction and plasticity. *Hear Res.* 2000; 147:104–112. [PubMed: 10962177]
- Muller CM, Scheich H. Contribution of GABAergic inhibition to the response characteristics of auditory units in the avian forebrain. *J Neurophysiol.* 1988; 59:1673–1689. [PubMed: 2900295]
- Norena A, Eggermont JJ. Comparison between local field potentials and unit cluster activity in primary auditory cortex and anterior auditory field in the cat. *Hear Res.* 2002; 166:202–213. [PubMed: 12062772]
- Ohl FW, Wetzel W, Wagner T, Rech A, Scheich H. Bilateral ablation of auditory cortex in Mongolian gerbil affects discrimination of frequency modulated tones but not of pure tones. *Learn Mem.* 1999; 6:347–362. [PubMed: 10509706]
- Ojima H, Murakami K. Intracellular characterization of suppressive responses in supragranular pyramidal neurons of cat primary auditory cortex in vivo. *Cereb Cortex.* 2002; 12:1079–1091. [PubMed: 12217972]
- Parkinson D, Kratz KE, Daw NW. Evidence for a nicotinic component to the actions of acetylcholine in cat visual cortex. *Exp Brain Res.* 1988; 73:553–568. [PubMed: 3224664]
- Phillips DP. Sensory representations, the auditory cortex, and speech perception. *Sem Hearing.* 1998; 19:319–332.
- Phillips DP, Farmer ME. Acquired word deafness, and the temporal grain of sound representation in the primary auditory cortex. *Behav Brain Res.* 1990; 40:85–94. [PubMed: 2285476]
- Phillips DP, Mendelson JR, Cynader MS, Douglas RM. Responses of single neurones in cat auditory cortex to time-varying stimuli: frequency-modulated tones of narrow excursion. *Exp Brain Res.* 1985; 58:443–454. [PubMed: 4007088]
- Porter JT, Cauli B, Tsuzuki K, Lambolez B, Rossier J, Audinat E. Selective excitation of subtypes of neocortical interneurons by nicotinic receptors. *J Neurosci.* 1999; 19:5228–5235. [PubMed: 10377334]
- Prusky GT, Shaw C, Cynader MS. Nicotine receptors are located on lateral geniculate nucleus terminals in cat visual cortex. *Brain Res.* 1987; 412:131–138. [PubMed: 3607444]
- Rauschecker JP, Tian B. Mechanisms and streams for processing of “what” and “where” in auditory cortex. *Proc Natl Acad Sci U S A.* 2000; 97:11800–11806. [PubMed: 11050212]

- Recanzone GH, Schreiner CE, Merzenich MM. Plasticity in the frequency representation of primary auditory cortex following discrimination training in adult owl monkeys. *J Neurosci.* 1993; 13:87–103. [PubMed: 8423485]
- Rogers SW, Gahring LC, Collins AC, Marks M. Age-related changes in neuronal nicotinic acetylcholine receptor subunit alpha4 expression are modified by long-term nicotine administration. *J Neurosci.* 1998; 18:4825–4832. [PubMed: 9634548]
- Role LW, Berg DK. Nicotinic receptors in the development and modulation of CNS synapses. *Neuron.* 1996; 16:1077–1085. [PubMed: 8663984]
- Ross SA, Wong JY, Clifford JJ, Kinsella A, Massalas JS, Horne MK, Scheffer IE, Kola I, Waddington JL, Berkovic SF, Drago J. Phenotypic characterization of an alpha 4 neuronal nicotinic acetylcholine receptor subunit knock-out mouse. *J Neurosci.* 2000; 20:6431–6441. [PubMed: 10964949]
- Sahin M, Bowen WD, Donoghue JP. Location of nicotinic and muscarinic cholinergic and mu-opiate receptors in rat cerebral neocortex: evidence from thalamic and cortical lesions. *Brain Res.* 1992; 579:135–147. [PubMed: 1320443]
- Sarter M, Givens B, Bruno JP. The cognitive neuroscience of sustained attention: where top-down meets bottom-up. *Brain Res Rev.* 2001; 35:146–160. [PubMed: 11336780]
- Schulze H, Langner G. Auditory cortical responses to amplitude modulations with spectra above frequency receptive fields: evidence for wide spectral integration. *J Comp Physiol [A].* 1999; 185:493–508.
- Sillito AM, Kemp JA. Cholinergic modulation of the functional organization of the cat visual cortex. *Brain Res.* 1983; 289:143–155. [PubMed: 6661640]
- Tan AY, Zhang LI, Merzenich MM, Schreiner CE. Tone-evoked excitatory and inhibitory synaptic conductances of primary auditory cortex neurons. *J Neurophysiol.* 2004; 92:630–643. [PubMed: 14999047]
- Tian B, Rauschecker JP. Processing of frequency-modulated sounds in the lateral auditory belt cortex of the rhesus monkey. *J Neurophysiol.* 2004; 92:2993–3013. [PubMed: 15486426]
- Velenovsky DS, Cetas JS, Price RO, Sinex DG, McMullen NT. Functional subregions in primary auditory cortex defined by thalamocortical terminal arbors: an electrophysiological and anterograde labeling study. *J Neurosci.* 2003; 23:308–316. [PubMed: 12514229]
- Wang J, Caspary D, Salvi RJ. GABA-A antagonist causes dramatic expansion of tuning in primary auditory cortex. *Neuroreport.* 2000; 11:1137–1140. [PubMed: 10790896]
- Wang J, McFadden SL, Caspary D, Salvi R. Gamma-aminobutyric acid circuits shape response properties of auditory cortex neurons. *Brain Res.* 2002; 944:219–231. [PubMed: 12106684]
- Wehr M, Zador AM. Balanced inhibition underlies tuning and sharpens spike timing in auditory cortex. *Nature.* 2003; 426:442–446. [PubMed: 14647382]
- Weinberger NM. Specific long-term memory traces in primary auditory cortex. *Nat Rev Neurosci.* 2004; 5:279–290. [PubMed: 15034553]
- Wetzel W, Ohl FW, Wagner T, Scheich H. Right auditory cortex lesion in Mongolian gerbils impairs discrimination of rising and falling frequency-modulated tones. *Neurosci Lett.* 1998; 252:115–118. [PubMed: 9756335]
- Whitfield IC, Evans EF. Responses of auditory cortical neurons to stimuli of changing frequency. *J Neurophysiol.* 1965; 28:655–672. [PubMed: 14347626]
- Winer JA, Sally SL, Larue DT, Kelly JB. Origins of medial geniculate body projections to physiologically defined zones of rat primary auditory cortex. *Hear Res.* 1999; 130:42–61. [PubMed: 10320098]
- Xiang Z, Huguenard JR, Prince DA. Cholinergic switching within neocortical inhibitory networks. *Nature.* 1998; 281:985–988.
- Zhang Y, Dyck RH, Hamilton SE, Nathanson NM, Yan J. Disrupted tonotopy of the auditory cortex in mice lacking M1 muscarinic acetylcholine receptor. *Hear Res.* 2005; 201:145–155. [PubMed: 15721569]
- Zoli M, Lena C, Picciotto MR, Changeux JP. Identification of four classes of brain nicotinic receptors using beta2 mutant mice. *J Neurosci.* 1998; 18:4461–4472. [PubMed: 9614223]

Figure 1.

Schematic (left) of hypothesis that spectral integration by neurons in A1 involves direct thalamocortical inputs carrying information about CF (and near-CF) stimuli and long-distance, intracortical “horizontal” projections carrying information about spectrally-distant (nonCF) stimuli. CF, characteristic frequency; MGv, ventral division of the medial geniculate body. Graph (right) illustrates hypothetical results of intracellular recording from neuron on left responding to tone stimuli, with suprathreshold EPSPs generating the classical receptive field flanked by subthreshold EPSPs elicited in response to higher and lower frequencies.

Figure 2.

Hypothesized effect of ACh on receptive fields due to modulation of thalamocortical and intracortical circuits (left). Cellular effects of ACh include those mediated by muscarinic ACh receptors (mAChRs)—including enhancement of postsynaptic excitability and presynaptic reduction of intracortical transmission—and effects mediated by nicotinic ACh receptors (nAChRs)—including enhancement of thalamocortical transmission by axonal and presynaptic receptors on thalamocortical axons, and excitation of cortical GABAergic interneurons. A potential result of these combined actions (right) is to reduce receptive field breadth and enhance responsiveness within the “sharpened” receptive field (red line).