CORRESPONDENCE

Self-Testing of Vaginal pH to Prevent Preterm Delivery: A Controlled Trial

by Prof. Dr. med. Eva-Maria Bitzer, Dipl.-Psych. Andrea Schneider, Paul Wenzlaff, Prof. Dr. med. Prof. Dr. h. c. Udo B. Hoyme, Dr. med. Elisabeth Siegmund-Schultze in volume 6/2011

Weaknesses in Study Design

We welcome the fact that even studies that did not yield positive results find their way to publication. However, to conclude from the absence of a positive result that no association exists between bacterial vaginosis (disturbance of the microbiological balance of the flora of the female genital tract) and preterm birth would be regrettable.

The negative result does not surprise me and corresponds to my own long years of experience. In addition to the substantial selection problem in this study, the following microbial causes are likely to have contributed to the negative result:

- Measuring the vaginal pH captures only the number of lactobacilli and not the number of potentially pathogenic bacteria in the vagina. The pH is also raised after ingestion of antibiotics, but no bacterial disturbance is present in such a scenario, and neither is there a risk of infection.
- It has been common knowledge for a long time that a disturbed vaginal flora and bacterial vaginosis, often accompanied by *E coli*, are among the risk factors for late miscarriage and preterm birth. In contrast to other risk factors, this cannot be detected and rectified by the patient but by her doctor.
- Although metronidazole and clindamycin are mentioned as guideline-recommended first-line agents of choice for bacterial vaginosis, as studies for these substances exist only for non-pregnant women, treatment with these substances during pregnancy in order to reduce the rate of preterm births is inappropriate (1, 2). Metronidazole is problematic during pregnancy because of its interaction with DNA, and clindamycin tends to trigger an abnormal selection of *E coli*.
- Improving the anogenital region in the long term by promoting the growth of lactobacilli by means of acidification and water-free ointments and reducing intestinal pathogens is the better method, in my experience (3). Sickness funds should commit to this treatment if they want to prevent suffering and save money.

DOI: 10.3238/arztebl.2011.0460a

REFERENCES

1. Donders GG, et al.: Predictive value for preterm birth of abnormal vaginal flora, bacterial vaginosis and aerobic vaginitis during the first trimester of pregnancy or why metronidazole is not a good option in pregnancy. 6th Eur Conf Eur Soc Inf Dis Obstet Gynecol (ESIDOG), Leuven/Belgien, 2008, Abstract Plen II-2.
2. Klebanoff MA, et al.: Failure of metronidazole to prevent preterm delivery among pregnant women with asymptomatic *Trichomonas vaginalis* infection. *N Engl J Med* 2001; 345: 487–93.
3. Petersen EE: Infektionen in Gynäkologie und Geburtshilfe. 5th edition. Stuttgart: Thieme-Verlag 2011.
4. Bitzer EM, Schneider A, Wenzlaff P, Hoyme UB, Siegmund-Schultze E: Self-testing of vaginal pH to prevent preterm delivery: A controlled trial. *Dtsch Arztebl Int* 2011; 108(6): 81–6.

Prof. Dr. med. Eiko E. Petersen
Freiburg, Germany
eiko.petersen@web.de

Conflict of interest statement

Professor Petersen has received honoraria for speaking and continuing medical education events from Bayer, Taurus, Essex Pharma, Kaymogyn, and Grünenthal.

Methodological Problems

When considering the common problem of publication bias it is to be welcomed that the evaluation results from this study were published.

In view of the fact that 1887 of 7469 pregnant women (25%) mentioned at least one abnormal test value, but only 803 (10.8%) sought a medical examination and only 642 (9%) were treated with whatever medications, it becomes obvious that a reduction in the preterm birth rate cannot always be achieved.

One of the study's problems is that it is impossible to figure out, even as a tendency, why vaginal pH self-testing is not efficacious. This is due to an accumulation of methodological problems in the intervention and evaluation.

Interventions have to be well planned and controlled with regard to their structures, processes, and outcomes.

The aim of planning and conducting interventions is to generate, in a case-control approach, intervention groups that are not selected by health or social status. In the reported study, the case group included a sample that was without exception self-selected.

In the context of the study, it was not possible to control the structures and processes; the intervention was conducted under the pregnant women's own direction. Whether and how the pH test glove was used was elicited only by means of a questionnaire. The control group was not questioned. Only by doing so it would have been possible, however, to recognize the selection in the groups and possibly to standardize these.

The present study does not raise any doubt in the individual medical benefits of the glove. However, any

further administration of the glove to study populations should always be accompanied by a robust evaluation.

DOI: 10.3238/arztebl.2011.0460b

REFERENCES

1. Bitzer EM, Schneider A, Wenzlaff P, Hoyme UB, Siegmund-Schultze E: Self-testing of vaginal pH to prevent preterm delivery: a controlled trial. *Dtsch Arztebl Int* 2011; 108(6): 81–6.

Dr. phil. Wolf Kirschner
Berlin, Germany

Prof. Dr. med. Klaus Friese
President
Deutsche Gesellschaft für Gynäkologie und Geburtshilfe, München, Germany
Klaus.friese@med.uni-muenchen.de

Dr. med. Albrecht Scheffler
Berlin, Germany

Conflict of interest statement

The authors declare that no conflict of interest exists.

Serious Shortcomings

The present study (1) has serious shortcomings. It is unsuitable for assessing the efficacy of vaginal pH self-testing in the prevention of preterm births. Important limitations are mentioned in the main part of the article but not in the abstract. The many “abstract only readers” will thus get misleading information.

In the study, it was not ensured:

- that a sufficiently high proportion of participants received the test kit in a timely fashion and then proceeded to take regular measurements. Only 1823 pregnant women took the measurements according to the trial protocol (this amounts to merely 9.7% of participants in the intervention group) (2). This was not mentioned in the actual publication;
- that the diagnostic and therapeutic approaches taken by the treating gynecologists were standardized and thus comparable;
- that in case of a pathological status confirmed by a physician, treatment was given in accordance with current guidelines.

The question needs to be asked why the study was even conducted in this form. Basically, the results are meaningless and may even be misleading. And why were we not even consulted ahead of time, when it was us who introduced the innovative self-test for vaginal pH for pregnant women in 1994 (3, 4) and have since then developed this further? We were not even mentioned as the initiators, which usually is common best practice among scientists.

The absence of essential prerequisites cannot be balanced by extensive statistics. However, in spite of the listed shortcomings, a minimally protective effect of pH self-testing was noted in association with birth weight. Therefore, it may be assumed that more notable effects would have been found in a correctly designed study.

DOI: 10.3238/arztebl.2011.0461a

REFERENCES

1. Bitzer EM, Schneider A, Wenzlaff P, Hoyme UB, Siegmund-Schultze E: Self-testing of vaginal pH to prevent preterm delivery: A controlled trial. *Dtsch Arztebl Int* 2011; 108(6): 81–6.
2. Schneider A, Rudolph A, Bitzer EM, Dörning H: Vermeidung von Frühgeburten. Gemeinsames Modellprojekt nach § 63 SGB V der Kaufmännischen Krankenkasse (KKH), Barmer Ersatzkasse, Techniker Krankenkasse (TK), Hamburg Münchener Krankenkasse (HMK). *Endbericht* 2009; 48.
3. Saling E, Fuhr N, Placht A, Schumacher E: „Selbst-Vorsorge-Aktion für Schwangere“ – ein wichtiger Bestandteil des Frühgeburtenvermeidungsprogrammes. In: Weidinger H, (eds.): *Geburtshilfe und Frauenheilkunde. Moderne Medizin zwischen Natur und High-Tech*. 7. Bayreuther Gespräch. Berlin, Wien: Blackwell Wissenschafts-Verlag 1995; 63–70.
4. Saling E, Fuhr N, Placht A, Schumacher E: A new efficient strategy for prevention of prematurity. In: Kurjak A, Latin V, Rippmann E (ed.): *Advances on the pathophysiology of pregnancy: CIC Edizioni Internazionali* 1995; 228–34.

Prof. Dr. med. Erich Saling
Jürgen Lütjhe

Institut für Perinatale Medizin, Berlin, Germany
info@saling-institut.de

Conflict of interest statement

Professor Saling developed the pH self-test for the purpose of avoiding preterm births. Mr Lütjhe declares that no conflict of interest exists.

In Reply:

We agree that a prospective randomized controlled trial with early recruitment of 64 000 pregnant women, monitoring of participants’ compliance, and guideline conformity among participating gynecologists would be the ideal setting for confirming the effect of pH self-testing on the rate of preterm births. Such a study would be a clinical study, and the statutory health insurers currently do not have the legal foundation to support such studies. Our investigation was done as a statutorily mandatory evaluation on the basis of §65 SGB V (social code, book V) in the context of a model project according to §63 SGB V, which dealt with the further development of healthcare services.

The limitations of the intervention and study design pointed out by our correspondents (among others, a self-selected control group, late recruitment of pregnant women, no recommendations on treatment and follow-up) reflect routine care in the context of which the statutory health insurers would probably provide pH self-testing as a standard service to pregnant women. It was therefore intentionally not part of the study protocol to stipulate possible subsequent diagnostic evaluation and resultant treatment (that is, a treatment intervention) in addition to the intervention “pH self-testing.” These should be provided by current routine care. Our evaluation therefore enables conclusions about the effectiveness in real healthcare settings under everyday routine conditions (1–3).

The strength of our study lies in the fact that we collected individual participants’ data for the total study population—the intervention group as well as the control group—with regard to the desired outcomes (preterm rate and birth weight) and for central patients’

characteristics (risk factors) that influence these outcomes. The study design of the model project enabled us to test the validity and robustness as well as possible biases in the results much better than for example Thuringia's prematurity prevention initiative (*Frühgeburtenvermeidungsaktion*). It should not be interpreted as a disadvantage that we made use of these opportunities and analyzed the comprehensive data with the appropriate statistical methods (4). Furthermore, we showed at least a trend towards reducing the proportion of neonates with a very low birth weight, who obviously present the real medical-social problem clientele.

We did not survey the control groups, as recommended by Kirschner et al. Since pregnant women could have been surveyed only after giving birth and a low response rate would have had to be expected in the intervention group as well as the control group, additional memory and selection related biases would have been introduced.

We thank Petersen for his explanations of the alternatives to pH self-testing.

DOI: 10.3238/arztebl.2011.0461b

REFERENCES

1. Bundesärztekammer: Definition der Versorgungsforschung. www.baek.de
2. Bitzer EM: Versorgungsforschung aus Public Health Perspektive – Forschungsansätze mit patientennahen Ergebnisparametern und Routinedaten von Krankenkassen. Hamburg: Dr. Kovacs, 2010.

3. Schäfer T: Wissenschaftliche Begleitung nach §68 SGB V – Grundzüge eines Evaluationskonzepts. in: Brennecke R (ed.). Sozialmedizinische Ansätze der Evaluation in Gesundheitswesen, Band 1: Grundlagen und Versorgungsforschung. Berlin, Heidelberg, New York: Springer 1992: 442–54.
4. Glaeske G, Augustin M, Abholz H, et al.: Epidemiologische Methoden für die Versorgungsforschung. Das Gesundheitswesen 2009; 71: 685–69.
5. Bitzer EM, Schneider A, Wenzlaff P, Hoyme UB, Siegmund-Schultze E: Self-testing of vaginal pH to prevent preterm delivery: A controlled trial. Dtsch Arztebl Int 2011; 108(6): 81–6.

Prof. Dr. med. Eva-Maria Bitzer

Dipl.-Psych. Andrea Schneider

Institut für Sozialmedizin, Epidemiologie und Gesundheitssystemforschung (ISEG)
Hannover, Germany
bitzer@iseg.org

Paul Wenzlaff

Zentrum für Qualität und Management im Gesundheitswesen
Niedersachsen, Germany

Prof. Dr. med. Udo B. Hoyme

Klinik für Frauenheilkunde und Geburtshilfe, Helios-Klinikum
Erfurt, Germany

Dr. med. Elisabeth Siegmund-Schultze

KKH-Allianz, Hannover, Germany

Conflict of interest statement

Professor Hoyme acts as a speaker and adviser for the statutory sickness funds supporting the study. In the current study he was the scientific adviser. He has received honoraria for advisory activities and travel expenses from Inverness (manufacturers of the glove).

The other authors declare that no conflict of interest exists.