Perspective

ASSOCIATE EDITOR: DAVID R. SIBLEY

The Highs and Lows of Cannabinoid Receptor Expression in Disease: Mechanisms and Their Therapeutic Implications

Lydia K. Miller and Lakshmi A. Devi

Department of Pharmacology and Systems Therapeutics, Mount Sinai School of Medicine, New York, New York

Abstract	461
I. Introduction: the cannabinoid receptors and their response to disease.	461
II. Mechanisms of cannabinoid receptor regulation.	464
III. Therapeutic implications	465
IV. Cannabinoid receptor gene structure	465
A. Cannabinoid receptor-1.	466
B. Cannabinoid receptor-2	466
V. High-throughput studies of transcription factor interactions with cannabinoid receptors 1 and 2 ...	468
VI. Conclusions and future directions	468
Acknowledgments	468
References	468

Abstract—Alterations in the endogenous cannabinoid system have been described in almost every category of disease. These changes can alternatively be protective or maladaptive, such as producing antinociception in neuropathic pain or fibrogenesis in liver disease, making the system an attractive therapeutic target. However, the challenge remains to selectively target the site of disease while sparing other areas, particularly mood and cognitive centers of the brain. Identifying regional changes in cannabinoid receptor-1 and -2 (CB₁R and CB₂R) expression is particularly important when considering endocannabinoid system-based therapies, because regional increases in cannabinoid receptor expression have been shown to increase potency and efficacy of exogenous agonists at

sites of disease. Although there have been extensive descriptive studies of cannabinoid receptor expression changes in disease, the underlying mechanisms are only just beginning to unfold. Understanding these mechanisms is important and potentially relevant to therapeutics. In diseases for which cannabinoid receptors are protective, knowledge of the mechanisms of receptor up-regulation could be used to design therapies to regionally increase receptor expression and thus increase efficacy of an agonist. Alternatively, inhibition of harmful cannabinoid up-regulation could be an attractive alternative to global antagonism of the system. Here we review current findings on the mechanisms of cannabinoid receptor regulation in disease and discuss their therapeutic implications.

I. Introduction: The Cannabinoid Receptors and Their Response to Disease

The endocannabinoid system is uniquely poised to respond locally to disease. Endocannabinoids are synthesized “on demand” from membrane phospholipids in response to increases in intracellular calcium (as occurs with neuronal activation or cell stress) and immediately released to act in paracrine fashion on nearby G_{i/o}-protein coupled receptors CB₁R¹ and CB₂R. Endocannabi-

Address correspondence to: Dr. Lakshmi A. Devi, Department of Pharmacology and Systems Therapeutics, Box 1603, Mount Sinai School of Medicine, One Gustave L. Levy Place, New York, NY 10029. E-mail: lakshmi.devi@mssm.edu

This article is available online at <http://pharmrev.aspetjournals.org>.
doi:10.1124/pr.110.003491.

TABLE 1
CB₁R and CB₂R expression changes in disease

Major findings in several disease categories have been selected; for more exhaustive review, see references cited in text.

Disease	CB ₁ R/CB ₂ R Regulation	Proposed Role of CB ₁ R/ CB ₂ R in Disease	Therapeutic Implications
Neuropathic pain	CB ₁ R and CB ₂ R up-regulated in peripheral and central sensory pathways in animal models of neuropathic pain (Siegling et al., 2001; Lim et al., 2003; Zhang et al., 2003; Walczak et al., 2005; Mitrirattanakul et al., 2006).	Inhibition of neurotransmitter release decreases hyperexcitability of sensory pathways (CB ₁ R); inhibition of inflammation (CB ₂ R) (for review, see Costigan et al., 2009).	CB ₁ R up-regulation enhances analgesic response to exogenous cannabinoids in animal model of neuropathic pain (Lim et al., 2003).
Neuroinflammation and brain injury	CB ₂ R up-regulated in microglia/macrophage-like cells of patients with MS or ALS (Yiangou et al., 2006) and in microglia of EAE mouse model of MS (Maresz et al., 2005). CB ₁ R up-regulated in rat brain after mild concussive head injury and NMDAR blockade (Hansen et al., 2001).	CB ₂ R expressed on T cells reduces inflammation in MS (Maresz et al., 2007). Neuroprotection by decreasing glutamate release/excitotoxicity, among other possible mechanisms (for review, see Mechoulam et al., 2002).	T-cell CB ₂ R is already highly activated by endocannabinoids in MS; increasing receptor number is thus important for increasing efficacy of CB ₂ R agonist (Maresz et al., 2007).
Cancer	Increased CB ₁ R and CB ₂ R expression in multiple human cancers (for review, see Sarfaraz et al., 2008). Down-regulated CB ₁ R in human colorectal tumors (Wang et al., 2008).	Cannabinoid receptor activation leads to apoptosis via cell cycle regulation and ER stress (for review, see Sarfaraz et al., 2008). Loss of CB ₁ R leads to enhanced colorectal tumor proliferation in mouse model of colorectal cancer (Wang et al., 2008).	Up-regulation of receptors may be crucial for antitumor effects of cannabinoids. For example, a nonselective agonist inhibits growth of prostate cancer cells with high expression of CB ₁ R and CB ₂ R (Sarfaraz et al., 2005), whereas Δ^9 -THC stimulates growth in breast cancer cells with low expression (McKallip et al., 2005). Increasing receptor expression where it is lost could be a novel therapeutic option (Wang et al., 2008).
Gastrointestinal	CB ₁ R up-regulated in mouse model of diarrhea (Izzo et al., 2001); CB ₁ R and CB ₂ R up-regulated in multiple models of colitis (Massa et al., 2004; Kimball et al., 2006).	CB ₁ R expressed on enteric neurons decreases intestinal motility, secretion, and visceral pain and promotes wound healing. CB ₂ R expressed on gut immune cells decreases inflammation (for review, see Izzo and Camilleri, 2008).	Up-regulation of CB ₁ R probably accounts for increased efficacy and potency of agonists in slowing intestinal transit in mouse model of diarrhea (Izzo et al., 2001).
Liver	Hepatic CB ₁ R and CB ₂ R up-regulated in cirrhosis in humans and rodent activated hepatic stellate cells (aHSCs) (Julien et al., 2005; Teixeira-Clerc et al., 2006; Jeong et al., 2008). Hepatic CB ₁ R also up-regulated in mice fed high-fat (Osei-Hyiaman et al., 2005; Jourdan et al., 2010; Quarta et al., 2010) and alcohol diets (Jeong et al., 2008).	Fibrogenic stimuli up-regulate hepatic CB ₁ R, which stimulates lipogenesis in hepatocytes (Osei-Hyiaman et al., 2005) and promotes fibrogenesis through aHSCs and myofibroblasts (Teixeira-Clerc et al., 2006). CB ₂ R activation leads to apoptosis of profibrogenic aHSCs and myofibroblasts (Julien et al., 2005).	Antagonism of CB ₁ R and agonism of CB ₂ R reduces proliferation of fibrogenic cells (Julien et al., 2005; Teixeira-Clerc et al., 2006). Antagonism of CB ₁ R decreases hepatic fatty acid synthesis in mice fed high-fat diet (Osei-Hyiaman et al., 2005). CB ₁ R antagonism also prevents high-fat diet-induced increases in hepatic CB ₁ R expression (Jourdan et al., 2010).
Metabolic	Adipose CB ₁ R up-regulated in Zucker rat model of obesity (Bensaid et al., 2003) and mice fed high-fat diet (Jourdan et al., 2010).	CB ₁ R stimulates lipogenesis and inhibits fatty acid oxidation in adipose cells and may increase insulin release from the pancreas (Matias et al., 2006; for review, see Kunos et al., 2008).	CB ₁ R up-regulation in adipocytes may account for increased efficacy of CB ₁ R antagonist in reducing weight in obese compared with lean Zucker rats (Vickers et al., 2003).

TABLE 1
Continued

Major findings in several disease categories have been selected; for more exhaustive review, see references cited in text.

Disease	CB ₁ R/CB ₂ R Regulation	Proposed Role of CB ₁ R/ CB ₂ R in Disease	Therapeutic Implications
Cardiovascular	Skeletal muscle CB ₁ R up-regulated in mice fed high-fat diet (Pagotto et al., 2006).	CB ₁ R up-regulation in skeletal muscle may contribute to insulin resistance in obesity (for review, see Kunos et al., 2008).	CB ₁ R antagonism prevents high-fat diet-induced increases in adipose CB ₁ R expression (Jourdan et al., 2010).
	CB ₁ R up-regulated in myocardium and aorta in rat model of hypertension (Batkai et al., 2004).	CB ₁ R lowers blood pressure by decreasing cardiac contractility and vascular resistance, but this effect is only seen in hypertensive, not normotensive animals (Batkai et al., 2004).	CB ₁ R up-regulation in the heart and vasculature may account for increased potency and efficacy of agonists in hypertensive animals (Batkai et al., 2004).
	CB ₂ R up-regulated in immune cells in atherosclerotic plaques in humans and mouse model (Steffens et al., 2005).	CB ₂ R decreases plaque progression, possibly by reducing infiltration of immune cells and cytokine release (Steffens et al., 2005).	
Psychiatric	CB ₁ R up-regulated in prefrontal cortex of depressed suicide victims (Hungund et al., 2004) and patients with schizophrenia (Dean et al., 2001).	CB ₁ R has both excitatory and inhibitory effects on synaptic transmission in the prefrontal cortex (Lafourcade et al., 2007; Chiu et al., 2010); the effect of its up-regulation in these diseases is not known.	Targeting the endocannabinoid system in these diseases will require better understanding of its role.

MS, multiple sclerosis; ALS, amyotrophic lateral sclerosis; EAE, experimental autoimmune encephalitis; NMDAR, *N*-methyl-D-aspartate receptor; ER, endoplasmic reticulum; Δ^9 -THC, tetrahydrocannabinol; aHSC, activated hepatic stellate cell.

noids are then rapidly cleared by cellular uptake and enzymatic degradation. Cannabinoid receptors work through a variety of signaling mechanisms to exert physiological and pathophysiological effects in different tissues. In neurons, where CB₁R expression is highest, stimulation of presynaptic CB₁R inhibits neurotransmitter release by stimulating potassium channels and inhibiting calcium channels (for review, see Howlett et al., 2002). In the liver, where CB₁R is normally expressed at low levels, stimulation of CB₁R leads to enhanced expression of acetyl-CoA carboxylase-1 and fatty acid synthase and thus increases lipogenesis (Osei-Hyiaman et al., 2005). CB₂R expression is highest in immune cells, where it seems to have several immunosuppressive effects, including inhibition of proinflammatory cytokine production (Maresz et al., 2007). Cannabinoid receptor signaling is discussed in detail in several recent reviews (Cabral and Griffin-Thomas, 2009; Turu and Hunyady, 2010).

In addition to increasing levels of endocannabinoids, the system often responds to stress by altering the expression of CB₁R and/or CB₂R. In some diseases, such as neuropathic pain and multiple sclerosis, increases in cannabinoid receptor expression are thought to reduce symptoms and/or inhibit progression of disease and thus serve a protective role (for review, see Pertwee, 2009). In

other diseases, alterations in receptor expression are maladaptive, examples being CB₁R up-regulation in liver fibrosis and down-regulation in colorectal cancer (Teixeira-Clerc et al., 2006; Wang et al., 2008). In both cases, regulation of cannabinoid receptor expression is of interest from a therapeutics perspective. Regional up-regulation of CB₁R correlates with enhanced potency and efficacy of agonists at sites of disease in several animal models, including intestinal inflammation and hypertension (Izzo et al., 2001; Batkai et al., 2004); a more causal relationship was suggested in neuropathic pain, where inhibition of CB₁R up-regulation reduced the analgesic effects of cannabinoids (Lim et al., 2003). Regional up-regulation of CB₁R is similarly thought to enhance efficacy of systemic antagonists in models of obesity (for review, see Vickers et al., 2003; Kunos et al., 2008). Such up-regulation should therefore increase the benefit-to-side-effect ratio of systemic agonists (for review, see Pertwee, 2009) and antagonists. In addition, identifying the lack of cannabinoid receptor up-regulation could be important. For example, prostate cancer cells that highly express cannabinoid receptors respond favorably to agonists (Sarfaraz et al., 2005), whereas breast cancer cells that express low levels of cannabinoid receptors show increased proliferation in response to Δ^9 -tetrahydrocannabinol (McKallip et al., 2005). These alterations in cannabinoid receptor expression have been extensively reviewed elsewhere (Di Marzo et al., 2004; Pertwee, 2005, 2009; Pacher et

¹Abbreviations: CB₁R, cannabinoid receptor-1; CB₂R, cannabinoid receptor-2; ChIP, chromatin immunoprecipitation; GR, glucocorticoid receptor; kb, kilobase(s); MAPK, mitogen-activated protein kinase; Trk, receptor tyrosine kinase; UTR, untranslated region.

al., 2006; Di Marzo, 2008; Izzo and Camilleri, 2008) and are briefly summarized in Table 1 to emphasize the global nature of these alterations and their therapeutic implications.

II. Mechanisms of Cannabinoid Receptor Regulation

Despite the growing list of diseases that show cannabinoid receptor expression changes, relatively little is known about the mechanisms underlying these changes. In Table 2, we summarize all current findings of which we are aware. These studies of mechanism span a number of diseases and vary in the level of detail, from investigating changes in cannabinoid receptor protein levels to promoter occupancy, but several themes are apparent across the disease models examined so far. For one, up-regulation is induced by factors that are released locally in response to disease, in accordance with the region-specific nature of these expression changes.

For example, spinal cord CB₁R protein increases were found to be mediated in part by the Trk/MAPK pathway in a rat model of neuropathic pain, suggesting a mechanism by which neurotrophic factors released locally by nerve injury (Ha et al., 2001) could modulate CB₁R expression (Lim et al., 2003). Spinal cord CB₁R increases in neuropathic pain were also found to be mediated by glucocorticoids (Wang et al., 2007). Although corticosteroids are increased systemically in the same model (Wang et al., 2004), localized CB₁R up-regulation is probably made possible by local increases in spinal glucocorticoid receptors (GR) (Wang et al., 2004). Localized mechanisms of CB₁R regulation were also observed in liver and immune cells. Retinoic acid, which is synthesized and stored by hepatic stellate cells, was found to increase CB₁R transcription in hepatocytes through retinoic acid receptor- γ (Mukhopadhyay et al., 2010). Cytokines, which are released locally in inflammation to regulate neighboring immune cells, have been impli-

TABLE 2
Mechanisms of CB₁R and CB₂R regulation in disease

Disease/System	Mechanism	Cell Type/ Animal Model	Implications
Liver steatosis	RAR- γ binds the -500 to +50 region of the CB ₁ R promoter and increases its expression (Mukhopadhyay et al., 2010).	Primary cultured mouse hepatocytes	Retinoic acid released from hepatic stellate cells induces CB ₁ R expression in hepatocytes, which in turn induces lipogenesis (Jeong et al., 2008).
Inflammation	Cannabinoids induce CB ₁ R mRNA expression in T cells by an IL-4- and CB ₂ R-dependent mechanism (Borner et al., 2007).	Jurkat T cells	CB ₁ R is expressed at very low levels in resting T cells; when induced, it seems to have anti-inflammatory effects (Molina-Holgado et al., 2003; Nakajima et al., 2006). These studies together suggest a feedback mechanism in T cells whereby cannabinoids induce expression of anti-inflammatory IL-4 via CB ₂ R, and IL-4 in turn increases CB ₁ R transcription (Borner et al., 2008).
	IL-4 directly increases CB ₁ R mRNA expression in T cells via STAT6 binding to a STAT6 element at nt -2769 upstream from human exon1 TSS (Borner et al., 2008).	Primary human T cells and Jurkat T cells	
	IFN γ and GM-CSF synergize to increase CB ₂ R mRNA expression in microglia (Maresz et al., 2005).	Primary cultured mouse microglial cells	Increases in proinflammatory IFN γ and GM-CSF could mediate the increased microglial expression of CB ₂ R seen in EAE model of MS (Maresz et al., 2005).
Neuropathic Pain	CB ₁ R protein up-regulation in the spinal cord is mediated by Trk and MAPK (Lim et al., 2003) and GR (Wang et al., 2007).	Rat chronic constriction injury model of neuropathic pain	Neurotrophic factors and corticosteroids released with nerve injury may mediate CB ₁ R up-regulation via the Trk/MAPK pathway and GR, respectively (Lim et al., 2003; Wang et al., 2007).
Cancer	Increased DNA methylation of CB ₁ R promoter leads to decreased expression of CB ₁ R (Wang et al., 2008).	Human colorectal cancer cells	Loss of CB ₁ R expression leads to enhanced tumor proliferation in a mouse model of colorectal cancer (Wang et al., 2008); first evidence of a role for epigenetics in CB ₁ R regulation.
Feeding behaviors and energy homeostasis	CNS-specific knockout of SF-1 leads to loss of CB ₁ R expression in the ventromedial hypothalamus. SF-1 directly increases CB ₁ R expression via SF-1 element at nt -101 within the mouse promoter (Kim et al., 2008).	CNS-specific SF-1 knockout mouse and various cell lines	CNS-specific SF-1 knockouts do not show the appetite-stimulating effects of CB ₁ R agonists. SF-1 regulation of CB ₁ R expression in the VMH is thus required for cannabinoid effects on food intake (Kim et al., 2008).
Neurodegenerative	Decreased striatal CB ₁ R levels in HD are due to decreased transcription (McCaw et al., 2004).	Mouse HD model	Striatal CB ₁ R modulates dopamine transmission, which is dysregulated in HD (see refs within McCaw et al., 2004).

RAR- γ , retinoic acid receptor- γ ; IL, interleukin; STAT, signal transducer and activator of transcription; TSS, transcription start site; IFN, interferon; GM-CSF, granulocyte-macrophage-colony-stimulating factor; EAE, experimental autoimmune encephalitis; MS, multiple sclerosis; CNS, central nervous system; SF-1, steroidogenic factor-1; nt, nucleotide(s); HD, Huntington's disease.

cated in up-regulation of both CB₁R and CB₂R in immune cells (Maresz et al., 2005; Börner et al., 2008).

Second, the regulatory factors identified so far are common to many physiological and pathophysiological processes and could thus be starting points for investigation of cannabinoid receptor regulation in other disease models. For example, retinoic acid has been implicated in cell survival, differentiation, axonal outgrowth, and immune regulation (for review, see Maden, 2007; Montrone et al., 2009; Noy, 2010), whereas neurotrophic factors have been implicated in neuronal survival and synaptic plasticity (for review, see Huang and Reichardt, 2001). A role for epigenetics, a widespread mechanism by which diseases cause long-lasting changes in gene expression by DNA and histone modifications, is also emerging. Down-regulation of CB₁R in human colorectal cancer cells has been attributed to methylation of the CB₁R promoter and leads to enhanced tumor proliferation in animal models (Wang et al., 2008).

Finally, autoregulation is an emerging and intriguing mechanism of cannabinoid receptor regulation. Endocannabinoids and exogenous cannabinoids have been implicated in CB₁R up-regulation in hepatocytes and T cells, respectively (Börner et al., 2007; Mukhopadhyay et al., 2010), and long-term CB₁R antagonism has been found to counter increases in hepatic and adipose CB₁R expression in response to high-fat diet (Jourdan et al., 2010). In addition, administration of a mixed CB₁R/CB₂R agonist increases expression of the CB₂A isoform (see section IV.B) in mouse cerebellum (Liu et al., 2009). Because endocannabinoids are often increased along with cannabinoid receptors in disease (Mitrirattanakul et al., 2006; Jeong et al., 2008), such autoinduction is probably common to many diseases. Understanding this autoregulation is also important from a therapeutics standpoint. Cannabinoid receptor antagonists could produce their effects through down-regulation of cannabinoid receptors in addition to blockade of cannabinoid receptor signaling pathways, whereas agonists could increase receptor expression in addition to stimulation of signaling pathways and thus amplify response to treatment.

III. Therapeutic Implications

The mechanisms of cannabinoid receptor regulation not only shed light on the pathophysiology of a disease but are also of interest from a therapeutics perspective. The challenge remains to selectively target the endocannabinoid system while sparing other areas, particularly mood and cognitive centers of the brain. As mentioned in section I, regional increases in cannabinoid receptor expression are thought to selectively enhance the effects of agonists at sites of disease, thus increasing their benefit-to-side-effect ratio (for review, see Pertwee, 2009). In diseases for which

cannabinoid receptors are protective, knowledge of the mechanisms of this receptor up-regulation could be used to design therapies to regionally enhance receptor expression and thus further optimize benefit-to-side effect ratio or eliminate the need for systemic agonists altogether. In neuropathic pain, for example, enhanced analgesic effect of cannabinoids has been linked to increases in spinal cord CB₁R expression (Lim et al., 2003). As mentioned in section I, this up-regulation is mediated in part by Trk/MAPK pathways and glucocorticoid receptors (Lim et al., 2003; Wang et al., 2007). Increasing spinal cord CB₁R expression by local administration of Trk agonists or glucocorticoids could further increase the efficacy and potency of exogenous cannabinoids and thus allow for lower, nonpsychotropic doses. Moreover, increases in endocannabinoid production in neuropathic pain (Mitrirattanakul et al., 2006) could have a saturating effect on cannabinoid receptors. Increasing receptor number could boost responsiveness to these increased endocannabinoids and thus bypass the need for exogenous agonists. Such a rationale has already been proposed for treatment of multiple sclerosis, in which CB₂R on T cells was found to be highly activated in the experimental autoimmune encephalomyelitis model, presumably by increased levels of endocannabinoids (Maresz et al., 2007). In other diseases, increasing cannabinoid receptor expression where it is lost could be an attractive therapeutic option. Loss of CB₁R as a result of promoter methylation in colorectal cancer, as seen in human cell lines, could be rescued using a demethylating agent in the hopes of decreasing tumor proliferation (Wang et al., 2008).

Alternatively, inhibition of cannabinoid receptor up-regulation where it is harmful could be an attractive alternative to systemic antagonism. For example, CB₁R antagonists have been proposed for treatment of liver fibrosis, which is marked by maladaptive up-regulation of CB₁R (Teixeira-Clerc et al., 2006). However, systemic use of the CB₁R antagonist rimonabant for “prevention of cardiovascular events” (CRESCENDO trial) was terminated because of adverse neuropsychiatric effects (Topol et al., 2010). A better alternative might be blockade of hepatic CB₁R up-regulation; the retinoic acid system has already been implicated and could thus be a possible target (Mukhopadhyay et al., 2010). Given that CB₁R is normally expressed at very low levels in the liver (Teixeira-Clerc et al., 2006) and thus probably does not serve much of a role in normal liver physiology, inhibition of its up-regulation in the liver is likely to be safe.

IV. Cannabinoid Receptor Gene Structure

Our understanding of cannabinoid receptor gene structure, essential for studying its direct regulation, is relatively recent. Experimentally determined gene

structures and promoters of CB₁R and CB₂R are summarized here and in Fig. 1.

A. Cannabinoid Receptor-1. In human, rat, and mouse, the CB₁R coding region is contained within one exon and shows significant homology across species; however, there seems to be considerable variation in the length of 5' untranslated region (UTR). Zhang et al. (2004) identified three additional upstream exons in human CB₁R using hippocampal RNA, giving a large (approximately 20 kb) 5' UTR characteristic of neuronally expressed genes. This 5'UTR can be alternatively spliced (CB₁A–D) or transcribed at different sites (CB₁A–D versus CB₁E) to yield five possible transcripts with region-specific expression in the brain. The 3-kb sequence upstream from the exon1 transcription start site showed significant promoter activity in various CB₁R-expressing neuroblastoma cell lines. 5' Flanking sequences of exon 3 yielded much lower promoter activity, consistent with the lower expression of transcript CB₁E. The more active promoter upstream of exon1 was further investigated in T cells, which normally express low levels of CB₁R (Börner et al., 2008). It is noteworthy that positive and negative regulatory regions mediating basal expression of CB₁R in resting T cells differed from findings in neuronal lines, suggesting cell-type specific CB₁R promoter regulation. A functional signal transducer and activator of transcription-6 (STAT6) element located –2769 upstream from the exon1 start site was also found to mediate interleukin-4-inducible CB₁R expression.

Mouse CB₁R gene structure was studied by McCaw et al. (2004) using RNA from striatum. In contrast to human CB₁R (namely the striatal transcripts isolated by Zhang et al., 2004), the mouse gene contains a shorter 5'UTR, with only one additional exon located upstream of the coding exon. This exon contains multiple transcription start sites (with the major sites at the beginning of exon1); however, the promoter activities of these regions were not examined. Later studies identified direct interactions of retinoic acid receptor- γ and steroidogenic factor-1 with the mouse CB₁R promoter (Kim et al., 2008; Mukhopadhyay et al., 2010); however, these studies did not specify which putative promoter was examined.

The originally cloned CB₁R from a rat cerebral cortex cDNA library (Matsuda et al., 1990) similarly consists of two exons; however, additional 5' exons have not been investigated. Homology between human and rodent 5'UTR has not been examined; in general, species comparison is more detailed for CB₂R (see next section).

B. Cannabinoid Receptor-2. Like CB₁R, the CB₂R coding region is contained in a single exon and is flanked by upstream noncoding exons in human, mouse, and, debatably, rat. Human CB₂R consists of three exons alternatively transcribed and spliced to yield isoforms CB₂A and -B (Liu et al., 2009). CB₂B, the first cloned cDNA, is transcribed from a promoter proximal to exon2

FIG. 1. CB₁R and CB₂R gene structures, promoters, and alternative transcripts. Transcripts are depicted below the complete gene structures and are not drawn to scale. Data from Zhang et al. (2004), human CB₁R; McCaw et al. (2004), mouse CB₁R; Matsuda et al. (1990), rat CB₁R; Liu et al. (2009), human, rat, and mouse CB₂R.

and expressed most highly in immune cells and tissues. The more recently identified CB₂A contains exon1 and exon3 and is generated from a promoter 5' proximal to exon1. In contrast to CB₂B, CB₂A is most highly expressed in testis and shows some expression in the brain.

Mouse CB₂R similarly consists of three exons alternatively transcribed by two promoters (Onaivi et al., 2006; Liu et al., 2009). In contrast to humans, however, both CB₂A and CB₂B are expressed predominantly in the spleen (Liu et al., 2009). Rat CB₂R gene structure seems more complex than that of human and mouse, although findings are conflicting. In Fig. 1, we present findings by

Liu et al. (2009) of CB₂A and -B isoforms transcribed from promoters flanking exon 1 and 2, like human and mouse CB₂R. Rat CB₂A and -B also showed expression patterns similar to those of mouse. In contrast, Brown et al. (2002) identified three coding exons in rat CB₂R using the same species and tissue (spleen).

Like CB₁R, human and rodent CB₂R differ in length of 5'UTR, with human CB₂R again being considerably longer. Moreover, the CB₂R 5'UTR showed little alignment between human-rodent (although rat-mouse 5'UTR aligned well, and there was significant homology in coding region across all species) (Liu et al., 2009). The lack of

TABLE 3

Transcription factors found to interact with CB₁R and CB₂R promoters using high-throughput ChIP (ChIP-Chip or ChIP-Seq) experiments

The selection of studies was found through the Chromatin Enrichment Analysis database (ChEA) (<http://amp.pharm.mssm.edu/lib/chea.jsp>) (Lachmann et al., 2010). Transcription factor functions are adapted from MetaCore database (<http://www.genego.com/metacore.php>).

Transcription Factor	References	Cell line	Function
CB ₁ R			
AR (androgen receptor)	Lin et al., 2009	Human prostate cancer PC3 cells	Activates transcription of androgen responsive genes to affect cell proliferation and differentiation.
SUZ12 (suppressor of zeste 12)	Boyer et al., 2006; Ku et al., 2008; Marson et al., 2008	Mouse ES cells	Component of the PRC2/EED-EZH2 complex, which represses target genes by methylating lysine residues on histone H3. Complex may also repress transcription by recruiting DNA methyltransferases.
EED (embryonic ectoderm development)	Boyer et al., 2006	Mouse ES cells	Also a component of the PRC2/EED-EZH2 complex (see above).
CREM (cAMP responsive element modulator)	Martianov et al., 2010	Mouse germ cells	Binds the CRE; isoforms are either activators or repressors.
JARID2 (jumonji, AT rich interactive domain 2)	Peng et al., 2009; Pasini et al., 2010	Mouse ES cells	Regulates histone methyltransferase complexes.
REST/NRSF (RE1-silencing transcription factor/ neuron restrictive silencer factor)	Abrajano et al., 2009	Mouse neurons	Binds NRSEs and represses neuronal gene expression via recruitment of the BHC (see below).
RCor1/CoREST (REST corepressor 1)	Abrajano et al., 2009	Mouse neurons	Component of the BHC that is recruited to NRSE sites by REST, where it deacetylates and demethylates histones and thus represses neuronal gene expression.
RNF2 (ring finger protein 2)	Boyer et al., 2006	Mouse ES cells	E3 ubiquitin-protein ligase that regulates monoubiquitination of histone H2A lysine residues, a repressive mark.
CUX-1 (cut-like homeobox 1)	Kedinger et al., 2009	Several human cancer cell lines	Part of the homeodomain family of DNA binding proteins; may regulate differentiation and cell cycle progression.
CB ₂ R			
EP300 (E1A binding protein p300)	Blow et al., 2010	Mouse embryonic heart tissue	Binds phosphorylated CREB and functions as a histone acetyltransferase. Important in cell proliferation and differentiation, and thought to have a role in the stimulation of hypoxia-induced genes.
ERG (v-ets erythroblastosis virus E26 oncogene homolog)	Wilson et al., 2010	Mouse HPC7 hematopoietic progenitor cells	Recruits SETDB1 histone methyltransferase to target genes.
STAT3 (signal transducer and activator of transcription 3)	Kwon et al., 2009	Mouse CD4 Ts	Activated by phosphorylation upon cell stimulation by cytokines and growth factors. Has broad functions, including cell growth and apoptosis.
GATA3 (GATA binding protein 3)	Kidder and Palmer, 2010	Mouse trophoblast stem cells	Regulates T-cell development.

CRE, cAMP response element; NRSE, neuron-restrictive silencer elements; BHC, BRAF-HDAC complex; REST, RE1-silencing transcription factor; CREB, cAMP response element-binding protein.

significant homology in 5'UTR and different tissue distribution of CB₂A and CB₂B isoforms in humans versus rodents (with the 2A isoform appearing in the brain only in humans) suggests differential CB₂R promoter regulation across species. Such differences in rodent and human CB₂R isoform expression should also be kept in mind when interpreting results from animal models. For example, the human CB₂A isoform could produce unwanted central nervous system effects because of its low expression in the brain; these effects would not be seen in mice (for review, see Campbell et al., 2001; Liu et al., 2009).

The production of these alternative CB₁R and CB₂R transcripts differing in 5'UTR in disease is worth investigating. These transcripts have already been shown to have region-specific expression; they could also differ in mRNA stability, subcellular localization and translational efficacies (see references in Zhang et al., 2004). The importance of the CB₁R 5'UTR in disease is suggested by the presence of single-nucleotide polymorphisms in 5' UTR introns and exons (TAG haplotype) that are associated with lower mRNA levels and with substance abuse (Zhang et al., 2004). It will also be important to identify the use of alternative promoters in disease when studying direct mechanisms of cannabinoid receptor regulation.

V. High-Throughput Studies of Transcription Factor Interactions with Cannabinoid Receptors 1 and 2

In addition to the directed studies of cannabinoid receptor promoter regulation (Table 2), high-throughput chromatin immunoprecipitation (ChIP) studies using various cell lines have identified a number of transcription factors that interact with the CB₁R and CB₂R promoters. We used the Chromatin Enrichment Analysis database of such ChIP experiments (Lachmann et al., 2010) to compile a list of transcription factors that interact with CB₁R and CB₂R (Table 3). Many of these transcription factors are implicated in DNA methylation and histone post-translational modifications, further supporting a role for epigenetic mechanisms in cannabinoid receptor regulation. These studies vary in their methods of defining target genes; often, regulatory regions are assigned based on a given distance (e.g., 50 kb) from a locus. Moreover, the binding of a transcription factor does not always mean that it has functional effects on transcription at a given locus (Chen et al., 2008).

In addition to identifying transcription factors that interact with the cannabinoid receptor promoters through such ChIP studies, identifying transcription factors downstream of cannabinoid receptor signaling will be important for further studies of cannabinoid receptor autoregulation. Along these lines, Bromberg et al. (2008) identified 33 transcription factors activated in Neuro2A neuroblastoma cells after stimulation with cannabinoid receptor agonist using an array of tran-

scription factor-binding site oligonucleotides. Taken together, transcription factors identified by these high-throughput studies can be starting points for further directed investigations of cannabinoid receptor regulation in disease.

VI. Conclusions and Future Directions

Alteration in cannabinoid receptor expression is appearing more and more to be a widespread response to disease. As investigation of the mechanisms underlying these changes continues, we will probably begin to see commonalities across diseases. Cytokines, growth factors, hormones, and other factors released in response to tissue injury and inflammation are thus rational starting points for further investigations of mechanism. Autoregulation of cannabinoid receptor expression will also likely be identified in other models. Findings of specific signaling pathways and transcription factors will hopefully be accompanied by identification of the epigenetic modifications ultimately underlying these expression changes.

Understanding these mechanisms of cannabinoid receptor regulation will hopefully expand our options for therapeutically targeting the endocannabinoid system. Selectively enhancing receptor expression could allow for lower doses of systemic agonists or eliminate the need for them altogether; selectively inhibiting expression could likewise avoid systemic antagonism. These improved endocannabinoid system-based therapies could have a lot to offer medicine.

Acknowledgments

This work was supported by National Institutes of Health National Institute on Drug Abuse [Grants DA08862, DA19521] (to L.A.D.) and National Institutes of Health National Institute General of General Medicine [Grant 1P50-GM071558-01A27398] (to L.A.D. and L.M.). We thank Ittai Bushlin, Maribel Lim, Dr. Raphael Rozenfeld, and Dr. Cristina Costantino for discussion and critical reading of the manuscript, and Alex Lachmann for help with Chromatin Enrichment Analysis database.

Authorship Contributions

Wrote or contributed to the writing of the manuscript: Miller and Devi.

References

- Abrajano JJ, Qureshi IA, Gokhan S, Zheng D, Bergman A, and Mehler MF (2009) REST and CoREST modulate neuronal subtype specification, maturation and maintenance. *PLoS One* 4:e7936.
- Bátkai S, Pacher P, Osei-Hyiaman D, Radaeva S, Liu J, Harvey-White J, Offertáler L, Mackie K, Rudd MA, Bukoski RD, et al. (2004) Endocannabinoids acting at cannabinoid-1 receptors regulate cardiovascular function in hypertension. *Circulation* 110:1996–2002.
- Bensaid M, Gary-Bobo M, Esclançon A, Maffrand JP, Le Fur G, Oury-Donat F, and Soubrié P (2003) The cannabinoid CB1 receptor antagonist SR141716 increases Acip30 mRNA expression in adipose tissue of obese fa/fa rats and in cultured adipocyte cells. *Mol Pharmacol* 63:908–914.
- Blow MJ, McCulley DJ, Li Z, Zhang T, Akiyama JA, Holt A, Plajzer-Frick I, Shoukry M, Wright C, Chen F, et al. (2010) ChIP-Seq identification of weakly conserved heart enhancers. *Nat Genet* 42:806–810.
- Börner C, Bedini A, Höllt V, and Kraus J (2008) Analysis of promoter regions regulating basal and interleukin-4-inducible expression of the human CB1 receptor gene in T lymphocytes. *Mol Pharmacol* 73:1013–1019.
- Börner C, Höllt V, Sebald W, and Kraus J (2007) Transcriptional regulation of the cannabinoid receptor type 1 gene in T cells by cannabinoids. *J Leukoc Biol* 81: 336–343.

- Boyer LA, Plath K, Zeitlinger J, Brambrink T, Medeiros LA, Lee TI, Levine SS, Wernig M, Tajonar A, Ray MK, et al. (2006) Polycomb complexes repress developmental regulators in murine embryonic stem cells. *Nature* **441**:349–353.
- Bromberg KD, Ma'ayan A, Neves SR, and Iyengar R (2008) Design logic of a cannabinoid receptor signaling network that triggers neurite outgrowth. *Science* **320**:903–909.
- Brown SM, Wager-Miller J, and Mackie K (2002) Cloning and molecular characterization of the rat CB₂ cannabinoid receptor. *Biochim Biophys Acta* **1576**:255–264.
- Cabral GA and Griffin-Thomas L (2009) Emerging role of the cannabinoid receptor CB₂ in immune regulation: therapeutic prospects for neuroinflammation. *Expert Rev Mol Med* **11**:e3.
- Campbell FA, Tramèr MR, Carroll D, Reynolds DJ, Moore RA, and McQuay HJ (2001) Are cannabinoids an effective and safe treatment option in the management of pain? A qualitative systematic review. *BMJ* **323**:13–16.
- Chen X, Xu H, Yuan P, Fang F, Huss M, Vega VB, Wong E, Orlov YL, Zhang W, Jiang J, et al. (2008) Integration of external signaling pathways with the core transcriptional network in embryonic stem cells. *Cell* **133**:1106–1117.
- Chiu CQ, Puente N, Grandes P, and Castillo PE (2010) Dopaminergic modulation of endocannabinoid-mediated plasticity at GABAergic synapses in the prefrontal cortex. *J Neurosci* **30**:7236–7248.
- Costigan M, Scholz J, and Woolf CJ (2009) Neuropathic pain: a maladaptive response of the nervous system to damage. *Annu Rev Neurosci* **32**:1–32.
- Dean B, Sundram S, Bradbury R, Scarr E, and Copolov D (2001) Studies on [³H]CP-55940 binding in the human central nervous system: regional specific changes in density of cannabinoid-1 receptors associated with schizophrenia and cannabis use. *Neuroscience* **103**:9–15.
- Di Marzo V (2008) Targeting the endocannabinoid system: to enhance or reduce? *Nat Rev Drug Discov* **7**:438–455.
- Di Marzo V, Bifulco M, and De Petrocellis L (2004) The endocannabinoid system and its therapeutic exploitation. *Nat Rev Drug Discov* **3**:771–784.
- Ha SO, Kim JK, Hong HS, Kim DS, and Cho HJ (2001) Expression of brain-derived neurotrophic factor in rat dorsal root ganglia, spinal cord and gracile nuclei in experimental models of neuropathic pain. *Neuroscience* **107**:301–309.
- Hansen HH, Schmid PC, Bittigau P, Lastres-Becker I, Berrendero F, Manzanares J, Ikonomidou C, Schmid HH, Fernández-Ruiz JJ, and Hansen HS (2001) Anandamide, but not 2-arachidonoylglycerol, accumulates during in vivo neurodegeneration. *J Neurochem* **78**:1415–1427.
- Howlett AC, Barth F, Bonner TI, Cabral G, Casellas P, Devane WA, Felder CC, Herkenham M, Mackie K, Martin BR, et al. (2002) International Union of Pharmacology. XXVII. Classification of cannabinoid receptors. *Pharmacol Rev* **54**:161–202.
- Huang EJ and Reichardt LF (2001) Neurotrophins: roles in neuronal development and function. *Annu Rev Neurosci* **24**:677–736.
- Hungund BL, Vinod KY, Kassir SA, Basavarajappa BS, Yalamanchili R, Cooper TB, Mann JJ, and Arango V (2004) Upregulation of CB₁ receptors and agonist-stimulated [³⁵S]GTPgammaS binding in the prefrontal cortex of depressed suicide victims. *Mol Psychiatry* **9**:184–190.
- Izzo AA and Camilleri M (2008) Emerging role of cannabinoids in gastrointestinal and liver diseases: basic and clinical aspects. *Gut* **57**:1140–1155.
- Izzo AA, Fezza F, Capasso R, Bisogno T, Pinto L, Iuvone T, Esposito G, Mascolo N, Di Marzo V, and Capasso F (2001) Cannabinoid CB₁-receptor mediated regulation of gastrointestinal motility in mice in a model of intestinal inflammation. *Br J Pharmacol* **134**:563–570.
- Jeong WI, Osei-Hyiaman D, Park O, Liu J, Bátkai S, Mukhopadhyay P, Horiguchi N, Harvey-White J, Marsicano G, Lutz B, et al. (2008) Paracrine activation of hepatic CB₁ receptors by stellate cell-derived endocannabinoids mediates alcoholic fatty liver. *Cell Metab* **7**:227–235.
- Jourdan T, Djaouti L, Demizieux L, Gresti J, Vergès B, and Degrace P (2010) CB₁ antagonism exerts specific molecular effects on visceral and subcutaneous fat and reverses liver steatosis in diet-induced obese mice. *Diabetes* **59**:926–934.
- Julien B, Grenard P, Teixeira-Clerc F, Van Nhieu JT, Li L, Karsak M, Zimmer A, Mallat A, and Lotersztajn S (2005) Antiinflammatory role of the cannabinoid receptor CB₂ in the liver. *Gastroenterology* **128**:742–755.
- Kedinger V, Sansregret L, Harada R, Vadnais C, Cadieux C, Fathers K, Park M, and Nepveu A (2009) p110 CUX1 homeodomain protein stimulates cell migration and invasion in part through a regulatory cascade culminating in the repression of E-cadherin and occludin. *J Biol Chem* **284**:27701–27711.
- Kidder BL and Palmer S (2010) Examination of transcriptional networks reveals an important role for TCFAP2C, SMARCA4, and EOMES in trophoblast stem cell maintenance. *Genome Res* **20**:458–472.
- Kim KW, Jo YH, Zhao L, Stallings NR, Chua SC Jr, and Parker KL (2008) Steroidogenic factor 1 regulates expression of the cannabinoid receptor 1 in the ventromedial hypothalamic nucleus. *Mol Endocrinol* **22**:1950–1961.
- Kimball ES, Schneider CR, Wallace NH, and Hornby PJ (2006) Agonists of cannabinoid receptor 1 and 2 inhibit experimental colitis induced by oil of mustard and by dextran sulfate sodium. *Am J Physiol Gastrointest Liver Physiol* **291**:G364–G371.
- Ku M, Koche RP, Rheinbay E, Mendenhall EM, Endoh M, Mikkelsen TS, Presser A, Nusbaum C, Xie X, Chi AS, et al. (2008) Genomewide analysis of PRC1 and PRC2 occupancy identifies two classes of bivalent domains. *PLoS Genet* **4**:e1000242.
- Kunos G, Osei-Hyiaman D, Liu J, Godlewski G, and Bátkai S (2008) Endocannabinoids and the control of energy homeostasis. *J Biol Chem* **283**:33021–33025.
- Kwon H, Thierry-Mieg D, Thierry-Mieg J, Kim HP, Oh J, Tunyaplin C, Carotta S, Donovan CE, Goldman ML, Tailor P, et al. (2009) Analysis of interleukin-21-induced Prdm1 gene regulation reveals functional cooperation of STAT3 and IRF4 transcription factors. *Immunity* **31**:941–952.
- Lachmann A, Xu H, Krishnan J, Berger SI, Mazloom AR, and Ma'ayan A (2010) ChEA: transcription factor regulation inferred from integrating genome-wide ChIP-X experiments. *Bioinformatics* **26**:2438–2444.
- Lafourcade M, Elezgarai I, Mato S, Bakiri Y, Grandes P, and Manzoni OJ (2007) Molecular components and functions of the endocannabinoid system in mouse prefrontal cortex. *PLoS One* **2**:e709.
- Lim G, Sung B, Ji RR, and Mao J (2003) Upregulation of spinal cannabinoid-1 receptors following nerve injury enhances the effects of Win 55,212–2 on neuropathic pain behaviors in rats. *Pain* **105**:275–283.
- Lin B, Wang J, Hong X, Yan X, Hwang D, Cho JH, Yi D, Utleg AG, Fang X, Schones DE, et al. (2009) Integrated expression profiling and ChIP-seq analyses of the growth inhibition response program of the androgen receptor. *PLoS One* **4**:e6589.
- Liu QR, Pan CH, Hishimoto A, Li CY, Xi ZX, Llorente-Berzal A, Viveros MP, Ishiguro H, Arinami T, Onaivi ES, et al. (2009) Species differences in cannabinoid receptor 2 (CNR2 gene): identification of novel human and rodent CB₂ isoforms, differential tissue expression and regulation by cannabinoid receptor ligands. *Genes Brain Behav* **8**:519–530.
- Maden M (2007) Retinoic acid in the development, regeneration and maintenance of the nervous system. *Nat Rev Neurosci* **8**:755–765.
- Maresz K, Carrier EJ, Ponomarev ED, Hillard CJ, and Dittel BN (2005) Modulation of the cannabinoid CB₂ receptor in microglial cells in response to inflammatory stimuli. *J Neurochem* **95**:437–445.
- Maresz K, Pryce G, Ponomarev ED, Marsicano G, Croxford JL, Shriver LP, Ledent C, Cheng X, Carrier EJ, Mann MK, et al. (2007) Direct suppression of CNS autoimmune inflammation via the cannabinoid receptor CB₁ on neurons and CB₂ on autoreactive T cells. *Nat Med* **13**:492–497.
- Marson A, Levine SS, Cole MF, Frampton GM, Brambrink T, Johnstone S, Guenther MG, Johnston WK, Wernig M, Newman J, et al. (2008) Connecting microRNA genes to the core transcriptional regulatory circuitry of embryonic stem cells. *Cell* **134**:521–533.
- Martianov I, Choukrallah MA, Krebs A, Ye T, Legras S, Rijkers E, Van Ijcken W, Jost B, Sassone-Corsi P, and Davidson I (2010) Cell-specific occupancy of an extended repertoire of CREM and CREB binding loci in male germ cells. *BMC Genomics* **11**:530.
- Massa F, Marsicano G, Hermann H, Cannich A, Monory K, Cravatt BF, Ferri GL, Sibaev A, Storr M, and Lutz B (2004) The endogenous cannabinoid system protects against colonic inflammation. *J Clin Invest* **113**:1202–1209.
- Matias I, Gonthier MP, Orlando P, Martiadis V, De Petrocellis L, Cervino C, Petrosino S, Hoareau L, Festy F, Pasquali R, et al. (2006) Regulation, function, and dysregulation of endocannabinoids in models of adipose and beta-pancreatic cells and in obesity and hyperglycemia. *J Clin Endocrinol Metab* **91**:3171–3180.
- Matsuda LA, Lolait SJ, Brownstein MJ, Young AC, and Bonner TI (1990) Structure of a cannabinoid receptor and functional expression of the cloned cDNA. *Nature* **346**:561–564.
- McCaw EA, Hu H, Gomez GT, Hebb AL, Kelly ME, and Donovan-Wright EM (2004) Structure, expression and regulation of the cannabinoid receptor gene (CB₁) in Huntington's disease transgenic mice. *Eur J Biochem* **271**:4909–4920.
- McKallip RJ, Nagarkatti M, and Nagarkatti PS (2005) Delta-9-tetrahydrocannabinol enhances breast cancer growth and metastasis by suppression of the antitumor immune response. *J Immunol* **174**:3281–3289.
- Mechoulam R, Spatz M, and Shohami E (2002) Endocannabinoids and neuroprotection. *Sci STKE* **2002**:re5.
- Mitrirattanakul S, Ramakul N, Guerrero AV, Matsuka Y, Ono T, Iwase H, Mackie K, Faull KF, and Spigelman I (2006) Site-specific increases in peripheral cannabinoid receptors and their endogenous ligands in a model of neuropathic pain. *Pain* **126**:102–114.
- Molina-Holgado F, Pinteaux E, Moore JD, Molina-Holgado E, Guaza C, Gibson RM, and Rothwell NJ (2003) Endogenous interleukin-1 receptor antagonist mediates anti-inflammatory and neuroprotective actions of cannabinoids in neurons and glia. *J Neurosci* **23**:6470–6474.
- Montone M, Martorelli D, Rosato A, and Dolcetti R (2009) Retinoids as critical modulators of immune functions: new therapeutic perspectives for old compounds. *Endocr Metab Immune Disord Drug Targets* **9**:113–131.
- Mukhopadhyay B, Liu J, Osei-Hyiaman D, Godlewski G, Mukhopadhyay P, Wang L, Jeong WI, Gao B, Duester G, Mackie K, et al. (2010) Transcriptional regulation of cannabinoid receptor-1 expression in the liver by retinoic acid acting via retinoic acid receptor-gamma. *J Biol Chem* **285**:19002–19011.
- Nakajima Y, Furuichi Y, Biswas KK, Hashiguchi T, Kawahara K, Yamaji K, Uchimura T, Izumi Y, and Maruyama I (2006) Endocannabinoid, anandamide in gingival tissue regulates the periodontal inflammation through NF-kappaB pathway inhibition. *FEBS Lett* **580**:613–619.
- Noy N (2010) Between death and survival: retinoic acid in regulation of apoptosis. *Annu Rev Nutr* **30**:201–217.
- Onaivi ES, Ishiguro H, Gong JP, Patel S, Perchuk A, Meozzi PA, Myers L, Mora Z, Tagliaferro P, Gardner E, et al. (2006) Discovery of the presence and functional expression of cannabinoid CB₂ receptors in brain. *Ann NY Acad Sci* **1074**:514–536.
- Osei-Hyiaman D, DePetrillo M, Pacher P, Liu J, Radaeva S, Bátkai S, Harvey-White J, Mackie K, Offertaler L, Wang L, et al. (2005) Endocannabinoid activation at hepatic CB₁ receptors stimulates fatty acid synthesis and contributes to diet-induced obesity. *J Clin Invest* **115**:1298–1305.
- Pacher P, Bátkai S, and Kunos G (2006) The endocannabinoid system as an emerging target of pharmacotherapy. *Pharmacol Rev* **58**:389–462.
- Pagotto U, Marsicano G, Cota D, Lutz B, and Pasquali R (2006) The emerging role of the endocannabinoid system in endocrine regulation and energy balance. *Endocr Rev* **27**:73–100.
- Pasini D, Cloos PA, Walfridsson J, Olsson L, Bukowski JP, Johansen JV, Bak M, Tommerup N, Rappsilber J, and Helin K (2010) JARID2 regulates binding of the Polycomb repressive complex 2 to target genes in ES cells. *Nature* **464**:306–310.
- Peng JC, Valouev A, Swigut T, Zhang J, Zhao Y, Sidow A, and Wysocka J (2009) Jarid2/Jumonji coordinates control of PRC2 enzymatic activity and target gene occupancy in pluripotent cells. *Cell* **139**:1290–1302.
- Pertwee RG (2005) The therapeutic potential of drugs that target cannabinoid receptors or modulate the tissue levels or actions of endocannabinoids. *AAPS J* **7**:E625–E654.

- Pertwee RG (2009) Emerging strategies for exploiting cannabinoid receptor agonists as medicines. *Br J Pharmacol* **156**:397–411.
- Quarta C, Bellocchio L, Mancini G, Mazza R, Cervino C, Brulke LJ, Fekete C, Latorre R, Nanni C, Bucci M, et al. (2010) CB(1) signaling in forebrain and sympathetic neurons is a key determinant of endocannabinoid actions on energy balance. *Cell Metab* **11**:273–285.
- Sarfaraz S, Adhami VM, Syed DN, Afaq F, and Mukhtar H (2008) Cannabinoids for cancer treatment: progress and promise. *Cancer Res* **68**:339–342.
- Sarfaraz S, Afaq F, Adhami VM, and Mukhtar H (2005) Cannabinoid receptor as a novel target for the treatment of prostate cancer. *Cancer Res* **65**:1635–1641.
- Siegling A, Hofmann HA, Denzer D, Mauler F, and De Vry J (2001) Cannabinoid CB(1) receptor upregulation in a rat model of chronic neuropathic pain. *Eur J Pharmacol* **415**:R5–R7.
- Steffens S, Veillard NR, Arnaud C, Pelli G, Burger F, Staub C, Karsak M, Zimmer A, Frossard JL, and Mach F (2005) Low dose oral cannabinoid therapy reduces progression of atherosclerosis in mice. *Nature* **434**:782–786.
- Teixeira-Clerc F, Julien B, Grenard P, Tran Van Nhieu J, Deveaux V, Li L, Serriere-Lanneau V, Ledent C, Mallat A, and Lotersztajn S (2006) CB1 cannabinoid receptor antagonism: a new strategy for the treatment of liver fibrosis. *Nat Med* **12**:671–676.
- Topol EJ, Bousser MG, Fox KA, Creager MA, Despres JP, Easton JD, Hamm CW, Montalescot G, Steg PG, Pearson TA, et al. (2010) Rimobabant for prevention of cardiovascular events (CRESCENDO): a randomised, multicentre, placebo-controlled trial. *Lancet* **376**:517–523.
- Turu G and Hunyady L (2010) Signal transduction of the CB1 cannabinoid receptor. *J Mol Endocrinol* **44**:75–85.
- Vickers SP, Webster LJ, Wyatt A, Dourish CT, and Kennett GA (2003) Preferential effects of the cannabinoid CB1 receptor antagonist, SR 141716, on food intake and body weight gain of obese (fa/fa) compared to lean Zucker rats. *Psychopharmacology (Berl)* **167**:103–111.
- Walczak JS, Pichette V, Leblond F, Desbiens K, and Beaulieu P (2005) Behavioral, pharmacological and molecular characterization of the saphenous nerve partial ligation: a new model of neuropathic pain. *Neuroscience* **132**:1093–1102.
- Wang D, Wang H, Ning W, Backlund MG, Dey SK, and DuBois RN (2008) Loss of cannabinoid receptor 1 accelerates intestinal tumor growth. *Cancer Res* **68**:6468–6476.
- Wang S, Lim G, Mao J, Sung B, Yang L, and Mao J (2007) Central glucocorticoid receptors regulate the upregulation of spinal cannabinoid-1 receptors after peripheral nerve injury in rats. *Pain* **131**:96–105.
- Wang S, Lim G, Zeng Q, Sung B, Ai Y, Guo G, Yang L, and Mao J (2004) Expression of central glucocorticoid receptors after peripheral nerve injury contributes to neuropathic pain behaviors in rats. *J Neurosci* **24**:8595–8605.
- Wilson NK, Foster SD, Wang X, Knezevic K, Schütte J, Kaimakis P, Chilarska PM, Kinston S, Ouwehand WH, Dzierzak E, et al. (2010) Combinatorial transcriptional control in blood stem/progenitor cells: genome-wide analysis of ten major transcriptional regulators. *Cell Stem Cell* **7**:532–544.
- Yiangou Y, Facer P, Durrenberger P, Chessell IP, Naylor A, Bountra C, Banati RR, and Anand P (2006) COX-2, CB2 and P2X7-immunoreactivities are increased in activated microglial cells/macrophages of multiple sclerosis and amyotrophic lateral sclerosis spinal cord. *BMC Neurol* **6**:12.
- Zhang J, Hoffert C, Vu HK, Groblewski T, Ahmad S, and O'Donnell D (2003) Induction of CB2 receptor expression in the rat spinal cord of neuropathic but not inflammatory chronic pain models. *Eur J Neurosci* **17**:2750–2754.
- Zhang PW, Ishiguro H, Ohtsuki T, Hess J, Carillo F, Walther D, Onaivi ES, Arinami T, and Uhl GR (2004) Human cannabinoid receptor 1: 5' exons, candidate regulatory regions, polymorphisms, haplotypes and association with polysubstance abuse. *Mol Psychiatry* **9**:916–931.