Published in final edited form as:

Eur Arch Paediatr Dent. 2008 March ; 9(1): 19–24.

Inheritance of Occlusal Topography: A Twin Study

C-Y. Su^{*}, P.M. Corby^{**}, M.A. Elliot^{*}, D.A. Studen-Pavlovich^{*}, D.N. Ranalli^{*}, B. Rosa^{***}, J. Wessel^{††}, N.J. Schork^{††}, T.C. Hart[†], and W.A. Bretz^{**}

^{*}Dept. Pediatric Dentistry, School of Dental Medicine, University of Pittsburgh Pittsburgh, PA

^{**}Departments of Cariology & Comprehensive Care, Periodontics and Implants, New York University College of Dentistry, New York, NY

[†]NIH/NIDCR, Division of Intramural Research, Bethesda, MD

^{††}Scripps Genomic Medicine and The Scripps Research Institute, San Diego, CA, USA

^{***}UNIMONTES, School of Dentistry, Department of Prosthodontics, Montes Claros, MG, Brazil

Abstract

Aim—This was to determine the relative contribution of genetic factors on the morphology of occlusal surfaces of mandibular primary first molars by employing the twin study model.

Methods—The occlusal morphology of mandibular primary first molar teeth from dental casts of 9 monozygotic (MZ) twin pairs and 12 dizygotic (DZ) twin pairs 4 to 7 years old, were digitized by contact-type three-dimensional (3D) scanner. To compare the similarity of occlusal morphology between twin sets, each twin pair of occlusal surfaces was superimposed to establish the best fit by using computerized least squared techniques. Heritability was computed using a variance component model, adjusted for age and gender.

Results—DZ pairs demonstrated a greater degree of occlusal morphology variance. The total amount of difference in surface overlap was 0.0508 mm (0.0018 inches) for the MZ (n=18) sample and 0.095 mm (0.0034 inches) for the DZ (n=24) sample and were not statistically significant (p=0.2203). The transformed mean differences were not statistically significantly different (p=0.2203). Heritability estimates of occlusal surface areas for right and left mandibular primary first molars were 97.5% and 98.2% (p<0.0001), respectively.

Conclusions—Occlusal morphology of DZ twin pairs was more variable than that of MZ twin pairs. Heritability estimates revealed that genetic factors strongly influence occlusal morphology of mandibular primary first molars.

Keywords

twins; occlusal anatomy; heritability; first primary mandibular molar

Introduction

Human teeth develop from tissues of both ectoderm and mesoderm in the sixth week of embryonic life [Pinkham et al., 1999]. Growth of the tooth germ proceeds after the appearance of the basal cell layer of the oral epithelium, and is followed by stages of proliferation, histodifferentiation, morphodifferentiation, and final apposition to calcification. Tooth size and shape occurs at the morphodifferentiation stage of the tooth

development which takes place at approximately the 18th week of embryonic development for permanent teeth [Pinkham et al., 1999]. Occlusal morphology is determined by temporal and spatial patterning events that include laying down an organic matrix that is subsequently reabsorbed and mineralized. Such morphodifferentiation is clearly under genetic control, but the extent to which environmental factors influence the process is unknown. Different teeth have a characteristic occlusal topography, and the variance in this topography can be measured and used to determine the relative similarity between individuals.

Twin studies offer the opportunity to dissect the relative contribution of genetic and environmental factors on a particular trait because MZ (identical) twins are 100% genetically similar whereas DZ twins are 50% genetically similar. If a particular trait is correlated highly between MZ twins and less correlated between DZ twins, it is reasonable to assume a genetic contribution to variation on that particular trait. Inheritability of a trait can be determined with heritability estimates by employing statistical models that quantify the degree of genetic versus environmental influences [Dempsey and Townsend, 2001].

A few twin studies have examined the role of genetics on tooth anatomy. In a study of 34 pairs of twins, it was shown that the mesiodistal dimension for all permanent teeth in MZ twins was more concordant than in DZ twins [Kabban et al., 2001]. Studies by Townsend and collaborators indicated a high phenotypic variation of intercuspal distance with moderate genetic influence [Townsend et al., 2003]. Morphological features such as the Carabelli's trait were found to have higher concordance in MZ twins than in dizygotic twins [Biggerstaff, 1973; Boraas et al., 1988]. Only one study has provided detailed mappings of the occlusal surface topography of first permanent molars of identical and fraternal twins and of singletons [Kabban et al., 2001]. That study, however, only examined 9 pairs of maxillary first molars and concluded that a significant genetic contribution to occlusal topography was apparent.

The purpose of the present study was to determine the relative contribution of genetic and environmental factors on the morphology of occlusal surfaces of first primary mandibular molars by employing the twin study model.

Material and Methods

Study Population

Twins were ascertained from a government-based health service registry in the city of Montes Claros, Brazil. Twin families were of a predominantly low socioeconomic status. The twin pairs participating in this study were in the range of 4 to 7 years old. The research protocol was approved by Institutional Review Boards of the University of Pittsburgh and of UNIMONTES, Montes Claros, Brazil.

Zygosity

DNA was extracted from peripheral venous blood samples. DNA marker loci were PCR amplified using standard methods. Amplification products were detected by blood kits (QIAamp, Qiagen, CA, USA) with an ABI-377 fluorescent sequencer and analyzed by GENESCAN 2.1 (Applied Biosystems, CA, USA) [Zang et al., 2001]. Zygosity was determined by genotyping all twins and their parents for 8 highly polymorphic DNA loci (on chromosomes 2, 7, 11, 17 and 20). Individuals discordant for one or more markers were considered dizygotic.

Dental Impressions/Casts

Dental impressions were obtained from study participants from mandibular and maxillary arches. The impressions were made using medium-body silicone impression material (3M-Espe, Selfeld, Germany). On the same day, the impressions were poured-up with a high precision dental plaster (BPB Formula, Suresnes, France). For the purpose of this study, for each twin set, only primary molars without occlusal caries were included in the analyses.

Occlusal Surfaces Available for Analysis

Based on the caries status of each participant (caries-free mandibular first primary molar), a total of 46 MZ twin pairs and 78 DZ twin pairs, and one DZ triplet were available for study analyses. The dental casts generated for each of these individuals were examined for surface defects such as air bubbles and/or fractured cusps which would impair our ability to generate accurate occlusal topographies. A total of 18 MZ (9 twin pairs) and 24 DZ (12 DZ pairs) casts were free of surface defects for both left and right mandibular first primary molars. Gender distribution of MZ twins included 6 pairs of females and 3 pairs of males and for DZ twins 7 same gender pairs and 5 pairs of discordant gender twins. Then the similarity of the occlusal morphology of mandibular first primary molars in these twin pairs was examined utilizing 3-D coordinate metrology.

3-D Coordinate Metrology

Prior to data collection, feasibility studies were conducted to determine resolution needs and scanning times. The time necessary for scanning one tooth was determined to be 30 minutes.

The mandibular first primary molar of each dental cast was digitized by a contact-type (touch probe) 3D scanner (PICZA, Roland, Japan) (Figure 1). The x pitch of the digitizer was set to 0.1524 mm, (0.006 inches) and the y pitch was also set to 0.1524 mm (0.006 inches). The upper z limit for scanning was set at the plane of 0.5 mm which was higher than the most upper height of the occlusal surface of a particular tooth. The optimum setting of the upper z limit allowed for a more efficient scanning process. Dr. PICZA software was utilized to control the PICZA scanner and to provide an interface to a computer database. After the images were obtained and stored in the computer, the images were exported as data point clouds for further analysis.

Occlusal Surface 3-D Analysis

To compare the similarity of the occlusal morphology between twin sets, each pair of occlusal surfaces was superimposed to establish the “best fit” using a least-squared technique. The task was accomplished by the function “manual register” of the software Geomagic Qualify 7(Raindrop Geomagic Inc., Durham, NC). The two superimposed images were then analyzed using the function of “3D comparison.” The “3D comparison” produced a colour scheme that demonstrated the deviations of occlusal surfaces’ topography as reflected by different designated colours (Figure 2a). Figure 2a is an example of a superimposed image of MZ mandibular molars. The results of 3D comparison also provided the values of standard deviation, average distance, and the percentage of outliers. Outliers were points that were considered to be too deviant from any neighbouring points to have any relevance to the segment being scanned (Figure 2b). The average distance provided by the 3D comparison quantifies the total difference between two overlapping surfaces. Figure 2b is a typical example of DZ superimposed images.

Total Surface Area Analysis

The assessment of the total surface area of the occlusal surface was analyzed using the surfacing software package Magics V6.3 (Materialise, Ann Arbor, MI). Each pair of the

registered images were superimposed to the best fit, and trimmed to the exact elliptical size using the Qualify software program. Each image was then imported under a STL file into the Magics program. The Magics program calculated the value that measured the total surface area produced by each inner cusp incline, groove and fossa (Figure 3).

Statistical Analysis

Frequency distributions, paired t-tests, and independent t-tests were computed using the JUMP software (SAS, Cary, NC) to assess occlusal morphology. The 3D average distance was squared (transformed) to account for positive and negative values canceling each other out. The transformed 3D comparison average distance data were used in the analysis. Paired t-tests were used to assess intra-examiner reliability for occlusal morphology comparisons in ten randomly selected twin paired samples that were analyzed at baseline and re-analyzed after a one month interval.

Heritability estimates were computed for surface area of mandibular first primary molars by using a variance component model implemented in the Sequential Oligogenic Linkage Analysis Routines (SOLAR) computer analysis package (AAVSO, Cambridge, MA), [Almasy and Blangero, 1998]. Heritability was defined in its narrow sense as the contribution of additive genetic factors on phenotypic expression. Likelihood ratio statistics were computed to assess the contribution of covariates to variation in each collected phenotype as well as to test for differences in heritability. Models were adjusted for age and gender.

Results

There were no statistically significant differences between baseline measurements and 1-month measurements for intra-examiner assessments ($p=0.708$) indicating adequate reproducibility of study measurements.

The DZ twin pairs demonstrated a higher percentage of outliers in comparison to the MZ pairs. For the MZ mandibular primary left first molars, one out of nine (11%) superimposed images exhibited outliers while DZ superimposed images presented in eight out of twelve instances (66%) ($p=0.001$). For the right molars, the corresponding numbers for outliers were 2/9 (22%) for MZ and 7/12 (58%) for DZ ($p=0.07$). The total amount of difference in surface overlap was 0.0508 mm (0.0018 inches) for the MZ ($n=18$) sample and 0.095 mm (0.0034 inches) for the DZ ($n=24$) sample and were not statistically significant ($p=0.2203$).

For the MZ sample, the data were further dichotomized into mandibular right and left first molars' average distance. The results for these analyses are shown in Table 1 and were not statistically significantly different. However, within the DZ data these differences for mandibular right first molars and for mandibular left first molars were statistically significantly different (Table 1).

The surface areas (in mm²) for the occlusal surfaces of mandibular right and left first molars were computed for all twins (Figure 3). Heritability estimates of occlusal surface areas for mandibular right and left first molars were, respectively, 97.5% and 98.2% (Table 2).

Discussion

Literature reports have established that there is a major genetic impact on tooth size and on tooth morphology. In humans studies of dental crown diameters provide evidence supporting significant contributions of additive genetic variation ranging from 56 to 92% of phenotype variance [Dempsey and Townsend, 2001].

Proposed genetic models include additive and dominant genetic effects. QTL mapping studies in animals provide evidence for genes of major effect that influence specific traits. Several attempts have been made by employing animal models to determine the genetic linkage to tooth morphology [Jernvall et al., 1994; Jernvall, 1995; Vaahtokari et al., 1996; Jernvall et al., 1998; Jernvall and Jung, 2000; Miletich and Sharpe, 2003]. One study located and examined specific quantitative trait loci (QTL) affecting the size and shape of mandibular molars in mice. Those results showed co-occurrence of QTL for molar shape, mandibular shape, and cranial dimensions in the mice, indicating genetic loci contribute to variance in these traits. Others have described the enamel knot as a central function in controlling growth and pattern of tooth cusps. Located at the tip of the tooth buds, the enamel knots act as the epithelial signaling centres to communicate with the mesenchymal tissues which appear to be crucial for correct tooth patterning and morphogenesis [Jernvall et al., 1994; Jernvall, 1995; Vaahtokari et al., 1996; Jernvall et al., 1998]. Recently, it has been demonstrated that a second set of signaling centres, known as secondary enamel knots, function to determine the multicuspid pattern of molar crowns [Jernvall and Jung, 2000; Miletich and Sharpe, 2003]. These studies have provided great insight into the molecular and genetic basis of tooth morphogenesis. In humans however evidence for these findings in animal models is virtually non-existent.

These findings are not directly relevant to the twin study. There is evidence that sequential gene expression is integral to tooth development. Mutational and animal knockout studies indicate that disruption of expression of these genes can disrupt normal tooth development. Occlusal development involves patterning [Cai et al., 2007] and this is likely under genetic control [Wright and Hart, 2002]. To date there has been little direct evidence in humans quantitating the genetic contribution to occlusal topography. The twin model permits assessment of genetic and environmental contribution to these traits.

The average distance value from 3D comparison is the sum of the difference between the two best-fitted superimposed images. In general, higher average distance values translate to greater differences between the superimposed images. The lower transformed average distance for the MZ group compared to the DZ group indicated a trend for greater similarity of occlusal topography for the MZ group (see results above).

A similar study on occlusal surface morphology has been performed on the occlusal surface of nine pairs of permanent molars [Kabban et al., 2001]. Three pairs were from MZ twins, four pairs from DZ twins and two from unrelated controls. Those results showed a marked trend of increasing inter-surface difference with the largest differences seen in the controls and the smallest in the MZ pairs thus confirming our results.

Evaluation of the average distance of the mandibular first molars of left and right quadrants indicated a significant difference within the DZ sample which converted to a greater similarity of surface topography for the right molars compared to the left molars. This trend was also observed for the MZ mandibular first of the right quadrant (Table 1). We are not able at this point to explain these findings.

The occlusal surface area of molars can provide information on the degree of similarities in occlusal morphology as well. For a given dimension of width and length, an increase in surface area indicates greater surface irregularities, which translates to greater cuspal height and deeper pits and fissures. The same interpretation can be extended to a smaller occlusal surface area where smaller cuspal height and shallower pits and fissures are observed. Heritability estimates for occlusal surface areas for the mandibular right and left first molars were very high (Table 2) suggesting that the trait of occlusal surface area for mandibular primary molars was influenced by genetic factors. No studies of the occlusal surface area of

primary molars have evaluated heritability estimates for these traits so comparisons of the results from this study are not possible.

Conclusion

The occlusal morphology of mandibular first primary molars of DZ twin pairs was more variable than that of MZ pairs. In addition, heritability estimates indicated a relatively high genetic contribution to observed variation of occlusal morphology based on surface area.

References

- Almasy L, Blangero J. Multipoint quantitative-trait linkage analysis in general pedigrees. *Am J Hum Genet.* 1998; 62:1198–1211. [PubMed: 9545414]
- Biggerstaff RH. Heritability of the Carabelli cusp in twins. *J Dent Res.* 1973; 52:40–44. [PubMed: 4509504]
- Boraas JC, Messer LB, Till MJ. A genetic contribution to dental caries, occlusion and morphology as demonstrated by twins reared apart. *J Dent Res.* 1988; 67:1150–55. [PubMed: 3165997]
- Cai J, Cho SW, Kim JY, et al. Patterning the size and number of tooth and its cusps. *Dev Biol.* 2007 Jan 9. [Epub].
- Dempsey PJ, Townsend GC. Genetic and environmental contributions to variation in human tooth size. *Heredity.* 2001; 86:685–93. [PubMed: 11595049]
- Jernvall J, Aberg T, Kettunen P, Keranen S, Thesleff S. The life history of an embryonic signaling center: BMP-4 induces p21 and is associated with apoptosis in the mouse tooth enamel knot. *Development.* 1998; 125:161–69. [PubMed: 9486790]
- Jernvall J, Jung HS. Genotype, phenotype, and developmental biology of molar tooth characters. *Am J Phys Anthropol.* 2000; 31:171–90. [PubMed: 11123840]
- Jernvall J. Mammalian molar cusp patterns: Developmental mechanisms of diversity. *Acta Zool Fennica.* 1995; 198:1–61.
- Jernvall J, Kettunen P, Karavanova I, Martin LB, Thesleff I. Evidence for the role of the enamel knot as a control center in mammalian tooth cusp formation: non-dividing cells express growth stimulating Fgf-4 gene. *Int J Dev Biol.* 1994; 38:463–469. [PubMed: 7848830]
- Kabban M, Fearn J, Jovanovski V, Zou L. Tooth size and morphology in twins. *Int J Paediatr Dent.* 2001; 11:577–586.
- Miletich I, Sharpe PT. Normal and abnormal dental development. *Human Mol Genet.* 2003; 12:R69–73. [PubMed: 12668599]
- Pinkham, JR.; Casamassimo, PS.; Fields, HW.; McTigue, DJ.; Nowak, A. *Pediatric Dentistry: Infancy through Adolescence.* 3. Philadelphia, PA: W.B. Saunders; 1999.
- Townsend G, Richards L, Hughes T. Molar intercuspal dimensions: genetic input to phenotypic variation. *J Dent Res.* 2003; 82:350–355. [PubMed: 12709500]
- Vahtokari A, Aberg T, Jernvall J, Keranen S, Thesleff I. The enamel knot as a signaling center in the developing mouse tooth. *Mech Dev.* 1996; 54:39–43. [PubMed: 8808404]
- Wright JT, Hart TC. The genome projects: implications for dental practice and education. *J Dent Educ.* 2002; 66:659–71. [PubMed: 12056771]
- Zhang Y, Lundgren T, Renvert S, et al. Evidence of a founder effect for four cathepsin C gene mutations in Papillon-Lefevre syndrome patients. *J Med Genet.* 2001; 38:96–101. [PubMed: 11158173]

Figure 1.
Picture of PICZA 3D Scanner with dental model place for scanning.

Figure 2.
a) Colour fringe plot by Qualify 7 with no outliers; b) Example of a case with outliers.

Figure 3.
Total surface area in mm² measurement by Magics Software.

Table 1

Average distance of superimposed images of first mandibular molars by zygosity and tooth type.

Zygosity	Transformed average distance	N	p-value
MZ right mandibular 1st molar	0.0135 mm 0.00054 inches	9	0.168 NS
MZ left mandibular 1st molar	0.0783 mm 0.00313 inches	9	
DZ right mandibular 1st molar	0.0450 mm 0.0018 inches	12	0.037
DZ left mandibular 1st molar	0.0125 mm 0.00497 inches	12	

Table 2

Heritability estimates for occlusal surface area of mandibular primary first molars.

Trait	Heritability (%)	Standard error	p-value	n	Log-likelihood
Occlusal surface area of right mandibular 1st molar (mm ²)	97.5	1.28	<.0001	42	-84.238
Occlusal surface area of left mandibular 1st molar (mm ²)	98.2	0.94	<.0001	42	-75.973