Published in final edited form as:

Psychol Med. 2011 March ; 41(3): 575–588. doi:10.1017/S0033291710001078.

Evidence for the Continuous Latent Structure of Mania in the Epidemiologic Catchment Area from Multiple Latent Structure and Construct Validation Methodologies

James J. Prisciandaro and John E. Roberts

Department of Psychology, University at Buffalo, The State University of New York

Abstract

Background—Although psychiatric diagnostic systems have conceptualized mania as a discrete phenomenon, appropriate latent structure investigations testing this conceptualization are lacking. In contrast to these diagnostic systems, several influential theories of mania have suggested a continuous conceptualization. The present study examined whether mania has a continuous or discrete latent structure using a comprehensive approach including taxometric, information-theoretic latent distribution modeling (ITLDM), and predictive validity methodologies in the Epidemiologic Catchment Area (ECA) study.

Methods—Eight dichotomous manic symptom items were submitted to a variety of latent structural analyses; including factor analyses, taxometric procedures, and ITLDM; in 10,105 ECA community participants. Additionally, a variety of continuous and discrete models of mania were compared in terms of their relative abilities to predict outcomes (i.e., health service utilization, internalizing and externalizing disorders, and suicidal behavior).

Results—Taxometric and ITLDM analyses consistently supported a continuous conceptualization of mania. In ITLDM analyses, a continuous model of mania demonstrated 6:52:1 odds over the best fitting latent class model of mania. Factor analyses suggested that the continuous structure of mania was best represented by a single latent factor. Predictive validity analyses demonstrated a consistent superior ability of continuous models of mania relative to discrete models.

Conclusions—The present study provided three independent lines of support for a continuous conceptualization of mania. The implications of a continuous model of mania are discussed.

Although psychiatric diagnostic systems (e.g., DSM-IV; APA, 1994) conceptualize mania and associated bipolar disorder phenomenology as categorical, there have been no appropriate latent structure investigations conducted to support or refute this conceptualization. In contrast to the *DSM*, most influential theories of mania and bipolar disorders have suggested a dimensional conceptualization of these constructs. For example, Kraepelin (1921) proposed that affective temperaments form the constitutional foundation for manic-depressive illness, and observed that non-disordered relatives of manic-depressive patients frequently exhibited these temperaments. Intellectual descendants of Kraepelin have subsequently provided empirical evidence to support the validity of a continuous conceptualization of manic-depressive illnesses (e.g., Judd & Akiskal, 2003, Merikangas et al., 2007). Aside from the theoretical importance of identifying the latent structure of mania, there are a number of concerns associated with potentially incorrectly conceptualizing mania

as a categorical construct; including statistical problems (e.g., attenuated statistical power; Cohen, 1983), spurious psychiatric comorbidity (Haslam, 2003), and inappropriate denial of treatment services.

Appropriate statistical methodologies for determining whether mania is a categorical or dimensional construct include taxometrics (Waller & Meehl, 1998) and information-theoretic latent distribution modeling (ITLDM; Markon & Krueger, 2006). Using multiple latent structure methodologies to investigate the continuous vs. discrete nature of a construct has long been recommended, but is rarely done in practice (Lenzenweger, 2004). Although no taxometric or ITLDM studies have examined mania or bipolar disorders, Meyer and Keller (2003) recently conducted a taxometric investigation of hyperthymic temperament. This study had a number of methodological issues (e.g., use of a single taxometric procedure), but nonetheless found evidence for a continuous latent structure.

Although ITLDM and taxometric analyses are crucial to determining whether mania is a categorical or dimensional construct, results from such analyses can be misleading due to flaws in study design (Grove, 1991) or to limitations of the analytic procedures (Ruscio et al., 2004). Consequently, researchers have argued that latent structural findings are provisional until their meaning has been clarified through construct validation (Waldman & Lilienfeld, 2001; Watson, 2003), and have provided examples of structural findings that did not survive such scrutiny (e.g., dissociative taxon; Watson, 2003; schizoid taxon; Nichols & Jones, 1985).

One approach to evaluating the construct validity of taxometric and ITLDM findings is to compare the relative abilities of multiple discrete and continuous models of mania to predict theoretically relevant constructs. If mania is continuous, it should demonstrate superior predictive ability relative to a discrete model because of the loss of statistical power associated with dichotomizing a continuous variable (Cohen, 1983). If alternatively, mania is discrete, then discrete models of the construct should out-predict continuous models because the extra variability afforded by continuous models would represent measurement error that would attenuate statistical relationships (Ruscio et al., 2006). Two previous studies have used this methodology to examine the construct validity of discrete and continuous models of depression (Aggen et al., 2005; Prisciandaro & Roberts, 2009). Both studies found that discrete models of depression did not significantly predict relevant outcomes once continuous models of depression were statistically controlled.

The present study represents the first to properly evaluate whether mania is a discrete or continuous construct. Given the various criticisms that have been raised regarding using taxometrics as a standalone methodology (e.g., lack of agreement over implementation decisions; see Prisciandaro & Roberts, 2010 for a discussion), the present investigation used an expanded, comprehensive approach (Prisciandaro & Roberts, 2010) to examine the latent structure of mania. In Part I of the present study, both taxometric and ITLDM analyses were conducted on the manic symptom data from the Epidemiologic Catchment Area (ECA; Eaton & Kessler, 1985). In Part II of the present study, an examination of the construct validity of taxometric and ITLDM findings was performed by testing the relative predictive abilities of various discrete and continuous models of mania.

Part I Method

Sample and Measure

The present study used the Epidemiologic Catchment Area (ECA; Eaton & Kessler, 1985) data. The ECA program obtained a representative group of over 20,000 individuals through probability sampling of five geographic catchment areas. Participants were administered a

structured diagnostic interview (i.e., the Diagnostic Interview Schedule; DIS; Robins et al., 1981) in the first wave of the ECA program, which included an assessment of manic symptoms (in accordance with DSM-III criteria; APA, 1980) over various recall periods (e.g., past 2 weeks, lifetime). Symptoms assessed from 2 weeks prior to participants' interviews were used because these data are minimally affected by long-term recall biases, and because it is clear that reported symptoms occurred during the same time period. The DIS contained nine dichotomous questions designed to assess symptoms of mania. These questions were collapsed into the 8 diagnostic criteria of a Manic Episode [elevated or expansive mood ("elevated mood")¹, increased activity or physical restlessness ("increased activity"), increased talkativeness or pressured speech ("talkativeness"), flight of ideas or racing thoughts ("racing thoughts"), inflated self-esteem or grandiosity ("grandiosity"), decreased need for sleep ("decreased sleep"), distractibility, excessive involvement in activities with high potential for painful consequences ("risky behavior"); APA, 1980] using syntax provided with the ECA data. Because of the probing structure of the DIS, an affirmative response to each symptom question indicated that the symptom did not only occur following drug or alcohol use. Test-retest reliability of diagnoses of Manic Episodes made using the DIS have been found to be acceptable ($\kappa = .56$; Semler et al., 1987). Concordance between diagnoses of Manic Episodes made by lay interviewers and psychiatrists using the DIS have also been found to be acceptable ($\kappa = .65$; Robins et al., 1981; Robins et al., 1982). Finally, concordance between DIS diagnoses of Manic Episodes and diagnoses based on semistructured clinical interviews have been found to be good in clinical populations ($\kappa = .63$; Helzer et al., 1985; $\kappa = .86$; Wittchen et al., 1985).

Although all ECA participants were administered all manic symptom questions, a subsample ($n = 10,105$) was selected from the ECA data due to methodological considerations. Included participants belonged to the community sample (institutionalized individuals were excluded because their inclusion could have produced a spurious taxon; Grove, 1991), were assessed for past 2-week symptoms, and completed the interview. The remaining sample contained more females than males (unweighted [weighted]: 58.6% [53.2%] vs. 41.4% [46.8%]), was predominantly White (unweighted [weighted]: 56.8% [54.1%], 25.8% [22.7%] Black, 13.3% [17.3%] Hispanic, 1.7% [2.7%] Asian, 0.8% [0.9%] American Indian, and 1.6% [0.5%] other) and ranged in age from 18 to 96 years old, with a mean age of 46 (unweighted; weighted = 41.9) years. The base rate of Manic Episodes was: lifetime (observed $n = 50$) 0.49% [unweighted; weighted = 0.46%]; past 2 weeks (observed $n = 15$) 0.15% [unweighted; weighted = 0.16%]. Missing data were negligible (2.2%) and were list-wise deleted (Kline, 2005).

Analytic Strategy

Taxometric analyses were conducted using Ruscio's taxometric programs (Ruscio et al., 2006) for the R platform (R Development Core Team, 2009). All other analyses were conducted using MPlus version 5.1 (Muthén & Muthén, 2007).

Preliminary dimensionality analyses—Incorrectly specifying mania as unidimensional vs. multidimensional can create difficulties interpreting ITLDM results (Markon & Krueger, 2006). To address this potential difficulty, factor analyses were conducted. The sample was randomly divided into two equal subsamples. Exploratory factor analysis (EFA) was conducted on the first subsample, using robust weighted least squares estimation (WLSMV; B. Muthén, 1989). A sample weight was applied to adjust for

¹In the ECA, only participants who endorsed ≥ 3 co-occurring symptoms of mania and denied elevated or expansive mood were asked a question regarding irritable mood. Because the question was not asked of all participants, irritable mood was not included in the present analyses.

systematic nonresponse and differential selection probabilities. The number of factors to extract was determined by parallel analysis (Horn, 1965) with 1,000 sets of random data. Results from the EFA model were used to construct a confirmatory factor analysis (CFA) model in the second subsample using WLSMV estimation. In addition to the abovementioned sample weight, information regarding the stratification and clustering of the data was also modeled (Muthén, 2004). CFA model fit was evaluated according to Hu and Bentler's (1999) guidelines ($CFI > .95$, $TLI > .95$, $RMSEA < .05$).

Taxometrics—Taxometric statistical methods have consistently demonstrated their ability to determine whether a construct has a discrete or continuous latent structure (Meehl, 1973; Meehl & Yonce, 1994, 1996; Ruscio, 2000). Rigorous taxometric investigations include multiple nonredundant procedures, additional consistency checks, and simulation techniques (Ruscio et al., 2006). Although the ECA data has a number of strengths for taxometrics (e.g., large sample size, unselected sample), the data also present a number of challenges. Specifically, the items available for the present analysis were dichotomous with sparse endorsement, and the hypothesized manic taxon had a very low base rate. Regarding the former concern, research has demonstrated that given large sample size (Ruscio, 2000) and proper implementation of analytic techniques (e.g., use of simulations and the inchworm consistency test; Ruscio & Marcus, 2007; Ruscio et al., 2004; Waller & Meehl, 1998), taxometric procedures are able to validly distinguish between discrete and continuous structures. Regarding the latter concern, simulation research has demonstrated that with sufficient inclusion of taxon members along with a large sample size, taxometric procedures can detect base rates as low as 0.1%-0.3% (Ruscio & Ruscio, 2004b). Ultimately, suitability for taxometric analysis was determined by simulating many sets of taxonic and dimensional data (Ruscio & Kaczetow, 2008); if taxometric results from simulated taxonic and dimensional data could be distinguished from one another, then the data were suitable for analysis.

MAXCOV (Meehl & Yonce, 1996) was conducted on all possible input/output indicator configurations. Summed input indicators were not used because recent evidence suggests that using summed input indicators for MAXCOV results in significantly less accurate results (Walters & Ruscio, 2009). Subsamples were created by dividing the sample into a large number of overlapping windows (i.e., 986 to $3,944$)², with 90% overlap to allow for the inchworm consistency test and to provide more interpretable results (Ruscio et al., 2006). 10 internal replications were implemented per run, and all MAXCOV curves were combined by averaging the covariance estimates for each subsample. MAMBAC (Meehl & Yonce, 1994) was also conducted. Summed input indicators were used, and the first and last cuts were made 25 cases from each input indicator's distributional tails (Ruscio & Ruscio, 2004a). Internal replications were implemented as described above. To improve interpretability (Ruscio et al., 2006), the number of cuts for MAMBAC was held equal to the maximum number of overlapping windows used for MAXCOV (i.e., 3,944). MAMBAC was repeated until all indicators had served as output, and curves were combined by averaging mean-difference estimates at each cut along the input indicators.

Supplementary consistency tests included base rate divergence (Ruscio & Ruscio, 2004b) and the case removal consistency test (Ruscio, 2000). The inchworm consistency test was also conducted by repeating MAXCOV analyses several times with each successive run

²To calculate the number of windows needed for the inchworm test, we determined the minimum subsample size needed for at least one window to contain more taxon than complement members ($n*2-1 = 25$, where n = number of individuals with past 2 week diagnoses of a Manic Episode). Then we calculated the overall number of windows needed to maintain the desired subsample size (i.e., 3,943; Formula 6.5 in Ruscio et al., 2006, p. 138). We divided this number by 4, and rounded to the nearest whole number, to determine the number of windows to be incrementally added to each successive step of the test (i.e., 986).

containing an increased number of overlapping windows (Waller & Meehl, 1998). Finally, the comparison of simulated taxonic and dimensional data and observed data was used as an interpretational tool as well as a consistency test (Ruscio & Kaczetow, 2008). Taxonic data were simulated using a modification to the base rate classification method (Ruscio, 2009): individuals with the highest total item scores were assigned to the taxon based on the observed base rate of past 2 week Manic Episodes in the present sample. One hundred sets of taxonic and dimensional data were generated and submitted to the same analyses as the research data. Results from the research data were compared to results from simulated taxonic and dimensional data both visually and using fit indices (i.e., comparison curve fit index; CCFI). CCFI values over .60 support taxonic structure, values under .40 support dimensional structure, and values between .40 and .60 are interpreted as ambiguous (Ruscio, Ruscio et al., 2007).

Information-theoretic latent distribution modeling (ITLDM)—ITLDM (Markon & Krueger, 2006) consists of estimating a variety of latent class (LCM) and latent trait (LTM) models using logistic modeling, and subsequently comparing their parsimony-adjusted fit using Bayesian Information Criteria (BIC). Specifically, (nominal) LCM models with between 2 and k classes (where $k = 8 =$ number of indicators); LTM (i.e., “discrete metrical”) models with between 2 and k latent values (with each model distributed according to a binomial distribution, $B[k-1, .5]$, rescaled to a mean of 0 and a standard deviation of 1); and a continuous normally distributed LTM model were estimated. A sample weight, as well as information regarding the stratification and clustering of the data, was included to properly model the complex sample design of the data. Comparisons between LTM models with few vs. many latent values evaluated the relative fit of discrete metrical and continuous models, respectively. Furthermore, comparisons between LTM and LCM models with the same number of values/classes evaluated whether the target construct consisted of ordered or unordered categories. For each comparison, the exponential of 0.5 times the negative BIC difference between the two models was interpreted as the posterior odds of one model over the other (Raftery, 1995). Descriptively, a BIC difference of 0-2 equals “weak” evidence, 2-6 equals “positive” evidence, 6-10 equals “strong evidence,” and > 10 equals “very strong” evidence in favor of the model with the lower BIC value (Raftery, 1995).

Part I Results

Preliminary dimensionality analyses

Parallel analysis suggested that only one interpretable factor could be extracted from the data. Thus, a one factor model was estimated in the first subsample, which provided good fit to the data ($\chi^2 = 9.79$, $df = 11$, $p = .55$; RMSEA = 0.00). The unidimensional CFA model constructed from these findings in the second subsample provided good fit to the data ($\chi^2 = 20.25$, $df = 11$, $p = .04$; CFI = 0.96, TLI = 0.96, RMSEA = 0.01). All factor loadings were statistically significant (mean loading = 0.58, mean $r^2 = .37$).

Taxometrics

Indicator validity was excellent (mean indicator validity = 5.26; Meehl, 1995) and within-class correlations were low (mean absolute value of $r = 0.04$; Meehl, 1995). MAXCOV and inchworm consistency test plots are presented in the top portion of Figure 1. MAXCOV results were all consistent with a dimensional solution: in all cases, the line representing results from the research data fit squarely within ± 1 SD from the average results for simulated dimensional data, and fell on or outside of the boundaries of ± 1 SD from the average results for simulated taxonic data. CCFI values confirmed this superior fit; CCFI ranged from .26 to .30 (mean CCFI = .28). Averaged MAMBAC results are presented in the leftmost bottom panel of Figure 1, alongside averaged results from simulated taxonic

(middle) and dimensional (far right) data. The averaged research curve equally resembles the simulated dimensional and taxonic curves. The associated CCFI value was also ambiguous (CCFI = 0.45).

Consistent with a dimensional interpretation, mean base rate estimates produced by MAXCOV (mean $SD = 0.30$) and MAMBAC ($SD = 0.43$) analyses were divergent. Further suggesting a dimensional interpretation, following targeted case removal, deviations from predicted base rate increases were substantial (MAXCOV deviation = 0.10; MAMBAC deviation = 0.77) and in the opposite direction than would be predicted by the presence of a taxon.³

Information-theoretic Latent Distribution Modeling

See Table 1 for ITLDM results. The two estimated models with the best relative fit to the data were the standard normal LTM (BIC = 9531.07) and the two-class LCM model (BIC = 9534.82). The BIC difference between the two-class LCM model and the standard normal LTM model (3.75) corresponded to “positive” evidence in favor of the LTM model. In other words, the odds of the LTM model over the two-class LCM model were 6.52:1.

To guard against the possibility of spurious continuous findings, all analyses were repeated for lifetime symptom data that contained a larger number, and higher base rate, of hypothesized taxon members. These analyses also supported a continuous conceptualization of mania.⁴ In sum, all taxometric and ITLDM results supported a dimensional interpretation of mania in the ECA data.

Part II Introduction

Part II of the present study evaluated the relative predictive abilities of discrete and continuous models of mania using the methodology detailed in Prisciandaro and Roberts (2009). Two sets of model comparisons were made across outcomes: 1) Between empirically derived discrete and continuous models from Part I of the present study and 2) between rationally selected discrete and continuous models. For the rationally selected continuous model, we created a single additive scale of mania with each manic symptom contributing 1 point. For the discrete model, we chose the predominant diagnostic model of mania: *DSM* Manic Episodes.

³Taxonic data can be simulated using a variety of methods, and the method of simulation chosen can significantly impact taxometric results. Thus, it is important to evaluate the robustness of taxometric results across different simulation methods. All taxometric analyses were re-conducted using an alternate method of taxonic data simulation. The base rate classification method (Ruscio, 2009) assigns individuals with the highest total indicator scores to the taxon group based on the mean base rate estimate from taxometric analyses of the research data. All MAXCOV and MAMBAC curves derived from the research data were more visually similar to curves derived from simulated dimensional data than to curves from simulated taxonic data. Obtained CCFI values from MAXCOV (mean CCFI = .15) and MAMBAC (CCFI = .04) analyses further supported a dimensional interpretation of the data. Overall, these supplementary results suggest that the present study's dimensional taxometric findings were not solely determined by the employed taxonic data simulation method. For further details, see www.buffalo.edu/~robertsj/psychmed.2010.supplement.pdf or www.drtprogram.cjb.net/psychmed.2010.supplement.pdf.

⁴Although past 2 week symptom data are preferable to lifetime data (and simulations suggested that the former could distinguish taxonic and dimensional structures), they are also potentially problematic because the base rate of Manic Episodes over a 2 week period is small (0.15%) and item endorsement is sparse. All analyses were repeated for lifetime symptom data to evaluate the robustness of obtained findings in data with a larger base rate of Manic Episodes (0.49%). Preliminary dimensionality analyses suggested a single latent factor. Twelve sets of MAXCOV analyses across three methods of taxonic data simulation unambiguously supported a continuous interpretation of the data (mean CCFI = .20). Two of three sets of MAMBAC analyses supported a continuous interpretation (mean CCFI = .22); the third set was ambiguous (CCFI = .46). ITLDM analyses suggested that a standard normal LTM model of mania provided the best relative fit to the data. The BIC difference (1.06) between this model and the second-best fitting model (LTM with 4 latent values) corresponded to “weak” evidence in favor of the standard normal model. In other words, the odds of the normal LTM model over the 4-valued LTM model were 1.7:1. Overall, these results support a continuous interpretation of the data, and suggest that the results presented from the past 2 week data were not unduly influenced by methodological concerns regarding an insufficient number of hypothesized taxon members in the sample. For further details, see www.buffalo.edu/~robertsj/psychmed.2010.supplement.pdf or www.drtprogram.cjb.net/psychmed.2010.supplement.pdf.

Three types of outcomes (i.e., dependent variables) were selected for predictive analyses: 1) Psychiatric health service utilization, which is elevated among individuals with bipolar disorders (Merikangas et al., 2007; Weissman et al., 1991); 2) Psychiatric disorders, which are highly comorbid with bipolar disorders (Kessler et al., 1997; McElroy et al., 2001); and 3) Suicidal behavior, which is elevated among individuals with bipolar disorders (Kallner et al., 2000; Sharma & Markar, 1994; Kessler et al., 1999).

The main criterion used to evaluate models' relative predictive abilities was whether the discrete model of mania predicted unique variance in outcomes once the continuous model was statistically controlled (and vice versa).

Part II Method

Sample

The sample for Part II of the present study was the same as for Part I. Missing data were negligible (4.2%) and were list-wise deleted (Kline, 2005).

Discrete and Continuous Models of Mania

Empirically derived models—The best fitting continuous and discrete models of mania from Part I ITLDM analyses were selected for subsequent predictive analyses (see Table 2 for model parameters). As reported in Part I, the best fitting continuous model was the unidimensional, standard normal latent trait model (LTM; BIC = 9531.07). All factor loadings were statistically significant (mean loading = 0.69, mean R^2 = 0.48). This model was represented as a manifest continuous variable for regression analyses by computing factor scores from the standard normal LTM model. The best fitting discrete model was the 2 class latent class model (LCM; BIC = 9534.82). Class 1 (n = 9474, “symptom free”) consisted of individuals with near-zero response probabilities across all symptoms (mean probability = 0.01) and class 2 (n = 406, “potentially symptomatic”) consisted of individuals with at least minimal response probabilities across all symptoms (mean probability = 0.16). This model was represented as a manifest discrete variable for regression analyses by calculating participants' posterior probabilities for each class, assigning participants to the class to which they most likely belonged, and creating a dichotomous variable to reflect these class assignments.

Rationally selected models—The rationally selected continuous model of mania was a single additive scale of manic symptoms, with each symptom contributing 1 point. The empirically derived and rationally selected dimensional models of mania were strongly associated (r = 0.97, p < .001). The rationally selected discrete model was past 2 week DSM-III diagnoses of Manic Episodes (n = 15). The association between the empirically derived and rationally selected categorical models of mania was minimal but statistically significant (r = .08, p < .001).

Outcome Variables

Outcome variables reflecting psychiatric diagnoses and suicidal behavior were assessed as part of the DIS interview. Health service utilization outcomes were assessed using a supplementary structured interview (Shapiro et al., 1985). Psychiatric diagnoses were assessed on a lifetime basis and were coded “0” = absent, “1” = present. Consistent with recommendations (e.g., Boyd et al., 1984; Meyers et al., 1984), DSM-III (APA, 1980) hierarchy rules were not observed. To reduce diagnostic outcomes, Krueger's (1999) dimensional model of common mental disorders was estimated using confirmatory factor analyses (CFA). Suicidal behavior variables were assessed on a lifetime basis and were coded “0” = no, “1” = yes. Items covered: 1) thoughts of death 2) desire to die 3) thoughts of

committing suicide; and 4) suicide attempts. Health service utilization variables (17 total) were assessed on a lifetime basis and were coded “0” = no, “1” = yes. Participants were asked if they had ever gone to a wide variety of people and places where “someone might get help for problems with emotions, nerves, drugs, alcohol, or their mental health,” including spiritual and natural healers, mental health and medical professionals, friends and relatives, and support groups.

Analytic Strategy

Factor Analyses—As described above, Krueger's (1999) model of common mental illnesses was estimated using CFA. Factor scores were computed for the two superordinate factors (internalizing and externalizing), and the resulting variables were used as outcomes in subsequent predictive analyses. Additionally, two separate unidimensional CFA models were estimated from the suicidal behavior and health service utilization variables, respectively; outcome variables were created by computing factor scores from each of these models. CFA analyses were conducted using robust weighted least squares estimation (WLSMV; Muthén, 1989), and each incorporated the ECA sample weight as well as information regarding the stratification and clustering of the data (Muthén, 2004).

Regression Analyses—Regression models were estimated using the %SREGSUB macro (SAS Institute Inc., 2002) in SAS version 9.1.3 (SAS Institute Inc., Cary, NC, USA), which allows for the analysis of subpopulations of complex survey data and includes information regarding the weighting, stratification, and clustering of the data into parameter and variance estimations. Separate sets of predictive comparisons were conducted for the empirically derived and the rationally selected models of mania. Within each set of comparisons, separate step-wise regression models were estimated for each of the 4 outcome variables. Step 1 included either a set of continuous or discrete predictors, whereas step 2 added the set of predictors not included in step 1. Each step-wise model was estimated twice per outcome: once with the continuous predictor entered at step 1, and again with the discrete predictor entered at step 1. The improvement in model fit from step 1 to step 2 represented the degree to which one model of mania provided unique predictive validity beyond that of the alternative model. Of particular interest were the unique relationships between predictor variables and outcomes in step 2 models. Betas involving predictors with meaningful scales of measurement (i.e., both of the rationally selected models and the discrete empirically derived model) were Y standardized. Alternatively, betas involving the continuous empirically derived model were fully (XY) standardized. Separate Bonferroni corrections were applied to significance tests of ΔR^2 and β to control experimentwise α inflation. With a desired α of 0.05 for each type of test, Bonferroni corrections suggested an α of 0.0016 for individual ΔR^2 (0.05/32) and β (0.05/32) significance tests.

Part II Results

Factor Analyses

Krueger's (1999) model of common mental illnesses provided good fit to the data ($\chi^2 = 134.02$, $df = 19$, $p < 0.001$, CFI = 0.95, TLI = 0.95, RMSEA = 0.02), and all factor loadings were statistically significant (mean loading = 0.74, mean $R^2 = 0.56$). Internalizing and externalizing factors were significantly correlated ($r = 0.43$, $p < .001$). The suicidal behavior CFA model provided acceptable fit to the data ($\chi^2 = 142.21$, $df = 2$, $p < .001$, CFI = 0.98, TLI = 0.95, RMSEA = 0.07), and all factor loadings were statistically significant (mean loading = 0.83, mean $R^2 = 0.70$). Finally, the health service utilization CFA model provided good fit to the data ($\chi^2 = 126.83$, $df = 26$, $p < .001$, CFI = 0.95, TLI = 0.97, RMSEA = 0.02), and all factor loadings were statistically significant (mean loading = 0.71, mean $R^2 = 0.51$). The four outcome variables that were created from these models (by computing factor scores

for all participants) had acceptable levels of skew and kurtosis (< 3 and < 10 , respectively; Kline, 2005).

Regression Analyses

Empirically derived models—Results regarding the empirically derived models' abilities to predict outcomes are presented in Table 3. As can be seen in the top half of Table 3, the inclusion of the discrete model at step 2 did not result in a statistically significant improvement in R^2 for any of the four outcomes. Additionally, the discrete model did not uniquely predict any of the four outcomes when the continuous model was simultaneously considered. As can be seen in the bottom half of Table 3, the inclusion of the continuous model at step 2 resulted in a statistically significant improvement in R^2 for each of the four outcomes (mean $\Delta R^2 = 0.02$, mean percent increase in $R^2 = 89\%$). Furthermore, the continuous model significantly uniquely predicted each of the four outcomes when the discrete model was simultaneously considered (mean $\beta = 0.20$). In sum, the continuous model demonstrated unambiguously superior predictive validity relative to the discrete model; the latter model had no incremental validity over the former model.

Rationally selected models—Results regarding the rationally selected models' abilities to predict outcomes are presented in Table 4. As can be seen in the top half of Table 4, the inclusion of the discrete model at step 2 led to a statistically significant improvement in R^2 for internalizing, externalizing, and suicidal behavior (mean $\Delta R^2 = 0.003$, mean percent increase in $R^2 = 7\%$), but not for health service utilization. However, the discrete model only uniquely predicted one of the four outcomes (suicidal behavior: $\beta = 0.16$) when the continuous model was simultaneously considered. As can be seen in the bottom half of Table 4, the inclusion of the continuous model at step 2 resulted in a statistically significant improvement in R^2 for each of the four outcomes (mean $\Delta R^2 = 0.04$, mean percent increase in $R^2 = 2,296\%$). Furthermore, the continuous model significantly uniquely predicted each of the four outcomes when the discrete model was simultaneously considered (mean $\beta = 0.07$). In sum, although the discrete model demonstrated some incremental predictive validity relative to the continuous model, the increase in R^2 associated with the inclusion of the discrete model was very small, and the discrete model only uniquely predicted one outcome when the continuous model was simultaneously considered.

Taken together, findings from Part II provide strong and consistent support for continuous models of mania, and weak and inconsistent support for discrete models of mania. To guard against the possibility of spurious continuous findings, all analyses were repeated for lifetime symptom data that contained a larger number, and higher base rate, of diagnosed Manic Episodes. These analyses also supported a continuous conceptualization of mania.⁵

⁵Because the base rates of manic symptoms and Manic Episodes in the present study's 2 week recall period were low, models of mania, especially the rationally selected discrete model, may have been limited in their ability to predict outcomes. Therefore, all predictive analyses were repeated for lifetime symptom data to evaluate the robustness of regression findings in data with a larger base rate of Manic Episodes ($0.49\% = 50$ potential taxon members). The inclusion of the empirically derived or rationally selected continuous model at step 2 resulted in a statistically significant improvement in R^2 for each of the four outcomes (empirically derived: mean $\Delta R^2 = 0.03$, mean percent increase in $R^2 = 41\%$; rationally selected: mean $\Delta R^2 = 0.09$, mean percent increase in $R^2 = 580\%$). Furthermore, in all cases, the continuous model significantly uniquely predicted each of the four outcomes when the discrete model was simultaneously considered (mean $\beta = 0.30$). The inclusion of the empirically derived discrete model at step 2 led to a statistically significant improvement in R^2 for health service utilization ($\Delta R^2 = 0.002$, percent increase in $R^2 = 2\%$), but not for internalizing, externalizing, and suicidal behavior. The inclusion of the rationally selected discrete model at step 2 led to a statistically significant improvement in R^2 for externalizing (mean $\Delta R^2 = 0.001$, mean percent increase in $R^2 = 1\%$), but not for internalizing, health service utilization, and suicidal behavior. However, in no cases did the discrete model uniquely predict any of the four outcomes when the continuous model was simultaneously considered. Taken together, findings from these supplementary analyses provide strong and consistent support for continuous models of mania, and weak and inconsistent support for discrete models of mania. For further details, see www.buffalo.edu/~robertsj/psychmed.2010.supplement.pdf or www.drtprogram.cjb.net/psychmed.2010.supplement.pdf.

Discussion

The present study is the first to examine whether mania is a discrete or continuous construct using appropriate methodologies. In an unselected epidemiologic sample of approximately 10,000 individuals, the manic symptom questions from the structured DIS interview were submitted to a wide variety of taxometric procedures and consistency tests, as well as to information-theoretic latent distribution modeling (ITLDM). The relative predictive validities of various continuous and discrete models of mania were also examined. Results all converged on a continuous solution.

These results provide support for prominent theories of bipolar disorders that conceptualize mania as an extreme variant of human functioning (e.g., Kraepelin, 1921; Depue et al., 1987). They are also consistent with studies demonstrating the validity of subthreshold mania (e.g., Merikangas et al., 2007). Taxometric studies that have investigated other bipolar mood constructs, such as depression (e.g., Ruscio & Ruscio, 2000; Prisciandaro & Roberts, 2005; Slade & Andrews, 2005) and hyperthymic temperament (Meyer & Keller, 2003) have also found these constructs to be continuous.⁶ On the other hand, the latent structures of a number of key bipolar mood constructs (e.g., mixed mania, mood cycling) have not yet been examined using appropriate statistical methodologies. Future studies should examine the reliability of the present study's findings, and also conduct similar investigations on related bipolar constructs.

The present study had a number of methodological strengths (e.g., large sample size, high indicator validity) that allowed it to overcome challenging aspects of the data (e.g., sparse, dichotomous data). Although unselected community samples are generally preferred in structural investigations because they minimize the likelihood of spurious dimensional or categorical findings (Ruscio et al., 2006), the present study's extremely low number of currently manic individuals potentially increased the likelihood of spurious dimensional findings. However, all analyses were repeated on lifetime manic symptom data, with an increased number of manic individuals, and these results converged on a continuous interpretation. Nevertheless, future research should evaluate the structure of mania in samples that are more likely to contain currently manic individuals.

The present study's support for a unidimensional representation of mania is not entirely consistent with previous factor analytic studies, which have found between 2 and 7 latent factors (Akiskal et al., 2003; Cassidy et al., 1998; Murphy & Beigel, 1974; Sato et al., 2002; Swann et al., 2001). However, these studies have extracted factors reflecting depression and psychosis, suggesting that symptoms unrelated to mania were included. Nonetheless, future factor analytic studies of *DSM* mania should include more than one item per *DSM* criterion to ensure sufficient coverage. Although the present study used indicators created from *DSM*-III (APA, 1980) criteria, present day Manic Episode criteria (APA, 1994) are nearly identical to those in *DSM*-III. Finally, because irritable mood was only assessed in a highly selected group of individuals in the ECA, the absence of irritable mood as a symptom indicator in the present study was an additional limitation; the results of the present study are thus more applicable to euphoric mania than to a conceptualization of mania that accepts irritable mood in the absence of euphoric/elevated mood.

If the results from the present study are supported by future investigations, this may argue for the use of a continuous representation of mania. If continuous conceptualizations of

⁶Though a small number of taxometric studies (e.g., Solomon et al., 2006; Ruscio, Zimmerman et al., 2007) have supported a discrete structure of depression, all available construct validity investigations have supported a continuous interpretation (Aggen et al., 2005; Prisciandaro & Roberts, 2009).

mania are more valid than discrete representations, categorical selection methods for research (e.g., selecting participants diagnosed with *DSM* Manic Episodes) would reduce statistical power (Cohen, 1983), prevent proper investigation of dimensional theoretical models of mania (e.g., Depue et al., 1987), disguise potential nonlinear relationships between mania and other variables (Ruscio et al., 2004), and create spurious statistically significant effects (Maxwell & Delaney, 1993). Instead, correlational research designs that sample individuals at all levels of manic symptom severity should be used. In order to sample individuals at all levels of symptom severity, continuous measures of manic symptoms must be further developed as existing measures may not reflect the true latent structure of mania.

As Akiskal and colleagues have demonstrated (e.g., Judd & Akiskal, 2003), the issue of clinical significance needs to be revisited for bipolar disorders; it is clear that subthreshold expressions of mania can result in pervasive impairment. The findings from the present study further support Akiskal and colleagues' reasoned arguments for expansion of the bipolar spectrum. It is important to recognize, however, that support for a continuous conceptualization of mania suggests that there are *no* points at which a certain number of symptoms or a certain level of symptom severity inherently denotes the presence of a manic disorder. Instead, a continuous conceptualization of mania necessitates that empirical associations between mania and important outcomes (e.g., functional impairment) form the primary basis for determining treatment resource allocation. For example, if the relationship between mania and impairment is nonlinear, treatment efforts could be directed at individuals whose levels of manic symptoms place them at the cusp of a substantial increase in functional impairment. If instead the relationship between mania and impairment is linear, a graded continuum of care could be provided to affected individuals such that different degrees of manic severity are met with concordant levels of treatment intensity (e.g., ranging from psychoeducation to hospitalization). Cost-benefit analyses could be used to determine which specific treatments are most effective at particular levels of severity. Before such a plan can be instituted, however, research is needed to determine the form of the relationship between mania and functional impairment, and to further investigate the cost-benefit ratios of specific treatments at varying levels of manic symptom severity.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank Craig Colder and Len Simms for their helpful comments on an earlier draft of this manuscript. We are also grateful to John Ruscio and Len Simms for their statistical advice.

References

- Aggen SH, Neale MC, Kendler KS. DSM criteria for major depression: Evaluating symptom patterns using latent-trait item response models. *Psychological Medicine*. 2005; 35:475–487. [PubMed: 15856718]
- Akiskal HS, Azorin JM, Hantouche EG. Proposed multidimensional structure of mania: Beyond the euphoric-dysphoric dichotomy. *Journal of Affective Disorders*. 2003; 73:7–18. [PubMed: 12507733]
- American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. 3rd. Washington, D.C.: Author; 1980.
- American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. 4th. Washington, D.C.: Author; 1994.

- Boyd JH, Burke JD Jr, Gruenberg E, Holzer CE III, Rae DS, George LK, Karno M, Stoltzman R, McEvoy L, Nestadt G. Exclusion criteria of DSM-III: A study of co-occurrence of hierarchy-free syndromes. *Archives of General Psychiatry*. 1984; 41:983–989. [PubMed: 6477056]
- Cassidy F, Forest K, Murry E, Carroll BJ. A factor analysis of the signs and symptoms of mania. *Archives of General Psychiatry*. 1998; 55:27–32. [PubMed: 9435757]
- Cohen J. The cost of dichotomization. *Applied Psychological Measurement*. 1983; 7:249–253.
- Depue, RA.; Krauss, S.; Spoont, MR. A two-dimensional threshold model of seasonal bipolar affective disorder. In: Magnusson, D.; Ohman, A., editors. *Psychopathology: An interactional perspective*. Orlando, FL: Academic Press; 1987. p. 95-123.
- Eaton, WW.; Kessler, LG. *Epidemiologic field method in psychiatry: The NIMH Epidemiologic Catchment Area program*. Orlando, FL: Academic Press; 1985.
- Grove, WM. Validity of taxometric inferences based on cluster analysis stopping rules. In: Cicchetti, D.; Grove, WM., editors. *Thinking clearly about psychology: Essays in honor of Paul E Meehl, Vol 2: Personality and psychopathology*. Minneapolis, MN: University of Minnesota Press; 1991. p. 313-329.
- Haslam N. The dimensional view of personality disorders: A review of the taxometric evidence. *Clinical Psychology Review*. 2003; 23:75–93. [PubMed: 12559995]
- Helzer JE, Robins LN, McEvoy LT, Spitznagel EL, Stoltzman RK, Farmer A, et al. A comparison of clinical and diagnostic interview schedule diagnoses. Physician reexamination of lay-interviewed cases in the general population. *Archives of General Psychiatry*. 1985; 42:657–666. [PubMed: 4015307]
- Horn JL. A rationale and test for the number of factors in factor analysis. *Psychometrika*. 1965; 30:179–185. [PubMed: 14306381]
- Hu L, Bentler PM. Cutoff criteria for fit indexes in covariance structure analysis: Conventional criteria versus new alternatives. *Structural Equation Modeling*. 1999; 6:1–55.
- Judd LL, Akiskal HS. The prevalence and disability of bipolar spectrum disorders in the US population: Re-analysis of the ECA database taking into account subthreshold cases. *Journal of Affective Disorders*. 2003; 73:123–131. [PubMed: 12507745]
- Kallner G, Lindelius R, Petterson U, Stockman O, Tham A. Mortality in 497 patients with affective disorders attending a lithium clinic or after having left it. *Pharmacopsychiatry*. 2000; 33:8–13. [PubMed: 10721878]
- Kessler RC, Borges G, Walters EE. Prevalence of and risk factors for lifetime suicide attempts in the National Comorbidity Survey. *Archives of General Psychiatry*. 1999; 56:617–626. [PubMed: 10401507]
- Kessler RC, Rubinow DR, Holmes C, Abelson JM, Zhao S. The epidemiology of DSM-III-R bipolar I disorder in a general population survey. *Psychological Medicine*. 1997; 27:1079–1089. [PubMed: 9300513]
- Kline, RB. *Principles and practice of structural equation modeling*. 2nd. NY: Guilford Publications; 2005.
- Kraepelin, E. *Manic-depressive insanity and paranoia*. Edinburgh: Livingstone; 1921.
- Krueger RF. The structure of common mental disorders. *Archives of General Psychiatry*. 1999; 56:921–926. [PubMed: 10530634]
- Lenzenweger MF. Consideration of the challenges, complications, and pitfalls of taxometric analysis. *Journal of Abnormal Psychology*. 2004; 113:10–23. [PubMed: 14992653]
- Markon KE, Krueger RF. Information-theoretic latent distribution modeling: Distinguishing discrete and continuous latent variable models. *Psychological Methods*. 2006; 11:228–243. [PubMed: 16953702]
- Maxwell SE, Delaney HD. Bivariate median splits and spurious statistical significance. *Psychological Bulletin*. 1993; 113:181–190.
- McElroy SL, Altshuler LL, Suppes T, Keck PE Jr, Frye MA, Denicoff KD, Nolen WA, Kupka RW, Leverich GS, Rochussen JR, Rush AJ, Post RM. Axis I psychiatric comorbidity and its relationship to historical illness variables in 288 patients with bipolar disorder. *American Journal of Psychiatry*. 2001; 158:420–426. [PubMed: 11229983]

- Meehl, PE. MAXCOV-HITMAX: A taxometric search method for loose genetic syndromes. In: Meehl, PE., editor. *Psychodiagnosis: Selected papers*. Minneapolis: University of Minnesota Press; 1973. p. 200-224.
- Meehl PE. Bootstraps taxometrics: Solving the classification problem in psychopathology. *American Psychologist*. 1995; 50:266–275. [PubMed: 7733538]
- Meehl PE, Yonce LJ. Taxometric analysis: I. Detecting taxonicity with two quantitative indicators using means above and below a sliding cut (MAMBAC procedure). *Psychological Reports*. 1994; 74:1059–1274.
- Meehl PE, Yonce LJ. Taxometric analysis: II. Detecting taxonicity using covariance of two quantitative indicators in successive intervals of a third indicator (MAXCOV procedure). *Psychological Reports*. 1996; 78:1091–1227.
- Merikangas KR, Akiskal HS, Angst J, Greenberg PE, Hirschfeld RMA, Petukhova M, Kessler RC. Lifetime and 12-month prevalence of bipolar spectrum disorder in the National Comorbidity Survey replication. *Archives of General Psychiatry*. 2007; 64:543–552. [PubMed: 17485606]
- Meyer TD, Keller F. Is there evidence for a latent class called “hypomanic temperament?”. *Journal of Affective Disorders*. 2003; 75:259–267. [PubMed: 12880938]
- Meyers JK, Weissman MM, Tischler GL, Holzer CE III, Leaf PJ, Orvaschel H, Anthony JC, Boyd JH, Burke JD Jr, Kramer M, Stoltzman R. Six-month prevalence of psychiatric disorders in three communities. *Archives of General Psychiatry*. 1984; 41:959–967. [PubMed: 6332591]
- Murphy DL, Beigel A. Depression, elation, and lithium carbonate responses in manic patient subgroups. *Archives of General Psychiatry*. 1974; 31:643–648. [PubMed: 4441238]
- Muthén BO. Dichotomous factor analysis of symptom data. *Sociological Methods & Research*. 1989; 18:19–65.
- Muthén, BO. *Mplus Technical Appendices*. Los Angeles, CA: Muthén & Muthén; 2004.
- Muthén, LK.; Muthén, BO. *Mplus User's Guide*. Fifth. Los Angeles, CA: Muthén & Muthén; 2007.
- Nichols DS, Jones RE Jr. Identifying schizoid-taxon membership with the Golden-Meehl MMPI items. *Journal of Abnormal Psychology*. 1985; 94:191–194. [PubMed: 3998287]
- Prisciandaro JJ, Roberts JE. A taxometric investigation of unipolar depression in the National Comorbidity Survey. *Journal of Abnormal Psychology*. 2005; 114:718–728. [PubMed: 16351392]
- Prisciandaro JJ, Roberts JE. A comparison of the predictive abilities of dimensional and categorical models of unipolar depression in the National Comorbidity Survey. *Psychological Medicine*. 2009; 39:1087–1096. [PubMed: 18845012]
- Prisciandaro JJ, Roberts JE. A comprehensive approach to testing structural questions. 2010 Manuscript in preparation.
- R Development Core Team. *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing; 2009.
- Raftery AE. Bayesian model selection in social research. *Sociological Methodology*. 1995; 25:111–196.
- Robins LN, Helzer JE, Croughan J, Ratcliff KS. National Institute of Mental Health Diagnostic Interview Schedule: Its history, characteristics, and validity. *Archives of General Psychiatry*. 1981; 38:381–389. [PubMed: 6260053]
- Robins LN, Helzer JE, Ratcliff KS, Seyfried W. Validity of the Diagnostic Interview Schedule, Version II: *DSM-III* diagnoses. *Psychological Medicine*. 1982; 12:855–870. [PubMed: 7156256]
- Ruscio J. Taxometric analysis with dichotomous indicators: The modified MAXCOV procedure and a case-removal consistency test. *Psychological Reports*. 2000; 87:929–939. [PubMed: 11191408]
- Ruscio J. Assigning cases to groups using taxometric results: An empirical comparison of classification techniques. *Assessment*. 2009; 16:55–70. [PubMed: 18607008]
- Ruscio, J.; Haslam, N.; Ruscio, AM. *Introduction to the taxometric method: A practical guide*. Mahwah, New Jersey: Lawrence Erlbaum Associates; 2006.
- Ruscio J, Kaczetow W. Simulating multivariate nonnormal data using an iterative algorithm. *Multivariate Behavioral Research*. 2008; 43:355–381.
- Ruscio J, Marcus DK. Detecting small taxa using simulated comparison data: A reanalysis of Beach, Amir, and Bau's (2005) data. *Psychological Assessment*. 2007; 19:241–246. [PubMed: 17563206]

- Ruscio J, Ruscio AM. Informing the continuity controversy: A taxometric analysis of depression. *Journal of Abnormal Psychology*. 2000; 109:473–487. [PubMed: 11016117]
- Ruscio J, Ruscio AM. A nontechnical introduction to the taxometric method. *Understanding Statistics*. 2004a; 3:151–193.
- Ruscio J, Ruscio AM. Clarifying boundary issues in psychopathology: The role of taxometrics in a comprehensive program of structural research. *Journal of Abnormal Psychology*. 2004b; 113:24–38. [PubMed: 14992654]
- Ruscio J, Ruscio AM, Keane TM. Using taxometric analysis to distinguish a small latent taxon from a latent dimension with positively skewed indicators: The case of Involuntary Defeat Syndrome. *Journal of Abnormal Psychology*. 2004; 113:145–154. [PubMed: 14992667]
- Ruscio J, Ruscio AM, Meron M. Applying the bootstrap to taxometric analysis: Generating empirical sampling distributions to help interpret results. *Multivariate Behavioral Research*. 2007; 42:349–386.
- Ruscio J, Zimmerman M, McGlinchey JB, Chelminski I, Young D. Diagnosing Major Depressive Disorder: XI. A taxometric investigation of the categorical-dimensional debate on the structure underlying *DSM-IV* symptoms. *Journal of Nervous and Mental Disease*. 2007; 195:10–19. [PubMed: 17220734]
- SAS Institute Inc. %SREGSUB macro provides additional capabilities for PROC SURVEYREG. 2002. Retrieved June 5, 2008, from <http://support.sas.com/kb/24/985.html#ref>
- Sato T, Bottlender R, Kleindienst N, Moller HJ. Syndromes and phenomenological subtypes underlying acute mania: A factor analytic study of 576 manic patients. *American Journal of Psychiatry*. 2002; 159:968–974. [PubMed: 12042185]
- Semler G, Wittchen HU, Joschke K, Zaudig M, Vongseiso T, Kaiser S, von Cranach M, Pfister H. Test-retest reliability of a standardized psychiatric interview (DIS/CIDI). *European Archives of Psychiatry and Neurological Sciences*. 1987; 236:214–222. [PubMed: 3582430]
- Shapiro, S.; Tischler, GL.; Cottler, L.; George, LK.; Amirkhan, JH.; Kessler, LG.; Skinner, EA. Health services research questions. In: Eaton, WW.; Kessler, LG., editors. *Epidemiologic field method in psychiatry: The NIMH Epidemiologic Catchment Area program*. Orlando, FL: Academic Press; 1985.
- Sharma R, Markar HR. Mortality in affective disorder. *Journal of Affective Disorders*. 1994; 31:91–96. [PubMed: 8071480]
- Slade T, Andrews G. Latent structure of depression in a community sample: A taxometric analysis. *Psychological Medicine*. 2005; 35:489–497. [PubMed: 15856719]
- Solomon A, Ruscio J, Seeley JR, Lewinsohn PM. A taxometric investigation of unipolar depression in a large community sample. *Psychological Medicine*. 2006; 36:973–985. [PubMed: 16700963]
- Swann AC, Janicak PL, Calabrese JR, Bowden CL, Dilsaver SC, Morris DD, Petty F, Davis JM. Structure of mania: Depressive, irritable, and psychotic clusters with different retrospectively-assessed course patterns of illness in randomized clinical trial participants. *Journal of Affective Disorders*. 2001; 67:123–132. [PubMed: 11869759]
- Waldman ID, Lilienfeld SO. Applications of taxometric methods to problems of comorbidity: Perspectives and challenges. *Clinical Psychology - Science & Practice*. 2001; 8:520–527.
- Waller, NG.; Meehl, PE. *Multivariate taxometric procedures: Distinguishing types from continua*. Thousand Oaks, CA: Sage Publications, Inc.; 1998.
- Walters GD, Ruscio J. To sum or not to sum: Taxometric analysis with ordered categorical assessment items. *Psychological Assessment*. 2009; 21:99–111. [PubMed: 19290770]
- Watson D. Investigating the construct validity of the dissociative taxon: Stability analyses of normal and pathological dissociation. *Journal of Abnormal Psychology*. 2003; 112:298–305. [PubMed: 12784840]
- Weissman, MA.; Bruce, ML.; Leaf, PJ.; Florio, LP.; Holzer, Charles, III. Affective Disorders. In: Robins, LN.; Regier, DA., editors. *Psychiatric Disorders in America: The Epidemiologic Catchment Area Study*. New York, NY: Free Press; 1991. p. 53-80.
- Wittchen HU, Semler G, von Zerssen D. A comparison of two diagnostic methods: Clinical ICD diagnoses vs. *DSM-III* and Research Diagnostic Criteria using the Diagnostic Interview Schedule (version 2). *Archives of General Psychiatry*. 1985; 42:677–684. [PubMed: 4015309]

MAXCOV

Figure 1.

MAXCOV (Top): Averaged maximum covariance (MAXCOV) results overlaying lines which represent ± 1 SD from the average results for simulated taxonic (left) and dimensional (right) data. Plots in each successive row contain an increased number of overlapping windows: 986, 1972, 2958, 3944. MAMBAC (Bottom): Averaged means above minus below a cut (MAMBAC) results from research data (far left), simulated taxonic data (center), and simulated dimensional (far right) data.

Table 1
Fit Criteria for Discrete and Continuous Models of Mania

	k	Criterion	
		ln(L)	BIC
Latent class models			
2 classes	17	-4689.23	9534.82
3 classes	26	-4681.21	9601.58
4 classes	35	-4672.09	9666.12
5 classes	44	-4665.76	9736.24
6 classes	53	-4663.99	9815.48
7 classes	62	-4659.92	9890.13
8 classes	71	-4653.03	9959.14
Latent trait models			
2 values	17	-4689.23	9534.82
3 values	18	-4688.93	9543.42
4 values	19	-4687.28	9549.32
5 values	20	-4687.54	9559.04
6 values	21	-4687.53	9568.22
7 values	22	-4687.33	9577.02
8 values	23	-4687.54	9586.64
Normal	16	-4691.95	9531.07

Note. k = number of parameters. BIC = Bayesian Information Criterion. ln(L) = log likelihood.

Table 2
Standard Normal LTM Factor Loadings and R-square Values, and Two Class LCM Item Response Probabilities for the 8 DSM Criteria of a Manic Episode

	Standard Normal LTM		Two Class LCM	
	Factor 1	R-Square	Class 1	Class 2
			"Symptom Free"	"P. Symptomatic"
Racing Thoughts	.81	.66	.00	.22
Talkativeness	.78	.60	.00	.17
Distractibility	.76	.57	.01	.28
Decreased Sleep	.74	.51	.01	.22
Risky Behavior	.64	.41	.00	.05
Elevated Mood	.62	.39	.00	.01
Grandiosity	.60	.37	.01	.17
Increased Activity	.55	.31	.01	.13

Note. LTM = Latent Trait Model. LCM = Latent Class Model. DSM = Diagnostic and Statistical Manual of Mental Disorders. P. Symptomatic = Potentially Symptomatic.

Table 3
Multiple Regression Analyses with Empirically derived Continuous or Discrete Models of Mania Alone Predicting Four Outcomes, Followed by Analyses with Both Models of Mania Simultaneously Predicting Outcomes

I. Addition of Discrete Predictors to Models with Continuous Predictors Only									
	Internalizing		Externalizing		Suicidal Beh.		Health Service Ut.		
	ΔR^2	β	ΔR^2	β	ΔR^2	β	ΔR^2	β	
Step 1									
Standard Normal LTM	.06*	.25*	.04*	.19*	.04*	.20*	.03*	.18*	
Step 2									
Standard Normal LTM	.00	.24*	.00	.20*	.00	.20*	.00	.18*	
Two Class LCM		.01		-.01		.01		.00	

II. Addition of Continuous Predictors to Models with Discrete Predictors Only									
	Internalizing		Externalizing		Suicidal Beh.		Health Service Ut.		
	ΔR^2	β	ΔR^2	β	ΔR^2	β	ΔR^2	β	
Step 1									
Two Class LCM	.03*	.17*	.02*	.12*	.02*	.14*	.02*	.12*	
Step 2									
Two Class LCM	.03*	.01	.02*	-.01	.02*	.01	.02*	.00	
Standard Normal LTM		.24*		.20*		.20*		.18*	

Note. Regression coefficients involving the Standard Normal LTM were XY standardized. Coefficients involving the Two Class LCM were Y standardized. LTM = Latent Trait Model. LCM = Latent Class Model. Internalizing = Internalizing Disorders. Externalizing = Externalizing Disorders. Suicidal Beh. = Suicidal Behavior. Health Service Ut. = Health Service Utilization. ΔR^2 = change in proportion of variance accounted for. β = regression coefficient.

* $p < .0016$.

Table 4
Multiple Regression Analyses with Rationally selected Continuous or Discrete Models of Mania Alone Predicting Four Outcomes, Followed by Analyses with Both Models of Mania Simultaneously Predicting Outcomes

I. Addition of Discrete Predictors to Models with Continuous Predictors Only									
	Internalizing		Externalizing		Suicidal Beh.		Health Service Ut.		
	ΔR^2	β	ΔR^2	β	ΔR^2	β	ΔR^2	β	
Step 1									
Manic Symptoms Scale	.06*	.08*	.03*	.06*	.04*	.07*	.03*	.06*	
Step 2									
Manic Symptoms Scale	.00*	.08*	.00*	.06*	.00*	.07*	.00	.06*	
Manic Episodes		.18		.22		.16*		.04	
II. Addition of Continuous Predictors to Models with Discrete Predictors Only									
	Internalizing		Externalizing		Suicidal Beh.		Health Service Ut.		
	ΔR^2	β	ΔR^2	β	ΔR^2	β	ΔR^2	β	
Step 1									
Manic Episodes	.00*	.23*	.01*	.26*	.00*	.21*	.00	.08	
Step 2									
Manic Episodes	.05*	.18	.03*	.22	.04*	.16*	.03*	.04	
Manic Symptoms Scale		.08*		.06*		.07*		.06*	

Note. LTM = Latent Trait Model. LCM = Latent Class Model. Internalizing = Internalizing Disorders. Externalizing = Externalizing Disorders. Suicidal Beh. = Suicidal Behavior. Health Service Ut. = Health Service Utilization. ΔR^2 = change in proportion of variance accounted for. β = Y standardized regression coefficient.

* $p < .0016$.