Published in final edited form as:

J Mech Behav Biomed Mater. 2011 October ; 4(7): 1167–1176. doi:10.1016/j.jmbbm.2011.03.027.

Active contraction of cardiac muscle: In vivo characterization of mechanical activation sequences in the beating heart

Alkiviadis Tsamis^a, Wolfgang Bothe^b, John-Peder Escobar Kvitting^b, Julia C. Swanson^b, D. Craig Miller^b, and Ellen Kuhl^{*,a,b,c}

^aDepartment of Mechanical Engineering, 496 Lomita Mall, Stanford, CA-94305, USA

^bDepartment of Cardiothoracic Surgery, 300 Pasteur Drive, Stanford, CA-94305, USA

^cDepartment of Bioengineering, 496 Lomita Mall, Stanford, CA-94305, USA

Abstract

Progressive alterations in cardiac wall strains are a classic hallmark of chronic heart failure. Accordingly, the objectives of this study are to establish a baseline characterization of cardiac strains throughout the cardiac cycle, to quantify temporal, regional, and transmural variations of active fiber contraction, and to identify pathways of mechanical activation in the healthy beating heart. To this end, we insert two sets of twelve radiopaque beads into the heart muscle of nine sheep; one in the anterior-basal and one in the lateral-equatorial left ventricular wall. During three consecutive heartbeats, we record the bead coordinates via biplane videofluoroscopy. From the resulting four-dimensional data sets, we calculate the temporally and transmurally varying Green-Lagrange strains in the anterior and lateral wall. To quantify active contraction, we project the strains onto the local muscle fiber directions. We observe that mechanical activation is initiated at the endocardium slightly after end diastole and progresses transmurally outward, reaching the epicardium slightly before end systole. Accordingly, fibers near the outer wall are in contraction for approximately half of the cardiac cycle while fibers near the inner wall are in contraction almost throughout the entire cardiac cycle. In summary, cardiac wall strains display significant temporal, regional, and transmural variations. Quantifying wall strain profiles might be of particular clinical significance when characterizing stages of left ventricular remodeling, but also of engineering relevance when designing new biomaterials of similar structure and function.

Keywords

biomechanics; cardiac mechanics; muscle; excitation; contraction; transmural strains

1. INTRODUCTION

Repairing the failing heart is a desirable but elusive goal. Despite tremendous scientific progress during the past decades, heart failure remains one of the most common, costly, disabling, and deadly medical conditions affecting more than 25 million people worldwide [23]. Dilation of the left ventricle, change in ventricular shape, and decrease in ejection

© 2011 Elsevier Ltd. All rights reserved.

*corresponding author, phone: +1.650.450.0855, fax: +1.650.725.1587, ekuhl@stanford.edu, URL: <http://biomechanics.stanford.edu>.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

fraction are clinical indicators of disease progression and strong predictors of survival [17]. Effective therapies for heart failure seek to attenuate these effects and reverse disease progression [30]. Particularly promising examples are biomaterial approaches that seek to support the ventricular wall to reduce stress and alleviate cardiomyocyte overstretch [12]. Research in biomaterials currently focuses on three mechanisms of wall support: The injection of living cells into the myocardium to create in situ engineered cardiac tissue [36]; the implantation of in vitro engineered tissue patches [2, 28]; and the implantation of synthetic patches or passive cardiac support devices around the heart [8]. The objective of this study is to provide guidelines for the structural and functional requirements of these biomaterial solutions. In particular, we seek to establish a baseline characterization of cardiac and fiber strains throughout the cardiac cycle, to quantify temporal, regional, and transmural variations of active muscle contraction, and to identify pathways of mechanical activation in the healthy beating heart. This baseline characterization of cardiac strains is also of clinical significance since it may help to judge strain alterations in response to cardiac disease.

For more than 30 years, the use of implanted radiopaque markers has provided considerable insight into the complex transmural dynamics of the healthy and diseased ventricular myocardium. In the first series of experiments, small lead spheres were inserted with a needle into the anterior left ventricular wall of five canine hearts to characterize the complex transmural deformation pattern using biplane cineradiography [16]. Ventricular wall strains were extracted from the experimentally measured marker coordinates using the field theories of continuum mechanics. While the data analysis was initially based on the assumption of infinitesimal deformations, subsequent studies demonstrated the need to extend the analysis to finite deformations [35]. Over the years, the bead insertion technique was refined and standardized to the method we have adopted within this study [5, 35]. One of the major breakthroughs enabled by the transmural marker technique was to precisely quantify how individual strain components change with depth, from the epicardium, the outer wall, to the endocardium, the inner wall. While the earlier studies used a discrete point-wise representation of strains, refined methods now use a continuous functional representation, either in the form of fitted bilinear cubic finite elements [29] or fitted polynomials [25], the method we adopt here. In combination with tissue histology, this functional representation naturally lends itself to the quantification of fiber and sheet strains which have significantly improved our understanding of the mechanics underlying coordinated pump function [1, 6, 9, 11].

Throughout the past two decades, marker-based strain characterization has been used both in canine [1, 11, 29, 35] and ovine [4, 5, 6, 7, 25] hearts, with generally similar results, however, strains being slightly larger in dogs than in sheep. The marker technique has been used successfully to identify the impact of acute changes in strain profiles, for example initiated by varying afterload [34] or by ventricular pacing [10]. However, the true beauty of this technique lies in its permanent nature allowing to precisely quantify chronic changes, for example in response to mitral regurgitation [4] or myocardial infarction, in its plain form, or treated with a passive support device [7]. Overall, this technique is relatively well-known in the surgical community but less so in the engineering community. One of the goals of this manuscript is therefore to discuss this technique in view of a modern continuum mechanics analysis which allows us to reconstruct the tensorial characterization of ventricular wall kinematics.

Most existing marker studies use the end diastolic state as the reference configuration and the end systolic state as the deformed configuration. Accordingly, they fail to give a complete picture of the spatio-temporal strain evolution across the ventricular wall. Using a novel time-aligned averaging technique to create a functional representation of the

underlying Green-Lagrange strains [32, 33], we reconstruct strain profiles throughout the entire wall throughout the entire cardiac cycle averaged over nine animals. This allows us to visualize the mechanical activation sequence across the ventricular wall. The resulting color-coded graphics demonstrate that maximum fiber contraction may in fact occur significantly after the end systolic state.

2. MATERIALS AND METHODS

We premedicate nine adult male dorsett-hybrid sheep with ketamine, anesthetize them with sodium thiopental, intubate and ventilate them, and maintain general anesthesia with inhalational isoflurane and supplemental oxygen. After performing a left thoracotomy, we measure the wall thickness at the anterior-basal and at the lateral-equatorial regions of the left ventricular wall using epicardial echocardiography. At each of these two regions, we insert three transmural columns of four bead sets each [35], see Figure 1. Using a depth-adjustable bead insertion trocar, we first insert three 0.7mm diameter beads in each column, evenly spaced between the endo- and epicardial surfaces. In addition, we sew a fourth 1.7 mm diameter bead onto the epicardial surface above each column. To access the left ventricular volume, we implant thirteen evenly spaced epicardial markers silhouetting the left ventricle. Before taking the animals to the catheterization laboratory for data acquisition, we place an implantable micromanometer pressure transducer in the left ventricular chamber and a micromanometer catheter into the ascending aorta for ventricular and aortic pressure measurements. During baseline conditions with the heart in normal sinus rhythm and the animals in the right lateral decubitus position, we record videofluoroscopic images of all beads and markers using biplane videofluoroscopy at a sampling frequency of 60Hz. We simultaneously record aortic pressure, left ventricular pressure, and electrocardiogram voltage signals. Using a semi-automated image processing and digitization software developed in our laboratory [31], we obtain three-dimensional coordinates $x(X, t)$ of the four by three beads of both bead sets and of the thirteen left ventricular markers offline from the acquired biplane images. The accuracy of our three-dimensional displacement reconstruction from biplane videofluoroscopy is 0.1 ± 0.3 mm [13].

3. THEORY AND CALCULATION

3.1. Cardiac Coordinates

To create a functional representation of the strain field in the anterior-basal and lateral-equatorial wall, we define a local Cartesian coordinate system $\{e_c, e_t, e_r\}$ aligned with the local circumferential, radial, and longitudinal directions, see Figure 1. We place its origin into the geometric center of three epicardial beads which define a plane tangential to the outer cardiac wall. The radial axis e_r is oriented normal to this tangential plane, with a positive outward direction. The circumferential axis e_c is introduced normal to the vector connecting the apex of the heart with the origin of the coordinate system and normal to the radial axis, pointing in the mathematically positive direction when viewing the heart from base to apex. The longitudinal axis e_l is defined normal to the radial and circumferential axes, $e_l = e_r \times e_c$, with a positive direction from apex to base, see Figure 1.

3.2. Temporal Interpolation and Data Analysis

To create an average strain representation in the anterior-basal and lateral-equatorial wall, we map all nine experimental data sets into four time intervals between End Diastole, ED, End Isovolumic Contraction, EIC, End Systole, ES, and End Isovolumic Relaxation, EIR, as previously described [32, 33]. We select these characteristic time points semi-manually from the individual pressure-volume relationships, with the volume defined in terms of the thirteen silhouetting epicardial markers. Based on different average interval lengths, these

four time intervals are further subdivided into five segments from ED to EIC, eleven segments from EIC to ES, seven segments from ES to EIR, and ten segments from EIR to ED, respectively. Within each of these four intervals, we perform a linear temporal interpolation between the experimentally acquired raw data points $\mathbf{x}(\mathbf{X}, t)$ to create temporally aligned data sets of hemodynamic and kinematic data at the corresponding five, eleven, seven, and ten segment points, thus representing the cardiac cycle through t_n discrete time points with $n = 1, \dots, 33$. Using these time-aligned data sets, we perform the strain calculations described in the following section to extract the means and standard deviations of the Green-Lagrange strain components $\mathbf{E}(\mathbf{X}, t)$ and fiber strains $E_{FF}(\mathbf{X}, t)$ at 33 discrete time points throughout the cardiac cycle.

3.3. Spatial Interpolation and Cardiac Strains

We assume that within the space enclosed by each bead set, the current placement $\boldsymbol{\varphi}(\mathbf{X}, t)$ of each material point $\mathbf{X}$ at each time t can be approximated by a time-dependent mapping $\mathbf{c}_I(t)$ of the spatial interpolation $N_I(\mathbf{X})$ in the reference configuration,

$$\boldsymbol{\varphi}(\mathbf{X}, t) = \sum_{I=1}^{n_{\text{apx}}} \mathbf{c}_I(t) N_I(\mathbf{X}) \quad (1)$$

where $I = 1, \dots, n_{\text{apx}}$ is the order of the approximation. In particular, at each of the $J = 1, \dots, n_{\text{pnt}}$ experimentally measured data points with $n_{\text{pnt}} = 12$, this implies that

$$\mathbf{x}_J(t) = \sum_{I=1}^{n_{\text{apx}}} \mathbf{c}_I(t) N_I(\mathbf{X}_J) \quad (2)$$

where $\mathbf{x} = [x_c, x_b, x_r]^t$ are the measured data points, $\mathbf{c} = [c_c, c_b, c_r]^t$ are the unknown coefficients at each discrete time point t associated with the circumferential, longitudinal, and radial directions, and N_I are the interpolation terms associated with the corresponding coordinates $\mathbf{X}$ of point J in the reference configuration, in our case, at end diastole. Following the literature [25], we choose a linear interpolation in the circumferential and longitudinal directions and a quadratic interpolation in the radial direction such that N_I can be expressed through the following $n_{\text{apx}} = 9$ terms,

$N_I = [X_c X_r^2, X_c X_r, X_c, X_l X_r^2, X_l X_r, X_l, X_r^2, X_r, 1]^t$. Since we acquire the coordinates of twelve data points while only applying an approximation with nine unknowns, i.e., $n_{\text{pnt}} = 12 > n_{\text{apx}} = 9$, the overall system for the unknown coefficients $\mathbf{c}_I$

$$\mathbf{x}(t)_{[3 \times 12]} = \mathbf{c}(t)_{[3 \times 9]} \cdot \mathbf{N}_{[9 \times 12]} \quad (3)$$

is overdetermined. In the above matrix representation of equation (2), we have introduced the matrix notation for the measured coordinates of the data points $\mathbf{x}_{[3 \times 12]} = [\mathbf{x}_1, \dots, \mathbf{x}_{12}]$, for the unknown coefficients $\mathbf{c}_{[3 \times 9]} = [\mathbf{c}_1, \dots, \mathbf{c}_9]$, and for the interpolation matrix $\mathbf{N}_{[9 \times 12]} = [N_I(\mathbf{X}_1), \dots, N_I(\mathbf{X}_{12})]^t$. We solve this system for the unknown coefficient matrix $\mathbf{c}(t)$ using the pseudo inverse of the interpolation matrix $\mathbf{N}$.

$$\mathbf{c}(t)_{[3 \times 9]} = \mathbf{x}(t)_{[3 \times 12]} \cdot \mathbf{N}_{[12 \times 9]}^\dagger \cdot [\mathbf{N}_{[9 \times 12]} \cdot \mathbf{N}_{[12 \times 9]}^\dagger]^{-1} \quad (4)$$

With the coefficients $\mathbf{c}(t)$ known at each discrete time point t , we can evaluate the nonlinear deformation map $\boldsymbol{\varphi}(\mathbf{X}, t) = \sum_{I=1}^{n_{\text{apx}}} \mathbf{c}_I(t) N_I(\mathbf{X})$ as introduced in equation (1). Accordingly, the deformation gradient $\mathbf{F} = \nabla \boldsymbol{\varphi}$ takes the following functional representation,

$$\mathbf{F}(\mathbf{X}, t) = \sum_{i=1}^{n_{\text{apx}}} \mathbf{c}_i(t) \otimes \nabla N_i(\mathbf{X}) \quad (5)$$

where $\nabla(\mathbf{o}) = [\partial_c(\mathbf{o}), \partial_l(\mathbf{o}), \partial_r(\mathbf{o})]^t$ denotes the spatial gradient with respect to the local cardiac coordinate system. The Green-Lagrange strain tensor in the cardiac coordinate system can then be expressed as follows.

$$\mathbf{E}(\mathbf{X}, t) = \frac{1}{2} [\mathbf{F}^t \cdot \mathbf{F} - \mathbf{I}] \quad (6)$$

Using regionally and transmurally varying fiber directions reported in the literature [15], we project the Green-Lagrange strain tensor onto the local fiber direction $\mathbf{f}(\mathbf{X})$

$$E_{\text{FF}}(\mathbf{X}, t) = \mathbf{f}(\mathbf{X}) \cdot \mathbf{E}(\mathbf{X}, t) \cdot \mathbf{f}(\mathbf{X}) \quad (7)$$

to characterize the local fiber strains $E_{\text{FF}}(\mathbf{X}, t)$. To illustrate the transmural variation of the cardiac strains and fiber strains, we evaluate equations (6) and (7) for varying depths, i.e., along the radial direction $\mathbf{X} = [0, 0, X_r]^t$.

4. RESULTS

4.1. Hemodynamics

Figure 2 shows the temporal evolution of the averaged left ventricular pressure LVP and left ventricular volume LVV over the cardiac cycle, left, and the resulting averaged pressure-volume loop, right. The left ventricular pressure exhibits the characteristic n-shape with small standard deviations. The left ventricular volume remains virtually unchanged during isovolumetric contraction from ED to EIC, with an end-diastolic volume of 57.4 ± 10.5 mL. It decreases during ejection from EIC to ES, reaching an end-systolic volume of 43.9 ± 9.3 mL. Average stroke volume and average ejection fraction are 15 mL and 26%, respectively. With average end-systolic and end-diastolic pressures of 83.1 ± 7.0 mmHg and 8.1 ± 2.2 mmHg, the average cardiac energy production per beat is 0.15 Joule, corresponding to the area enclosed by the pressure-volume loop.

4.2. Spatio-Temporal Evolution of Cardiac Strains in the Anterior Wall

Figure 3 summarizes the temporal evolution of the average cardiac strains in the anterior-basal left ventricular midwall, i.e., at 50% depth, at which the fiber direction is approximately aligned with the circumferential direction. The circumferential strains E_{CC} displayed on the top left are slightly tensile after end diastole with a maximum value of $+2.0 \pm 1.7\%$ at end isovolumetric contraction, then become compressive and decrease towards their minimum value of $-7.9 \pm 3.8\%$ at end isovolumetric relaxation, before returning back to zero at end diastole. The radial strains E_{RR} displayed on the bottom left behave in the opposite way. The wall thins slightly from end diastole towards $-3.8 \pm 3.5\%$ at end isovolumetric contraction, then thickens during systole towards $+11.3 \pm 6.4\%$ at end isovolumetric relaxation, before returning back to zero at end diastole.

Figure 4 displays the spatio-temporal evolution of the average cardiac strains across the anterior-basal left ventricular wall over three consecutive beats, where red indicates positive, i.e., tensile strains, and blue indicates negative, i.e., compressive strains. Mechanical activation characterized through the transition from red to blue in the circumferential strain

E_{CC} begins at the endocardium, the inner wall, at 90% depth, and then progresses towards the epicardium, the outer wall, at 10% depth. Mechanical deactivation, however, progresses vice versa, beginning at the epicardium and progressing towards the endocardium. This implies that the cells that are *activated first* are *deactivated last*. Mechanical deactivation proceeds faster than mechanical activation as indicated by the steeper slope in the transition from blue to red. This mechanical excitation pattern closely resembles the electrical excitation pattern starting at the endocardium, progressing gradually outward towards the epicardium, as the activation wave spreads across the heart [14]. This observation is in excellent agreement with the electrophysiological response of endocardial cells, close to the Purkinje fiber system, which display a longer action potential duration than epicardial cells [22, 27], meaning they are electrically activated first and deactivated last. Temporal strain variations throughout the cardiac cycle are larger in the endocardium than in the epicardium for both circumferential E_{CC} and radial E_{RR} strains.

4.3. Spatio-Temporal Evolution of Cardiac Strains in the Lateral Wall

Figure 5 displays the temporal evolution of the average cardiac strains at the lateral-equatorial left ventricular midwall, i.e., at 50% depth. For all six strain components, the general trend in the lateral-equatorial wall is similar to the anterior-basal wall, displayed in Figure 3. There are some minor differences in the circumferential strains E_{CC} at end isovolumetric relaxation, which seem to be less pronounced in the lateral midwall at $-5.1 \pm 6.0\%$ than in the anterior midwall at $-7.9 \pm 3.8\%$. However, the longitudinal extension E_{LL} at end isovolumetric contraction seems to be more pronounced in the lateral midwall at $+4.9 \pm 4.4\%$ than in the anterior midwall at $+0.8 \pm 3.8\%$.

Figure 6 depicts the spatio-temporal evolution of the average cardiac strains across the lateral-equatorial left ventricular wall, where, again, red indicates positive, i.e., tensile strains, and blue indicates negative, i.e., compressive strains. The general trend seems to be similar to that of the anterior wall, displayed in Figure 5. Overall, transmural strain variations in the lateral wall seem to be less pronounced than those in the anterior wall, except for the change in the radial strain which is similar to that of the anterior wall. In addition, mechanical activation of the lateral-equatorial wall seems to be *slightly delayed* as compared to that of the anterior-basal wall. This manifests itself in both circumferential E_{CC} and radial E_{RR} strains which are mechanically activated only at end systole. This mechanical activation delay agrees nicely with the electrical activation delay of the lateral equatorial wall with respect to the anterior basal wall, which has been reported to be one of the regions of early electrical activation [14].

4.4. Spatio-Temporal Evolution of Fiber Strains in the Anterior and Lateral Wall

Figure 7 illustrates the temporal evolution of the average fiber strains E_{FF} , both in the anterior-basal and lateral-equatorial left ventricular wall. The general trends are similar in the anterior and lateral wall. Transmurally, the maximum fiber stretch E_{FF} occurs at end isovolumetric contraction at $+3.5 \pm 4.1\%$ close to the epicardium, decreasing transmurally towards the endocardium, where the maximum fiber stretch E_{FF} is less than $+0.5 \pm 0.9\%$. Transmurally, the maximum fiber contraction E_{FF} occurs between end systole and end isovolumetric relaxation at $-8.0 \pm 2.5\%$ close to the endocardium and in the midwall, decreasing transmurally towards the epicardium, where maximum fiber contraction E_{FF} is only $-6.2 \pm 4.6\%$. Overall, this indicates that fiber strains display a *significant transmural heterogeneity*, being more contractile in the inner wall than in the outer wall and more contractile at the anterior site than at the lateral site. Temporally, maximum fiber contraction is found between end systole and isovolumetric relaxation in the endocardium, progressing across the wall and reaching the epicardium at end isovolumetric relaxation. In this sense, the mechanical activation sequence nicely follows the electrical activation sequence, from

endocardium to epicardium, from the anterior-basal to the lateral equatorial wall, reported in the literature [14]. In general, fiber contraction seems to be more pronounced and more coordinated in the anterior-basal than in the lateral-equatorial wall.

Last, Figure 8 summarizes the spatio-temporal evolution of the average fiber strains E_{FF} , both in the anterior-basal and lateral-equatorial wall. It confirms the general trends observed in Figure 7. Fiber shortening, represented through blue colors, seems to be more coordinated in the anterior than in the lateral wall. Fiber pre-stretch, indicated through red colors, spans the entire anterior-basal wall and the upper half of the lateral-equatorial wall. Some small pre-stretched regions in the endocardium of the lateral-equatorial wall might be attributed to the tension exerted by the papillary muscles located in this area. Fiber strain profiles are transmurally homogeneous at end diastole, see the almost vertical orange zero-strain lines near end diastole in Figures 8, top and bottom. This indicates that *fiber relaxation occurs almost simultaneously across the wall* both in the anterior and lateral regions. However, fiber strain profiles are transmurally heterogeneous during systole, see the inclined orange zero-strain lines from end isovolumetric contraction towards end systole in Figures 8, top and bottom. This indicates that *fiber contraction is initiated at the endocardium* slightly after end diastole *and progresses transmurally outward*, reaching the epicardium slightly before end systole. Fiber relaxation, indicated through the transition from blue to orange, proceeds faster than the fiber contraction, indicated through the transition from orange to blue. Accordingly, *fibers near the outer wall are in contraction for approximately half of the cardiac cycle while fibers near the inner wall are in contraction almost throughout the entire cardiac cycle*. Roughly speaking, in Figures 8, blue contractile regions with $E_{FF} < 0$ are more pronounced close to the endocardium, while red tensile regions with $E_{FF} > 0$ are more pronounced close to the epicardium. This effect is more pronounced in the anterior-basal than in the lateral-equatorial wall.

5. DISCUSSION

5.1. Limitations

A limitation inherent to the proposed marker technique is that it is invasive in nature and therefore not applicable to humans. Although ovine hearts behave relatively similar to human hearts, we need to be careful when extrapolating our findings to cardiac wall dynamics in humans. In fact, the overall goal of this research is to utilize our videofluoroscopic marker technique to validate common non-invasive imaging techniques used in human patients in the clinic such as tissue Doppler imaging and speckle tracking echocardiography. Another potential limitation might be that we perform the data acquisition right after the surgical procedure itself. The implantation of myocardial markers and transmural beads is an invasive procedure associated with local left ventricular wall trauma which might potentially lead to acute changes in ventricular wall dynamics. This might explain our relatively low ejection fraction and our relatively low strain values when compared with canine models reported in the literature [35], which, however, agree relatively well with our previous results in ovine models [7]. To quantify these acute effects, we have previously acquired marker and bead coordinates immediately after implantation and several weeks post surgery. While hemodynamics were unfavorably impacted by the surgical intervention, differences in absolute strain values were only minor, and general trends remained unchanged [26]. Lastly, it could be viewed as a limitation that the results of our computational analysis are sensitive to the chosen reference configuration. After experimenting with different reference configurations, we have chosen end diastole as the strain free reference state. While the choice of the reference state obviously influences the absolute values of the individual strain components, general trends remain unaltered. Different reference configurations basically induce a vertical shift of the strain vs time curves, but the strain amplitudes remain virtually unchanged. To extract fiber strains, we

have used generic fiber orientation maps generated by tissue histology averaged over five animals. It would, of course, be more appropriate to utilize subject-specific fiber orientation maps when projecting the strain tensor onto local fiber directions.

5.2. Clinical Significance

Left ventricular wall strains and active fiber contraction are believed to be important indicators for left ventricular remodeling. Altered wall strains can result in the production of cytokines and reactive oxygen species stimulating cardiomyocyte apoptosis and extracellular matrix disruption which, in turn, might trigger global left ventricular dilation and remodeling. This study therefore aimed at establishing a baseline characterization of cardiac and fiber strains throughout the cardiac cycle to quantify temporal, regional, and transmural variations of active fiber contraction in the healthy beating heart. Our observed mechanical excitation patterns are in excellent qualitative agreement with the electrical excitation patterns reported in the literature [14]. They complement our in vitro database of strains and stresses in healthy cardiac muscle tissue [3]. The identification of deviations from these baseline values is clinically important to guide the optimal timing of clinical interventions, e.g., in asymptomatic severe mitral regurgitation [4] or in progressive heart failure [20]. The permanent nature of the implanted beadsets renders the proposed marker technology an ideal methodology to quantify chronic kinematic changes during progressive growth and remodeling [21].

5.3. Engineering Relevance

The past two decades have seen a clear trend towards patient-specific computational treatment planning in cardiovascular disease. This manuscript has established a baseline characterization of ventricular wall dynamics that might be useful to calibrate and validate mechanistic material models and computational simulation tools for passive cardiac relaxation [19], active cardiac contraction [18], and potential viscous effects [24]. Lastly, and maybe most importantly, current therapies for heart failure seek to support the ventricular wall to reduce cardiomyocyte overstretch by using synthetic or living tissue engineered solutions [12]. This study provides useful guidelines for the structural and functional requirements of these biomaterials and may therefore be relevant to tissue engineering in general.

6. CONCLUSIONS

In summary, this study demonstrates that cardiac wall strains display significant temporal, regional, and transmural variations. Maximum fiber contraction was found to vary from $-8.0 \pm 2.5\%$ and $-5.4 \pm 3.1\%$ in the anterior and lateral endocardium to $-6.2 \pm 4.6\%$ and $-4.6 \pm 4.2\%$ in the anterior and lateral epicardium. Mechanical activation is initiated at the endocardium progressing transmurally outwards towards the epicardium, whereas mechanical deactivation occurs almost simultaneously across the wall. Fibers near the outer wall are therefore in contraction for approximately half of the cardiac cycle while fibers near the inner wall are in contraction almost throughout the entire cardiac cycle. We believe that this work provides a valuable baseline data set to judge strain alterations in response to cardiac disease and treatment, to validate alternative non-invasive imaging modalities, to calibrate and validate material models and computational tools for active cardiac contraction, and to engineer synthetic or living biomaterials with similar structure and function.

Acknowledgments

This work was supported by the US National Institutes of Health grants R01 HL 29589 and R01 HL67025 (Dr Miller); the US National Science Foundation CAREER award CMMI-0952921 (Dr Kuhl); the Swiss National Science Foundation Fellowship PBELP2-130913 (Dr Tsamis); the Western States Affiliate American Heart Association Fellowship (Dr Swanson); the US-Norway Fulbright Foundation, the Swedish Heart-Lung Foundation, and the Swedish Society for Medical Research (Dr Kvitting); and the Deutsche Herzstiftung, Frankfurt, Germany, Research Grant S/06/07 (Dr Bothe).

References

1. Arts T, Costa KD, Covell JW, McCulloch AD. Relating myocardial laminar architecture to shear strain and muscle fiber orientation. *Am H Physiol Heart Physiol*. 2001; 280:H2222–2229.
2. Böl M, Reese S, Parker KK, Kuhl E. Computational modeling of muscular thin films for cardiac repair. *Comp Mech*. 2009; 43:535–544.
3. Böl M, Abilez OJ, Assar AN, Zarins CK, Kuhl E. In vitro / in silico characterization of active and passive stresses in cardiac muscle. *Int J Multiscale Comp Eng*. 2011 in press.
4. Carhäll C, Nguyen TC, Itoh A, Ennis DB, Bothe W, Liang D, Ingels NB, Miller DC. Alterations in transmural myocardial strain: An early marker of left ventricular dysfunction in mitral regurgitation? *Circulation*. 2008; 118:S256–S262. [PubMed: 18824764]
5. Cheng A, Langer F, Rodriguez F, Criscione JC, Daughters GT, Miller DC, Ingels NB. Transmural cardiac strains in the lateral wall of the ovine left ventricle. *J Physiol Heart Circ Physiol*. 2005; 288:H1546–H1556.
6. Cheng A, Langer F, Rodriguez F, Criscione JC, Daughters GT, Miller DC, Ingels NB. Transmural sheet strains in the lateral wall of the ovine left ventricle. *J Physiol Heart Circ Physiol*. 2005; 289:H1234–H1241.
7. Cheng A, Nguyen TC, Malinowski M, Ennis DB, Daughters GT, Miller DC, Ingels NB. Transmural left ventricular shear strain alterations adjacent to and remote from infarcted myocardium. *J Heart Valve Dis*. 2006; 15:209–218. [PubMed: 16607903]
8. Cheng A, Nguyen TC, Malinowski M, Langer F, Liang D, Daughters GT, Ingels NB, Miller DC. Passive ventricular constraint prevents transmural shear strain progression in left ventricle remodeling. *Circulation*. 2006; 114:I79–I86. [PubMed: 16820650]
9. Cheng A, Nguyen TC, Malinowski M, Daughters GT, Miller DC, Ingels NB. Heterogeneity of left ventricular wall thickening mechanisms. *Circulation*. 2008; 118:713–721. [PubMed: 18663088]
10. Coppola BA, Covell JW, McCulloch AD, Omens JH. Asynchrony of ventricular activation affects magnitude and timing of fiber stretch in late-activated regions of the canine heart. *Am J Physiol Heart Physiol*. 2007; 293:H754–H761.
11. Costa KD, Takayama Y, McCulloch AD, Covell JW. Laminar fiber architecture and three-dimensional systolic mechanics in canine ventricular myocardium. *Am H Physiol Heart Physiol*. 1999; 276:595–607.
12. Christman KL, Lee RJ. Biomaterials for the treatment of myocardial infarction. *J Am College of Cardiol*. 2006; 48:907–913.
13. Daughters GT, Sanders WJ, Miller DC, Schwarzkopf A, Mead CW, Ingels NB. A comparison of two analytical systems for 3-D reconstruction from biplane videograms. *Proc of IEEE Computers in Cardiology*. 1989; 15:79–82.
14. Durrer D, van Dam RT, Freud GE, Janse MJ, Meijler FL, Arzbaeher RC. Total excitation of the isolated human heart. *Circulation*. 1970; 41:899–912. [PubMed: 5482907]
15. Ennis DB, Nguyen TC, Riboh JC, Wingström L, Harrington KB, Daughters GT, Ingels NB, Miller DC. Myofiber angle distributions in the ovine left ventricle do not conform to computationally optimized predictions. *J Biomech Eng*. 2008; 41:3219–3224.
16. Fenton TR, Cherry JM, Klassen GA. Transmural myocardial deformation in the canine left ventricular wall. *Am J Physiol*. 1978; 235:H523–H530.
17. Gaudron P, Eilles C, Kubler I, Ertl G. Progressive left ventricular dysfunction and remodeling after myocardial infarction. *Circulation*. 1993; 87:755–763. [PubMed: 8443896]

18. Göktepe S, Kuhl E. Electromechanics of cardiac tissue: A unified approach to the fully coupled excitation-contraction problem. *Comp Mech.* 2010; 45:227–243.
19. Göktepe S, Acharya SNS, Wong J, Kuhl E. Computational modeling of passive myocardium. *Int J Num Meth Biomed Eng.* 2011; 27:1–12.
20. Göktepe S, Abilez OJ, Kuhl E. A generic approach towards finite growth with examples of athlete's heart, cardiac dilation, and cardiac wall thickening. *J Mech Phys Solids.* 2010; 58:1661–1680.
21. Göktepe S, Abilez OJ, Parker KK, Kuhl E. A multiscale model for eccentric and concentric cardiac growth through sarcomerogenesis. *J Theor Bio.* 2010; 265:433–442. [PubMed: 20447409]
22. Göktepe S, Wong J, Kuhl E. Atrial and ventricular fibrillation - Computational simulation of spiral waves in cardiac tissue. *Arch Appl Mech.* 2010; 80:569–580.
23. Lloyd-Jones D, Adams RJ, Brown TM, Carnethon M, Dai S, De Simone G, Ferguson TB, Ford E, Furie K, Gillespie C, Go A, Greenlund K, Haase N, Hailpern S, Ho PM, Howard V, Kissela B, Kittner S, Lackland D, Lisabeth L, Marelli A, McDermott MM, Meigs J, Mozaffarian D, Mussolino M, Nichol G, Roger VL, Rosamond W, Sacco R, Sorlie P, Stafford R, Thom T, Wasserthiel-Smoller S, Wong ND, Wylie-Rosett J. Heart disease and stroke statistics - 2010 Update. A report from the American Heart Association. *Circulation.* 2010; 121:e46–e215. [PubMed: 20019324]
24. Katsnelson LB, Nikitina LV, Chemla D, Solovyova O, Coirault C, Lecarpentier Y, Markhasin VS. Influence of viscosity on myocardial mechanical activity: A mathematical model. *J Theor Bio.* 2004; 230:385–405. [PubMed: 15302547]
25. Kindberg K, Carhäll C, Karlsson M, Nguyen TC, Cheng A, Langer F, Rodriguez F, Daughters GT, Miller DC, Ingels NB. Transmural strains in the ovine left ventricular lateral wall during diastolic filling. *J Biomech Eng.* 2007; 131:061004-1–061004-8. [PubMed: 19449958]
26. Kindberg K, Karlsson M, Ingels NB, Criscione JC. Nonhomogeneous strain from sparse marker arrays for analysis of transmural myocardial mechanics. *J Biomech Eng.* 2009; 129:603–610. [PubMed: 17655482]
27. Kotikanyadanam M, Göktepe S, Kuhl E. Computational modeling of electrocardiograms - A finite element approach towards cardiac excitation. *Int J Num Meth Biomed Eng.* 2010; 26:524–533.
28. Leong KF, Chua CK, Sudarmadji N, Yeong WY. Engineering functionally graded tissue engineering scaffolds. *J Mech Beh Biomed Mat.* 2008; 1:14–152.
29. McCulloch AD, Omens JH. Non-homogeneous analysis of three-dimensional transmural finite deformation in canine ventricular myocardium. *J Biomechanics.* 1991; 24:539–548.
30. McGee EC, Gillinov AM, McCarthy PM. Reverse ventricular remodeling: mechanical options. *Curr Opin Cardiol.* 2006; 21:215–220. [PubMed: 16601460]
31. Niczyporuk MA, Miller DC. Automatic tracking and digitization of multiple radiopaque myocardial markers. *Comp Biomed Res.* 1991; 24:129–142.
32. Rausch MK, Bothe W, Kvitting JP, Göktepe S, Miller DC, Kuhl E. In vivo dynamic strains of the ovine anterior mitral valve leaflet. *J Biomechanics.* in press. 10.1016/j.jbiomech.2011.01.020
33. Rausch MK, Bothe W, Kvitting JP, Swanson JC, Ingels NB, Miller DC, Kuhl E. Characterization of mitral valve annular dynamics in the beating heart. *Ann Biomed Eng.* in press. 10.1007/s10439-011-0272-y
34. Takayama Y, Costa KD, Covell JW. Contribution of laminar myofiber architecture to load-dependent changes in mechanics of LV myocardium. *Am J Physiol Heart Physiol.* 2002; 282:H1510–H1520.
35. Waldman LK, Fung YC, Covell JW. Transmural myocardial deformation in the canine left ventricle. Normal in vivo three-dimensional finite strains. *Circ Res.* 1985; 57:152–163. [PubMed: 4006099]
36. Wollert KC, Meyer GP, Lotz J, Ringes-Lichtenberg S, Lippolt P, Breidenbach C, Fichtner S, Korte T, Hornig B, Messinger D, Arseniev L, Hertenstein B, Ganser A, Drexler H. Intracoronary autologous bone-marrow cell transfer after myocardial infarction: the BOOST randomised controlled clinical trial. *Lancet.* 2004; 364:141–148. [PubMed: 15246726]

Figure 1. Locations of transmural beadsets in the anterior-basal and lateral-equatorial left ventricular wall, left. Left ventricle with both beadsets silhouetted by thirteen epicardial markers, right. C , R , and L denote the axes of the local circumferential-radial-longitudinal coordinate system.

Figure 2.

Temporal evolution of left ventricular pressure and volume, left, and average pressure-volume loop, right. The left ventricular pressure exhibits the characteristic n-shape with small standard deviations. The left ventricular volume varies between 57.4 ± 10.5 mL at ED and 43.9 ± 9.3 mL at ES. Average stroke volume and average ejection fraction are 15 mL and 26%, respectively. Means and standard deviations are displayed for $n=9$ animals to the top of the curve. LVP left ventricular pressure, LVV left ventricular volume, ED end diastole, EIR end isovolumetric relaxation, ES end systole, and EIC end isovolumetric contraction.

Figure 3.

Temporal evolution of cardiac strains at 50% depth of the anterior-basal left ventricular wall. Maximum circumferential contraction E_{CC} is $-7.9 \pm 3.8\%$ and maximum wall thickening E_{RR} is $+11.3 \pm 6.4\%$. Means and standard deviations of components of the Green Lagrange strains are displayed for $n=9$ animals to the top of the curves. C circumferential, R radial, and L longitudinal direction. ED end diastole, EIR end isovolumetric relaxation, ES end systole, and EIC end isovolumetric contraction.

Figure 4.

Spatio-temporal evolution of cardiac strains in the anterior-basal left ventricular wall averaged over $n=9$ animals. Components of the Green Lagrange strains are displayed across the ventricular wall with 10% depth indicating the epicardium, the outer wall, and 90% depth indicating the endocardium, the inner wall. Strains are displayed for three consecutive heartbeats. C circumferential, R radial, and L longitudinal direction. ED end diastole, EIR end isovolumetric relaxation, ES end systole, and EIC end isovolumetric contraction.

Figure 5.

Temporal evolution of cardiac strains at 50% depth of the lateral-equatorial left ventricular wall. Maximum circumferential contraction is $-7.9 \pm 3.8\%$ and maximum wall thickening is $+10.5 \pm 5.2\%$. Means and standard deviations of components of the Green Lagrange strains are displayed for $n=9$ animals to the top of the curves. C circumferential, R radial, and L longitudinal direction. ED end diastole, EIR end isovolumetric relaxation, ES end systole, and EIC end isovolumetric contraction.

Figure 6.

Spatio-temporal evolution of cardiac strains in the lateral-equatorial left ventricular wall averaged over $n=9$ animals. Components of the Green Lagrange strains are displayed across the ventricular wall with 10% depth indicating the epicardium, the outer wall, and 90% depth indicating the endocardium, the inner wall. Strains are displayed for three consecutive heartbeats. C circumferential, R radial, and L longitudinal direction. ED end diastole, EIR end isovolumetric relaxation, ES end systole, and EIC end isovolumetric contraction.

Figure 7.

Temporal evolution of fiber strains at 20%, 50%, and 80% depth of the anterior-basal and lateral-equatorial left ventricular wall. Maximum fiber contraction varies from $-8.0 \pm 2.5\%$ and $-5.4 \pm 3.1\%$ in the anterior and lateral endocardium, i.e., at 80% depth, to $-6.2 \pm 4.6\%$ and $-4.6 \pm 4.2\%$ in the anterior and lateral epicardium, i.e., at 20% depth. Means and standard deviations are displayed for $n=9$ animals to the top of the curves. ED end diastole, EIR end isovolumetric relaxation, ES end systole, and EIC end isovolumetric contraction.

Figure 8.

Spatio-temporal evolution of fiber strains in the anterior-basal and lateral-equatorial left ventricular wall averaged over $n=9$ animals. Fiber strains are displayed across the ventricular wall with 10% depth indicating the epicardium, the outer wall, and 90% depth indicating the endocardium, the inner wall. Strains are displayed for three consecutive heartbeats. ED end diastole, EIR end isovolumetric relaxation, ES end systole, and EIC end isovolumetric contraction.