Substrate specificity of SpoIIGA, a signal-transducing aspartic protease in *Bacilli*

Received December 10, 2010; accepted January 20, 2011; published online February 28, 2011

Daisuke Imamura¹, Ritsuko Kuwana¹,
Lee Kroos², Michael Feig²,
Hiromu Takamatsu^{1,*} and Kazuhito Watabe¹

¹Faculty of Pharmaceutical Sciences, Setsunan University, Osaka 573-0101, Japan; and ²Department of Biochemistry and Molecular Biology, Michigan State University, East Lansing, MI, USA

*Hiromu Takamatsu, Faculty of Pharmaceutical Sciences, Setsunan University, Hirakata, Osaka 573-0101, Japan.
Tel/Fax: +81 72 866 3114, email: takamatu@pharm.setsunan.ac.jp

SpoIIGA is a novel type of membrane-associated aspartic protease that responds to a signal from the forespore by cleaving Pro- σ^E in the mother cell during sporulation of *Bacillus subtilis*. Very little is known about how SpoIIGA recognizes Pro- σ^E . By co-expressing proteins in *Escherichia coli*, it was shown that charge reversal substitutions for acidic residues 24 and 25 of Pro- σ^E , and for basic residues 245 and 284 of SpoIIGA, impaired cleavage. These results are consistent with a model predicting possible electrostatic interactions between these residues; however, no charge reversal substitution for residue 245 or residue 284 of SpoIIGA restored cleavage of Pro- σ^E with a charge reversal substitution for residue 24 or residue 25. *Bacillus subtilis* SpoIIGA cleaved Pro- σ^E orthologs from *Bacillus licheniformis* and *Bacillus halodurans*, but not from *Bacillus cereus*. A triple substitution in the pro-sequence of *B. cereus* Pro- σ^E allowed cleavage by *B. subtilis* SpoIIGA, indicating that residues distal from the cleavage site contribute to substrate specificity. Co-expression of SpoIIGA and Pro- σ^E orthologs in different combinations suggested that *B. licheniformis* SpoIIGA has a relatively narrow substrate specificity as compared with *B. subtilis* SpoIIGA, whereas *B. cereus* SpoIIGA and *B. halodurans* SpoIIGA appear to have broader substrate specificity.

Keywords: Aspartic protease/*Bacillus subtilis*/sigma factor/signal transduction/sporulation.

Abbreviations: BC, *Bacillus cereus*; BH, *Bacillus halodurans*; BL, *Bacillus licheniformis*; BS, *Bacillus subtilis*; F2, double FLAG tag; GFP, green fluorescent protein; H6, 6 histidine tag; HIV-1, human immunodeficiency virus type 1; IPTG, isopropyl β -D-thiogalactopyranoside; RNAP, RNA polymerase; VMD, Visual Molecular Dynamics; WT, wild type.

The Gram-positive bacterium *Bacillus subtilis* undergoes sporulation in response to starvation. Sporulation involves a highly ordered program of gene expression and morphological change (1). The

first morphological change of sporulation is the appearance of an asymmetrically positioned septum that divides the cell into a larger mother cell and a smaller forespore compartment (Fig. 1A, left). The first compartment-specific transcription factor to become active is σ^F in the forespore. This subunit of RNA polymerase (RNAP) directs transcription of at least 47 genes (2, 3), including *spoIIR*, whose product is believed to be secreted from the forespore into the space between the two membranes of the polar septum, and signals activation of σ^E in the mother cell (4–6) (Fig. 1B, middle). The σ^E is synthesized as an inactive precursor, Pro- σ^E , and is activated by proteolytic cleavage that removes its N-terminal 27 residues (7, 8). The gene (*spoIIGB* or *sigE*) encoding Pro- σ^E is co-transcribed in an operon with *spoIIGA* (9), whose product is necessary for the processing of Pro- σ^E to σ^E . The σ^E RNAP transcribes over 270 genes (2, 10–12). SpoIIGA and Pro- σ^E are expressed predominantly in the mother cell (13) (Fig. 1). SpoIIGA is predicted to have an N-terminal domain with five transmembrane segments embedded in the mother cell membrane (14) and a C-terminal aspartic protease domain in the mother cell cytoplasm (15). SpoIIGA dimerizes to form the catalytic site of the aspartic protease (see Fig. 3 which appears later). Pro- σ^E associates with the mother cell membrane via its N-terminal pro-sequence (16, 17) and is cleaved (Fig. 1, arrowhead) by SpoIIGA, releasing active σ^E into the mother cell. The products of certain genes under σ^F control in the forespore and other genes under σ^E control in the mother cell bring about further morphological change, resulting in the formation of an endospore (1).

Since *in vitro* processing of Pro- σ^E has not been achieved yet, it remained possible that SpoIIGA and SpoIIR modify the activity of an unidentified protein that directly cleaves Pro- σ^E . Recently, we reconstituted Pro- σ^E processing in *Escherichia coli* (Fig. 1C, right) and provided evidence that SpoIIR and SpoIIGA are necessary and sufficient for accurate processing of Pro- σ^E in a heterologous host, and that SpoIIGA is a novel type of signal-transducing aspartic protease that cleaves Pro- σ^E to σ^E (15). This *E. coli* system is also a powerful tool for mutagenesis to analyse Pro- σ^E processing. Pro- σ^E was co-expressed in *E. coli* with C-terminally double-FLAG-tagged SpoIIR (SpoIIR-F2) and with C-terminally tagged SpoIIGA (SpoIIGA-GFP-F2). SpoIIR is expected to be secreted to the periplasm where it could interact with SpoIIGA's inner membrane-embedded N-terminal domain, stimulating SpoIIGA's C-terminal domain to cleave inner membrane-associated Pro- σ^E , releasing σ^E to the cytoplasm (15) (Fig. 1). Expression of both

Fig. 1 Model of Pro- σ^E processing during sporulation of *B. subtilis* and in *E. coli*. (Left) Asymmetric septation during *B. subtilis* sporulation generates a smaller forespore compartment and a larger mother cell compartment. (Middle) An expanded view of the septum depicts proteins involved in Pro- σ^E processing. SpoIIR is produced in the forespore and is believed to cross the forespore membrane and interact with SpoIIGA, activating Pro- σ^E processing (4–6). (Right) SpoIIR, SpoIIGA and Pro- σ^E are produced under T7 RNAP control in *E. coli*. (Middle) An expanded view of the *E. coli* outer membrane, periplasm and inner membrane.

SpoIIR-F2 and SpoIIGA-GFP-F2 was necessary and sufficient for cleavage of Pro- σ^E to σ^E in *E. coli*. Modelling and mutational analysis of the C-terminal half of SpoIIGA provided evidence that it forms a dimer similar to the human immunodeficiency virus type 1 (HIV-1) protease and related retroviral aspartic proteases. However, unlike the retroviral proteases, SpoIIGA has an N-terminal, membrane-embedded domain (14, 18). SpoIIGA-GFP-F2 expressed in *E. coli* was membrane associated and appeared to form inactive dimers or oligomers, that upon interaction between their N-terminal domains and SpoIIR-F2 on the periplasmic side of the inner membrane was proposed to cause a conformational change in the cytoplasmic C-terminal domains of SpoIIGA-GFP-F2, allowing formation of active aspartic protease dimer capable of cleaving membrane-associated Pro- σ^E (15). According to this model, SpoIIGA is unique since previously described membrane-associated aspartic proteases have two catalytic aspartate residues in a single-polypeptide chain, and they have not been proposed to transduce a signal across a membrane. Very little is known about how SpoIIGA recognizes its substrate, Pro- σ^E . A previous study showed that E25 of Pro- σ^E is important for processing in *B. subtilis* (19). Also, the N-terminal 28 residues of Pro- σ^E were shown to be sufficient for processing of a Pro- σ^E - σ^K fusion protein (20), and the N-terminal 52–55 amino acids of Pro- σ^E were sufficient for processing when fused to proteins unrelated to σ -factors (16). These results suggested that some, if not all, of the information necessary for recognition by the SpoIIGA protease resides in the pro-sequence of Pro- σ^E (residues 1–27). Here we investigated substrate specificity of SpoIIGA in *Bacilli*.

Experimental Procedures

Plasmids

Plasmids used in this study are described in Supplementary Table SI, and primers used in their construction are listed in Supplementary Table SII. All cloned PCR products and all genes subjected to mutagenesis (QuikChange kit, Stratagene) were sequenced to confirm that no undesired mutations were present.

Co-transformation

Plasmids bearing different antibiotic-resistance genes were co-transformed into *E. coli* strain BL21(DE3) (Novagen) as described previously (15).

Induction of gene expression

Escherichia coli BL21(DE3) can be induced to synthesize T7 RNAP by addition of isopropyl β -D-thiogalactopyranoside (IPTG). This strain, bearing plasmids with genes fused to the T7 RNAP promoter, was grown in Luria–Bertani medium containing 50 μ g/ml kanamycin sulphate and/or 50 μ g/ml ampicillin overnight at 37°C with shaking. A 50 μ l of culture was transferred to 5 ml of fresh Luria–Bertani medium with antibiotics, and incubation was continued at 37°C with shaking at 140 r.p.m. until the culture reached 0.5 of OD₆₆₀. To induce gene expression, IPTG (0.5 mM) was added, and incubation with shaking was continued for 2 h since cleavage was shown previously to reach a maximum at this time (15).

Western blot analysis

Escherichia coli cells from cultures (1 ml) induced as described above were collected by centrifugation (12,000g). Samples were prepared by resuspending cells in 100 μ l of sample buffer [50 mM Tris–HCl, pH 6.8, 2% SDS, 10% (v/v) glycerol and 0.015% bromophenol blue] and boiling for 3 min. Samples were separated on SDS–12% polyacrylamide gels with Tris–Tricine electrode buffer (0.1 M Tris, 0.1 M Tricine and 0.1% SDS) and electroblotted to Immobilon-P membranes (Millipore). Membranes were incubated in TBS [20 mM Tris–HCl (pH 7.5), 0.5 M NaCl] with 5% non-fat dry milk for 1 h at room temperature with shaking to block non-specific antibody binding and then incubated overnight at room temperature with shaking in monoclonal antibody against σ^E (a gift from W. Haldenwang) at 1:1,000 dilution into TBS with 2% non-fat dry milk to detect pro- σ^E and σ^E . Horseradish peroxidase-conjugated goat anti-mouse IgG (Promega) were used at 1:1,000 dilution for 1 h at room temperature with shaking serving as the secondary antibodies. Horseradish peroxidase-conjugated antibodies to FLAG (Sigma) were used at 1:5,000 dilution for 1 h at room temperature with shaking to detect SpoIIR-F2 and SpoIIGA-GFP-F2. Chemiluminescent detection was performed according to the manufacturer's instructions (ECL kit; Amersham). Signals were captured using a charge-coupled-device camera (ATTO CoolSaver AE-6955; ATTO Corporation, Tokyo, Japan).

Results and discussion

Importance of conserved, charged residues close to the cleavage site in Pro- σ^E

An alignment of *B. subtilis* Pro- σ^E and its orthologs revealed that the pro-sequence is not as conserved as the downstream sigma factor domain (Supplementary

Fig. S1). However, E25 in the pro-sequence of *B. subtilis* Pro- σ^E appears to be conserved in *Bacilli* and some other species, although not in *Clostridia*. Likewise, D24 and K22 in the pro-sequence of *B. subtilis* Pro- σ^E appear to be highly conserved in *Bacilli*. In the HIV-1 protease, it is proposed that the opening of flaps exposes the electronegative active site, then, a neutral or positively charged substrate could potentially be guided into a conformation optimal for binding (21). We reasoned that one or more of the conserved, charged residues in pro-sequence of Pro- σ^E might interact electrostatically with a residue of opposite charge in SpoIIIGA to help position the substrate for cleavage. To test this hypothesis, we engineered substitutions in *B. subtilis* Pro- σ^E and in SpoIIIGA-GFP-F2, and we co-expressed mutant forms of these proteins together with SpoIIR-F2 in *E. coli* as described previously (15). Expression of SpoIIR-F2 from pID50 and its derivatives was normal (Supplementary Fig. S2). As noted previously (15), pID50 expresses Pro- σ^E with its methionine at residue 17 changed to isoleucine (M17I) to avoid initiation of translation at this position, which would interfere with the analysis of cleavage by SpoIIIGA-GFP-F2 (expressed from pID48) because the species produced from the alternative start codon co-migrates with σ^E . Also, the Pro- σ^E is C-terminally tagged with 6-histidine residues (H6). Neither the H6 tag nor the M17I substitution impairs cleavage (15), and we refer to pro- σ^E (M17I)-H6 as the wild-type substrate here.

We engineered three charge reversals in Pro- σ^E (K22D, D24K and E25K) by subjecting pID50 to site-directed mutagenesis, and the *sigE* gene was sequenced to ensure that only the desired mutation

was created. We chose to make charge reversal substitutions because we hoped to make compensatory charge reversals in SpoIIIGA (see below). As expected, the wild-type substrate was cleaved to σ^E -H6 by SpoIIIGA-GFP-F2 (Fig. 2A, lane 1) but not by the D183A mutant form (Fig. 2A, lane 2) (15). D183 is the putative catalytic aspartate residue of SpoIIIGA and was used as a negative control for SpoIIIGA activity in this study. The K22D substitution in Pro- σ^E -H6 had little or no effect on cleavage, indicating that charge reversal at residue 22 is tolerated (Fig. 2A, lanes 3 and 4). However, the D24K and E25K substitutions in Pro- σ^E -H6 prevented cleavage (Fig. 2A, lanes 5–8). The failure of the E25K mutant form of Pro- σ^E to be cleaved in the *E. coli* system is consistent with its failure to be cleaved by wild-type SpoIIIGA during *B. subtilis* sporulation (19). The inability of SpoIIIGA to cleave the D24K and E25K mutant forms of Pro- σ^E shows that charge reversals are not tolerated at these positions. These results are consistent with a model predicting that D24 and/or E25 might interact electrostatically with a residue of opposite charge in SpoIIIGA (see below).

Effects of changes in SpoIIIGA on cleavage of Pro- σ^E

Based on a structural model of SpoIIIGA in complex with a short segment of Pro- σ^E (15), we predicted that R245 of SpoIIIGA might interact with D24 and/or E25 of Pro- σ^E because the residues are in close proximity (Fig. 3) and of opposite charge. R245 is predicted to be located in a flap in each subunit of the putative SpoIIIGA dimer. The corresponding flaps in the HIV-1 protease dimer (the template for the structural model) control access of the substrate to the active site.

Fig. 2 Effects of substitutions in Pro- σ^E and in SpoIIIGA on cleavage. (A) Effects of charge reversals in conserved residues in the pro-sequence of Pro- σ^E . *Escherichia coli* bearing pID50 to produce SpoIIR-F2 and Pro- σ^E -H6 (WT), or a derivative of pID50 to produce SpoIIR-F2 and Pro- σ^E -H6 with the indicated substitution, and pID48 to produce active SpoIIIGA-GFP-F2 (WT), or pID53 to produce inactive SpoIIIGA(D183A)-GFP-F2, were induced with IPTG for 2 h. Intervening lanes were deleted from the image. (B) Effects of substitutions in SpoIIIGA on cleavage of wild-type and mutant Pro- σ^E . As described for panel A, *E. coli* bearing pID50 or a derivative, and pID48 or a derivative, were induced. For both panels, samples (10 μ l) were subjected to western blot analysis with antibodies against FLAG (upper blot) or σ^E (lower blot).

Fig. 3 Model of SpoIIGA in complex with a short segment of Pro- σ^E . Only the C-terminal domain of SpoIIGA is shown, with the two chains of the putative dimer coloured black and white (arrows and helices indicate β -sheet and α -helix secondary structures, respectively), except van der Waals surfaces are shown for the putative catalytic aspartates (D183) and residues hypothesized to play a role in substrate recognition (R245 and K284) or flap function (P259) for each chain. Only a short segment of Pro- σ^E is shown (residues 22–33), in stick representation coloured yellow, except van der Waals surfaces are shown for K22, D24 and E25. The left part shows a 'side view' and the right part shows a 'top view' of the complex. The 'side view' is the orientation typically shown for HIV-1 protease. Note that both the N-terminus (i.e. the point of attachment to the N-terminal membrane domain) and C-terminus of each SpoIIGA chain is at the bottom in the 'side view', which is upside down relative to the depiction in Fig. 1. This image was made with Visual Molecular Dynamics (VMD) software support. VMD is developed with NIH support by the Theoretical and Computational Biophysics group at the Beckman Institute, University of Illinois at Urbana-Champaign.

Alignment of *B. subtilis* SpoIIGA with its orthologs showed that R245 appears to be highly conserved in *Bacilli*, although not in other species (Supplementary Fig. S3). We engineered an R245D substitution in *B. subtilis* SpoIIGA-GFP-F2 and co-expressed it with SpoIIR-F2 and wild-type or mutant forms of Pro- σ^E -H6 in *E. coli*. The R245D substitution abolished the ability of SpoIIGA-GFP-F2 to cleave wild-type Pro- σ^E -H6 (Fig. 2B, lane 3), consistent with the idea that R245 interacts with the substrate. However, SpoIIGA(R245D)-GFP-F2 did not cleave the D24K or E25K mutant form of Pro- σ^E -H6 (Fig. 2B, lanes 8 and 13). Perhaps R245 of SpoIIGA interacts with another residue in Pro- σ^E or the particular substitutions we chose are incompatible with the predicted interaction. In addition to possibly interacting with Pro- σ^E , R245 of SpoIIGA is predicted to form a salt bridge with D190 that might be important for structural stability in the substrate binding cavity (15), so the R245D substitution might disrupt that interaction, although the altered protein accumulates normally (Fig. 2B, lanes 3, 8 and 13).

A K284D substitution in SpoIIGA-GFP-F2 was tested in the same way as the R245D substitution. Based on our structural model (15), K284 is predicted to be in a loop near the substrate, so it might interact with D24 and/or E25 (Fig. 3). However, this seemed less likely than an interaction between D24 and/or E25 of Pro- σ^E and R245 of SpoIIGA, since K284 is not conserved among orthologs of SpoIIGA in *Bacilli* (Supplementary Fig. S3). The K284D substitution in SpoIIGA-GFP-F2 reduced its activity towards wild-type Pro- σ^E -H6 (Fig. 2B, lane 5) and did not permit cleavage of the D24K and E25K mutant forms of Pro- σ^E -H6 (Fig. 2B, lanes 10 and 15).

The effects of the R245D and K284D substitutions in SpoIIGA on cleavage of wild-type Pro- σ^E provide the first evidence that R245 and K284 are important

for SpoIIGA function. Although our results do not provide evidence that R245 or K284 of SpoIIGA interact with D24 or E25 of Pro- σ^E , neither do they rule out such interactions.

Since SpoIIGA(P259L) had been identified as a suppressor of Pro- σ^E (E25K) previously in sporulating *B. subtilis* (22), we also tested the effects of a P259L substitution in SpoIIGA-GFP-F2 using our *E. coli* system. In *B. subtilis* during sporulation, SpoIIGA(P259L) cleaves both wild-type Pro- σ^E and the E25K mutant form (22) [Pro- σ^E (D24K) was not tested] suggesting that SpoIIGA(P259L) has broadened substrate specificity. It is conceivable that the broadened substrate specificity of SpoIIGA(P259L) is due to increased mobility of flaps that in the HIV-1 protease control access of the substrate to the active site because P259 is located at the base of the flap (Fig. 3). Surprisingly, the P259L substitution in SpoIIGA-GFP-F2 abolished its activity on wild-type Pro- σ^E (Fig. 2B, lane 4) and failed to permit cleavage of the D24K and E25K mutant forms of Pro- σ^E (Fig. 2B, lanes 9 and 14). There are many differences between sporulating *B. subtilis* and the *E. coli* system that might account for the different results (Fig. 1), including the composition of the membrane into which SpoIIGA is inserted. We speculate that in *E. coli*, SpoIIGA(P259L) cannot close the flaps tightly enough to bind the substrate and permit cleavage. We note that P259 appears to be highly conserved in SpoIIGA orthologs (Supplementary Fig. S3).

Ability of *B. subtilis* SpoIIGA to cleave Pro- σ^E orthologs

To further investigate how SpoIIGA recognizes Pro- σ^E , we took advantage of the natural variation in the pro-sequence among Pro- σ^E orthologs (Supplementary Fig. S1). We cloned the *sigE* genes that code for Pro- σ^E orthologs from *Bacillus licheniformis*, *Bacillus cereus* and *Bacillus halodurans*, which

Fig. 4 Cleavage of Pro- σ^E orthologs by *B. subtilis* SpoIIIGA. (A) Alignment of the N-terminal region of Pro- σ^E orthologs tested for cleavage. The N-terminal sequences of Pro- σ^E from *B. subtilis* (BS), *B. licheniformis* (BL), *B. cereus* (BC) and *B. halodurans* (BH) were aligned using ClustalW (23). An arrow indicates the cleavage site in *B. subtilis* Pro- σ^E . Underlining indicates differences from the *B. subtilis* sequence. Bold case letters indicate residues subjected to substitutions in the BC pro-sequence to corresponding BS residues in Supplementary Fig. S4B. At the bottom, asterisks, double dots and dots indicate identical, conserved and semi-conserved residues, respectively. (B) Cleavage of Pro- σ^E orthologs by *B. subtilis* SpoIIIGA. *Escherichia coli* bearing pID50, pID129, pID121 or pID122 to produce *B. subtilis* SpoIIR-F2 and Pro- σ^E -H6 from *B. subtilis* (BS), *B. licheniformis* (BL), *B. cereus* (BC) or *B. halodurans* (BH), respectively, and pID48 to produce active *B. subtilis* SpoIIIGA-GFP-F2, or pID53 to produce inactive *B. subtilis* SpoIIIGA(D183A)-GFP-F2, were induced with IPTG for 2 h. Intervening lanes were deleted from the image. (C) Cleavage of triply substituted *B. cereus* Pro- σ^E by *B. subtilis* SpoIIIGA. *E. coli* bearing pID121 or pID176 to produce *B. subtilis* SpoIIR-F2 and Pro- σ^E -H6 from *B. cereus* [Pro- σ^E (BC)] or its derivative [Pro- σ^E (BC, F6L, Y7R, V9T)], respectively, and pID48 to produce active *B. subtilis* SpoIIIGA-GFP-F2, pID53 to produce inactive *B. subtilis* SpoIIIGA(D183A)-GFP-F2, or pID126 to produce SpoIIIGA(BC)-GFP-F2 were induced with IPTG for 2 h. For (B and C), samples (10 μ l) were subjected to western blot analysis with antibodies against FLAG (upper blot) or σ^E (lower blot).

differ in their pro-sequence at 2, 10 and 13 positions, respectively, from *B. subtilis* Pro- σ^E (Fig. 4A). Each *sigE* gene was cloned downstream of a T7 RNAP promoter in a plasmid that also expresses SpoIIR-F2 from a T7 RNAP promoter. Expression of Pro- σ^E -H6 from *B. licheniformis*, Pro- σ^E (BL)-H6, in *E. coli* generated a smaller species (data not shown), presumably because methionine at position 17 served as an alternative translation initiation codon (Fig. 4A), as observed previously for *B. subtilis* Pro- σ^E -H6 (15). Therefore, an M17I substitution was made in Pro- σ^E (BL)-H6 (analogous to the one made in *B. subtilis* Pro- σ^E -H6). Expression of Pro- σ^E -H6 from *B. cereus*,

Fig. 5 Combinations of SpoIIIGA and Pro- σ^E orthologs. *Escherichia coli* bearing pID50 (lanes 1–4), pID129 (lanes 5–8), pID121 (lanes 9–12) or pID122 (lanes 13–16) to produce *B. subtilis* SpoIIIR-F2 and Pro- σ^E -H6 orthologs, and pID48 (lanes 1, 5, 9 and 13) to produce SpoIIIGA(BS)-GFP-F2, pID125 (lanes 2, 6, 10 and 14) to produce SpoIIIGA(BL)-GFP-F2, pID126 (lanes 3, 7, 11 and 15) to produce SpoIIIGA(BC)-GFP-F2, or pID173 (lanes 4, 8, 12 and 16) to produce SpoIIIGA(BH) were induced with IPTG for 2 h. Samples (10 μ l) were subjected to western blot analysis with antibodies against FLAG (upper blot) or σ^E (lower blot).

Pro- σ^E (BC)-H6, or *B. halodurans*, Pro- σ^E (BH)-H6, in *E. coli* did not produce a smaller species (Fig. 4B, lanes 6 and 8). Interestingly, upon co-expression in *E. coli*, *B. subtilis* SpoIIIGA-GFP-F2 cleaved Pro- σ^E (BL)-H6 and Pro- σ^E (BH)-H6, but not Pro- σ^E (BC)-H6 (Fig. 4B). These results suggest that the Pro- σ^E (BC) pro-sequence is not recognized by *B. subtilis* SpoIIIGA.

To demonstrate that Pro- σ^E (BC) could be cleaved by its cognate enzyme, we cloned the gene encoding, the SpoIIIGA ortholog from *B. cereus*, SpoIIIGA(BC), and engineered expression in *E. coli*. Although SpoIIIGA(BC)-GFP-F2 was not expressed as highly as *B. subtilis* SpoIIIGA-GFP-F2 in *E. coli*, Pro- σ^E (BC)-H6 was readily cleaved by SpoIIIGA(BC)-GFP-F2 (Fig. 5, lane 11). As shown in Supplementary Fig. S4A, this cleavage depended on co-expression of *B. subtilis* SpoIIR-F2, indicating that the *B. subtilis* signalling protein stimulates SpoIIIGA(BC)-GFP-F2.

To investigate the inability of *B. subtilis* SpoIIIGA to cleave Pro- σ^E (BC), we compared the pro-sequences of the Pro- σ^E orthologs (Supplementary Fig. S1). We considered F6, Y7, V9, V15 and T23 of Pro- σ^E (BC) to be the most likely residues to explain its inability to be cleaved by *B. subtilis* SpoIIIGA (Fig. 4A). The five residues of Pro- σ^E (BC)-H6 were changed individually to those of *B. subtilis*, creating F6L, Y7R, V9T, V15L and T23S mutant forms of Pro- σ^E (BC)-H6, which were tested for cleavage by *B. subtilis* SpoIIIGA-GFP-F2 and by SpoIIIGA(BC)-GFP-F2 (Supplementary Fig. S4B). None of the individual substitutions enabled Pro- σ^E (BC)-H6 to be cleaved by *B. subtilis* SpoIIIGA-GFP-F2. We considered the possibility that several residues of Pro- σ^E (BC) need to be changed simultaneously in order to allow cleavage by *B. subtilis*

SpoIIGA. Noting that F6, Y7 and V9 are highly diverged from the corresponding residues in *B. subtilis* Pro- σ^E , we hypothesized that these residues are incompatible with cleavage by *B. subtilis* SpoIIGA, because this enzyme had been shown to cleave *B. subtilis* Pro- σ^E lacking residues 2–11 (19), so this region is not required for substrate recognition. Therefore, we constructed triply substituted Pro- σ^E (BC)-H6 in which F6, Y7 and V9 were changed to the corresponding residues in *B. subtilis* Pro- σ^E . The resulting Pro- σ^E (BC, F6L, Y7R, V9T)-H6 was reproducibly cleaved by *B. subtilis* SpoIIGA-GFP-F2 (Fig. 4C, lane 4), although not as efficiently as by SpoIIGA(BC)-GFP-F2 (Fig. 4C, lane 6). Probably, more residues of Pro- σ^E (BC) would need to be changed to allow full cleavage by *B. subtilis* SpoIIGA. In any case, the result with triply-substituted Pro- σ^E (BC)-H6 indicates that residues distal from the cleavage site can inhibit cleavage by SpoIIGA and therefore contribute to the specificity of the cleavage reaction.

Specificity of SpoIIGA orthologs

To explore the specificity of SpoIIGA orthologs further, we cloned the putative proteases from *B. halodurans* and *B. licheniformis*. We tested the cleavage of all combinations of SpoIIGA and Pro- σ^E from species used in this study, using co-expression in *E. coli*, and in each case *B. subtilis* SpoIIR-F2 was also co-expressed (Fig. 5). The *B. halodurans* ortholog, SpoIIGA(BH), was not tagged with GFP-F2 because SpoIIGA(BH)-GFP-F2 was inactive (data not shown), so we were not able to verify expression of SpoIIGA(BH). However, all four Pro- σ^E substrates were cleaved in *E. coli* engineered to co-express SpoIIGA(BH) (Fig. 5, lanes 4, 8, 12 and 16). Likewise, SpoIIGA(BC)-GFP-F2 cleaved all four substrates (Fig. 5, lanes 3, 7, 11 and 15). As noted above, *B. subtilis* SpoIIGA(BS)-GFP-F2 cleaved all except Pro- σ^E (BC)-H6 (Fig. 5, lanes 1, 5, 9 and 13). Interestingly, SpoIIGA(BL)-GFP-F2 cleaved only its cognate substrate (Fig. 5, lanes 2, 6, 10 and 14). These results suggest that SpoIIGA(BL) has a relatively narrow substrate specificity as compared with SpoIIGA(BS), whereas SpoIIGA(BC) and SpoIIGA(BH) appear to have broader substrate specificity. The four enzymes are nearly identical in the region around the catalytic aspartate [D183 of SpoIIGA(BS)] and all have isoleucine or valine at critical positions [I294 and I295 of SpoIIGA(BS)] in a β -sheet secondary structure predicted to position the catalytic aspartate (Supplementary Fig. S3), so it is unlikely that these regions account for differences in substrate specificity. On the other hand, SpoIIGA(BS) and SpoIIGA(BL) differ at a few positions, and SpoIIGA(BC) and SpoIIGA(BH) differ greatly in three regions predicted to interact with substrate; the flap, loop and C-terminal regions corresponding to residues 242–257, 281–292 and 302–309, respectively, of SpoIIGA(BS) (Supplementary Fig. S3). These differences might account for the observed differences in substrate specificity.

Our results shed new light on residues in Pro- σ^E and in SpoIIGA that are important for cleavage, and demonstrate that SpoIIGA orthologs exhibit different

ranges of substrate specificity. Our results show for the first time that residues in the N-terminal one-third of the pro-sequence of Pro- σ^E contribute to substrate specificity. The experimental determination of the 3D structure of a SpoIIGA-Pro- σ^E complex and further mutational analysis will be required to clarify the mechanism of substrate recognition by SpoIIGA and its orthologs. As shown in Supplementary Fig. S4A, the cleavage of Pro- σ^E (BC)-H6 by SpoIIGA(BC)-GFP-F2 depended on co-expression of *B. subtilis* SpoIIR-F2, indicating that the *B. subtilis* signalling protein stimulates SpoIIGA(BC)-GFP-F2, presumably by direct interaction, as *B. subtilis* SpoIIR-F2 was shown to interact directly with *B. subtilis* SpoIIGA-GFP-H6 when the proteins were co-expressed in *E. coli* (15). Our finding that *B. subtilis* SpoIIR stimulates SpoIIGA (BC), and likely SpoIIGA(BL) and SpoIIGA(BH), may not be surprising since an alignment of SpoIIR orthologs (Supplementary Fig. S5) shows higher similarity and identity than for SpoIIGA orthologs (Supplementary Fig. S3). In particular, SpoIIR orthologs are highly similar in a region near their C-termini, and all contain the identical sequence NWWCV (Supplementary Fig. S5), which might be important for interaction with SpoIIGA.

Supplementary Data

Supplementary Data are available at *JB* online.

Acknowledgements

We are grateful for the gift of monoclonal antibody against σ^E from Dr William G. Haldenwang (University of Texas Health Science Center, San Antonio, TX, USA).

Funding

The Science Research Promotion Fund; Grant-in-Aid for Scientific Research (C) (20580089); Grant-in-Aid for Young Scientists (B) (19780066) from the Japanese Society for the Promotion of Science; The Michigan Agricultural Experiment Station; National Science Foundation (0447799 to M.F.); The Alfred P. Sloan Foundation (to M.F.); and National Institutes of Health (GM43585 to L.K.).

Conflict of interest

None declared.

References

1. Kroos, L. (2007) The *Bacillus* and *Myxococcus* developmental networks and their transcriptional regulators. *Annu. Rev. Genet.* **41**, 13–39
2. Steil, L., Serrano, M., Henriques, A.O., and Völker, U. (2005) Genome-wide analysis of temporally regulated and compartment-specific gene expression in sporulating cells of *Bacillus subtilis*. *Microbiology* **151**, 399–420
3. Wang, S.T., Setlow, B., Conlon, E.M., Lyon, J.L., Imamura, D., Sato, T., Setlow, P., Losick, R., and Eichenberger, P. (2006) The forespore line of gene expression in *Bacillus subtilis*. *J. Mol. Biol.* **358**, 16–37
4. Hofmeister, A.E., Londoño-Vallejo, A., Harry, E., Stragier, P., and Losick, R. (1995) Extracellular signal protein triggering the proteolytic activation of a

- developmental transcription factor in *B. subtilis*. *Cell* **83**, 219–226
5. Karow, M.L., Glaser, P., and Piggot, P.J. (1995) Identification of a gene, *spoIIR*, that links the activation of σ^E to the transcriptional activity of σ^F during sporulation in *Bacillus subtilis*. *Proc. Natl. Acad. Sci. USA* **92**, 2012–2016
 6. Londoño-Vallejo, J.A. and Stragier, P. (1995) Cell-cell signaling pathway activating a developmental transcription factor in *Bacillus subtilis*. *Genes Dev.* **9**, 503–508
 7. LaBell, T.L., Trempy, J.E., and Haldenwang, W.G. (1987) Sporulation-specific σ factor σ^{29} of *Bacillus subtilis* is synthesized from a precursor protein, P^{31} . *Proc. Natl. Acad. Sci. USA* **84**, 1784–1788
 8. Miyao, A., Theeragool, G., Takeuchi, M., and Kobayashi, Y. (1993) *Bacillus subtilis* *spoVE* gene is transcribed by σ^E -associated RNA polymerase. *J. Bacteriol.* **175**, 4081–4086
 9. Kenney, T.J. and Moran, C.P. Jr. (1987) Organization and regulation of an operon that encodes a sporulation-essential sigma factor in *Bacillus subtilis*. *J. Bacteriol.* **169**, 3329–3339
 10. Eichenberger, P., Fujita, M., Jensen, S.T., Conlon, E.M., Rudner, D.Z., Wang, S.T., Ferguson, C., Haga, K., Sato, T., Liu, J.S., and Losick, R. (2004) The program of gene transcription for a single differentiating cell type during sporulation in *Bacillus subtilis*. *PLoS Biol.* **2**, e328
 11. Eichenberger, P., Jensen, S.T., Conlon, E.M., van Ooij, C., Silvaggi, J., González-Pastor, J.E., Fujita, M., Ben-Yehuda, S., Stragier, P., Liu, J.S., and Losick, R. (2003) The σ^E regulon and the identification of additional sporulation genes in *Bacillus subtilis*. *J. Mol. Biol.* **327**, 945–972
 12. Feucht, A., Evans, L., and Errington, J. (2003) Identification of sporulation genes by genome-wide analysis of the σ^E regulon of *Bacillus subtilis*. *Microbiology* **149**, 3023–3034
 13. Fujita, M. and Losick, R. (2002) An investigation into the compartmentalization of the sporulation transcription factor σ^E in *Bacillus subtilis*. *Mol. Microbiol.* **43**, 27–38
 14. Stragier, P., Bonamy, C., and Karmazyn-Campelli, C. (1988) Processing of a sporulation sigma factor in *Bacillus subtilis*: how morphological structure could control gene expression. *Cell* **52**, 697–704
 15. Imamura, D., Zhou, R., Feig, M., and Kroos, L. (2008) Evidence that the *Bacillus subtilis* SpoIIIGA protein is a novel type of signal-transducing aspartic protease. *J. Biol. Chem.* **283**, 15287–15299
 16. Ju, J., Luo, T., and Haldenwang, W.G. (1997) *Bacillus subtilis* Pro- σ^E fusion protein localizes to the forespore septum and fails to be processed when synthesized in the forespore. *J. Bacteriol.* **179**, 4888–4893
 17. Ju, J. and Haldenwang, W.G. (2003) Tethering of the *Bacillus subtilis* σ^E proprotein to the cell membrane is necessary for its processing but insufficient for its stabilization. *J. Bacteriol.* **185**, 5897–5900
 18. Londoño-Vallejo, J.A. (1997) Mutational analysis of the early forespore/mother-cell signalling pathway in *Bacillus subtilis*. *Microbiology* **143**, 2753–2761
 19. Peters, H.K. 3rd, Carlson, H.C., and Haldenwang, W.G. (1992) Mutational analysis of the precursor-specific region of *Bacillus subtilis* σ^E . *J. Bacteriol.* **174**, 4629–4637
 20. Carlson, H.C., Lu, S., Kroos, L., and Haldenwang, W.G. (1996) Exchange of precursor-specific elements between Pro- σ^E and Pro- σ^K of *Bacillus subtilis*. *J. Bacteriol.* **178**, 546–549
 21. Scott, W.R. and Schiffer, C.A. (2000) Curling of flap tips in HIV-1 protease as a mechanism for substrate entry and tolerance of drug resistance. *Structure* **8**, 1259–1265
 22. Peters, H.K. III and Haldenwang, W.G. (1994) Isolation of a *Bacillus subtilis* *spoIIIGA* allele that suppresses processing-negative mutations in the Pro- σ^E gene (*sigE*). *J. Bacteriol.* **176**, 7763–7766
 23. Larkin, M.A., Blackshields, G., Brown, N.P., Chenna, R., McGettigan, P.A., McWilliam, H., Valentin, F., Wallace, I.M., Wilm, A., Lopez, R., Thompson, J.D., Gibson, T.J., and Higgins, D.G. (2007) Clustal W and Clustal X version 2.0. *Bioinformatics* **23**, 2947–2948