Published in final edited form as:

Cardiol Young. 2011 February ; 21(1): 94–96. doi:10.1017/S1047951110001368.

Paroxysmal complete atrioventricular block in a patient with pulmonary atresia and intact ventricular septum

Akash R. Patel, David Goldberg, and Maully Shah

Division of Cardiology, Department of Pediatrics, The Children's Hospital of Philadelphia, University of Pennsylvania School of Medicine, Pennsylvania, United States of America

Abstract

Sudden death in pulmonary atresia and intact ventricular septum with right ventricular dependent coronary circulation is a well-established complication, and is thought to be caused by myocardial ischaemia. We report a case of paroxysmal complete atrioventricular block that raises the possibility of an additional mechanism of sudden death.

Keywords

External cardiac ambulatory telemetry; congenital cardiac disease; Fontan procedure

Case report

A 4-year-old boy with pulmonary atresia and intact ventricular septum presented with syncope over a year after he underwent a fenestrated extracardiac Fontan procedure. Previously, several near-syncope episodes occurred with no clear aetiology. Prior Holter monitoring revealed no sinus pauses or atrioventricular block. External cardiac ambulatory telemetry was placed for further evaluation. Two weeks later, the patient experienced a syncopal episode while walking and the telemetry recording of this event revealed paroxysmal complete atrioventricular block. There was an abrupt transition from sinus rhythm to complete atrioventricular block and an asystolic pause of 9.6 seconds followed by junctional escape beats and eventual return to sinus rhythm (Fig 1).

The patient was admitted to the cardiac intensive care unit for observation and evaluation. A 15-lead electrocardiogram showed normal sinus rhythm with T-wave inversion and Q-waves in the inferior leads, which were unchanged from previous electrocardiograms. The echocardiogram was unchanged from baseline and demonstrated normal ventricular function with trivial mitral insufficiency and no aortic insufficiency. Based on this event, an epicardial single-chamber ventricular pacemaker was placed and the patient was discharged home without further episodes.

The patient's past medical history is noteworthy for development of right ventricular dependent coronary circulation. A cardiac catheterisation at 2 days of age revealed a diminutive right coronary system. The right coronary artery filled not only from the aorta but also from several coronary cameral fistulae. No areas of discrete stenosis were

© Cambridge University Press, 2010

Correspondence to: Dr A. R. Patel, MD, Division of Cardiology, Department of Pediatrics, The Children's Hospital of Philadelphia, University of Pennsylvania School of Medicine, 34th Street and Civic Center Blvd, Philadelphia, Pennsylvania 19104, United States of America. Tel: 215 590 2230; Fax: 215 590 5825; patela@email.chop.edu.

The authors have not disclosed any potential conflicts of interest.

identified. Repeat angiography prior to the Fontan procedure showed no right coronary artery originating from the right sinus of valsalva, suggesting acquired coronary ostial occlusion.

Discussion

The spectrum of disease in pulmonary atresia and intact ventricular septum is quite variable depending on the adequacy of right-sided structures, degree of tricuspid regurgitation, and extent of coronary abnormalities that can lead to variations in management and surgical approach.^{1,2} Currently, pulmonary atresia and intact ventricular septum have a reported actuarial survival of approximately 81% at 5 years, which is comparable to other cardiac conditions undergoing staged univentricular palliation.^{3,4} Previous studies, however, suggest increased early mortality when associated with significant coronary artery abnormalities.^{1,5}

There are 31–68% of patients with pulmonary atresia with intact ventricular septum who have associated coronary artery abnormalities including coronary-cameral fistulae, coronary stenoses, and coronary atresia.^{3,5} In the setting of coronary obstruction, the coronary perfusion becomes dependent upon the right ventricle. This so-called right ventricular dependent coronary circulation can lead to acute myocardial ischaemia, if the right ventricle is decompressed as part of surgical palliation or occurs over time due to progression of coronary arterial stenoses.⁵

To date, there has been no evidence that the underlying conduction system, coronary pattern, and coronary distribution are abnormal in this disease state.^{6–8} This may be significant, however, because the atrioventricular nodal artery, the coronary branch that supplies the atrioventricular node, arises from the right coronary artery 90% of the time.⁹ In this setting, the blood supply to the atrioventricular node could be significantly diminished. An analogous situation occurs in adults following myocardial infarction in the right coronary distribution. An increased incidence of atrioventricular block has been described in these patients.¹⁰

We postulate that patients with pulmonary atresia and intact ventricular septum with right ventricular dependent coronary circulation may be at risk for paroxysmal atrioventricular block due to diminished coronary perfusion to the atrioventricular node. This may be an underappreciated mechanism of sudden death in this population.

Acknowledgments

This study received no financial support.

References

1. Hanley FL, Sade RM, Blackstone EH, et al. Outcomes in neonates with pulmonary atresia with intact ventricular septum. A multi-institutional study. *J Thorac Cardiovasc Surg.* 1993; 105:406–407. [PubMed: 8445920]
2. Asburn DA, Blackstone EH, Wells WJ, et al. Determinants of mortality and type of repair in neonates with pulmonary atresia and intact ventricular septum. *J Thorac Cardiovasc Surg.* 2004; 127:1000–1008. [PubMed: 15052196]
3. Guleserian KJ, Armsby LB, Thiagarajan RR, del Nido PJ, Mayer JE. Natural history of pulmonary atresia with intact ventricular septum and right-ventricle-dependent coronary circulation managed by the single-ventricle approach. *Ann Thorac Surg.* 2006; 81:2250–2258. [PubMed: 16731162]
4. Rychik J, Levy H, Gaynor JW, Decamp WM, Spray TL. Outcome after operations for pulmonary atresia with intact ventricular septum. *J Thorac Cardiovasc Surg.* 1998; 11:924–931. [PubMed: 9832682]

5. Freedom RM, Anderson RH, Perrin D. The significance of ventriculo-coronary arterial connections in the setting of pulmonary atresia with an intact ventricular septum. *Cardiol Young*. 2005; 15:447–468. [PubMed: 16164782]
6. Ansari A, Goltz D, McCarthy KP, Cook A, Yen Ho S. The conduction system in hearts with pulmonary atresia and intact ventricular septum. *Ann Thorac Surg*. 2003; 75:1502–1505. [PubMed: 12735569]
7. Sandor GG, Cook AC, Sharland GK, Ho SY, Potts JE, Anderson RH. Coronary arterial abnormalities in pulmonary atresia with intact ventricular septum diagnosed during fetal life. *Cardiol Young*. 2002; 12:436–444. [PubMed: 15773446]
8. Daubney PEF, Delany DJ, Anderson RH, et al. Pulmonary atresia with intact ventricular septum: range of morphology in a population-based study. *J Am Coll Cardiol*. 2002; 39:1670–1679. [PubMed: 12020496]
9. Pejkovic B, Krajnc I, Anderbuber F, Kosutic D. Anatomical aspects of the arterial blood supply to the sinoatrial and atrioventricular nodes of the human heart. *J Int Med Res*. 2008; 36:691–698. [PubMed: 18652764]
10. Mehta SR, Eikelboom JW, Natarajan MK, et al. Impact of right ventricular involvement on mortality and morbidity in patients with inferior myocardial infarction. *J Am Coll Cardiol*. 2001; 37:37–43. [PubMed: 11153770]

Figure 1.
The rhythm was sinus with abrupt transition to complete atrioventricular block and an asystolic pause of 9.6 seconds followed by junctional escape beats and eventual return to sinus rhythm.