Published in final edited form as:

Crit Rev Biochem Mol Biol. 2011 August ; 46(4): 284–294. doi:10.3109/10409238.2011.577052.

Revealing Histone Variant Induced Changes Via Quantitative Proteomics

Anna M. Arnaudo^{1,#}, Rosalynn C. Molden^{2,#}, and Benjamin A. Garcia^{1,2,3,*}

¹Department of Molecular Biology, Princeton University, Princeton NJ 08544

²Department of Chemistry, Princeton University, Princeton NJ 08544

³Quantitative and Computational Biology Program, Princeton University, Princeton, NJ 08544

Abstract

Histone variants are isoforms of linker and core histone proteins that differ in their amino acid sequences. These variants have distinct genomic locations and post-translational modifications, thus increasing the complexity of the chromatin architecture. Biological studies of histone variants indicate that they play a role in many processes including transcription, DNA damage response, and the cell cycle. The small differences in amino acid sequence and the diverse post-translational modification states that exist between histone variants make traditional analysis using immunoassay methods challenging. In recent years, a number of mass spectrometric techniques have been developed to identify and quantify histones at the whole protein or peptide levels. In this review we discuss the biology of histone variants and methods to characterize them using mass spectrometry-based proteomics.

Introduction

The nucleosome, the basic repeating unit of chromatin, consists of DNA wrapped around an octamer of core histone proteins, two copies each of H2A, H2B, H3 and H4. Linker histone H1 may also be present, and contribute to chromatin structure. The presence of covalent histone post-translational modifications (PTMs) and the incorporation of histone sequence variants alter the composition of the nucleosome (Figure 1). Most PTMs occur on the N-terminal tails of histone proteins and include methylation (mono-, di-, tri-), phosphorylation, acetylation, and ubiquitination. The observation that certain histone modifications are associated with active genes and others with repressed genes lead to the hypothesis that information contained in histone PTMs forms a “histone code,” read by numerous effector proteins to influence chromatin structure and downstream events such as transcription (Strahl and Allis, 2000). Histone variants affect a variety of chromatin-related processes, are localized to different areas of the genome, and have unique modification patterns; they are proposed to form an extra layer of the histone code (Hake and Allis, 2006).

Histone PTMs and variants impact a variety of biological processes, including transcription, DNA damage response, cell cycle, viral infection, stem cell pluripotency, and fertility. Chromosome condensation and proper segregation during mitosis are correlated with histone H3 phosphorylation at serine 10 and serine 28, implicating these PTMs in cell cycle

*Address correspondence to: Benjamin A. Garcia 415 Schultz Laboratory, Dept. of Molecular Biology, Princeton University, Princeton, NJ 08544. Telephone: 609-258-8854; Fax: 609-258-8854; bagarcia@princeton.edu.

[#]Both authors contributed equally

Declaration of interest

The authors declare no conflict of interest.

regulation (Garcia et al., 2005). Histone deacetylase inhibitors (HDACi), drugs that increase global histone acetylation by blocking deacetylase activity, have been used to induce pluripotent stem cells (Huangfu et al., 2008b, Huangfu et al., 2008a) and to reactivate latent human immunodeficiency virus (HIV) (Van Lint et al., 1996), thus providing evidence for histone acetylation's role in these processes. Additionally, histone H2A variant H2A.X is required for male fertility in mice; in its absence, spermatogenesis halts at the pachytene phase (meiosis I) resulting in loss of mature sperm production and infertility (Celeste et al., 2002). These examples highlight the biological importance of select histone variants and PTMs. Many of the other variants and PTMs remain to be characterized.

Histone variants and their PTMs often need to be quantified across different conditions in order to determine their functions in the cell. The reliable identification and quantification of histone PTMs is challenging because histones may be extensively modified and similar in structure and molecular weight. Histone variant analysis is equally difficult because variants can differ in sequence by as little as one amino acid. The two main methods currently used to study histone variants and their PTMs are immunoblotting and mass spectrometry. Immunoblotting is very sensitive but it is not very quantitative and it is a laborious task that provides information only about a single modification or a subset of modifications in a given sample. In addition, many histone modifications and variants are similar in structure and sequence, making the specificity and cross-reactivity of antibodies a problem (Fuchs et al., 2011, Egelhofer et al., 2011). Epitope occlusion is a concern because modifications are also often closely spaced, e.g. Histone H3 Lysine 9 is acetylated and Serine 10 is phosphorylated during mitosis (Hirota et al., 2005). In contrast, liquid-chromatography mass spectrometry (LC-MS) can be used to identify large numbers of proteins, their modifications and the locations of their modifications in a just couple of hours. In addition, there are many mass spectrometric techniques for the quantification of modified proteins. This review will focus on the progress that has been made in applying quantitative LC-MS techniques to study histone variants and their modifications.

Histone Overview

Histone variants differ from canonical histones in their genomic organization and gene expression. Canonical histone genes are intronless and occur in clusters, facilitating joint gene regulation (Albig and Doenecke, 1997). Their mRNAs lack a polyA tail, instead having a dyad sequence that forms a 3' stem loop structure. Their expression is replication-dependent, which means that their expression is linked to the S-phase of the cell cycle. Expression increases as much as 35-fold in S-phase to allow for sufficient histone protein incorporation during DNA synthesis (Harris et al., 1991). The number of canonical genes that code for each histone family member can exceed ten genes (Marzluff et al., 2002). These canonical genes can produce different protein sequences, drastically increasing variability of the histone population. The only exception is histone H4, whose 12 genes all produce the same sequence. Canonical histone proteins (H3, H4, H2A, and H2B) are small, ranging in size from 11–15 kDa, and consist of a lysine rich N-terminal tail, a globular histone fold domain, and a C-terminal tail. Different combinatorial PTMs to these core proteins add to the complexity.

In contrast to canonical histone genes, histone variants are replication-independent with expression occurring throughout the cell cycle. Genes encoding histone variants typically lay outside of clusters, can contain introns, and occur in low copy number. Like standard genes, mRNA produced from variant genes possess a polyA tail (Doenecke et al., 1997). Histone variants have protein sequences, modifications, structures and biological functions distinct from their respective canonical family members.

Mass Spectrometric Analysis of Histones

Most commonly, acid extraction is used to isolate histones from mammalian cell nuclei. This method gives reasonably pure histones for analysis (>90% purity), thus decreasing the sample complexity and diminishing the need for complicated separation strategies seen in whole cell proteomics (Shechter et al., 2007). Several alternatives also exist for isolating histones under non-denaturing conditions, as histones can be extracted from nuclei after micrococcal nuclease digestion using high salt concentrations or by hydroxyapatite chromatography (Benson et al., 2006, von Holt et al., 1989). Different protocols for isolating nuclei and histones are required for plants and some fungi (including *Saccharomyces cerevisiae*) since they have tough to lyse cell walls, and typically the histone yield and purity are observed to be somewhat lower for these eukaryotic systems. Nevertheless, the histone isolation procedures are essentially the same as for mammalian cells, and can be very easily implemented and executed with very little issues (Ide and Saunders, 1981, Edmondson et al., 1996, Waterborg et al., 1987). After extraction there are different processing steps depending on the type of mass spectrometric (MS) analysis and the method of quantification. There are three main MS analysis techniques used to analyze histones: Bottom Up, Middle Down and Top Down proteomics. These methods are compared in Figure 2, with a specific emphasis on differences in their utility during histone variant analysis.

Bottom Up Proteomics

Bottom Up proteomics is probably the most widely used form of MS analysis for histones and PTMs. For this method, histones are first subjected to proteolytic digestion, and then the resulting peptides are de-salted and loaded onto a C8 or C18 column for reverse phase LC-MS analysis. As the peptides elute, they are subjected to MS analysis and fragmentation, known as MS/MS analysis. The most common type of fragmentation for Bottom Up analysis is collision-activated dissociation (CAD), which breaks the peptide at the amide bond to create a series of b- and y-type ions. The difference in mass between ions in the series can be used to determine the peptide sequence and the positions of PTMs.

Trypsin, a protease that cleaves after arginine and lysine, is the enzyme most commonly used for digestion during Bottom Up MS analysis. However, histone proteins are arginine and lysine-rich, especially in their N-terminal tails where modifications are prevalent, so the standard tryptic digest creates ions that are too small to be retained on an HPLC column and thus cannot be detected using LC-MS analysis. The protease ArgC, which only cleaves after arginines, can be used in the place of trypsin to create longer peptides for LC-MS analysis but ArgC is not as robust as trypsin. Another alternative to a pure tryptic digest is to chemically derivatize lysines using propionic anhydride or acetic anhydride, blocking their cleavage by trypsin (Garcia et al., 2007a, Smith et al., 2002, Bonaldi et al., 2004). The derivatization method is efficient, in addition, propionic or acetic anhydride derivatization decreases the charge on peptides, increasing their retention on reverse phase columns and improving chromatographic resolution.

Bottom Up proteomics has limitations when it comes to analyzing histone variants and combinations of histone modifications. During ArgC or tryptic digests, information about the modifications that occur in combination is lost because of the small size of the resultant peptides (5–20 amino acids in length). In addition, Bottom Up proteomics cannot be used to distinguish between many of the histone variants and their modified states because of small differences in their sequences. For example, there is no unique peptide that distinguishes all of the H2B variants from each other so they cannot be quantified using Bottom Up MS (see Figure 2). Offline separation techniques can be used to purify histone variants prior to

Bottom Up analysis. For example, histones H3.1, H3.2 and H3.3 have many tryptic peptides in common and can't be distinguished from a mixture of digested histones, but they can be separated prior to digestion and LC-MS analysis using reversed phase HPLC (RP-HPLC). Alternatively, whole protein mass spectrometry (Top Down MS) or mass spectrometry of larger histone peptides (Middle Down MS) can be used to gain more information about histone variants and combinatorial modifications.

Top Down proteomics

Top Down mass spectrometry is the only method that can distinguish between all of the histone isoforms and their PTMs. Pre-fractionated whole histones are identified using Top Down MS by their intact mass and fragmentation spectra (Kelleher, 2004). Using this method even isobaric histones, such as H2B1H and H2B1K, can be identified based on their fragment ions. Analysis of H2B variants from bulk H2B is presented in Figure 2. Histones are very basic proteins and are therefore highly charged and generate multiply charged fragment ions. As a consequence, non-ergodic fragmentation methods such as electron transfer dissociation (ETD) or electron coupled dissociation (ECD) and high-resolution instruments such as the LTQ-Orbitrap or FT-ICR (Syka et al., 2004, Zubarev et al., 1998) are required for Top Down histone analysis.

Despite the advantages of Top Down MS, it is not utilized as often as Bottom Up MS for histone analysis because it is not as sensitive: many of the low-level histone variants and modifications that can be identified and quantified using Bottom Up methods are not readily detected in Top Down spectra. Another disadvantage of Top Down analysis is that it does not lend itself well to high-throughput analysis because it is not easily coupled to online separations. Some advances have been made in combining online separations with Top Down MS for the analysis of histone mixtures (Tian et al., 2010, Eliuk et al., 2010), but further improvements need to be made before Top Down methods are used to routinely identify and quantify histone variants and their PTMs.

Middle Down Proteomics

Middle Down proteomics combines the advantages of Top Down and Bottom Up methods. Middle Down MS cannot distinguish between as many histone variants as Top Down mass spectrometry; however it is much more sensitive and can be used with online chromatography (see Figure 2 for a comparison). Large histone peptides (greater than 3000 Da) are generated using Glu-C or Asp-N digests, separated using online or offline HPLC, and then fragmented using ETD or ECD for identification. Middle Down proteomics is used to profile coexisting histone modifications and can distinguish between more histone variants than Bottom Up MS. So far Middle Down MS has been applied to H4 as well as H3, H2A and H2B variants and their modifications (Garcia et al., 2007b, Young et al., 2009, Bonenfant et al., 2006).

Quantification Strategies

Multiple proteomic approaches have been developed to allow for protein quantification ranging in scale from the whole cell proteome to single protein analyses. These approaches can be broken down into four general categories - "label-free" quantification, chemical labeling, metabolic labeling and quantification by standard peptide (Bantscheff et al., 2007). Variations on all of these techniques have been utilized in studying chromatin histone post-translational modifications, histone variants, and histone turnover as shown in Figure 3, and discussed in more detail below.

Label-free Quantification

Label-free quantification methods establish relative protein quantities without incorporation of any labels or use of standard peptides, thus abrogating the cost and sample preparation required for labeling or peptide based strategies and allowing quantification across multiple samples or experimental conditions. A common label-free technique is spectral counting, which bases quantification on the number of MS/MS spectra acquired for each protein (Liu et al., 2004). In spectral counting the total number of MS/MS spectra corresponding to each peptide is compared across experimental conditions. The quality and reproducibility of the data can be affected by many factors, such as peptide ionization efficiency, settings for data dependent acquisition, and chromatographic separation (Washburn et al., 2003, Liu et al., 2004, Ott et al., 2007). The large scale nature of some spectral counting experiments utilize multi-dimensional separation strategies (e.g. strong cation exchange coupled to reverse phase) to better resolve peptide mixtures and allow for analysis of thousands of proteins-including histones (Washburn et al., 2003). Spectral counting of human HEK 293 cells, for example, revealed that treatment with cytokine tumor necrosis factor alpha (TNF-alpha) causes an increase in expression of H2A variants H2A.Z and macroH2A. This increase in H2A variants coupled with a decrease in DNA replication machinery implicates TNF-alpha in a cell cycle arrest event as part of the innate immune response (Ott et al., 2007). Such studies indicate that spectral counting may be a way to assess changes in histone variants across whole cells or tissue samples.

An alternative label-free strategy to spectral counting utilizes peak area from an extracted ion chromatogram to estimate peptide abundance. These studies typically rely on the Bottom Up strategies discussed above. The relative abundance of a covalently modified peptide is established by comparing the peak area of that modified form to the total peak area for all forms of the peptide. In *Drosophila*, it was found by Bottom Up LC-MS quantification of offline RP-HPLC purified H3 variants that histone H3.3 is enriched for acetylation and K4 methylation, suggesting a role for H3.3 in gene activation (McKittrick et al., 2004). A more stream-lined analysis of histone PTMs can be done by combining propionylation with trypsin digestion of unfractionated histones (Plazas-Mayorca et al., 2009).

Top Down mass spectrometry is not commonly used in conjunction with online LC analysis so an alternative label-free method has been developed for whole protein analysis. The relative amounts of different modified histone forms are measured by comparing the intensities of abundant fragment ions. The Kelleher lab found that the relative fragment ion intensity of modified histone H4 peptides was the same as the relative concentration of the peptides in solution, within 5% error margins (Pesavento et al., 2006). This technique has been applied to measure the relative abundances of PTMs across H3 variants in rat brains. A slight enrichment in the activating mark K4 methylation in H3.3 and a slight enrichment of the repressive mark K27 trimethylation in H3.2 were found (Garcia et al., 2007c).

Chemical Derivatization Based Quantification

The post-harvest derivatization of histones for Bottom Up using anhydrides or methyl and ethyl esters facilitates relative quantification between experimental conditions. An initial round of anhydride or ester derivatization blocks monomethylated and unmodified lysines then, following tryptic digestion, a second round of derivatization with esters or anhydrides block the C-terminus or N-terminus of the peptides respectively. Heavy isotope labeled esters or anhydrides, such as D10-propionic anhydride, can be used in the second round of derivatization to add one label per peptide. Mixing equal amounts of labeled and unlabeled propionylated histone samples from two different conditions facilitates relative quantification. A disadvantage of chemical derivatization is that samples are mixed at the

peptide level, and results can be skewed if sample loss occurs early on in sample preparation (see Figure 3). Ester derivatization requires peptides to be dried, increasing sample loss and making it a less desirable method compared to propionic anhydride derivatization.

A number of studies have utilized differential chemical derivatization to quantify changes in PTMs or variants (DiMaggio and Garcia, 2010). Most recently, propionic anhydride labeling was used to observe the effect of G9a/Glp1 methyltransferase knockdown on global histone modifications. G9a/Glp1 methyltransferases are responsible for dimethylation of H3K9 and their knockdown resulted in a decrease of this modification. Additionally, H3K14 acetylation, H3K79 and H3K36 methylation levels increased in a H3 variant specific manner-suggesting cross-talk between these modifications (Plazas-Mayorca et al.).

Metabolic Stable Isotope Labeling

Metabolic labeling is a relative quantification technique that relies on the replacement of a standard light isotope with a “heavy” isotope (e.g. ^{15}N , ^{13}C , D, or ^{18}O) in cellular proteins. The heavy isotope is incorporated into the protein using a heavy-labeled metabolite that is added to media during cell culture. In a typical metabolic labeling experiment comparing cells grown under two biological conditions, cells under one condition receive heavy-isotope labeled media while cells grown under the other condition receive normal media. Equal numbers of cells from each condition are mixed and processed for mass spectrometric analysis. Proteins from different conditions are distinguished by a known mass shift in their spectra. Relative protein amounts can be quantified using the area under the curve in the extracted ion chromatograms for heavy and light-labeled peptides or by relative fragment ion intensities. The main advantage of metabolic labeling is that the labeled cells and unlabeled cells are mixed as a first step, so proteins have identical losses for all of the proceeding processing steps (Figure 3). The main disadvantage of metabolic labeling is that it has to be done in tissue culture. Two types of metabolic labeling are used for histone quantification: stable isotope labeling of amino acids in cell culture (SILAC), and labeling of metabolites that become incorporated into the histone modifications.

SILAC was developed by the Mann group in 2002 and since has been used in numerous histone studies (Ong et al., 2002). It was used recently to investigate partitioning of H3 variants during DNA replication (Xu et al., 2010). In this study, FLAG epitope-tagged H3.1 and H3.3 variants were grown in HeLa cells. The cells were arrested and switched into media containing $^{13}\text{C}_6^{15}\text{N}_2$ -lysine so that all new histones after cell cycle arrest would become labeled. After immunopurification of the tagged variants, and Bottom Up analysis of the immunopurified histones, they discovered that H3.1–H4 tetramers do not split into dimers during replication dependent nucleosome assembly, but H3.3–H4 dimers do split during replication dependent assembly. They present this as support for the hypothesis that histone modifications in newly formed nucleosomes are copied from histones in preexisting nucleosomes.

SILAC can also be used in combination with Top Down mass spectrometry. It was used to study changes in levels of different canonical HeLa cell H2A isoforms across the cell cycle by following the level of incorporation of ^{13}C , ^{15}N labeled valine and arginine into the H2A isoforms after cell synchronization (Boyne et al., 2006). This study confirmed that the majority of canonical H2A isoforms are produced during S-phase, and showed that levels of different H2A isoforms did not vary much across the cell cycle.

Metabolites can also be used to label specific modifications. Jenuwein and coworkers used $^{13}\text{CD}_3$ -methionine to measure levels of H3K9 methylation after inducing the expression Jmjd2b H3K9 de-methylase in murine cells (Fodor et al., 2006). Later, the Garcia Lab used heavy-methionine labeling in addition to $^{13}\text{C}_6^{15}\text{N}_2$ -lysine labeling to study

methylation turnover in site-specific manner and to compare methylation turnover to histone turnover (Zee et al., 2010a). They found that activating methylation marks such as H3K4 generally turned over faster than repressive marks such as H3K9. They also found that methyl marks had faster turnover than the histones themselves, demonstrating that histones are dynamically modified. In a later paper they demonstrated that PTM turnover between variants was fairly constant and that all histone variants, with the exception of histone H2A variants, turned over at approximately the same rate (Zee et al., 2010b).

Absolute quantification by use of peptide standards

Both relative and absolute quantification of histones and their PTMs are possible through the use of synthetic, isotopically labeled standard peptides. Peptides are synthesized to have the same sequence and modifications as the digested form of the histones of interest. A known concentration of the standard peptide is spiked into samples, then the intensity of the analyte to the standard is compared. Alternatively the absolute quantity of the analyte can be determined by making a calibration curve of peptide standard concentration versus ion intensity. This method is not entirely accurate because analyte losses occur in processing steps before the standard peptide is added; however, it can be used in combination with reaction monitoring MS techniques to monitor very low levels of analyte. Standard peptide quantification is not commonly used for analysis of histones because standard peptides are expensive and only one PTM or variant can be quantified at one time.

There are a few examples of standard peptides being used to quantify histone PTMs. Darwanto and coworkers used peptide standards to quantify relative levels of H3K79 methylation and H2B ubiquitination in wild-type U937 lymphoma cells and in U937 cells expressing the methyltransferase hDot1 with a deleted methyl-binding domain. They found that a decrease in H3K79 methylation led to an increase in H2B ubiquitination suggesting cross-talk between the modifications (Darwanto et al., 2010). They used standard peptides for this study because ubiquitinated H2B is not very abundant and would not be easily detected using less sensitive techniques.

Function of Histone Variants

Histone H3 Family

Histone H3 has three major variants as well as a testis specific variant (H3t) and a variant that is localized to centromeres (CENP-A). The three main variants of H3 are surprisingly similar in sequence with H3.1 only differing from H3.2 by a change in Cysteine 96 to Serine and H3.3 differing from H3.1 by only 5 residues. Despite the small differences in sequence homology, the variants have large differences in their expression, localization in chromatin, and modification state. H3.3 is expressed in a replication independent manner, while the canonical variants (H3.1 and H3.2) are only expressed during S phase (Stein et al., 1984). In addition, different chaperones recognize and assemble H3.1 and H3.3 into nucleosomes in a replication dependent and independent manner respectively (Tagami et al., 2004). Compared to H3.1 and H3.2, H3.3 is usually localized to heterochromatin and enriched for histone modifications that are associated with gene activation (McKittrick et al., 2004, Ahmad and Henikoff, 2002). H3.1 and H3.2 do not show specific areas of localization in the genome and are generally considered to be identical; however one study suggests that H3.2 is modified with marks associated with gene repression while H3.1 shows both activating and repressive marks (Ahmad and Henikoff, 2002, Hake et al., 2006).

Quantitative studies of H3 variant modifications have been accomplished using Bottom Up, Top Down and Middle Down techniques (Hake et al., 2006, Thomas et al., 2006, Garcia et al., 2007c). Fractionation of H3 variants is necessary for complete analysis by mass

spectrometry. The H3 variants are easily resolved using reverse-phase HPLC and different modified forms of each variant can be further separated using cation-exchange chromatography or hydrophilic interaction chromatography (HILIC). The H3 variants need to be separated before digestion for Bottom Up analysis because they have nearly identical N-terminal tails. Top and Middle Down analysis can also be improved by separating H3 variants and their modifications because there are thousands of combinations of modifications associated with each H3 variant. The most successful separation method used in combination with Top and Middle Down mass spectrometry is weak cation-exchange HILIC, which successfully separates isobaric species based on the number and position of modifications. This technique was used to identify over 200 modified forms of H3.2 (Young et al., 2009).

Histone H2A Family

The H2A family of histones is one of the most sequence divergent families, with 20 unique sequences in humans. Canonical human histone H2A is encoded for by 16 genes within gene clusters and has 12 unique sequences. Sequence diversity comes mainly from divergent C-terminal tails, but the biological relevance of this diversity remains unknown. Phosphorylation of S1 and acetylation sites within the N-terminal have been reported for canonical H2A (Bonenfant et al., 2006, Boyne et al., 2006). Monoubiquitination of H2A at C-terminal residue K119 has been implicated in gene silencing (Wang et al., 2004). The H2A family variants include H2A.X, H2A.Bbd, H2A.Z and macroH2A; these variants differ in sequence and have diverse functions.

Variant H2A.X is most similar in sequence to canonical but has a divergent C-terminal tail. Unlike canonical genes the mRNA for H2A.X can have either a polyA tail or the stem loop dyad structure, indicating that it can undergo both replication dependent and independent transcription (Mannironi et al., 1989). H2A.X can be acetylated, ubiquitinated and phosphorylated. In response to DNA damage, H2A.X is phosphorylated at serine 139 and localizes to DNA double strand breaks, helping to recruit proteins to carry out DNA repair (Rogakou et al., 1998, Ikura et al., 2007). Acetylation and ubiquitination of H2A.X also function in this process with acetylation at lysine 5 serving as a prerequisite for ubiquitination and subsequent release of H2A.X from DNA damage sites (Ikura et al., 2007).

A second variant of the H2A family is H2A.Bbd, whose name is derived from the fact that this variant is excluded from the inactive X and therefore Barr body deficient. This along with the finding that it associates with acetylated histone H4 has implicated H2A.Bbd in transcriptional activation (Chadwick and Willard, 2001). Recent reports also suggest a role for H2A.Bbd in spermatogenesis (Ishibashi et al.). The H2A.Bbd protein sequence is short and arginine rich in comparison to canonical. In vitro work demonstrates that these sequence differences alter the nucleosome core particle. Nucleosomes containing H2A.Bbd contain 128bp of DNA as opposed to the traditional 146, indicating that H2A.Bbd incorporation impacts chromatin structure (Doyen et al., 2006).

Histone H2A.Z is highly conserved and is essential for development in higher eukaryotes (Draker and Cheung, 2009). Chromatin immunoprecipitation experiments show that H2A.Z localizes to the promoters of genes and in mammals, localization to the transcription start site correlates with gene activation (Barski et al., 2007). In yeast, it was found that H2A.Z is present in transcriptionally active areas near the telomeres and silent mating locus and prevents spreading of heterochromatin from the silent mating locus. These findings suggest that H2A.Z is important for maintaining active chromatin in regions near silent chromatin (Meneghini et al., 2003). Acetylation of H2A.Z is thought to be linked to its role in gene activation. These facts implicate H2A.Z in gene activation, however, there is evidence to

indicate H2A.Z can also act in gene repression (Draker and Cheung, 2009, Millar et al., 2006, Mehta et al., 2010).

MacroH2A is the largest H2A variant containing an approximately 30kDa macro domain at the C-terminus. Human macroH2A is encoded for by two genes macroH2A2 and macroH2A1, which has two splice variants- macroH2A1.1 and macroH2A1.2. The ability of macroH2A1.1 to bind metabolites of NAD⁺ gives this splice variant potential for unique functions. MacroH2A is generally enriched at transcriptionally silent areas both on autosomes and on the inactive X. However, recent ChIP-Chip experiments show that macroH2A1 can be found at a subset of active genes (Gamble et al.). Mass spectrometric analysis reveals that macroH2A can be ubiquitinated, phosphorylated and acetylated. Ubiquitination at K115 has been implicated in X-inactivation, while phosphorylation of S137 is enriched during mitosis (Chu et al., 2006, Bernstein et al., 2008).

Histone H2B Family

Human H2B has 17 isoforms (Figure 1), which are encoded by 25 genes. Most H2B sequences vary by just a few amino acids and are encoded by genes located in the main histone gene clusters. The two H2B proteins that differ significantly from other H2B sequences are the testis-specific H2B proteins H2B1A and H2BFWT (Zalensky et al., 2002, Churikov et al., 2004). H2B variants have not been studied in as much depth as other histone variants, possibly because of their sequence similarity, so their biological significance is not well understood. H2B variants are not as extensively modified as H3 or H4, and aside from a few lysine acetylation sites other modified forms are very low in abundance. H2B isoforms have been identified using Bottom Up, Middle Down and Top Down mass spectrometry (Bonenfant et al., 2006, Siuti et al., 2006). Neither technique gives a complete picture of all H2B isoforms because H2B isoforms have too much sequence homology to be identified from digest peptides, and Top Down mass spectrometry is not sensitive enough to detect lower-level variants. Mass spectrometry has also been used to detect H2B modifications, including some modifications specific to the testis-specific variants (Lu et al., 2009, Wyrick and Parra, 2009, Bonenfant et al., 2006).

Histone H1 Family

Histone H1 is commonly referred to as the linker histone. A single copy of this histone has been proposed to bind near the entry/exit site of DNA on the nucleosome stabilizing the 30nm fiber and impacting chromatin structure and DNA linker length (Bustin et al., 2005, Woodcock et al., 2006). The amount of histone H1 present for each core nucleosome has found to differ between cell types. In higher eukaryotes, histone H1 contains a globular domain, a short N-terminal tail and a longer lysine rich C-terminal tail. Sequence divergence between histone H1 isoforms occurs mainly in the N- and C-terminal regions of the protein and is high, with mammals having as many as 11 isoforms. These isoforms are generally categorized based on the timing of expression or tissue specificity (Izzo et al., 2008). The replication dependent isoforms include histone H1.1–H1.5 and H1.t, while replication independent variants include H1.X, H1.t2, H1LS1, H1oo and H1.0 (which is sometimes referred to as H1degrees). Histones H1.0– H1.5 are expressed in somatic cells, with H1.1 restricted to certain tissue types (Godde and Ura, 2008, Daujat et al., 2005). Histone H1.t, H1.t2 and H1LS1 are expressed in testes specific tissues and histone H1oo is oocyte specific (Godde and Ura, 2008). Histone H1.X has mainly been reported in tissue culture cells (Wisniewski et al., 2007). In addition to sequence diversity, mass spectrometric analysis and other studies have revealed a high number of post-translational modifications present on different linker histone variants including lysine methylation, phosphorylation, acetylation, ubiquitination, formylation, and ADP ribosylation (Wisniewski et al., 2007, Garcia et al., 2004, Poirier and Savard, 1980, Talasz et al., 2009, Talasz et al., 1996, Weiss et al.). The

high number of isoforms and their unique modification patterns cause some to suggest that in combination this diversity results in a linker code- akin to the histone code.

Multiple studies of these isoforms reveal some interesting insights into function and potential for functional diversity and redundancy (Godde and Ura, 2008, Izzo et al., 2008). Deletion of testes specific histone H1.t from mice resulted in deposition of other isoforms and normal spermatogenesis (Lin et al., 2000). Mice can also survive lacking copies of the somatic isoforms H1.0, H1.2, H1.3 or H1.4, but deletion of histone H1.2, H1.3 and H1.4 resulted in embryonic lethality (Fan et al., 2005). These results suggest that there is some functional redundancy, but there are cases where individual isoforms have been assigned unique functions. For example, histone H1.4 has been implicated in transcriptional repression (Kuzmichev et al., 2004). Reverse phase separation of histone H1 from bulk Hela histone followed by digestion, propionylation and tandem mass spectrometry revealed methylation at K26 and phosphorylation at S27 of histone H1.4 (Garcia et al., 2004). Methylation at K26 allows for binding of heterochromatin protein 1 (HP1), which functions in heterochromatin formation and gene silencing, while pS27 prevents binding (Daujat et al., 2005). Histone H1.4 is enriched in heterochromatic regions as compared to euchromatic regions- a unique localization pattern in comparison to some other H1 variants (Izzo et al., 2008). Overall these findings suggest that histone H1.4K26me facilitates HP1 binding, chromatin condensation and gene silencing, while phosphorylation of neighboring S27 can ameliorate these effects.

Phosphorylation is the most well characterized modification on histone H1 isoforms. In general, phosphorylation levels on H1 change over the course of the cell cycle with a maximum amount occurring during mitosis. Interestingly, the level of phosphorylation differs on specific variants in mice and rats (Talaszi et al., 1996). Hydrophobic interaction chromatography followed by Top Down analysis confirm these results in human cells showing that H1.2 and H1.4 from synchronized Hela cells have a single and double phosphorylation site respectively during interphase, while these numbers increase to four and six respectively during mitosis (Zheng et al.). Phosphorylation in the C-terminal domain is thought to ease binding of H1 to DNA, facilitating dynamic changes to chromatin organization (Lever et al., 2000). Distinct phosphorylation events on H1 isoforms have been proposed to have various functions including stabilizing condensed chromatin during mitosis (Talaszi et al., 2009) and promoting transcription and DNA replication (Talaszi et al., 2009, Zheng et al.).

Acknowledgments

B.A.G is supported by an the American Society for Mass Spectrometry Research award sponsored by the Waters Corp., an NJCCR SEED grant, an NSF Faculty CAREER award, an NSF grant (CBET-0941143), an Agilent Thought Leader award, and an NIH Innovator award (DP2OD007447) from the Office Of The Director, NIH.

References

- Ahmad K, Henikoff S. The Histone Variant H3.3 Marks Active Chromatin by Replication-Independent Nucleosome Assembly. *Molecular Cell*. 2002; 9:1191–1200. [PubMed: 12086617]
- Albig W, Doenecke D. The human histone gene cluster at the D6S105 locus. *Hum Genet*. 1997; 101:284–294. [PubMed: 9439656]
- Bantscheff M, Schirle M, Sweetman G, Rick J, Kuster B. Quantitative mass spectrometry in proteomics: a critical review. *Anal Bioanal Chem*. 2007; 389:1017–1031. [PubMed: 17668192]
- Barski A, Cuddapah S, Cui K, Roh TY, Schones DE, Wang Z, Wei G, Chepelev I, Zhao K. High-resolution profiling of histone methylations in the human genome. *Cell*. 2007; 129:823–837. [PubMed: 17512414]

- Benson LJ, Gu Y, Yakovleva T, Tong K, Barrows C, Strack CL, Cook RG, Mizzen CA, Annunziato AT. Modifications of H3 and H4 during chromatin replication, nucleosome assembly, and histone exchange. *J Biol Chem.* 2006; 281:9287–9296. [PubMed: 16464854]
- Bernstein E, Muratore-Schroeder TL, Diaz RL, Chow JC, Changolkar LN, Shabanowitz J, Heard E, Pehrson JR, Hunt DF, Allis CD. A phosphorylated subpopulation of the histone variant macroH2A1 is excluded from the inactive X chromosome and enriched during mitosis. *Proc Natl Acad Sci U S A.* 2008; 105:1533–1538. [PubMed: 18227505]
- Bonaldi T, Imhof A, Regula JT. A combination of different mass spectroscopic techniques for the analysis of dynamic changes of histone modifications. *Proteomics.* 2004; 4:1382–1396. [PubMed: 15188406]
- Bonenfant D, Coulot M, Towbin H, Schindler P, Van Oostrum J. Characterization of histone H2A and H2B variants and their post-translational modifications by mass spectrometry. *Mol Cell Proteomics.* 2006; 5:541–552. [PubMed: 16319397]
- Boyne MT 2nd, Pesavento JJ, Mizzen CA, Kelleher NL. Precise characterization of human histones in the H2A gene family by top down mass spectrometry. *J Proteome Res.* 2006; 5:248–253. [PubMed: 16457589]
- Bustin M, Catez F, Lim JH. The dynamics of histone H1 function in chromatin. *Mol Cell.* 2005; 17:617–620. [PubMed: 15749012]
- Celeste A, Petersen S, Romanienko PJ, Fernandez-Capetillo O, Chen HT, Sedelnikova OA, Reina-San-Martin B, Coppola V, Meffre E, Difilippantonio MJ, Redon C, Pilch DR, Oлару A, Eckhaus M, Camerini-Otero RD, Tessarollo L, Livak F, Manova K, Bonner WM, Nussenzweig MC, Nussenzweig A. Genomic instability in mice lacking histone H2AX. *Science.* 2002; 296:922–927. [PubMed: 11934988]
- Chadwick BP, Willard HF. A novel chromatin protein, distantly related to histone H2A, is largely excluded from the inactive X chromosome. *J Cell Biol.* 2001; 152:375–384. [PubMed: 11266453]
- Chu F, Nusinow DA, Chalkley RJ, Plath K, Panning B, Burlingame AL. Mapping post-translational modifications of the histone variant MacroH2A1 using tandem mass spectrometry. *Mol Cell Proteomics.* 2006; 5:194–203. [PubMed: 16210244]
- Churikov D, Siino J, Svetlova M, Zhang KL, Gineitis A, Bradbury EM, Zalensky A. Novel human testis-specific histone H2B encoded by the interrupted gene on the X chromosome. *Genomics.* 2004; 84:745–756. [PubMed: 15475252]
- Darwanto A, Curtis MP, Schrag M, Kirsch W, Liu P, Xu G, Neidigh JW, Zhang K. A modified "cross-talk" between histone H2B Lys-120 ubiquitination and H3 Lys-79 methylation. *J Biol Chem.* 2010; 285:21868–21876. [PubMed: 20442396]
- Daujat S, Zeissler U, Waldmann T, Happel N, Schneider R. HP1 binds specifically to Lys26-methylated histone H1.4, whereas simultaneous Ser27 phosphorylation blocks HP1 binding. *J Biol Chem.* 2005; 280:38090–38095. [PubMed: 16127177]
- DiMaggio PA, Garcia BA. Mass Spectrometry Based Proteomics for Interrogating the Histone Code. *Current Proteomics.* 2010; 7:177–187.
- Doenecke D, Albig W, Bode C, Drabent B, Franke K, Gavenis K, Witt O. Histones: genetic diversity and tissue-specific gene expression. *Histochem Cell Biol.* 1997; 107:1–10. [PubMed: 9049636]
- Doyen CM, Montel F, Gautier T, Menoni H, Claudet C, Delacour-Larose M, Angelov D, Hamiche A, Bednar J, Faivre-Moskalenko C, Bouvet P, Dimitrov S. Dissection of the unusual structural and functional properties of the variant H2A.Bbd nucleosome. *EMBO J.* 2006; 25:4234–4244. [PubMed: 16957777]
- Draker R, Cheung P. Transcriptional and epigenetic functions of histone variant H2A.Z. *Biochem Cell Biol.* 2009; 87:19–25. [PubMed: 19234520]
- Edmondson DG, Smith MM, Roth SY. Repression domain of the yeast global repressor Tup1 interacts directly with histones H3 and H4. *Genes Dev.* 1996; 10:1247–1259. [PubMed: 8675011]
- Egelhofer TA, Minoda A, Klugman S, Lee K, Kolasinska-Zwierz P, Alekseyenko AA, Cheung MS, Day DS, Gadel S, Gorchakov AA, Gu T, Kharchenko PV, Kuan S, Latorre I, Linder-Basso D, Luu Y, Ngo Q, Perry M, Rechtsteiner A, Riddle NC, Schwartz YB, Shanower GA, Vielle A, Ahringer J, Elgin SC, Kuroda MI, Pirrotta V, Ren B, Strome S, Park PJ, Karpen GH, Hawkins RD, Lieb JD.

- An assessment of histone-modification antibody quality. *Nat Struct Mol Biol.* 2011; 18:91–93. [PubMed: 21131980]
- Eliuk SM, Maltby D, Panning B, Burlingame AL. High Resolution Electron Transfer Dissociation Studies of Unfractionated Intact Histones from Murine Embryonic Stem Cells Using On-line Capillary LC Separation. *Molecular & Cellular Proteomics.* 2010; 9:824–837. [PubMed: 20133344]
- Fan Y, Nikitina T, Zhao J, Fleury TJ, Bhattacharyya R, Bouhassira EE, Stein A, Woodcock CL, Skoultchi AI. Histone H1 depletion in mammals alters global chromatin structure but causes specific changes in gene regulation. *Cell.* 2005; 123:1199–1212. [PubMed: 16377562]
- Fodor BD, Kubicek S, Yonezawa M, O'Sullivan RJ, Sengupta R, Perez-Burgos L, Opravil S, Mechtler K, Schotta G, Jenuwein T. Jmjd2b antagonizes H3K9 trimethylation at pericentric heterochromatin in mammalian cells. *Genes Dev.* 2006; 20:1557–1562. [PubMed: 16738407]
- Fuchs SM, Krajewski K, Baker RW, Miller VL, Strahl BD. Influence of combinatorial histone modifications on antibody and effector protein recognition. *Curr Biol.* 2011; 21:53–58. [PubMed: 21167713]
- Gamble MJ, Frizzell KM, Yang C, Krishnakumar R, Kraus WL. The histone variant macroH2A1 marks repressed autosomal chromatin, but protects a subset of its target genes from silencing. *Genes Dev.* 24:21–32. [PubMed: 20008927]
- Garcia BA, Barber CM, Hake SB, Ptak C, Turner FB, Busby SA, Shabanowitz J, Moran RG, Allis CD, Hunt DF. Modifications of human histone H3 variants during mitosis. *Biochemistry.* 2005; 44:13202–13213. [PubMed: 16185088]
- Garcia BA, Busby SA, Barber CM, Shabanowitz J, Allis CD, Hunt DF. Characterization of phosphorylation sites on histone H1 isoforms by tandem mass spectrometry. *J Proteome Res.* 2004; 3:1219–1227. [PubMed: 15595731]
- Garcia BA, Mollah S, Ueberheide BM, Busby SA, Muratore TL, Shabanowitz J, Hunt DF. Chemical derivatization of histones for facilitated analysis by mass spectrometry. *Nature Protocols.* 2007a; 2:933–938.
- Garcia BA, Pesavento JJ, Mizzen CA, Kelleher NL. Pervasive combinatorial modification of histone H3 in human cells. *Nat Methods.* 2007b; 4:487–489. [PubMed: 17529979]
- Garcia BA, Siuti N, Thomas CE, Mizzen CA, Kelleher NL. Characterization of neurohistone variants and post-translational modifications by electron capture dissociation mass spectrometry. *International Journal of Mass Spectrometry.* 2007c; 259:13p.
- Godde JS, Ura K. Cracking the enigmatic linker histone code. *J Biochem.* 2008; 143:287–293. [PubMed: 18234717]
- Hake SB, Allis CD. Histone H3 variants and their potential role in indexing mammalian genomes: the "H3 barcode hypothesis". *Proc Natl Acad Sci U S A.* 2006; 103:6428–6435. [PubMed: 16571659]
- Hake SB, Garcia BA, Duncan EM, Kauer M, Dellaire G, Shabanowitz J, Bazett-Jones DP, Allis CD, Hunt DF. Expression Patterns and Post-translational Modifications Associated with Mammalian Histone H3 Variants. *Journal of Biological Chemistry.* 2006; 281:10p.
- Harris ME, Bohni R, Schneiderman MH, Ramamurthy L, Schumperli D, Marzluff WF. Regulation of histone mRNA in the unperturbed cell cycle: evidence suggesting control at two posttranscriptional steps. *Mol Cell Biol.* 1991; 11:2416–2424. [PubMed: 2017161]
- Hirota T, Lipp JJ, Toh BH, Peters JM. Histone H3 serine 10 phosphorylation by Aurora B causes HP1 dissociation from heterochromatin. *Nature.* 2005; 438:1176–1180. [PubMed: 16222244]
- Huangfu D, Maehr R, Guo W, Eijkelenboom A, Snitow M, Chen AE, Melton DA. Induction of pluripotent stem cells by defined factors is greatly improved by small-molecule compounds. *Nat Biotechnol.* 2008a; 26:795–797. [PubMed: 18568017]
- Huangfu D, Osafo K, Maehr R, Guo W, Eijkelenboom A, Chen S, Muhlestein W, Melton DA. Induction of pluripotent stem cells from primary human fibroblasts with only Oct4 and Sox2. *Nat Biotechnol.* 2008b; 26:1269–1275. [PubMed: 18849973]
- Ide GJ, Saunders CA. Rapid Isolation of Yeast Nuclei. *Current Genetics.* 1981; 4:85–90.
- Ikura T, Tashiro S, Kakino A, Shima H, Jacob N, Amunugama R, Yoder K, Izumi S, Kuraoka I, Tanaka K, Kimura H, Ikura M, Nishikubo S, Ito T, Muto A, Miyagawa K, Takeda S, Fishel R,

- Igarashi K, Kamiya K. DNA damage-dependent acetylation and ubiquitination of H2AX enhances chromatin dynamics. *Mol Cell Biol.* 2007; 27:7028–7040. [PubMed: 17709392]
- Ishibashi T, Li A, Eirin-Lopez JM, Zhao M, Missiaen K, Abbott DW, Meistrich M, Hendzel MJ, Ausio J. H2A.Bbd: an X-chromosome-encoded histone involved in mammalian spermiogenesis. *Nucleic Acids Res.* 38:1780–1789. [PubMed: 20008104]
- Izzo A, Kamieniarz K, Schneider R. The histone H1 family: specific members, specific functions? *Biol Chem.* 2008; 389:333–343. [PubMed: 18208346]
- Kelleher NL. Top-down proteomics. *Anal Chem.* 2004; 76:197A–203A. [PubMed: 14697051]
- Kuzmichev A, Jenuwein T, Tempst P, Reinberg D. Different EZH2-containing complexes target methylation of histone H1 or nucleosomal histone H3. *Mol Cell.* 2004; 14:183–193. [PubMed: 15099518]
- Lever MA, Th'ng JP, Sun X, Hendzel MJ. Rapid exchange of histone H1.1 on chromatin in living human cells. *Nature.* 2000; 408:873–876. [PubMed: 11130728]
- Lin Q, Sirotkin A, Skoultchi AI. Normal spermatogenesis in mice lacking the testis-specific linker histone H1t. *Mol Cell Biol.* 2000; 20:2122–2128. [PubMed: 10688658]
- Liu H, Sadygov RG, Yates JR 3RD. A model for random sampling and estimation of relative protein abundance in shotgun proteomics. *Anal Chem.* 2004; 76:4193–4201. [PubMed: 15253663]
- Lu S, Xie YM, Li X, Luo J, Shi XQ, Hong X, Pan YH, Ma X. Mass spectrometry analysis of dynamic post-translational modifications of TH2B during spermatogenesis. *Mol Hum Reprod.* 2009; 15:373–378. [PubMed: 19346237]
- Mannironi C, Bonner WM, Hatch CL. H2A.X, a histone isoprotein with a conserved C-terminal sequence, is encoded by a novel mRNA with both DNA replication type and polyA 3' processing signals. *Nucleic Acids Res.* 1989; 17:9113–9126. [PubMed: 2587254]
- Marzluff WF, Gongidi P, Woods KR, Jin J, Maltais LJ. The human and mouse replication-dependent histone genes. *Genomics.* 2002; 80:487–498. [PubMed: 12408966]
- Mckittrick E, Gafken PR, Ahmad K, Henikoff S. Histone H3.3 is enriched in covalent modifications associated with active chromatin. *Proc Natl Acad Sci U S A.* 2004; 101:1525–1530. [PubMed: 14732680]
- Mehta M, Braberg H, Wang S, Lozsa A, Shales M, Solache A, Krogan NJ, Keogh MC. Individual lysine acetylations on the N terminus of *Saccharomyces cerevisiae* H2A.Z are highly but not differentially regulated. *J Biol Chem.* 2010; 285:39855–39865. [PubMed: 20952395]
- Meneghini MD, Wu M, Madhani HD. Conserved histone variant H2A.Z protects euchromatin from the ectopic spread of silent heterochromatin. *Cell.* 2003; 112:725–736. [PubMed: 12628191]
- Millar CB, Xu F, Zhang K, Grunstein M. Acetylation of H2AZ Lys 14 is associated with genome-wide gene activity in yeast. *Genes Dev.* 2006; 20:711–722. [PubMed: 16543223]
- Ong SE, Blagoev B, Kratchmarova I, Kristensen DB, Steen H, Pandey A, Mann M. Stable isotope labeling by amino acids in cell culture, SILAC, as a simple and accurate approach to expression proteomics. *Mol Cell Proteomics.* 2002; 1:376–386. [PubMed: 12118079]
- Ott LW, Resing KA, Sizemore AW, Heyen JW, Cocklin RR, Pedrick NM, Woods HC, Chen JY, Goebel MG, Witzmann FA, Harrington MA. Tumor Necrosis Factor- α - and interleukin-1-induced cellular responses: coupling proteomic and genomic information. *J Proteome Res.* 2007; 6:2176–2185. [PubMed: 17503796]
- Pesavento JJ, Mizzen CA, Kelleher NL. Quantitative Analysis of Modified Proteins and Their Positional Isomers by Tandem Mass Spectrometry: Human Histone H4. *Analytical Chemistry.* 2006; 78:4271–4280. [PubMed: 16808433]
- Plazas-Mayorca MD, Bloom JS, Zeissler U, Leroy G, Young NL, Dimaggio PA, Krugylak L, Schneider R, Garcia BA. Quantitative proteomics reveals direct and indirect alterations in the histone code following methyltransferase knockdown. *Mol Biosyst.* 6:1719–1729. [PubMed: 20577673]
- Plazas-Mayorca MD, Zee BM, Young NL, Fingerman IM, Leroy G, Briggs SD, Garcia BA. One-pot shotgun quantitative mass spectrometry characterization of histones. *J Proteome Res.* 2009; 8:5367–5374. [PubMed: 19764812]
- Poirier GG, Savard P. ADP-ribosylation of pancreatic histone H1 and of other histones. *Can J Biochem.* 1980; 58:509–515. [PubMed: 6791796]

- Rogakou EP, Pilch DR, Orr AH, Ivanova VS, Bonner WM. DNA double-stranded breaks induce histone H2AX phosphorylation on serine 139. *J Biol Chem.* 1998; 273:5858–5868. [PubMed: 9488723]
- Shechter D, Dormann HL, Allis CD, Hake SB. Extraction, purification and analysis of histones. *Nat Protoc.* 2007; 2:1445–1457. [PubMed: 17545981]
- Siuti N, Roth MJ, Mizzen CA, Kelleher NL, Pesavento JJ. Gene-specific characterization of human histone H2B by electron capture dissociation. *J Proteome Res.* 2006; 5:233–239. [PubMed: 16457587]
- Smith CM, Haimberger ZW, Johnson CO, Wolf AJ, Gafken PR, Zhang Z, Parthun MR, Gottschling DE. Heritable chromatin structure: mapping "memory" in histones H3 and H4. *Proc Natl Acad Sci U S A.* 2002; 99 Suppl 4:16454–16461. [PubMed: 12196632]
- Stein, GS.; Stein, JL.; Marzluff, WF. *Histone Genes.* New York: John Wiley and Sons; 1984.
- Strahl BD, Allis CD. The language of covalent histone modifications. *Nature.* 2000; 403:41–45. [PubMed: 10638745]
- Syka JE, Coon JJ, Schroeder MJ, Shabanowitz J, Hunt DF. Peptide and protein sequence analysis by electron transfer dissociation mass spectrometry. *Proc Natl Acad Sci U S A.* 2004; 101:9528–9533. [PubMed: 15210983]
- Tagami H, Ray-Gallet D, Almouzni G, Nakatani Y. Histone H3.1 and H3.3 complexes mediate nucleosome assembly pathways dependent or independent of DNA synthesis. *Cell.* 2004; 116:51–61. [PubMed: 14718166]
- Talasaz H, Helliger W, Puschendorf B, Lindner H. In vivo phosphorylation of histone H1 variants during the cell cycle. *Biochemistry.* 1996; 35:1761–1767. [PubMed: 8639656]
- Talasaz H, Sarg B, Lindner HH. Site-specifically phosphorylated forms of H1.5 and H1.2 localized at distinct regions of the nucleus are related to different processes during the cell cycle. *Chromosoma.* 2009; 118:693–709. [PubMed: 19609548]
- Thomas CE, Kelleher NL, Mizzen CA. Mass Spectrometric Characterization of Human Histone H3: A Bird's Eye View. *Journal of Proteome Research.* 2006; 5:240–247. [PubMed: 16457588]
- Tian Z, Zhao R, Tolic N, Moore RJ, Stenoien DL, Robinson EW, Smith RD, Pasa-Tolic L. Two-dimensional liquid chromatography system for online top-down mass spectrometry. *Proteomics.* 2010; 10:3610–3620. [PubMed: 20879039]
- Van Lint C, Emiliani S, Ott M, Verdin E. Transcriptional activation and chromatin remodeling of the HIV-1 promoter in response to histone acetylation. *EMBO J.* 1996; 15:1112–1120. [PubMed: 8605881]
- Von Holt C, Brandt WF, Greyling HJ, Lindsey GG, Retief JD, Rodrigues JD, Schwager S, Sewell BT. Isolation and characterization of histones. *Methods Enzymol.* 1989; 170:431–523. [PubMed: 2549339]
- Wang H, Wang L, Erdjument-Bromage H, Vidal M, Tempst P, Jones RS, Zhang Y. Role of histone H2A ubiquitination in Polycomb silencing. *Nature.* 2004; 431:873–878. [PubMed: 15386022]
- Washburn MP, Ulaszek RR, Yates JR 3RD. Reproducibility of quantitative proteomic analyses of complex biological mixtures by multidimensional protein identification technology. *Anal Chem.* 2003; 75:5054–5061. [PubMed: 14708778]
- Waterborg JH, Winicov I, Harrington RE. Histone Variants and Acetylated Species from the Alfalfa Plant *Medicago-Sativa*. *Archives of Biochemistry and Biophysics.* 1987; 256:167–178. [PubMed: 3606121]
- Weiss T, Hergeth S, Zeissler U, Izzo A, Tropberger P, Zee BM, Dundr M, Garcia BA, Daujat S, Schneider R. Histone H1 variant-specific lysine methylation by G9a/KMT1C and Glp1/KMT1D. *Epigenetics Chromatin.* 3:7. [PubMed: 20334638]
- Wisniewski JR, Zougman A, Kruger S, Mann M. Mass spectrometric mapping of linker histone H1 variants reveals multiple acetylations, methylations, and phosphorylation as well as differences between cell culture and tissue. *Mol Cell Proteomics.* 2007; 6:72–87. [PubMed: 17043054]
- Woodcock CL, Skoultchi AI, Fan Y. Role of linker histone in chromatin structure and function: H1 stoichiometry and nucleosome repeat length. *Chromosome Res.* 2006; 14:17–25. [PubMed: 16506093]

- Wyrick JJ, Parra MA. The role of histone H2A and H2B post-translational modifications in transcription: a genomic perspective. *Biochim Biophys Acta*. 2009; 1789:37–44. [PubMed: 18675384]
- Xu M, Long C, Chen X, Huang C, Chen S, Zhu B. Partitioning of histone H3–H4 tetramers during DNA replication-dependent chromatin assembly. *Science*. 2010; 328:94–98. [PubMed: 20360108]
- Young NL, Dimaggio PA, Plazas-Mayorca MD, Baliban RC, Floudas CA, Garcia BA. High throughput characterization of combinatorial histone codes. *Mol Cell Proteomics*. 2009; 8:2266–2284. [PubMed: 19654425]
- Zalensky AO, Siino JS, Gineitis AA, Zalenskaya IA, Tomilin NV, Yau P, Bradbury EM. Human testis/sperm-specific histone H2B (hTSH2B). Molecular cloning and characterization. *J Biol Chem*. 2002; 277:43474–43480. [PubMed: 12213818]
- Zee BM, Levin RS, Bo X, Leroy G, Wingreen NS, Garcia BA. In Vivo Residue-specific Histone Methylation Dynamics. *Journal of Biological Chemistry*. 2010a; 285:10p.
- Zee BM, Levin RS, Dimaggio PA, Garcia BA. Global turnover of histone post-translational modifications and variants in human cells. *Epigenetics Chromatin*. 2010b; 3:22. [PubMed: 21134274]
- Zheng Y, John S, Pesavento JJ, Schultz-Norton JR, Schiltz RL, Baek S, Nardulli AM, Hager GL, Kelleher NL, Mizzen CA. Histone H1 phosphorylation is associated with transcription by RNA polymerases I and II. *J Cell Biol*. 189:407–415.
- Zubarev RA, Kelleher NL, McLafferty FW. Electron capture dissociation of multiply charged protein cations. A nonergodic process. *Journal of the American Chemical Society*. 1998; 120:3265–3266.

Figure 1.

All of the histone variants contain a highly conserved histone fold domain and vary mainly in their C and N-terminal sequences. Shown above is a schematic comparing histone variant sequences. Boxes represent the histone fold domain and orange lines represent site-specific sequence variations. Histones that are in different shades of the same color are from the same histone family but have large differences in sequence.

Figure 2.

(A) A sequence alignment H2B1K and H2B1H demonstrate the extent of sequence similarity between H2B isoforms. (B) Side by side comparison of a mixture of H2B sequences analyzed using Bottom Up, Middle Down and Top Down mass spectrometry. Top Down: The Top Down mass spectra were collected using direct infusion of purified H2B and fragmented using ETD on an LTQ-Orbitrap. Isobaric H2B1K and H2B1H species could be differentiated after MS/MS analysis. Middle Down: Middle Down spectra were collected after online RP-HPLC separation of Glu-C digested H2B and fragmented using ETD. A peak corresponding to the N-terminal tail of H2B1H was chosen for MS/MS analysis, however the N-terminal tail of H2B1H is identical to that of H2B1O and H2B2F. Seven out of 14 somatic H2B variants can be uniquely identified using Middle Down MS. Bottom Up: Bottom Up spectra were collected after separation of peptides generated from a tryptic digest of H2B and analyzed using CAD fragmentation. Four out of 14 H2B variants can be uniquely identified using this method.

Figure 3. Histone Quantification Methods. Depicted above are (A.) chemical derivatization, (B.) metabolic labeling, (C.) standard peptides and (D.) label-free quantification. For the labeling based strategies (A.,B., C.), the processing step where label incorporation occurs is shown.