Published in final edited form as:

Med Care Res Rev. 2010 October ; 67(5 Suppl): 163S–197S. doi:10.1177/1077558710374621.

The Use of Quality Improvement and Health Information Technology Approaches to Improve Diabetes Outcomes in African-American and Hispanic Patients

Arshiya A. Baig, MD, MPH¹, Abigail E. Wilkes, MPH¹, Andrew M. Davis, MD, MPH¹, Monica E. Peek, MD, MPH¹, Elbert S. Huang, MD, MPH¹, Douglas S. Bell, MD, PhD^{2,3}, and Marshall H. Chin, MD, MPH¹

¹Department of Medicine, University of Chicago, Chicago, IL

²RAND Corporation, Santa Monica, CA

³Department of Medicine, David Geffen School of Medicine at UCLA, Los Angeles, CA

Abstract

Differences in rates of diabetes-related lower extremity amputations represent one of the largest and most persistent health disparities found for African-Americans and Hispanics compared to whites in the United States. Since many minority patients receive care in under-resourced settings, quality improvement (QI) initiatives in these settings may offer a targeted approach to improve diabetes outcomes in these patient populations. Health information technology (health IT) is widely viewed as an essential component of health care QI and may be useful in decreasing diabetes disparities in under-resourced settings. This article reviews the effectiveness of health care interventions utilizing health IT to improve diabetes process of care and intermediate diabetes outcomes in African-American and Hispanic patients. Health IT interventions have addressed patient, provider, and system challenges in the provision of diabetes care but require further testing in minority patient populations to evaluate their effectiveness in improving diabetes outcomes and reducing diabetes-related complications.

Keywords

diabetes; health information technology; health disparities; quality improvement; under resourced settings

Corresponding author: Arshiya Baig, MD, MPH, 5841 S. Maryland Ave, MC 2007, Chicago, IL 60637. Phone: 773.834.4760. Fax: 773. 834.2238. abaig@uchicago.edu.
Abigail E. Wilkes: 5841 S. Maryland Ave, MC 2007, Chicago, IL 60637. Phone: 773.834.2605. Fax: 773. 834.2238. Email: awilkes@medicine.bsd.uchicago.edu
Andrew M. Davis: 5841 S. Maryland Ave, MC 2007, Chicago, IL 60637. Phone: 773.834.5921. Fax: 773. 702.2468. Email: amd@uchicago.edu
Monica E. Peek: 5841 S. Maryland Ave, MC 2007, Chicago, IL 60637. Phone: 773.702.2083. Fax: 773. 834.2238. Email: mpeek@medicine.bsd.uchicago.edu
Elbert S. Huang: 5841 S. Maryland Ave, MC 2007, Chicago, IL 60637. Phone: 773.834.3419. Fax: 773. 834.2238. Email: ehuan@medicine.bsd.uchicago.edu
Douglas S. Bell: UCLA School of Medicine, Med-GIM & HSR, 911 Broxton Plaza 1st Floor, Box 951736, Los Angeles, CA 90095-1736. Phone: (310) 794-3168. Fax: (310) 794-0732. Email: Dbell@mednet.ucla.edu
Marshall H. Chin: 5841 S. Maryland Ave, MC 2007, Chicago, IL 60637. Phone: 773.702.4769. Fax: 773. 834.2238. Email: mchin@medicine.bsd.uchicago.edu

Background

Chronic conditions are the leading cause of disparities in health (Wong, Martin, Shapiro, Boscardin, & Ettner, 2002; Centers for Disease Control and Prevention (CDC), 2008). Minorities and low-income individuals suffer from the highest rates of chronic conditions, such as diabetes (CDC, 2008; National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK), 2008). Mexican-Americans and African-Americans are almost twice as likely as non-Hispanic whites to have diabetes (CDC, 2008). Considering the projected increase of diabetes prevalence among Hispanics and African-Americans, mitigating these disparities is crucial (Narayan, Boyle, Geiss, Saaddine, & Thompson, 2006).

Of particular concern, the projected increase in diabetes incidence among these ethnic groups may fuel growing disparities in diabetes complications if the disease is not controlled equally across ethnic groups (Agency for Healthcare Research and Quality (AHRQ), 2007). Diabetes puts individuals at risk for several major microvascular and macrovascular complications, including chronic kidney disease, diabetic retinopathy, coronary artery disease, peripheral vascular disease, and lower extremity amputations. Despite the availability of effective medications and well-published evidence-based treatments, only one-third of Hispanics and African-Americans with diabetes achieve optimal glycemic control (American Diabetes Association, 2008; Intensive blood-glucose control, 1998; Kirk, Ralph, Bell, Passmore, & Bonds, 2006; Koro, Bowlin, Bourgeois, & Fedder 2004; Saydah, Cowie, Eberhardt, Rekenire, & Narayan, 2007; Tight blood pressure control, 1998). Hispanics and African-Americans with diabetes also have lower rates of blood pressure and lipid control compared to whites (Collins et al., 2007; Duru et al., 2009; Rosamond et al., 2007). The suboptimal control of blood glucose, blood pressure, and lipids places these individuals at higher risk for complications of diabetes.

Racial and ethnic disparities in lower extremity amputation (LEA) rates among people with diabetes have been well documented in numerous population-based studies (Dillingham, Pezzin, & Mackenzie, 2002; Morrissey et al., 2007; Regenbogen, Gawande, Lipsitz, Greenberg, & Jha, 2009). The 2008 National Healthcare Disparities Report (AHRQ, 2009) reported that African-Americans were 2.3 times as likely as whites to be hospitalized for LEA. Similarly, the 2007 National Healthcare Disparities Report (AHRQ, 2008) found that Hispanics were 2.9 times as likely as whites to be hospitalized for LEA from diabetes. According to AHRQ, the difference in rates of LEA hospitalizations for African-Americans and Hispanics compared to whites is one of the largest disparities seen in health care quality for these groups and is worsening.

Quality improvement initiatives have shown promise in improving diabetes quality of care and may be critical in decreasing diabetes disparities (Peek, Cargill, & Huang, 2007). QI initiatives that address salient intermediate outcomes in diabetes—such as control of glycosylated hemoglobin (HbA1c), blood pressure, and lipids and improved processes of care, such as foot examinations and medication intensification—may be central to decreasing disparities in LEA (Schade & Hannah, 2007; Selvin, Wattanakit, Steffes, Coresh, & Sharrett, 2006). QI initiatives in all health care settings provide an opportunity to address disparities in diabetes care and quality, but QI initiatives in under-resourced settings may present a particularly promising opportunity, since many minority patients receive care in rural health centers, county hospitals, or safety net clinics (Bach, Pham, Schrag, Tate, & Hargraves, 2004; Jha, Orav, Li, & Epstein, 2007; Jha, Orav, Zheng, & Epstein, 2008). Physicians in these settings often face resource constraints and report greater difficulties in obtaining access for their patients to subspecialists, high-quality diagnostic imaging, and nonemergency admissions to the hospital (Bach et al., 2004; Jha et al., 2007; Jha et al., 2008). Finding innovative ways to improve diabetes processes of care and intermediate

outcomes in under-resourced settings may be a powerful way to address disparities in diabetes complications (Bach et al., 2004).

Health information technology (health IT) is widely viewed as an essential component of health care quality improvement. QI interventions using health IT have been found to improve adherence to guideline-based care, enhance surveillance and monitoring, decrease medication errors, and decrease utilization of care (Chaudhry et al., 2006). Many health IT interventions, such as electronic medical records (EMR), computerized prompts, population management (including reports and feedback), specialized decision support, electronic scheduling, and personal health records have been shown to improve diabetes processes of care and intermediate outcomes. (Dorr et al., 2007) Implementation of health IT has become a national priority, but adoption still remains low (American Recovery and Reinvestment Act of 2009; DesRoches et al., 2008; Foxhall, 2009). Community health centers and small private practices lag in implementing health IT due to barriers prevalent in these settings (Jha, DesRoches, & Campbell, 2009; Shields et al., 2007). Encouraging health IT adoption as a part of QI initiatives to improve diabetes care in under-resourced settings may provide a special opportunity to reduce diabetes disparities.

New Contribution

To our knowledge, no systematic literature review has examined the use of health IT in quality improvement initiatives to reduce diabetes health disparities. Additionally, few studies examine the use of health IT in under-resourced settings. We systemically review health IT interventions to examine their impact on diabetes outcomes in African-American and Hispanic patient populations. Our review focuses on intermediate diabetes outcomes and processes of care, which are on the causal pathway to lower extremity amputations. For this study, we review health care interventions that target diabetes disparities in African-American and Hispanic patients and discuss the use of health IT for enhancing these QI efforts. We aim to (1) update from 2006 to 2009 a previous systematic review of QI initiatives to improve diabetes outcomes in minority patients; (2) systematically review QI interventions that utilized health IT to improve diabetes disparities from 2000 to 2009; and (3) give recommendations on the use of health IT to reduce diabetes disparities.

Conceptual Framework

Berwick's (2002) model summarizes the lessons from the Institute of Medicine's "Crossing the Quality Chasm" report. Using this model, we can organize QI initiatives from Level A to Level D and comment on QI strategies that target the patient experience (A), microsystems (B), organizations that house these microsystems (C), and policy or regulatory factors (D). We used Berwick's model to classify the types of QI initiatives previously reviewed in decreasing diabetes disparities (Peek et al., 2007).

Peek et al. (2007) published a systematic review of QI interventions in health care settings that aimed to reduce diabetes disparities in minority populations. The authors categorized interventions into those that focused on patients, providers, and health care systems. Among patient-oriented QI initiatives, those that utilized interpersonal relationships and social networks and were culturally tailored had the most success in improving diabetes outcomes. Provider QI initiatives that incorporated in-person feedback were also successful. Organizational changes—such as the use of case management, community health workers, and non-physician providers—were effective at improved rates of making and keeping appointments; overcoming social, cultural and linguistic barriers; and providing clinical care through treatment algorithms. The use of medical assistance programs increased prescription adherence and decreased hospitalizations. Disease management systems—such as patient

registries, practice guidelines, case management, and tracking and monitoring of patients—improved diabetes processes and outcomes.

Similar to the goal of the Peek et al. (2007) study, we aimed to identify efforts to reduce diabetes disparities in Hispanic and African-American populations in under-resourced settings. In our study, under-resourced settings included federally qualified health centers, rural clinics, public hospitals, and public clinics. We begin by using the Berwick (2002) model to describe the challenges these settings face in reducing diabetes disparities and to identify promising points of intervention for health IT use at the patient, provider, and system levels. The health IT applications described here were not necessarily applied in under-resourced settings, however they are meant to serve as examples of what may be possible.

Patient Centered Challenges

Studies with African-American and Latino patients with diabetes have demonstrated that medication adherence, health care cost, and lack of health insurance serve as barriers to receiving effective diabetes care. Reichsman, Warner, & Cella, 2009; Rosal, Benjamin, & Pekow, 2008). Some patients also have poor access to diabetes education in under-resourced settings, in particular information that is culturally tailored, in their language, and easy to comprehend (Reichsman et al., 2009). Poor communication, language barriers, lack of trust in the health care system, and lack of provider cultural competence each adversely affect patient satisfaction (Reichsman et al., 2009; Rosal et al., 2008). Patient directed interventions, therefore, can help address these challenges by improving diabetes knowledge, encouraging provider communication, strengthening social support from peers, and providing access to care.

Health IT may improve diabetes care through patient directed interventions. Patient satisfaction, self-management, self-empowerment, and diabetes knowledge have been enhanced by patient-focused health IT applications, such as online, interactive personal health records, secure messaging with physicians, and video visits (Halamka, Mandl, & Tang 2008; Katz, Nissan, & Moyer 2004; Wakefield et al., 2008). Patients can schedule their appointments online while receiving social support and reminders from peers on a diabetes Web site (Eysenbach, 2008). Patient communication may be enhanced through secure messaging with physicians while simultaneously decreasing the cost of frequent visits for patients and providers. Patient directed interventions also could be designed to specifically target those with low health literacy or English as a second language. Trust in the health care system also may be improved using an interactive personal health record that promotes transparency and secure messaging with providers.

Provider Centered Challenges

Clinicians in safety net clinics note many financial, cultural, and psychosocial factors that impact the quality of diabetes care (Reichsman et al., 2009). Providers report inadequate time to counsel patients on lifestyle modifications, limited resources for dietary and physical activity counseling, limited documentation of diabetes-related risk factors, limited availability of medical records, and inconsistent delivery of diabetes prevention strategies (such as dietary or weight loss counseling and pharmacological interventions) to patients in under-resourced settings (Rosal et al., 2008). Diabetes care may be enhanced through health IT applications that address these provider related challenges.

Safety net providers using health IT have reported benefit from increased efficiency in providing care through the use of EMRs (Kim, Chen, Keith, Yee, & Krushei, 2009). EMRs can also be used for clinical documentation, data storage and tracking, and results

management (Davis, Doty, Shea, & Stremikis, 2009). Computerized order entry, standardized histories, and clinical decision support can increase appropriate medication prescribing and reduce adverse drug interactions, and they have the potential to provide better adherence to prescribing guidelines (Davis et al., 2009).

System Centered Challenges

Organizational and policy level challenges specific to under-resourced health care settings also exist. These challenges include long waits for appointments, poor access to subspecialists, difficulty in the formation of multidisciplinary teams, and lack of access to funding for infrastructure upgrades. Other policy level and regulatory barriers to QI implementation include lack of reimbursement for non-visit care and coordination, drug costs, and lack of the presence and/or accessibility to community health workers and community outreach programs.

Health IT applications can address systems-based challenges to improve diabetes quality of care. Health care systems can track disparities through health information exchanges (HIEs), regional health information systems, and disease registries. Telemedicine and video visits may improve the timeliness of care and decrease long waits for appointments in under-resourced settings. Improved communication through secure messaging or social networking sites can facilitate formation of multidisciplinary teams. Text messaging systems that provide messaging among clinics, patients, and outreach workers may enhance the presence and accessibility of community health care workers in community level interventions.

Methods

Systematic QI Intervention Search

The review article, "Diabetes Health Disparities, A Systematic Review of Health Care Interventions," by Peek et al. (2007) described a detailed electronic database search and hand search of health system-based diabetes QI interventions targeting minority populations from 1985 to 2006. We updated this review with studies published from 2006 to 2009.

To update this review, we searched for diabetes QI intervention articles published from January 2006 to March 2009 using electronic databases Cochrane, CINAHL, ACP Journal Club, Psycinfo, and Medline. We used prespecified Medical Subject Headings (MeSH) and keywords to identify evaluation studies (evaluation studies, clinical Trials; effectiveness; improvement; performance) designed to address health care delivery (delivery of health care, integrated; quality of health care; health services accessibility) among African-Americans and Hispanics (African-Americans; Hispanic Americans; Mexican Americans; Latinos) and among adults and children with diabetes (diabetes mellitus, type 2; diabetes mellitus, type 1; diabetes complications; peripheral vascular disease). We also utilized reference lists from recently published QI intervention articles and major review articles. We supplemented our electronic search with a hand search of issues from selected journals with a high likelihood of publishing diabetes QI studies (Diabetes Care, Diabetes Educator, Journal of General Internal Medicine, and Medical Care) published within the preceding 2 years.

Studies were required to include at least a 50% minority sample population, be based in the United States, and focus on improving diabetes treatment processes and outcomes. We defined "minority" as African-American or Hispanic American/Latino. We excluded studies that focused on diabetes prevention and gestational diabetes. We included all study types in our review (e.g. randomized controlled trials, pre-post, pilot, cross-sectional). We included QI interventions whether they used health IT or not.

From the Cochrane, CINAHL, ACP Journal Club, and Psycinfo databases we found 63 articles, but none met our inclusion criteria. Our Medline search returned 43 articles, and we retained one after application of inclusion/exclusion criteria. From our hand search, we found 18 articles that met exclusion/inclusion criteria.

Systematic Health Information Technology (Health IT) Search

While some of the articles from Peek et al. (2007) and our update search contained aspects of health IT, we conducted a supplementary health IT electronic database search from 2000 to 2009. For our systematic search of QI interventions that utilized health IT, we added keyword and MeSH terms for health IT (biomedical technology; medical informatics applications; data collection) and removed all other search terms except those referring to diabetes and peripheral vascular disease (diabetes mellitus, type 2; diabetes mellitus, type 1; diabetes complications; peripheral vascular disease) using the Medline database only and restricted the search to articles published between January 2000 and March 2009. This search yielded 286 articles, and after applying our inclusion/exclusion criteria, we kept none. In addition, a hand search of issues of selected journals with a high likelihood of publishing health IT interventions (e.g., HealthCare Informatics and the Journal of the American Medical Informatics Association) published within the previous 2 years was conducted. References from health IT review articles, including Peek et al.'s (2007) review, were investigated for eligibility for our review. Our hand search resulted in 31 articles; 10 met our inclusion/exclusion criteria.

The QI update from 2006 to 2009 yielded 18 articles. The expanded health IT QI search from 2000 to 2009, yielded 10 articles, with one overlapping article with the QI update (Shea et al., 2009) Since the overlapping article reported on the use of health IT in a diabetes QI intervention, we report that study's findings in the health IT results section.

Data Abstraction

We used a validated instrument to guide our abstraction (Zaza et al., 2000). We abstracted data from each study into a table that included the objective of the improvement strategy, the population sampled, the setting, a description of the form of health IT used, and the findings. We documented baseline and follow-up rates of diabetes processes of care (proportion of patients receiving HbA1c testing, lipid testing, blood pressure measurement, foot examination, and eye examination), health outcomes (HbA1c, total cholesterol, low density lipoprotein (LDL) cholesterol, high density lipoprotein (HDL) cholesterol, triglycerides, systolic blood pressure, and diastolic blood pressure), and diabetes complication rates (blindness, rates of myocardial infarctions, kidney failure, peripheral vascular disease, and lower extremity amputation hospitalizations) when noted in the article. In addition, each study in the table was given a quality score by two members of the research team using an adapted questionnaire based on the Downs and Black (1998) guidelines (27 point scale: 0 worst, 27 best). We report the average of the two quality scores.

Results

We report the findings from our review in two sections: (1) QI systematic review and 2) health IT systematic review. The QI review describes the 18 articles we found in our search for diabetes QI interventions from 2006 to 2009. We present an update on QI initiatives targeted at improving diabetes processes of care and outcomes in minority patient populations and highlight new findings and changes from Peek et al. (2007). The health IT review reports the findings from the 10 articles that used health IT in their diabetes QI interventions from 2000 to 2009.

Quality Improvement Systematic Review

Eighteen articles used QI initiatives to improve diabetes processes of care and outcomes among African-American and Hispanic patient populations, not including health IT-oriented initiatives (Babamoto et al., 2009; Cramer et al., 2007; Davidson, Ansari, & Karlan, 2007; Gold et al., 2008; Ingram et al., 2007; Joshu, Rangel, Garcia, Brownson, & O'Toole, 2007; King et al., 2006; Liebman, Heffernan, & Sarvela, 2007; Lujan, Ostwald, & Ortiz, 2007; Mahotiere, Ocepok-Welikson, Daley, & Byssainthe, 2006; Mauldon, Melkus, & Cagganello 2006; Miller et al., 2006; Sixta & Ostwald, 2008; Steinhardt, Mamerow, Brown, & Jolly, 2009; Thom, Tirado, Woon, & McBride, 2006; Thompson, Horton, & Flores 2007; Utz et al., 2008; Wagner, Pizzimenti, Daniel, Pandya, & Hardigan, 2008). Table 1 summarizes QI interventions from Peek, et al. (2007) and from the 2006–2009 update that have demonstrated improvements in a range of diabetes processes of care and clinical outcomes.

Patient oriented—Of the 18 articles, 6 reported interventions that were patient oriented (Cramer et al., 2007; Liebman et al., 2007; Mauldon et al., 2006; Steinhardt et al., 2009; Utz et al., 2008; Wagner et al., 2008). Of these, five were culturally tailored for their specific population (Cramer et al., 2007; Liebman et al., 2007; Mauldon et al., 2006; Steinhardt et al., 2009; Utz et al., 2008). Studies emphasizing cognitive-behavioral education and self-care management and those that adapted the Diabetes Prevention Program demonstrated improvements in diabetes metabolic outcomes (Cramer et al., 2007; Mauldon et al., 2006). Interventions based on enhancing patient resilience improved a range of patient outcomes (Steinhardt et al., 2009).

Provider oriented—We found two new studies that tested provider directed interventions (Miller et al., 2006; Thom et al., 2006). Thom's study tested provider cultural competency training but failed to show any statistically significant clinical improvements in diabetes outcomes. When treatment algorithms factoring in HbA1c and random blood sugar were available at the time of patients' visits, providers were more likely to intensify medical therapy and lower patient HbA1c levels (Miller et al., 2006).

System oriented—We found nine studies that focused on health system interventions for improving diabetes care for minorities (Babamoto et al., 2009; Davidson et al., 2007; Gold et al., 2008; Ingram et al., 2007; Joshu et al., 2007; King et al., 2006; Lujan et al., 2007; Sixta & Ostwald, 2008; Thompson et al., 2007). Confirming Peek's work (2007), we also found that the use of community health workers and promotoras facilitated improvements in diabetes intermediate outcomes, processes of care, knowledge, and social support (Babamoto et al., 2009; Ingram et al., 2007; Joshu et al., 2007; Lujan et al., 2007; Sixta & Ostwald 2008; Thompson et al., 2007). Our update also provided evidence that promotoras can successfully reach out to patients with diabetes on the fringes of the traditional health care system (Ingram et al., 2007). All six community health worker studies were conducted in majority Mexican-American populations. The use of nonphysician providers reduced diabetes-related urgent visits and emergency room visits (Davidson et al., 2007). The use of a centralized diabetes specialist in guiding off-site primary treatment of diabetes demonstrated trends in improvements in HbA1c, lipids, and blood pressure that were not statistically significant (King et al., 2006). Interventions utilizing multidisciplinary teams improved patients' HbA1c levels (Gold et al., 2008).

Multi-target interventions—We found one study that used a multi-target approach for reducing the disparity in the rate of biennial lipid profiles between African-Americans and white Medicare beneficiaries with diabetes in New York City (Mahotiere et al., 2006).

Health Information Technology Systematic Review

We identified 8 studies with 10 articles that utilized health IT to address diabetes quality improvement in minority populations (Bray, Roupe et al., 2005; Bray, Thompson, Wynn, Cummings, & Whetstone, 2005; Cherry, Moffatt, Rodriguez, & Dryden, 2002; Chin et al, 2004; Gerber et al., 2005; Levetan, Dawn, Robbins, & Ratner, 2002; Phillips et al., 2005; Shea et al, 2006; Shea et al., 2009; Ziemer et al., 2006). Table 2 describes these studies in detail.

Patient oriented—We identified two studies that utilized health IT in patient-oriented QI initiatives (Gerber et al., 2005; Levetan, Dawn, Robbins, & Ratner, 2002).

The Levetan et al. (2002) study evaluated the impact of computer-generated personalized goals on HbA1c among a majority African-American patient population and had a quality score of 17. This randomized controlled trial found that those who received the personalized goal report who had a baseline HbA1c $\geq 7.0\%$ had an absolute reduction of 1.69% in HbA1c versus 0.77% in HbA1c among control subjects at 6 months follow-up ($p=0.03$).

The Gerber et al. (2005) randomized controlled trial evaluated a clinic-based multimedia intervention using computer kiosks for diabetes education targeting individuals with low health literacy in five inner city, publicly run health centers with a mix of Latino and African-American patients (quality score 20). This study found no changes in HbA1c, weight, blood pressure, diabetes knowledge, self-efficacy, or self-reported medical care at the end 1 year.

The Levetan et al. (2002) study demonstrated that a personalized report of diabetes values and goals may hold promise for improving HbA1c, but a clinic needs to have an EMR already in use and a software algorithm to conduct this intervention. Gerber's work targeted those with lower health literacy and less computer experience but had problems with computer malfunction. The Gerber et al. (2005) study highlighted the additional challenges of appropriate technical support, testing, and user skills for implementing advanced IT interventions in safety net settings.

Provider oriented—We identified two articles that utilized health IT in provider-oriented QI initiatives from the Improving Primary Care of African-Americans with Diabetes (IPCAAD) study (Phillips et al., 2005; Ziemer et al., 2006). IPCAAD was a randomized controlled trial that evaluated computerized reminders and in-person feedback from an endocrinologist on patient diabetes outcomes and provider behavior regarding medication intensification. The study was conducted in a municipal hospital primary care clinic within a large academic center serving primarily African-American patients and had a 3-year follow-up.

The Phillips et al. (2005) study found that improvement in HbA1c was significantly greater for patients who saw providers receiving in-person feedback plus reminders compared to the providers in the control group (-0.6% vs. -0.2% , $p<0.02$). It also found improvements in systolic blood pressure within the feedback plus reminder and feedback only groups but not in the reminders only or control arm. LDL within groups improved in all intervention arms. This study had a quality score of 18.

In the Ziemer et al. (2006) study, medication intensification occurred more frequently when providers received in-person feedback with or without reminders compared to the group that received reminders alone or usual care (52% vs. 35% , $p<0.001$) at 3 years. Adequate medication intensification was associated with a significant fall in HbA1c (-0.19% , $p<0.001$). This study had a quality score of 16.5.

The IPCAAD studies demonstrated that electronic clinical reminders can modify provider prescribing behavior and improve glycemic control. Reminders were more effective when the providers received in-person feedback from an endocrinologist. Considering the lack of access to subspecialty care in under-resourced settings, this intervention may not be feasible to implement in safety net settings.

System oriented—We found five studies that published six articles assessing organization/system-based interventions (Bray, Roupe et al., 2005; Bray, Thompson, Wynn, Cummings, & Whetstone, 2005; Cherry et al., 2002; Chin et al., 2004; Shea et al., 2006; Shea et al., 2009)

The Health Resources and Services Administration's Health Disparities Collaboratives is a QI model that used plan-do-study-act cycles, the MacColl Institute's Chronic Care Model, and learning sessions to improve diabetes outcomes. In the first year of the intervention in 19 Midwestern health centers, 37% of the community health centers chose to implement an electronic registry to track patient examination and laboratory data (Chin et al., 2004). Several key processes of care improved at 1 year with the intervention, including rates of HbA1c measurement (80% to 90%, adjusted odds ratio 2.1, 95% CI 1.6–2.8), foot examination (40% to 64%; OR: 2.7, 95% CI 1.8–4.1), and lipid assessments (55% to 66%, OR: 1.6, 95% CI 1.1–2.3). HbA1c did not significantly improve. Staff from the sites that developed a registry found it useful but burdensome and requested more technical support. This study had a quality score of 13. There have been follow-up studies to Chin's initial evaluation. A national study of the Health Disparities Collaboratives found improvement in diabetes processes of care but not in intermediate outcomes compared to matched control clinics at 1- to 2-year follow-up. (Landon et al., 2007) Follow-up at 4 years of diabetic patients in 16 Midwestern and West Central health centers in the Health Disparities Collaboratives found improvements in processes of care and HbA1c and LDL cholesterol values (Chin et al., 2007).

Bray's studies also describe interventions using an electronic registry called the Cardiovascular/Diabetes Electronic Management System (CVDEMS) (Bray, Roupe et al., 2005; Bray, Thompson et al., 2005). CVDEMS is an electronic patient care registry system that allows office staff to enter demographic information regarding each patient into a clinic population registry. Providers can query detailed patient and laboratory information and obtain summary reports. The two Bray et al. (2005) studies explored the feasibility, cost-effectiveness, and efficacy of using diabetes care management, group visits, visit reminders, and CVDEMS. The intervention was implemented in solo or small group primary care practices with a majority rural African-American patient population with diabetes. This study used a pre-post design with control patients from a separate clinical site. HbA1cs were similar at baseline for both the intervention (8.2 ± 2.6) and control groups (8.3 ± 2.6), but mean HbA1c was significantly lower in the intervention group (7.1 ± 2.3) versus the control group (8.6 ± 2.4 ; $p < 0.05$) at 1 year. Patients receiving the intervention experienced a 61% reduction in HbA1c, while the control group increased HbA1c levels. Documentation of lipid panels (55% to 76%) and foot examinations (12% to 54%) improved within the intervention group at 1 year. Provider productivity also improved from 20.17 to 31.55 average daily encounters. Some clinical staff thought the CVDEMS was duplicative of the medical record, and the leadership began to move toward implementation of an EMR in the practices. These studies had an average quality score of 11.5.

Two studies evaluated telemedicine as a way to bridge the gap between office visits for patients and access to subspecialty care in under-resourced settings (Cherry et al., 2002; Shea et al., 2006). Cherry's study (2002) tested a Web-based patient interface technology that prompted patients to answer daily self-management questions. It also allowed care managers

to follow-up with patients at highest calculated risk for hospitalization or adverse outcomes. It used a pre-post design with 1-year follow-up and was conducted in a Mexican border community. The study found a reduction in inpatient admission (-32% , $p<0.07$) and emergency room visits (-34% , $p<0.06$). Patients also had improvement in mental health quality of life ($p<0.03$) from baseline to follow-up. The home technology used a telephone line and was reported to be easy to use, but it required a partnership with a health care system that utilized an EMR. This study had a quality score of 10.5.

Shea et al. (2006, 2009) reported two randomized controlled trials that evaluated the effect of nurse guided telemedicine case management and treatment algorithms using a home telemedicine unit (HTU) on patient clinical outcomes in the Informatics for Diabetes Education and Telemedicine (IDEATel) Project. The IDEATel Project provided patients with an HTU which consisted of a Web-enabled computer with modem. The HTU allowed synchronous videoconferencing over standard telephone lines, electronic transmission of finger stick glucose and blood pressure readings, secure Web-based messaging with nurse case managers, and the ability to review clinical data and access Web-based education materials (Blanchet, 2008; Shea et al., 2006). Patients could take pictures of skin and feet and send those images to their nurse case managers. The study population included primarily African-American and Latino patients with diabetes who were Medicare recipients living in federally designated medically underserved areas of New York State.

The IDEATel project demonstrated significant decreases in HbA1c, blood pressure and LDL at 1- and 5-year follow-up (Shea et al., 2006, 2009). There were net reductions in HbA1c (-0.18% , $p=0.006$), systolic blood pressure (-3.4 mmHg, $p=0.001$), and LDL cholesterol (-9.5 mg, $p<0.001$) in the intervention group after 1 year of follow-up (Shea et al., 2006). After 5 years of follow-up, there were net reductions in HbA1c (0.29% , $p=0.001$), systolic blood pressure (-4.32 mmHg, $p=0.02$) and LDL (-3.84 mg/dL, $p<0.001$) in the intervention group compared to the control group, but there was no difference in mortality between groups (Shea et al., 2009). The 2006 and 2009 studies had quality scores of 21 and 20.5, respectively.

Although the Shea studies demonstrated improved diabetes outcomes, some challenges were noted. The HTU was expensive, and patients had difficulty learning how to use it. The intervention also required additional personnel for case management and to set up the technology in the clinic. There was a high attrition rate after 5 years (Shea et al., 2009). Cost analysis demonstrated that IDEATel did not reduce Medicare costs for health services; the cost per person per year was over \$8,000, which was excessive compared to other non-health IT programs that had similar clinical impacts (Moreno, Dale, Chen, & Magee, 2009).

These system oriented health IT interventions suggest that electronic registries and telemedicine may hold promise in improving diabetes outcomes. The telemedicine studies reduced health care utilization and improved diabetes clinical outcomes but were expensive and required additional personnel for case management.

Discussion

Our review provides an update on QI initiatives targeted at improving diabetes disparities and a review of health IT interventions to improve diabetes outcomes in minority patient populations. The 2006–2009 QI update presented many findings similar to studies published from 1985–2006, with some novel interventions. We found that QI projects utilizing cognitive-behavioral education, self-care management, and resilience training tended to be more successful in improving diabetes outcomes (Cramer et al., 2007; Mauldon et al., 2006; Steinhardt et al., 2009). Treatment algorithms for providers improved diabetes clinical

outcomes, though cultural competency training of providers did not lead to improvement in patient outcomes (Miller et al., 2006; Thom et al., 2006). The use of community health workers demonstrated improvements in diabetes intermediate outcomes, processes of care, knowledge, and social support and in outreach to patients who were on the fringe of the health care system (Babamoto et al., 2009; Ingram et al., 2007; Joshi et al., 2007; Lujan et al., 2007; Sixta & Ostwald 2008; Thompson et al., 2007). Nonphysician providers reduced diabetes-related urgent visits and emergency room visits (Davidson et al., 2007). From our review, we found mixed evidence from health IT studies designed to improve diabetes outcomes in minority patients.

Lessons Learned from Health IT Initiatives

In our review of diabetes QI initiatives using health IT, patient level interventions included personalized diabetes reports and computer-based multimedia diabetes education. These interventions reported mixed results. Personalized reports of diabetes values and goals can improve HbA1c, but a clinic needed to have an EMR already in use and a software algorithm to conduct this intervention (Levetan et al., 2002). Another study aimed to involve patients with lower health literacy and those with less computer experience, but it showed no improvements in diabetes outcomes, and it reported problems with computer malfunction (Gerber et al., 2005). Patient directed health IT interventions required sophisticated technology and support from case managers or other health care personnel and did not consistently improve diabetes outcomes (Gerber et al., 2005; Levetan et al., 2002).

We found few studies that tested provider oriented health IT interventions. The IPCAAD study tested the effect of clinical decision support systems for physicians in intensifying patient's medication regimens (Phillips et al., 2005; Ziemer et al., 2006). Medication intensification occurred more frequently when providers received in-person feedback regarding medication changes. The study also found improvements in HbA1c in patients who were in the intervention arm. These clinical reminders were more effective when the providers received in-person feedback from an endocrinologist, which may not be feasible to implement in all health care settings.

Registries facilitated tracking and analysis of data and improved diabetes outcomes in some studies. The Chin et al. (2004) and Bray, et al. (2005) studies demonstrate how registries can be used to track data and be integrated within more multi-targeted interventions to improve diabetes outcomes. Yet, the registries were found to be burdensome and duplicative of an EMR system.

Other systems oriented health IT interventions used telemedicine to bridge the gap between office visits for patients and demonstrated improvements in diabetes clinical outcomes (Cherry et al., 2002; Shea et al, 2006, 2009). From the patient perspective, the home technology units were difficult to use (Shea et al., 2006, 2009). Telemedicine interventions show promise, but the cost of the technology and need for additional support staff may overshadow the benefit.

Limitations of Current Work

Although we found many studies of the use of health IT to improve diabetes outcomes that were conducted in the past 9 years, there were many limitations to the literature. Given the heterogeneous designs and varying patient populations and settings, we could not compare effect sizes across studies; however the data may suggest trends in diabetes outcomes among racial/ethnic minority patients when using particular interventions. We found only a few health IT interventions that were tested in minority patient populations. We also did not find many studies that were conducted in under-resourced settings, such as public hospitals or

rural clinics. Study follow-up periods were generally short, usually about 12 months. A previous review of health IT have found similar limitations (Jackson, Bolen, Brancati, Batts-Turner, & Gary, 2006). Additionally, few studies reported cost data, thus making cost-effectiveness difficult to assess. Although we made an effort to search multiple databases and conduct hand searches, there may be diabetes QI literature that we did not identify. In addition, some organizations that have conducted interventions to improve diabetes care may not have published their findings. Our review was also limited by publication bias because positive findings tend to be published more often than negative findings.

Conclusion and Future Directions

Health care is rapidly becoming digital. New technologies may be harnessed to improve diabetes disparities, yet there are many gaps in the literature on the role of health IT in decreasing diabetes disparities. From our review of the literature, we identify opportunities for policy implementation and areas that need further research. Health IT policy should encourage: (1) the development of innovative health IT interventions that target diabetes processes and outcome measures; (2) rigorous evaluations of these interventions; and (3) support for under-resourced settings to strengthen their infrastructure to implement health IT.

Development of Targeted Health IT Interventions to Improve Diabetes Processes and Outcomes

Culturally tailored interventions—Culturally tailored, low literacy health IT interventions need to be developed to improve patient access to health information. Few studies have targeted non-English speakers and those with low literacy (Glasgow et al., 2005; Lorig, Ritter, & Laurent, 2006). People who speak and read little English or no English or have lower reading skills may face undue burden in using advanced technologies. Web sites and other health IT applications often assume a level of English language comprehension and a baseline computer skill level that may place immigrants, non-native English speakers, and those with limited literacy at a disadvantage. Designing health IT interventions that use voice recognition, television-based Internet, or touch screen systems may hold promise for these populations (Eng et al., 1998). Testing and modifying such technologies with these groups will be crucial to their application in reducing diabetes disparities (Gibbons, 2005).

Use of innovative technologies—Innovative technologies need to be integrated into health IT interventions. We found no studies that used mobile phones, social networking sites, YouTube, or Twitter to improve access or improve diabetes outcomes among minority populations or in under-resourced settings. We found no studies that utilized health IT integrated into community health worker programs to address diabetes disparities. Health information technologies may be used to increase social support among patients, encourage behavioral changes, and enhance community health worker programs.

Communication and care coordination—Health IT can be used to enhance provider-to-provider communication and improve care coordination. Health IT may be used to alleviate the fragmentation of health care for minority patients in under-resourced settings through improved communication between providers and access to subspecialists. Although we found many examples of telemedicine and e-referrals that extended the reach of subspecialists, few studies were conducted in minority patient populations (Kim et al., 2009; Shea et al., 2009). More studies are needed that aim to enhance communication across providers and subspecialists in addressing poor subspecialty access for patients and improving continuity of care among patients who may have fragmented care.

Cultural competency training—Health IT interventions that address patient-provider communication and cultural competency training need to be developed. Health IT interventions may be used in under-resourced settings to facilitate communication with culturally diverse patient populations in innovative ways. We found no studies that used health IT applications to enhance patient communication with providers, facilitated shared decision-making, or emphasized culturally competent interactions with patients.

Patient centered behavioral interventions—Health IT innovations can be used to design novel behavioral interventions and enhance current patient centered interventions. In under-resourced settings, patients may not have access to diabetes behavioral management programs. Through the use of health IT innovations—such as mobile phone technologies, computerized simulations, and interactive Web-based programs—individuals with diabetes may be able to monitor their physical activity, weight loss, and other physiological markers and receive prompts to encourage self-management. Other health IT innovations, such as personalized health records (PHRs) and social network applications, may enhance access to providers, provide consistency in care, empower patients, and provide peer support (Barrera, Glasgow, McKay, Boles, & Feli, 2002; Zrebiec, 2005). PHRs that are portable and interoperable with EMRs can be of great benefit to patients in under-resourced settings who may have fragmented care. There is also a need for more low literacy, culturally relevant patient applications. PHRs may benefit disadvantaged populations by enabling consistent care, electronic access to information on health conditions, and enhanced communication with primary care providers (Gibbons, 2005; Ueckert, Goerz, Ataian, Tessmann, & Prokosch, 2003). Many applications exist for PHRs, but they have not been tested in minority patient populations or under-resourced settings (Weitzman, Kaci, & Mandl, 2009; see also MiVIA at <https://www.mivia.org>).

Enhanced data tracking—There needs to be further development and testing of interventions that enhance data tracking and that are interoperable between many health systems. Regional HIEs are developing, but we found no studies that tested the impact of these systems in decreasing diabetes disparities. Health systems to track and report data on race/ethnicity and other social determinants of health are a basic need. (Ulmer, McFadden, & Nerenz, 2009). These systems can also facilitate disease tracking, allow quicker access to patient information, and improve continuity of care. Designing interventions that are interoperable between many health systems is crucial. Interoperable systems can enhance sharing of data among providers, laboratories, and health systems.

Rigorous Evaluations of Current Health IT Interventions and Their Impact on Diabetes Disparities

Effectiveness of health IT interventions and cost data—More studies are needed that examine the effectiveness of health IT interventions in under-resourced settings that serve minority patients. We need to understand the feasibility, acceptability, and effectiveness of these technological interventions for reducing disparities in diabetes. More studies are also needed to examine implementation cost and the cost-effectiveness of these interventions, especially in under-resourced settings where financial barriers may inhibit the use of health IT innovations. Understanding the true costs of implementation and any savings over time will allow better planning for under-resourced settings to implement health IT (Goldzweig, Towfigh, Maglione, & Shekelle, 2009; Harris, Haneuse, Martin, & Ralston, 2009; Huang et al., 2007; Moreno et al., 2009; Welch et al., 2007). Although financial incentives are necessary to prevent safety net clinics from lagging in health IT implementation, analyses of long term costs are also crucial to better understand the payoff of the initial investment in health IT for these settings. Longer follow up may also give better insight into the effect of health IT interventions because fine-tuning is necessary in the

initial set-up phases. Additionally, more studies are needed that measure reduction in diabetes disparities across populations in addition to improvement of diabetes outcomes in minority patients.

Testing of EMR capabilities—Studies are needed to test the ability of EMRs to track and monitor patients, provide feedback, assess medication adherence, and improve diabetes outcomes. EMRs can provide the capacity for computerized provider order entry and clinical decision support (Aarts & Koppel, 2009). EMRs can also automate the collection and movement of information that would enhance patient care and monitor quality of diabetes care. Tracking race data allows safety net clinics to identify trends among their patients and may facilitate recruitment of minority patients who are underrepresented in clinical trials. The studies we found using EMRs in minority patients did not fully use all of the EMRs' components to assess diabetes outcomes.

Multi-targeted intervention evaluation—More studies are needed that assess multi-targeted health IT interventions. Also, more studies are needed that integrate all aspects of health IT and harness their potential. Through the utilization of all the aspects of health IT—such as EMR, clinical decision support, computerized order entry, and interoperability with regional HIEs—we can understand how to fully utilize health IT innovations to reduce diabetes disparities.

Under-Resourced Settings Need Support to Build Infrastructure to Implement HIT Interventions

Multi-payer projects—Multi-payer pilot projects and financial incentives are needed to encourage health IT implementation in under-resourced settings that may face many financial barriers in implementing innovative health IT applications. These barriers may be overcome by providing external financial incentives, demonstration grants, and collaborations (Miller & Sim, 2004; Simon, Rundall, & Shortell, 2007). The American Recovery and Reinvestment Act (2009) provides financial incentives for providers and hospitals to use certified EMRs and subsidies for physicians with high volumes of Medicaid patients (Blumenthal, 2009). Other incentives may be necessary for under-resourced settings to implement EMRs (Blumenthal, 2009). Non-visit care, such as phone communication and electronic mail, is not generally reimbursed, so strong incentives exist for providers to delay EMR implementation. Rewarding practices for publishing performance reports and mandating specific quality improvement tactics or IT applications could hasten implementation (Miller & Sim, 2004; Simon, Rundall, & Shortell 2007). Financial resources need to be available for under-resourced settings to partner with local organizations, businesses, and government organizations to install EMRs, participate in regional HIEs, and use innovative health IT applications to improve patient outcomes (Goroll, Simon, Tripathi, Ascenzo, & Bates, 2009; McDonald et al., 2005; Mostashari, Tripathi, & Kendall, 2009). Pilot grants to fund implementation projects and other financial incentives, such as payment for improved performance and for non-visit care, will be critical in under- resourced settings implementing QI interventions using health IT.

Assessing human capital needs—It is important to assess human capital needs to implement and maintain health IT interventions in under-resourced settings. Although some of the studies we reviewed commented on the need for nursing staff, case managers, or staff to assist in maintaining health IT interventions, no studies reported the type of staff and number of staff needed to support health IT interventions in under-resourced settings (Bray, Thompson et al., 2005; Chin et al., 2007). Studies in EMR process measurement have noted the importance of a health IT champion, yet this concept still needs testing in under-resourced settings and in addressing diabetes (Zandieh et al., 2008). Chronic disease

management requires a team of physicians, nurses, case managers, and other allied health professionals. Hiring nurse educators, nurse case managers, and case managers to triage clinical decision alerts or receive alerts on missed appointments or elevated laboratory values may be one way to decrease the burden of “reminder fatigue” for providers (Dixon & Samarth, 2009). Clinical reminders can provide better information at the time of clinical decision-making for individual patients, but these reminders need to be supported by other personnel.

Best practices for under-resourced settings—Best practices in how community health centers and nonprofit organizations can partner with for-profit organizations and state and federal governments to install health IT are needed. Demonstration projects and evaluations can identify critical implementation lessons and disseminate best practices. Descriptions of processes undertaken to encourage adoption of health IT technologies and how to integrate them into current work flow are necessary to plan and execute the implementation of health IT innovations in under-resourced settings. Actively engaging providers in deciding what functions, clinical reminders, and alerts are most valuable is essential for integrating EMRs into the current infrastructure and avoiding provider fatigue and resistance to the EMR (Ko et al., 2007).

Technological support—Technological support needs to be implemented for under-resourced settings because these organizations may not have adequate personnel trained in health IT or a physician IT champion who can ensure a smooth transition to EMR. The American Recovery and Reinvestment Act of 2009 described establishment of regional centers to assist providers seeking to adopt and become meaningful users of health IT. These centers will be key in offering technologic support to rural and other under-resourced health care settings in the implementation and maintenance of health IT infrastructure (Centers for Medicare & Medicaid Services, 2009).

The rapid advancement of health IT offers unprecedented opportunities for policymakers, clinicians, and health systems to address disparities in diabetes care and outcomes. Integrating innovations in health IT with QI initiatives is likely to be a powerful way to reduce diabetes disparities.

Acknowledgments

Funding for this report was provided by the Agency for Healthcare Research and Quality (AHRQ) through contract P233200900421P. In addition, this project was completed with support from AHRQ through contract HHSN263200500063293B. We thank Jessica Watkins for her assistance in retrieving articles for the review. Dr. Baig's effort was also supported by a National Center on Minority Health and Health Disparities Loan Repayment Grant. Drs. Baig, Chin, and Huang, and Ms. Wilkes were supported by the National Institutes of Diabetes and Digestive and Kidney Diseases (NIDDK) Diabetes Research and Training Center (P60 DK20595). Dr. Peek is supported by career development awards from the Robert Wood Johnson Foundation (RWJF) Harold Amos Medical Faculty Development program and the NIDDK (K23 DK075006-01). Dr. Chin is supported by a Midcareer Investigator Award in Patient-Oriented Research from the NIDDK (K24 DK071933).

References

- Aarts J, Koppel RL. Implementation of computerized physician order entry in seven countries. *Health Affairs (Millwood)*. 2009; 28(2):404–414.
- Agency for Healthcare Research and Quality. Rockville, MD: U.S. Department of Health and Human Services, Agency for Healthcare Research and Quality; 2008. National healthcare disparities report 2007.
- Agency for Healthcare Research and Quality. Rockville, MD: U.S. Department of Health and Human Services, Agency for Healthcare Research and Quality; 2009. National healthcare disparities report 2008.

- American Diabetes Association. Standards of medical care in diabetes--2008. *Diabetes Care*. 2008; 31 Suppl 1:12–54.
- [Accessed April 14, 2010] American Recovery and Reinvestment Act of 2009 (Public Law 111-5). Available at http://frwebgate.access.gpo.gov/cgi-bin/getdoc.cgi?dbname=111_cong_bills&docid=f:h1enr.pdf
- Babamoto KS, Sey KA, Camilleri AJ, Karlan VJ, Catalasan J, Morisky DE. Improving diabetes care and health measures among Hispanics using community health workers: Results from a randomized controlled trial. *Health Education & Behavior*. 2009; 36(1):113–126. [PubMed: 19188371]
- Bach PB, Pham HH, Schrag D, Tate RC, Hargraves JL. Primary care physicians who treat blacks and whites. *New England Journal of Medicine*. 2004; 351(6):575–584. [PubMed: 15295050]
- Barrera M Jr, Glasgow RE, McKay HG, Boles SM, Feil EG. Do Internet-based support interventions change perceptions of social support?: An experimental trial of approaches for supporting diabetes self-management. *American Journal of Community Psychology*. 2002; 30(5):637–654. [PubMed: 12188054]
- Berwick DM. A user's manual for the IOM's 'Quality Chasm' report. *Health Affairs (Millwood)*. 2002; 21(3):80–90.
- Blanchet KD. Telehealth and diabetes monitoring. *Telemedicine and E-Health*. 2008; 14(8):744–746. [PubMed: 18954243]
- Blumenthal D. Stimulating the adoption of health information technology. *West Virginia Medical Journal*. 2009; 105(3):28–29. [PubMed: 19456037]
- Bray P, Roupe M, Young S, Harrell J, Cummings DM, Whetstone LM. Feasibility and effectiveness of system redesign for diabetes care management in rural areas: The eastern North Carolina experience. *The Diabetes Educator*. 2005; 31(5):712–718. [PubMed: 16203855]
- Bray P, Thompson D, Wynn JD, Cummings DM, Whetstone L. Confronting disparities in diabetes care: The clinical effectiveness of redesigning care management for minority patients in rural primary care practices. *Journal of Rural Health*. 2005; 21(4):317–321. [PubMed: 16294654]
- Centers for Disease Control and Prevention. Atlanta, GA: Author; 2008. National diabetes fact sheet: General information and national estimates on diabetes in the United States, 2007.
- Centers for Medicare & Medicaid Services. [cited 5 April 2010] Fact sheets: Details For: Medicare and Medicaid Health Information Technology: Title IV of the American Recovery and Reinvestment Act 2009. 2009. Available from <http://www.cms.hhs.gov/apps/media/press/factsheet.asp?Counter=3466&intNumPerPage>
- Chaudhry B, Wang J, Wu S, Maglione M, Mojica W, Roth E, et al. Systematic review: Impact of health information technology on quality, efficiency, and costs of medical care. *Annals of Internal Medicine*. 2006; 144(10):742–752. [PubMed: 16702590]
- Cherry JC, Moffatt TP, Rodriguez C, Dryden K. Diabetes disease management program for an indigent population empowered by telemedicine technology. *Diabetes Technology & Therapeutics*. 2002; 4(6):783–791.
- Chin MH, Cook S, Drum ML, Jin L, Guillen M, Humikowski CA, et al. Improving diabetes care in midwest community health centers with the health disparities collaborative. *Diabetes Care*. 2004; 27(1):2–8. [PubMed: 14693957]
- Chin MH, Drum ML, Guillen M, Rimington A, Levie JR, Kirchhoff AC, et al. Improving and sustaining diabetes care in community health centers with the health disparities collaboratives. *Medical Care*. 2007; 45(12):1135–1143. [PubMed: 18007163]
- Collins TC, Beyth RJ, Nelson DB, Petersen NJ, Suarez-Almazor ME, Bush RL, et al. Process of care and outcomes in patients with peripheral arterial disease. *Journal of General Internal Medicine*. 2007; 22(7):942–948. [PubMed: 17453264]
- Cramer J, Steven R, Sibley RF, Donald P, Bartlett L, Kahn S, et al. An adaptation of the diabetes prevention program for use with high-risk, minority patients with type 2 diabetes. *The Diabetes Educator*. 2007; 33(3):503–508. [PubMed: 17570881]
- Davidson MB, Ansari A, Karlan VJ. Effect of a nurse-directed diabetes disease management program on urgent care/emergency room visits and hospitalizations in a minority population. *Diabetes Care*. 2007; 30(2):224–227. [PubMed: 17259485]

- Davis K, Doty MM, Shea K, Stremikis K. Health information technology and physician perceptions of quality of care and satisfaction. *Health Policy*. 2009; 90(2–3):239–246. [PubMed: 19038472]
- DesRoches CM, Campbell EG, Rao SR, Donelan K, Ferris TG, Jha A, et al. Electronic health records in ambulatory care. A national survey of physicians. *New England Journal of Medicine*. 2008; 359(1):50–60. [PubMed: 18565855]
- Dillingham TR, Pezzin LE, Mackenzie EJ. Racial differences in the incidence of limb loss secondary to peripheral vascular disease: A population-based study. *Archives of Physical Medicine and Rehabilitation*. 2002; 83(9):1252–1257. [PubMed: 12235605]
- Dixon, BE.; Samarth, A. Findings from the AHRQ Health IT Portfolio. Rockville, MD: Agency for Healthcare Research and Quality; 2009. Innovations in using health IT for chronic disease management. (AHRQ Publication No. 09-0029-EF) Available at http://healthit.ahrq.gov/portal/server.pt/gateway/PTARGS_0_3882_840586_0_0_18/09-0029-EF_cdm.pdf
- Dorr D, Bonner LM, Cohen AN, Shoai RS, Perrin R, Chaney E, et al. Informatics systems to promote improved care for chronic illness: A literature review. *Journal of the American Medical Informatics Association*. 2007; 14(2):156–163. [PubMed: 17213491]
- Downs SH, Black N. The feasibility of creating a checklist for the assessment of the methodological quality both of randomized and non-randomized studies of health care interventions. *Journal of Epidemiology and Community Health*. 1998; 52(6):377–384. [PubMed: 9764259]
- Duru OK, Gerzoff RB, Selby JV, Brown AF, Ackermann RT, Karter AJ, et al. Identifying risk factors for racial disparities in diabetes outcomes: The Translating Research into Action for Diabetes Study. *Medical Care*. 2009; 47(6):700–706. [PubMed: 19480090]
- Eng TR, Maxfield A, Patrick K, Deering MJ, Ratzan SC, Gustafson DH. Access to health information and support. A public highway or a private road? *Journal of the American Medical Association*. 1998; 280(15):1371–1375. [PubMed: 9794322]
- Eysenbach G. Medicine 2.0: Social networking, collaboration, participation, apomediation, and openness. *Journal of Medical Internet Research*. 2008; 10(3):e22. [PubMed: 18725354]
- Foxhall, K. [cited 22 April 2010] Health IT stimulus bill zips to House floor. *Government Health IT*. 2009 23 1. 2009 Available from <http://www.govhealthit.com/newsitem.aspx?tid=79&nid=69532>
- Gerber BS, Brodsky IG, Lawless KA, Smolin LI, Arozullah AM, Smith EV, et al. Implementation and evaluation of a low-literacy diabetes education computer multimedia application. *Diabetes Care*. 2005; 28(7):1574–1580. [PubMed: 15983303]
- Gibbons MC. A historical overview of health disparities and the potential of eHealth solutions. *Journal of Medical Internet Research*. 2005; 7(5):e50. [PubMed: 16403714]
- Glasgow RE, Nutting PA, King DK, Nelson CC, Cutter G, Gaglio B, et al. Randomized effectiveness trial of a computer-assisted intervention to improve diabetes care. *Diabetes Care*. 2005; 28(1):33–39. [PubMed: 15616230]
- Gold R, Yu K, Liang LJ, Adler F, Balingit P, Luc P, et al. Synchronous provider visit and self-management education improves glycemic control in Hispanic patients with long-standing type 2 diabetes. *The Diabetes Educator*. 2008; 34(6):990–995. [PubMed: 18849465]
- Goldzweig CL, Towfigh A, Maglione M, Shekelle PG. Costs and benefits of health information technology: New trends from the literature. *Health Affairs (Millwood)*. 2009; 28(2):w282–w293.
- Goroll AH, Simon SR, Tripathi M, Ascenzo C, Bates DW. Community-wide implementation of health information technology: The Massachusetts eHealth Collaborative experience. *Journal of the American Medical Informatics Association*. 2009; 16(1):132–139. [PubMed: 18952937]
- Halamka JD, Mandl KD, Tang PC. Early experiences with personal health records. *Journal of the American Medical Informatics Association*. 2008; 15(1):1–7. [PubMed: 17947615]
- Harris LT, Haneuse SJ, Martin DP, Ralston JD. Diabetes quality of care and outpatient utilization associated with electronic patient-provider messaging: A cross-sectional analysis. *Diabetes Care*. 2009; 32(7):1182–1187. [PubMed: 19366959]
- Huang ES, Zhang Q, Brown SE, Drum ML, Meltzer DO, Chin MH. The cost-effectiveness of improving diabetes care in U.S. federally qualified community health centers. *Health Services Research*. 2007; 42(6 Pt 1):2174–2193. discussion 2294–2323. [PubMed: 17995559]

- Ingram M, Torres E, Redondo F, Bradford G, Wang C, O'Toole ML. The impact of promotoras on social support and glycemic control among members of a farmworker community on the US-Mexico border. *The Diabetes Educator*. 2007; 33 Suppl 6:172S–178S. [PubMed: 17620398]
- Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33). UK Prospective Diabetes Study (UKPDS) Group. *Lancet*. 1998; 352(9131):837–853. [PubMed: 9742976]
- Jackson CL, Bolen S, Brancati FL, Batts-Turner ML, Gary TL. A systematic review of interactive computer-assisted technology in diabetes care. *Interactive information technology in diabetes care. Journal of General Internal Medicine*. 2006; 21(2):105–110. [PubMed: 16390512]
- Jha AK, DesRoches CM, Campbell EG. Use of electronic health records in U.S. hospitals. *New England Journal of Medicine*. 2009; 360(16):1628–1638. [PubMed: 19321858]
- Jha AK, Orav EJ, Li Z, Epstein AM. Concentration and quality of hospitals that care for elderly black patients. *Archives of Internal Medicine*. 2007; 167(11):1177–1182. [PubMed: 17563027]
- Jha AK, Orav EJ, Zheng J, Epstein AM. The characteristics and performance of hospitals that care for elderly Hispanic Americans. *Health Affairs (Millwood)*. 2008; 27(2):528–537.
- Joshu CE, Rangel L, Garcia O, Brownson CA, O'Toole ML. Integration of a promotor-led self-management program into a system of care. *The Diabetes Educator*. 2007; 33 Suppl 6:151S–158S. [PubMed: 17620395]
- Katz SJ, Nissan N, Moyer CA. Crossing the digital divide: Evaluating online communication between patients and their providers. *American Journal of Managed Care*. 2004; 10(9):593–598. [PubMed: 15515991]
- Kim Y, Chen AH, Keith E, Yee HF, Krushei MB. Not perfect, but better: Primary care providers' experiences with electronic referrals in a safety net health system. *Journal of General Internal Medicine*. 2009; 24(5):614–619. [PubMed: 19308334]
- King A, Wolfe G, Healy S, Lineham T, Harrison S, Ponzio C, et al. Exploring methods to extend the reach of diabetes specialist expertise into primary care clinics. *The Diabetes Educator*. 2006; 32(4):525–530. [PubMed: 16873590]
- Kirk JK, Ralph B, Bell RA, Passmore LV, Bonds DE. Disparities in HbA1c levels between African-American and non-Hispanic white adults with diabetes. *Diabetes Care*. 2006; 29(9):2130–2136. [PubMed: 16936167]
- Ko Y, Abarca J, Malone DC, Dare DC, Geraets D, Houranieh A, et al. Practitioners' views on computerized drug-drug interaction alerts in the VA system. *Journal of the American Medical Informatics Association*. 2007; 14(1):56–64. [PubMed: 17068346]
- Koro CE, Bowlin SJ, Bourgeois N, Fedder DO. Glycemic control from 1988 to 2000 among U.S. adults diagnosed with type 2 diabetes: A preliminary report. *Diabetes Care*. 2004; 27(1):17–20. [PubMed: 14693960]
- Landon BE, Hicks LS, O'Malley AJ, Lieu TA, Keegan T, McNeil BJ, et al. Improving the management of chronic disease at community health centers. *New England Journal of Medicine*. 2007; 356(9):921–934. [PubMed: 17329699]
- Levetan CS, Dawn KR, Robbins DC, Ratner RE. Impact of computer-generated personalized goals on HbA(1c). *Diabetes Care*. 2002; 25(1):2–8. [PubMed: 11772893]
- Liebman J, Heffernan D, Sarvela P. Establishing diabetes self-management in a community health center serving low-income Latinos. *The Diabetes Educator*. 2007; 33 Suppl 6:132S–138S. [PubMed: 17620392]
- Lorig KR, Ritter PH, Laurent DD. Internet-based chronic disease self-management: A randomized trial. *Medical Care*. 2006; 44(11):964–971. [PubMed: 17063127]
- Lujan J, Ostwald SK, Ortiz M. Promotora diabetes intervention for Mexican Americans. *The Diabetes Educator*. 2007; 33(4):660–670. [PubMed: 17684167]
- Mahotiery T, Oceppek-Welikson K, Daley MB, Byssainthe JP. A program to reduce the disparity in the rate of biennial lipid profiles between African-American and white Medicare beneficiaries with diabetes mellitus in New York City. *Journal of Community Health*. 2006; 31(4):263–288. [PubMed: 16894826]

- Mauldon M, Melkus GD, Cagganello M. Tomando Control: A culturally appropriate diabetes education program for Spanish-speaking individuals with type 2 diabetes mellitus. Evaluation of a pilot project. *The Diabetes Educator*. 2006; 32(5):751–760. [PubMed: 16971708]
- McDonald CJ, Overhage JM, Barnes M, Schadow G, Blevins L, Dexter PR, et al. The Indiana network for patient care: A working local health information infrastructure. An example of a working infrastructure collaboration that links data from five health systems and hundreds of millions of entries. *Health Affairs (Millwood)*. 2005; 24(5):1214–1220.
- Miller CD, Ziemer DC, Kolm P, El-Kebbi IM, Cook CB, Gallina DL, et al. Use of a glucose algorithm to direct diabetes therapy improves A1C outcomes and defines an approach to assess provider behavior. *The Diabetes Educator*. 2006; 32(4):533–545. [PubMed: 16873591]
- Miller RH, Sim I. Physicians' use of electronic medical records: Barriers and solutions. *Health Affairs (Millwood)*. 2004; 23(2):116–126.
- Moreno L, Dale SB, Chen AY, Magee CA. Costs to Medicare of the Informatics for Diabetes Education and Telemedicine (IDEATel) home telemedicine demonstration: Findings from an independent evaluation. *Diabetes Care*. 2009; 32(7):1202–1204. [PubMed: 19366971]
- Morrissey NJ, Giacobelli J, Egorova N, Gelijns A, Moskowitz A, McKinsey J, et al. Disparities in the treatment and outcomes of vascular disease in Hispanic patients. *Journal of Vascular Surgery*. 2007; 46(5):971–978. [PubMed: 17980283]
- Mostashari F, Tripathi M, Kendall M. A tale of two large community electronic health record extension projects. *Health Affairs (Millwood)*. 2009; 28(2):345–356.
- Narayan KM, Boyle JP, Geiss LS, Saaddine JB, Thompson TJ. Impact of recent increase in incidence on future diabetes burden: U.S., 2005–2050. *Diabetes Care*. 2006; 29(9):2114–2116. [PubMed: 16936162]
- National Institute of Diabetes and Digestive and Kidney Diseases. Bethesda, MD: Author. U.S. Department of Health and Human Services, National Institutes of Health; 2008. National diabetes statistics, 2007 fact sheet.
- Peek ME, Cargill A, Huang ES. Diabetes health disparities: A systematic review of health care interventions. *Medical Care Research and Review*. 2007; 64 5 Suppl:101–156.
- Phillips LS, Ziemer DC, Doyle JP, Barnes CS, Kolm P, Branch WT, et al. An endocrinologist-supported intervention aimed at providers improves diabetes management in a primary care site: Improving primary care of African Americans with diabetes (IPCAAD) 7. *Diabetes Care*. 2005; 28(10):2352–2360. [PubMed: 16186262]
- Regenbogen SE, Gawande AA, Lipsitz SR, Greenberg CC, Jha AK. Do differences in hospital and surgeon quality explain racial disparities in lower-extremity vascular amputations? *Annals of Surgery*. 2009; 250(3):424–431. [PubMed: 19652590]
- Reichsman A, Warner JJ, Cella P. Opportunities for improved diabetes care among patients of safety net practices: A safety net providers' strategic alliance study. *Journal of the National Medical Association*. 2009; 101(1):4–11. [PubMed: 19245066]
- Rosal ME, Benjamin E, Pekow P. Opportunities and challenges for diabetes prevention at two community health centers. *Diabetes Care*. 2008; 31(2):247–254. [PubMed: 17989311]
- Rosamond W, Flegal K, Friday G, Furie K, Go A, Greenlund K, et al. Heart disease and stroke statistics--2007 update: A report from the American Heart Association Statistics Committee and Stroke Statistics Subcommittee. *Circulation*. 2007; 115(5):e69–e171. [PubMed: 17194875]
- Saydah S, Cowie C, Eberhardt MS, De Rekeneire N, Narayan KM. Race and ethnic differences in glycemic control among adults with diagnosed diabetes in the United States. *Ethnicity & Diabetes*. 2007; 17(3):529–535.
- Schade CP, Hannah KL. Quality of ambulatory care for diabetes and lower-extremity amputation. *American Journal of Medical Quality*. 2007; 22(6):410–417. [PubMed: 18006421]
- Selvin E, Wattanakit K, Steffes MW, Coresh J, Sharrett AR. HbA1c and peripheral arterial disease in diabetes: The Atherosclerosis Risk in Communities study. *Diabetes Care*. 2006; 29(4):877–882. [PubMed: 16567831]
- Shea S, Weinstock RS, Starren J, Teresi J, Palmas W, Field L, et al. A randomized trial comparing telemedicine case management with usual care in older, ethnically diverse, medically underserved

- patients with diabetes mellitus. *Journal of the American Medical Informatics Association*. 2006; 13(1):40–51. [PubMed: 16221935]
- Shea S, Weinstock RS, Teresi JA, Palmas W, Starren J, Cimino JJ, et al. A randomized trial comparing telemedicine case management with usual care in older, ethnically diverse, medically underserved patients with diabetes mellitus: 5 year results of the IDEATel study. *Journal of the American Medical Informatics Association*. 2009; 16(4):446–456. [PubMed: 19390093]
- Shields AE, Shin P, Leu MG, Levy DE, Betancourt RM, Hawkins D, et al. Adoption of health information technology in community health centers: Results of a national survey. *Health Affairs*. 2007; 26(5):1373–1383. [PubMed: 17848448]
- Simon JS, Rundall TG, Shortell SM. Adoption of order entry with decision support for chronic care by physician organizations. *Journal of the American Medical Informatics Association*. 2007; 14(4): 432–439. [PubMed: 17460136]
- Sixta CS, Ostwald S. Texas-Mexico border intervention by promotores for patients with type 2 diabetes. *The Diabetes Educator*. 2008; 34(2):299–309. [PubMed: 18375779]
- Steinhardt MA, Mamerow MM, Brown SA, Jolly CA. A resilience intervention in African American adults with type 2 diabetes: A pilot study of efficacy. *The Diabetes Educator*. 2009; 35(2):274–284. [PubMed: 19204102]
- Thom DH, Tirado MD, Woon TL, McBride MR. Development and evaluation of a cultural competency training curriculum. *BioMed Central Medical Education*. 2006; 6:38. [PubMed: 16872504]
- Thompson JR, Horton C, Flores C. Advancing diabetes self-management in the Mexican American population: A community health worker model in a primary care setting. *The Diabetes Educator*. 2007; 33 Suppl 6:159S–165S. [PubMed: 17620396]
- Tight blood pressure control and risk of macrovascular and microvascular complications in type 2 diabetes: UKPDS 38. UK Prospective Diabetes Study Group. *British Medical Journal*. 1998; 317(7160):703–713. [PubMed: 9732337]
- Ueckert F, Goerz M, Ataian M, Tessmann S, Prokosch HU. Empowerment of patients and communication with health care professionals through an electronic health record. *International Journal of Medical Informatics*. 2003; 70(2–3):99–108. [PubMed: 12909161]
- Ulmer, C.; McFadden, B.; Nerenz, DR., editors. Subcommittee on Standardized Collection of Race/Ethnicity Data for Healthcare Quality Improvement, Board on Health Care Services, Institute of Medicine. Washington, DC: National Academies Press; 2009. Race, ethnicity, and language data: Standardization for health care quality improvement.
- Utz SW, Williams IC, Jones R, Hinton I, Alexander G, Yan G, et al. Culturally tailored intervention for rural African Americans with type 2 diabetes. *Diabetes Educator*. 2008; 34(5):854–865. [PubMed: 18832290]
- Wagner H, Pizzimenti JJ, Daniel K, Pandya N, Hardigan PC. Eye on diabetes: A multidisciplinary patient education intervention. *Diabetes Educator*. 2008; 34(1):84–89. [PubMed: 18267994]
- Wakefield BJ, Ward MM, Holman JE, Ray A, Scherubel M, Burns TL, et al. Evaluation of home telehealth following hospitalization for heart failure: A randomized trial. *Telemedicine Journal and E-Health*. 2008; 14(8):753–761. [PubMed: 18954244]
- Weitzman ER, Kaci L, Mandl KD. Acceptability of a personally controlled health record in a community-based setting: Implications for policy and design. *Journal of Medical Internet Research*. 2009; 11(2):e14. [PubMed: 19403467]
- Welch WP, Bazarko D, Ritten K, Burgess Y, Harmon R, Sandy LG. Electronic health records in four community physician practices: Impact on quality and cost of care. *Journal of the American Medical Informatics Association*. 2007; 14(3):320–328. [PubMed: 17329734]
- Wong MD, Martin F, Shapiro W, Boscardin J, Ettner SL. Contribution of major diseases to disparities in mortality. *New England Journal of Medicine*. 2002; 347(20):1585–1592. [PubMed: 12432046]
- Zandieh SO, Yoon-Flannery K, Kuperman GJ, Langsam DJ, Hyman D, Kaushal R. Challenges to EHR implementation in electronic- versus paper-based office practices. *Journal of General Internal Medicine*. 2008; 23(6):755–761. [PubMed: 18369679]
- Zaza S, Wright-De Agüero LK, Briss PA, Truman PI, Hopkins DP, Hennessy MH, et al. Data collection instrument and procedure for systematic reviews in the Guide to Community Preventive

- Services Task Force on Community Preventive Services. American Journal of Preventive Medicine. 2000; 18 1 Suppl:44–74. [PubMed: 10806979]
- Ziemer DC, Doyle JP, Barnes CS, Branch WT Jr, Cook CB, El-Kebbi IM, et al. An intervention to overcome clinical inertia and improve diabetes mellitus control in a primary care setting: Improving primary care of African Americans with diabetes (IPCAAD) 8. Archives of Internal Medicine. 2006; 166(5):507–513. [PubMed: 16534036]
- Zrebiec JF. Internet communities: Do they improve coping with diabetes? The Diabetes Educator. 2005; 31(6):825–836. [PubMed: 16288090]

Table 1

Overview of Health Care System Quality Improvement Initiatives in Improving Diabetes Disparities

Level	Characteristics of QI that are associated with improved diabetes outcomes	Evidence exists to support these outcomes can be impacted			Limitations and challenges
Patient	<ul style="list-style-type: none">Utilize interpersonal (rather than computer-based) skillsUse social networks (family members, peer support groups, one-on-one interactive education, community health workers)Use culturally tailored interventionsEmphasize cognitive-behavioral education, self-care management, and adaptations of the Diabetes Prevention Program (DPP)Focus on improving patient resilience to stressorsProvide multidisciplinary patient diabetes education	<ul style="list-style-type: none">Lower HbA1cDietary habitsPhysical activityWeight changeDepressive symptoms (short-term)Physician trustHealth knowledge	<ul style="list-style-type: none">Health behaviorsDiabetes self-empowermentDiabetes self-managementLow density lipidsSystolic blood pressureIncreased diabetes knowledge	<ul style="list-style-type: none">Need broader expansion into racial/ethnic populationsLack of interventions targeting patients on the fringe of the health system	
Provider	<ul style="list-style-type: none">Use of in-person feedback instead of computerized decision-support to sustain provider behavior change and health outcomesEnsure timely availability of treatment algorithms factoring in HbA1c and random blood sugar at patient appointments	<ul style="list-style-type: none">Glycemic controlProcess measures: eye examinations, microalbumin testing, HbA1c monitoring, foot care, exercise counseling, lipid testing, influenza vaccines, nutrition educationHypertension and cholesterolProvider intensification of glucose managementMedication intensification leading to improved HbA1c		<ul style="list-style-type: none">Lack of intensive provider training in diabetes careLack of provider communication or shared decision making trainingComputer reminders can be easily ignoredSustained change may require the use of continued medical education and	

Level	Characteristics of QI that are associated with improved diabetes outcomes	Evidence exists to support these outcomes can be impacted			Limitations and challenges
					practice guidelines
Healthcare Organization	<ul style="list-style-type: none">Use of case management and community health workers (CHWs) and non-physician providers: RN/CHW can act as a patient adjunct to primary care team to assist with case management and overcome social, cultural, and linguistic barriers; act as powerful change agents	<ul style="list-style-type: none">Keeping appointments with physiciansHbA1c testingGlycemic controlBlood pressureDyslipidemiaDelayed onset of retinopathyDecreased hospitalizations	<ul style="list-style-type: none">Pharmacist-led medication management: HbA1c levels, increased aspirin use, lowered systolic blood pressure, diabetes knowledge, patient satisfactionLow density lipidsAnnual foot examinationsDiabetes knowledge	<ul style="list-style-type: none">Lack of partnering with communitiesLack of support for safety net providers caring for disproportionate share of racial/ethnic minoritiesLack of collaboration with national programsPharmacist-directed care is labor intensivePhysician resistanceLimited insurance reimbursementLack of Internet access and dependence on fax and telephone communication	
	<ul style="list-style-type: none">Use of medical (or medication) assistance programs (MAPs)	<ul style="list-style-type: none">Diabetes and dyslipidemia controlNumber of patients on medication for dyslipidemia and hypertension	<ul style="list-style-type: none">Increase prescription adherenceDecrease hospitalizations, and reduce hospital costsHbA1c, LDL, triglyceride levels	<ul style="list-style-type: none">Only used in pharmaceutical company-sponsored medications	
		<ul style="list-style-type: none">Use of disease management systems <i>Elements:</i> 1) identification of population with disease (diabetes registry); 2) guidelines for performance standards for care; 3) management of identified people; 4)	<ul style="list-style-type: none">Better diabetes controlBetter processes of care: increased screening for neuropathy, dyslipidemia, and microalbuminuria		<ul style="list-style-type: none">No reviewed studies had all four system components

Level	Characteristics of QI that are associated with improved diabetes outcomes	Evidence exists to support these outcomes can be impacted	Limitations and challenges
	tracking and monitoring		
Multi-target Interventions	<ul style="list-style-type: none"> • Use of multidisciplinary teams • Ensure that health care and academic institutions work as part of a broad community coalition • Focus on community involvement 	<ul style="list-style-type: none"> • Eliminated disparities in HbA1c testing, eye examinations, lipid profiles, and microalbumin testing • LDL testing and control • Lower extremity amputations 	<ul style="list-style-type: none"> • At least annual testing of HbA1c and serum creatinine • Annual screenings for hypertension and obesity

Table 2

Quality Improvement Interventions with Health IT Focus in Minority Populations, 2000-present

Reference	Health IT Focus	Study Objective(s)	Study Design	Population Characteristics	Results Summary	Cost of Health IT Implementation	Health IT Use Lessons Learned	Quality Score (out of 27)
Chin et al. (2004)	Electronic registry implemented in 37% of community health centers; used to follow up on examination and lab data	Evaluation of the Diabetes Health Disparities Collaborative with the aim to reduce health disparities and improve diabetes care quality in 19 community health centers	Pre-post design (1yr: 1998-99) Plan-do-study-act cycles using patient self-management tools, flowsheets, DM registry, group community health center visits regarding health system design All health centers were asked to perform at least two HbA1c tests at least 3 months apart over the year for 90% of their target population	19 Midwest community health centers (CHC) Chart review of 969 patients 33% African American 22% Hispanic	Improved processes of care including HbA1c measurement, foot examination, lipid assessments * Mean value of HbA1c improved, a drop of 0.2%. Quality improvement initiatives in CHCs can significantly improve diabetes care/process measures and may improve diabetes control Overall, the collaborative was considered worthwhile and successful.	No cost information reported	<ul style="list-style-type: none"> Only 37% of CHCs chose to use an electronic patient registry Staff found registry useful but burdensome Needed more technical support Difficulty developing registry Delay in finding an efficient and common electronic patient registry Delayed identification of target population 	13
Bray, Roupe, et al. (2005); Bray, Thompson, et al. (2005)	Electronic patient care registry system called the Cardiovascular/Diabetes Electronic Management System (CVDEMS)	Explore feasibility, cost-effectiveness, and efficacy of diabetes care management, group visits and electronic registry	Pre-post design (1yr) with control patients in separate practice Intervention: RN provided weekly intensive diabetes case management, 4 session group visit DM education program, DM registry and visit reminder system Control: Usual care	5 solo or small group primary care fee-for-service practices in rural North Carolina 314 patient participants 72% African American	Increased provider productivity on group visit days (20,17 to 31.55 average daily encounter rate) 61% of patients in intervention group had a reduction in HbA1c, while control group increased HbA1c. Improved processes of care including lipid panels and foot examinations improved in the intervention group No significant difference in weight or blood pressure Overall, intervention improved percentage of patients achieving	Substantial increase in office productivity, as measured by patient encounters Magnitude of increase in charges is sufficient to offset 75% of direct personnel costs on nurse case manager Demonstrated potential for sustainability	<ul style="list-style-type: none"> Pilot experience with CVDEMS showed much info was duplicative of the medical record, as a result, leadership began moving toward full implementation of electronic medical record. New system will allow for provider reminders and decision support. 	11.5

Reference	Health IT Focus	Study Objective(s)	Study Design	Population Characteristics	Results Summary	Cost of Health IT Implementation	Health IT Use Lessons Learned	Quality Score (out of 27)
Gerber et al. (2005)	Computer kiosks in waiting room, education with video testimonials, audio feedback, without text or complex navigation	Evaluate a clinic-based multimedia intervention for diabetes education among a low health literacy population	Randomized controlled trial (1 year) Intervention: Computer training (information, skills, support) kiosk in waiting room Control: Multiple choice computer quiz about diabetes	Five public clinics in Chicago 244 subjects enrolled; complete data for 183 subjects 29% African American 66% Latino	diabetes management goals. No significant difference in change in HbA1c, weight, blood pressure, knowledge, self-efficacy, or self-reported medical care Increased perceived susceptibility to diabetes complications in intervention group *	No cost information reported	<ul style="list-style-type: none"> Digital divide--lower health literacy correlated with less computer use, despite equal access to the technology Touch-screen monitors take the place of mouse and keyboard which may remove some barriers for older, lower literacy, less computer experienced individuals Optimal computer location was uncertain; limited privacy of waiting rooms; waiting for computer availability when multiple patients want to log in; computer malfunctions Staff assistance improved participant acceptance Multimedia audiovisual prompting necessary; repeated staff prompting important for repeated sessions 	20
Levetan et al. (2002)	Computer-generated 11" X 17" color poster depicting an individual's A1c, goals, and steps to achieve goals Individual report was generated from	Evaluate a system that provides uniquely formatted and personalized reports of diabetes status and goals on changes in HbA1c levels	Randomized controlled trial (6 months) All participants received diabetes education 3 months before enrollment	Participants were identified and enrolled from a group completing the American Diabetes Association educational	63% of patients in control group and 69% of patients in the intervention group experienced a decline in HbA1c Among patients with baseline HbA1c $\geq$	No cost information reported	<ul style="list-style-type: none"> Intervention required no effort on the part of the provider Visual nature of the poster may have contributed to 	17

Reference	Health IT Focus	Study Objective(s)	Study Design	Population Characteristics	Results Summary	Cost of Health IT Implementation	Health IT Use Lessons Learned	Quality Score (out of 27)
Phillips et al. (2005)	a Microsoft Access-based decision support system that collected patient enrollment questionnaire and matched it against established diabetes, cardiovascular, nutrition, and exercise guidelines	Target provider clinical inertia using computer reminders to improve diabetes patient clinical outcomes Part of the Improving Primary Care of African Americans with Diabetes (IPCAAD) study	Intervention: Computer-generated individually tailored poster and wallet card; along with one phone call from health educator Control: Standard care	program between October 1998–April 1999 150 patients with diabetes who completed a diabetes education program during the 3-month period before study enrollment 83% African American (control group) 89% African American (intervention group)	7.0%, the control group experienced a 0.77% absolute reduction and the intervention group had a 1.69% absolute reduction* Overall, participants in the intervention group significantly lowered their HbA1c compared to the control group	No cost information reported	success of patients with less than college education	18
Ziener et al. (2006)	Hard-copy computerized reminders that document critical values, notify when evaluations are due, and provide individualized recommendations for therapy	Evaluate whether computerized reminders vs. face-to-face feedback from an	Controlled trial (3 years); residents were randomized to intervention groups Three intervention groups: 1) residents received computerized reminders with patient-specific management recommendations, 2) residents received individual 5 minute face-to-face feedback every 2 weeks from endocrinologist, 3) both reminders and feedback Control: Standard care	Municipal hospital primary care clinic in large academic medical center 345 internal medicine residents serving 4,138 patients 94% patients African American	Change in HbA1c was greater for the reminder group (Δ -0.6%) compared to control group (Δ -0.2%)* Improved systolic blood pressure in feedback plus reminder and feedback only groups compared to reminder only and control groups LDL improved in all intervention arms compared to control In-person endocrinologist feedback (especially with reminders) modestly enhanced diabetes control and is more effective than reminders alone	No cost information reported	Printed reminders were easy to ignore while face-to-face feedback required the provider's attention	16.5

Reference	Health IT Focus	Study Objective(s)	Study Design	Population Characteristics	Results Summary	Cost of Health IT Implementation	Health IT Use Lessons Learned	Quality Score (out of 27)
	medications, and recommendations for diabetes management	endocrinologist improve provider clinical inertia defined as "did nothing/did anything/did enough" Part of the Improving Primary Care of African Americans with Diabetes (IPCAAD)	Three intervention groups: 1) residents received computerized reminders with patient-specific management recommendations, 2) residents received individual 5 min face-to-face feedback every 2 weeks from endocrinologist, 3) both reminders and feedback Control: Standard care When patient glucose levels exceeded 150mg/dL, provider behavior was characterized as "did nothing," "did anything" (intensification of therapy), or "did enough" (met intensification recommendations)	345 internal medicine residents serving 4,138 patients total 4,038 patients with glucose levels high enough to trigger intensification of therapy 94% of patients African American	declined throughout the 3-year period Therapy intensification increased when providers received in-person feedback with or without reminders (52%) compared to the group that received reminders alone or usual care (35%) Adequate medication intensification was associated with fall in HbA1c Therapy intensification among the reminder only group and control group did not remain above baseline levels		"forced" to respond to them	
Shea et al. (2006)	Telemedicine Home telemedicine unit (HTU) developed specifically for Informatics for Diabetes Education and Telemedicine (IDEATel) project Designed for videoconferencing, remote monitoring of glucose and blood pressure, accessing Web portal to clinical data, Web-messaging with nurse case managers, and diabetes education Web site	Improve HbA1c, blood pressure, and lipid levels utilizing telemedicine	Randomized controlled trial (1 year follow up) Intervention: RN telemedicine case management and treatment algorithms Control: Usual care	Urban patients identified and enrolled through Columbia University Medical Center Rural patients identified and enrolled through State University of New York 1,665 Medicare beneficiaries residing in federally designated medically underserved areas 15% African American 36% Latino	Intervention group experienced improvements in: HbA1c (net compared to control group of -0.18%)* Systolic blood pressure (net -3.4mmHg) and diastolic blood pressure (net -1.9mmHg)* Total (net -11.06mg/dL) and LDL lipid levels (net -9.5mg/dL)*	Burden on the health care delivery system- cost of technology and personnel for case management Cost to project for HTU devices was \$3,425 Increase in Medicare claims in intervention group (increased use of services) Reimbursement models may not cover telemedicine visits	Physical size of the HTU and difficulty in learning to use it, responsible for most dropouts HTU could not be installed in some homes due to poor telephone line quality Providers still preferred traditional communication with nurse case managers despite access to secure Web-based messaging Providers must be licensed in the state	21

Reference	Health IT Focus	Study Objective(s)	Study Design	Population Characteristics	Results Summary	Cost of Health IT Implementation	Health IT Use Lessons Learned	Quality Score (out of 27)
Shea et al. (2009)	Telemedicine HTU developed specifically for IDEATel project Designed for videoconferencing, remote monitoring of glucose and blood pressure, accessing Web portal to clinical data, Web-messaging with nurse case managers,	To examine the effectiveness of telemedicine intervention to achieve clinical management goals in older, ethnically diverse, medically underserved patients with diabetes	Randomized controlled trial (5 year follow up) Intervention: RN telemedicine case management and treatment algorithms Control: Usual care	Urban patients identified and enrolled through Columbia University Medical Center Rural patients identified and enrolled through State University of New York 1,665 Medicare beneficiaries (793	Intervention group experienced improvements in: HbA1c: -0.29%* Systolic blood pressure: -4.32 mmHg* Diastolic blood pressure: -2.64 mmHg* LDL :-3.84mg/dL*	No cost information reported Costs to Medicare for this program are reported elsewhere. (Moreno et al., 2009)	where telemedicine is accessed ▪ Digital divide (disproportionate experience using technology) was not a large barrier to use ▪ Comfort with technology and belief in benefit from it played an important role in enrollment (Palmas et al., 2006) ▪ Most frequent reason for participation was belief that technology could help them (52% urban, 42% rural) (Palmas et al., 2006) ▪ 22% of rural non-participants were not comfortable with the technology (Palmas et al., 2006) ▪ 71% of urban non-participants did not believe technology could help them; 52% uncomfortable with it (Palmas et al., 2006) ▪ High attrition rate	20.5

Reference	Health IT Focus	Study Objective(s)	Study Design	Population Characteristics	Results Summary	Cost of Health IT Implementation	Health IT Use Lessons Learned	Quality Score (out of 27)
Cherry et al. (2002)	and diabetes education Web site	To determine the impact of a Web-based patient interface technology as part of a diabetes disease management program	Pre-post design (1year: 2000–2001) Participants had Health Buddy appliance installed on home telephone line, were prompted to answer self-management questions daily; care managers followed up with patients at highest calculated risk for hospitalization or adverse outcomes	after 5 years) residing in federally designated medically underserved areas 15% African American 36% Latino	Hazard ratio 1.01--mortality was not different between groups, though power was limited	Saved approx \$747/member/year compared to 1999 comparative sample of people with diabetes	Health Hero technology platform: flexibility, simplicity, cost timeliness, cost	10.5
	Telemedicine Diabetes Disease Management Program featuring the use of Health Hero iCare Desktop Buddy appliance Health Hero iCare Desktop allows for integrated patient enrollment, scheduling, monitoring tools, a secure Web site; used by Mercy's telephone support staff			Mercy Health Center in Laredo, Texas Indigent border residents with diabetes; economically disadvantaged 169 patients with diabetes identified through hospital discharge data, physician private practices, local health department diabetic clinics, regional clinic registries and outreach programs	Intervention reduced: Inpatient hospitalizations (~32%) Post-discharge care visits (~44%) Outpatient visits (~49%) * ER encounters (~34%) Improved perceived quality of life assessed by Medical Outcomes Study 12-item Short Form survey: mental component improved 2.8 points, * physical component improved 2.1 points 94% of participants self-reported regular medication adherence (up from 34%)	Health Buddy reportedly easy to use, high levels of perceived value	Health Hero technology platform: flexibility, simplicity, cost timeliness, cost	

* Indicates a statistically significant finding ($p \leq 0.05$) Note: CHC = community health center; CVDEMS = cardiovascular/diabetes electronic management system; DM = diabetes mellitus; ER = emergency room; health IT = health information technology; HTU = home telemedicine unit; IDEATel = Informatics for Diabetes Education and Telemedicine; IPCAAD = Improving Primary Care of African Americans with Diabetes; LDL = low density lipoprotein; RN = registered nurse; Palmas, W., Teresi, J., Morin, P., Wolff, L.T., Field, L., Eimicke, J.P., Capps, L., Prigollini, A., Orbe I., Weinstock, R.S., & Shea S. (2006). Recruitment and enrollment of rural and urban medically underserved elderly into a randomized trial of telemedicine case management for diabetes care. *Telemedicine Journal and e-Health*, 12(5), 601-7.