Published in final edited form as:

Chemistry. 2010 November 15; 16(43): 12797–12800. doi:10.1002/chem.201002102.

The Development of a General Strategy for the Synthesis of Tyramine-Based Natural Products by Using Continuous Flow Techniques

Dr. Srinivas Achanta, Dr. Virginie Liautard, Robert Paugh, and Prof. Michael G. Organ*

Department of Chemistry, York University 4700 Keele Street, Toronto, ON, M3J 1P3 (Canada)

Keywords

flow chemistry; Heck reaction; microwave chemistry; tyramine

The movement of synthetic chemistry from a traditional batch format (e.g., round-bottom flask or kettle reactor) to a flowed platform offers many advantages.^[1] In smaller reaction channels, the chemical transformation itself can benefit from improved mass and thermal transfer leading to enhanced kinetics and cleaner product mixtures with fewer byproducts. The latter benefits are significantly affected by the introduction of both a space and time element to the reaction process. For example, starting materials and/or catalysts can be kept separate and only brought together at the point of reaction and every resultant minute plug of reaction mixture thus formed is continuously moved away from infusing reactants eliminating many side reactions. Operationally, the movement of synthesis to a flowed platform allows for real-time reaction monitoring coupled with instantaneous changes of reaction parameters that vastly truncates reaction process optimization. Finally, once optimal conditions are achieved, larger quantities of product can be obtained by simply flowing using those parameters for the necessary time (or in parallel reaction tubes) to accrue the desired amount of product, a concept termed ‘scale-out’.

We have been developing microwave-assisted, continuous-flow organic synthesis (MACOS) and applying it in single-step reactions including those promoted by thin-metal films that line the reaction tube.^[2] Most recently we have investigated by using the scale-out strategy with MACOS to produce multigram quantities of benzo-fused sultams.^[3] Here we are furthering the development of this platform to multi-step transformations for the production of tyramine-based natural products entirely by flow^[4] (see Scheme 1 for target structures).^[5–8] We are attracted to the members of this family of natural products for their inherent biological activity,^[9–13] but additionally they can themselves serve as templates for the diversity-oriented synthesis of analogues. Indeed, family members contain alkylating sites, a Michael acceptor for addition reactions, and in some cases halides for substitution reactions (e.g., cross-coupling). As such, it would be beneficial to demonstrate the capability to prepare them potentially on at least a gram scale using a scale-out synthetic approach. The retrosynthetic route that we developed to approach all of the members of this natural product family by flow revolves around a Heck/alkylation sequence (Scheme 1). Unsaturated amides **11** were prepared readily by simple acylation of either tyramine itself (**12**, R⁴=R⁵= H) or the appropriate derivative with acryloyl chloride.^[14]

Heck reactions are generally run at relatively high temperatures (>100°C) for several hours. However, for the flow sequence to be effective, the reaction has to reach a high level of completion during the time in which it is receiving microwave irradiation while it is traversing the reaction capillary; this is in the order of minutes. To explore how the Heck reaction could be made to perform under flow conditions,^[15] we first examined a simplified set of starting materials (Table 1). As a control, we performed the reaction first under batch conditions in an oil bath (Table 1, entry 1) and found that 18 h were required to achieve full conversion. Knowing that the flowing reaction only resides for a few minutes in the microwave irradiation zone, we irradiated it with enough power to create temperatures well in excess of 100°C (Table 1, entries 2–7). With optimization of concentration, flow rate, and temperature, excellent levels of conversion were attained using MACOS. Using the conditions outlined in Table 1, entry 5, compounds **1**, **2**, **5**, and **6** were prepared from their corresponding Heck precursors (**10** and **11**, where compound **11** was used directly from the acylation procedure) in 32, 22, 48, and 46% yield, respectively (Scheme 2).^[16]

From the above applied results, some shortcomings of the Heck conditions were realized. Most importantly, yield dropped off sharply, which was found to be primarily the result of the target products undergoing a second Heck reaction.^[17] With some investigation, it was found that the excess tryamine from the acylation reaction (used crude in the next step) was present as the HCl salt and was being deprotonated by the triethylamine that was originally used for the subsequent Heck reaction. The resultant tyramine free base (in excess) then underwent a Michael reaction with compound **11**, the initially formed desired product, thus removing it from the reaction mixture; the aryl iodide, now in excess due to the depletion of **11**, underwent a second undesired Heck coupling. This problem was overcome by the use of equimolar quantities of tryamine and acryloyl chloride and only a slight excess of Hunig's base in the Heck reaction, which now led to good yields of analogues **1** and **3** (see bottom of Scheme 2). The addition of tetrabutylammonium bromide and Hunig's base, which has a soluble conjugate acid, were key to this yield improvement and instrumental to producing larger quantities (vide infra).

With the first two steps worked out, we extended the flow process to include a subsequent alkylation step to approach **7** and aplysamine 6 (**8**) (Scheme 3).^[18] Acylation proceeded cleanly and the crude material was used directly, as usual, in the Heck reaction. Coupling of mono bromide **11a** proceeded uneventfully and in excellent recovery. However, unexpectedly, the dibromide (**11b**) underwent reduction of one of the bromides. To probe the origin of this undesired reaction, we first repeated it with identical reactions conditions only in batch, and the same result was attained. Thus, the side reaction is not a consequence of flow. Clearly, the propensity for reduction diminishes when one of the bromides is removed as we never observed reduction in the case of **11a**. Perhaps the hydroxyl group in **17** makes the ring electron-rich enough to avoid oxidative addition of a single bromide under these reaction conditions, but the presence of the second bromide tips the balance and activates the ring. The next question is the hydride source for this reduction. We suspected the phenol and to test this we protected it as a methyl ether; this halted reduction confirming the phenol as the source of the problem.

Carrying on with the preparation of the final tyramine analogues, we set out to alkylate the phenol position. The reaction was thought to work best with a soluble base that had a soluble conjugate acid that could facilitate kinetic deprotonation of the phenol. With only minor optimization, potassium trimethylsilanoxide was loaded into one syringe along with **16** or **17**, while the alkyl bromide was loaded into a second syringe. Substitution proceeded cleanly providing the corresponding alkylated products **19** and **20**. It was determined that deprotection with TFA could not be carried out in DMF. Thus, the solvent had to be swapped out to CH₂Cl₂ meaning that the two steps could not be flowed directly from one to

the next. Nevertheless, deprotection also proceeded nicely at room temperature providing desired products **7** and **8** with excellent recovery (Scheme 3).

As mentioned, we were developing a process to make the compounds in this natural product series with an eye on the potential to produce larger quantities of final product. To this end, compounds **1** and **2** were prepared on a 1.02- and 1.64-gram scale using the corresponding two-step sequence described in Scheme 2. Compounds **7** and **8** (aplysamine 6) were prepared by using the four-step protocol detailed in Scheme 3 in final quantities of 1.5 and 0.63 grams, respectively.

In summary, we have developed a general synthetic method for the preparation of a number of tyramine-derived natural products and analogues that is based on a combination of room-temperature alkylation and deprotection processes with microwave-heated Heck coupling. The protocol has been demonstrated to be suitable for scale-out, meaning that we can prepare, in principle, any quantity of these compounds we require without the need for process reoptimization. Applications of multistep, flow synthesis of natural products of progressing complexity are underway in our laboratory.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by NSERC (Canada), the ORDCF (Ontario) and the NIH Center for Chemical Methodologies and Library Development at the University of Kansas (P50 GM069663).

References

1. For recent reviews on flow chemistry, see: a) Razzaq T, Kappe CO. Chem Asian J. 2010; 5:1274–1289. [PubMed: 20411525] b) Garcia-Verdugo E, Luis SV. Chemical Reactions and Processes under Flow Conditions. The Royal Society of Chemistry Cambridge 2010:44–85. c) Lin WY, Wang Y, Wang S, Tseng HR. Nano Today. 2009; 4:470–481. [PubMed: 20209065] d) Wiles C, Watts P. Future Medicinal Chemistry. 2009; 1:1593–1612. [PubMed: 21425981]
2. a) Shore G, Yoo WJ, Li CJ, Organ MG. Chem Eur J. 2010; 16:126–133. b) Shore G, Organ MG. Chem Eur J. 2008; 14:9641–9646. c) Shore G, Organ MG. Chem Commun. 2008:838–840. d) Shore G, Morin S, Mallik D, Organ MG. Chem Eur J. 2008; 14:1351–1356. e) Bremner S, Organ MG. J Comb Chem. 2007; 9:14–16. [PubMed: 17206827] f) Shore G, Morin S, Organ MG. Angew Chem. 2006; 118:2827–2832. Angew Chem Int Ed. 2006; 45:2761–2766. g) Organ MG, Comer E. Chem Eur J. 2005; 11:7223–7227. h) Comer E, Organ MG. J Am Chem Soc. 2005; 127:8160–8167. [PubMed: 15926844]
3. Ullah F, Samarakoon T, Rolfe A, Kurtz RD, Hanson PR, Organ MG. Chem Eur J. 2010; 16:10959–10962.
4. For examples of natural products synthesis conducted in flow, see: a) Hopkin MD, Baxendale IR, Ley SV. Chem Commun. 2010; 46:2450–2452. b) Herath A, Dahl R, Cosford NDP. Org Lett. 2010; 12:412–415. [PubMed: 20038130] c) Sahoo HR, Kralj JG, Jensen KF. Angew Chem Int Ed. 2010; 49:5704–5708.
5. For the isolation and biological activity of aplysamine 6, see: Buchanan MS, Carroll AR, Fechner GA, Boyle A, Simpson M, Addepalli R, Avery VM, Hooper JNA, Cheung T, Chen H, Quinn R. J Nat Prod. 2008; 71:1066–1067. [PubMed: 18393464]
6. For the isolation of ailanthamide, see: Chen JJ, Chung CY, Hwang TL, Chen JF. J Nat Prod. 2009; 72:107–111. [PubMed: 19128011]
7. For the isolation of beecheyamide, see: Cheng MJ, Wu CC, Tsai IL, Chen IS. J Chin Chem Soc. 2004; 51:1065–1072.

8. For the isolation of N-trans-feruloyl tyramine and N-trans-coumaroyl tyramine, see: a) Okuyama T, Shibata S, Hoson M, Kawada T, Osada H, Noguchi T. *Plant Methods*. 1986;171–174. b) Yoshihara T, Yamaguchi K, Takamatsu S, Sakamura S. *Agri Biol Chem*. 1981; 45:2593–2598.
9. For cytotoxicity studies involving tyramines, see: Zhao G, Hui Y, Rupprecht K, McLaughlin JL. *J Nat Prod*. 1992; 55:347–356. [PubMed: 1593281]
10. For use of tyramines as antioxidants and cytoprotectives, see: Tomosaka H, Chin YW, Salim AA, Keller WJ, Chai H, Kinghorn DA. *Phytother Res*. 2008; 22:979–981. [PubMed: 18389483]
11. For the use of tyramines to cleave DNA, see: Ma J, Jones SH, Hecht SM. *Bioorg Med Chem*. 2004; 12:3885–3889. [PubMed: 15210155]
12. For the use of tyramines to inhibit acetylcholinesterase, see: Kim DK, Lee K. *Arch Pharmacol Res*. 2003; 26:735–738.
13. For the use of tyramines to inhibit nitric oxide synthase, see: Kim Y, Han MS, Lee S, Kim J, Kim YC. *Phytother Res*. 2003; 17:983–985. [PubMed: 13680842]
14. Tyramine, or one of its analogues (1.0 equiv) was reacted with acryloyl chloride (1.0 equiv) and Hunig's base (1.1 equiv) in DMF. The crude amide was used directly in the subsequent Heck reaction.
15. For other examples of the Heck reaction performed by using flow techniques, see: a) Nikbin N, Ladlow M, Ley SV. *Org Proc Res Dev*. 2007; 11:458–462. b) Mennecke K, Kirschning A. *Beilstein J Org Chem*. 2009; 5(21)
16. For complete experimental details and full spectral characterization of all new compounds, see the Supporting Information.
17. For the structure of the double-Heck product and full spectral characterization, see the Supporting Information.
18. During the course of our studies, the first synthesis of aplysamine 6 was reported, see: Ullah N, Arafeh KM. *Tetrahedron Lett*. 2009; 50:158–160.

- 1, $R^1 = H$, $R^2 = H$, $R^3 = H$, $R^4 = H$, $R^5 = H$, $R^6 = H$ (*N-trans*-coumaroyl tyramine)
- 2, $R^1 = H$, $R^2 = H$, $R^3 = CH_3$, $R^4 = H$, $R^5 = H$, $R^6 = CH_3$ (Ailanthamide)
- 3, $R^1 = H$, $R^2 = OCH_3$, $R^3 = H$, $R^4 = H$, $R^5 = H$, $R^6 = H$ (*N-trans*-feruloyl tyramine)
- 4, $R^1 = CH_3$, $R^2 = OCH_3$, $R^3 = CH_3$, $R^4 = H$, $R^5 = H$, $R^6 = H$
- 5, $R^1 = CH_3$, $R^2 = OCH_3$, $R^3 = CH_3$, $R^4 = H$, $R^5 = H$, $R^6 = CH_3$ (Beecheyamide)
- 6, $R^1 = CH_3$, $R^2 = Br$, $R^3 = H$, $R^4 = H$, $R^5 = H$, $R^6 = H$
- 7, $R^1 = CH_3$, $R^2 = Br$, $R^3 = H$, $R^4 = H$, $R^5 = H$, $R^6 = CH_2CH_2CH_2NH_2$
- 8, $R^1 = CH_3$, $R^2 = Br$, $R^3 = H$, $R^4 = Br$, $R^5 = Br$, $R^6 = CH_2CH_2CH_2NH_2$ (Aplysamine 6)

Scheme 1.

Synthetic approach to the tyramine-based family of natural products and analogues.

Scheme 2. Heck reaction between fragments **10** and **11**.^[16]

Scheme 3.

Full protocol to prepare brominated tyramine derivatives **7** and **8**.^[16]

Table 1

Optimization of the Heck reaction.

Entry	14 [equiv]	Conc. of 13 ^[a] [M]	Time or flow rate ^[b]	Temp. [°C] ^[c]	Conversion [%] ^[d]
1	1.5	0.6	18 h	90	100
2	1.8	0.6	20	100	76
3	1.8	0.6	20	135	85
4	1.8	0.6	15	125	94
5	1.5	0.6	20	140	92
6	1.8	0.9	20	110	65
7	1.8	1.5	20	140	complex mixture

^[a]Concentration is based on **14**.

^[b]The experiment in entry 1 is a batch reaction performed in an oil bath for 18 h, all other entries are flowed reactions and the rate listed is in μLmin^{-1} .

^[c]The temperature listed in entry 1 is of the oil bath. All other temperatures are the temperature recorded off the surface of the reaction capillary by the internal IR sensor of the Biotage Initiator Synthesizer.

^[d]Percent conversion is determined by taking an aliquot from the eluent stream from the capillary (crude material) and calculating the ratio of the peaks of the starting materials and product in the proton NMR spectrum.