Published in final edited form as:

J Acquir Immune Defic Syndr. 2010 September 1; 55(1): 65–72. doi:10.1097/QAI.0b013e3181d916db.

HIV risk behavior in treatment-seeking opioid-dependent youth: Results from a NIDA Clinical Trials Network multi-site study

Christina S. Meade, PhD¹, Roger D. Weiss, MD², Garrett M. Fitzmaurice, ScD², Sabrina A. Poole, MS³, Geetha A. Subramaniam, MD⁴, Ashwin A. Patkar, MD¹, Hilary S. Connery, MD², and George E. Woody, MD³

¹Department of Psychiatry & Behavioral Sciences, Duke University Medical Center, Durham, NC

²Department of Psychiatry, Harvard Medical School and McLean Hospital, Belmont, MA

³Department of Psychiatry, University of Pennsylvania and Treatment Research Institute, Philadelphia, PA

⁴National Institute on Drug Abuse, Bethesda, MD

Abstract

Objective—To assess baseline rates of and changes in HIV drug and sexual risk behavior as a function of gender and treatment in opioid-dependent youth.

Methods—150 participants were randomly assigned to extended buprenorphine/naloxone therapy for 12 weeks (BUP) or detoxification for 2 weeks (DETOX); all received drug counseling for 12 weeks. HIV risk was assessed at baseline and 4-, 8-, and 12-week follow-ups. Behavioral change was examined using generalized estimating equations.

Results—Baseline rates of past-month HIV risk for females/males were 51%/45% for injection drug use (IDU) (ns), 77%/35% for injection risk ($p < .001$), 82%/74% for sexual activity (ns), 14%/24% for multiple partners (ns), and 68%/65% for unprotected intercourse (ns). IDU decreased over time ($p < .001$), with greater decreases in BUP versus DETOX ($p < .001$) and females versus males in BUP ($p < .05$). Injection risk did not change for persistent injectors. Sexual activity decreased in both genders and conditions ($p < .01$), but sexual risk did not.

Conclusions—Overall IDU and sexual activity decreased markedly, particularly in BUP patients and females, but injection and sexual risk behaviors persisted. While extended buprenorphine/naloxone therapy appears to have favorable effects on HIV risk behavior in opioid-dependent youth, risk reduction counseling may be necessary to extend its benefits.

Keywords

HIV risk; opioid dependence; youth; gender differences; buprenorphine treatment; injection drug use

INTRODUCTION

Over the past decade, there has been a marked increase in illicit opioid use, including heroin and prescription analgesics, among adolescents in the United States,¹ with a substantial proportion of initiators progressing to dependence within 1 year.² Opioid dependence is

Corresponding author and request for reprints to: Christina S. Meade, PhD, Duke Global Health Institute, Box 90519, Durham, NC 27708, Tel. 919.613.6549, Fax. 919.613.6215, christina.meade@duke.edu.

Trial registration: ClinicalTrials.gov #NCT00078130; IND #64,578

often a chronic and relapsing medical condition, and the prognosis for opioid-dependent youth appears poor.^{3, 4} Among the many adverse consequences of opioid dependence is exposure to infectious diseases, including HIV.

In 2006, 34% of new HIV infections occurred in persons 13–29 years old – more than in any other age group,⁵ and opioid-dependent youth may be at particularly high risk. Over half of young heroin users inject the drug and are at risk for contracting HIV through sharing of syringes, injection paraphernalia, and drug solution.^{6–9} Non-injection drug use also contributes to the spread of HIV through its association with sex risk behavior. Previous research has found that heroin^{10–13} and prescription opioid^{8, 14} users engage in high rates of sex risk behaviors, including multiple partners, unprotected intercourse, and sex trading. However, only one of these studies focused on youth.⁸ A number of developmental factors make adolescents more vulnerable to HIV risk behavior than adults. These include hormonal and physical changes of puberty,^{15, 16} shifting relationship patterns, including development of romantic relationships, and increasing influence of peer norms,^{17–20} a heightened sense of invulnerability,^{21, 22} cognitive immaturity leading to impulsive decision-making,²³ tenuous affect regulation,^{16, 24, 25} and underlying brain changes associated with increased risk-taking propensity.^{26–28} Since results from adult samples may not generalize to adolescents, it is critical to identify the rates and patterns of HIV risk behaviors among opioid-dependent youth.

Females now account for a quarter of new HIV/AIDS diagnoses, with high-risk heterosexual contact and injection drug use (IDU) accounting for nearly all of these infections.²⁹ Studies conducted among adult drug users suggest that women are more likely to engage in HIV risk behaviors, including sharing needles and other injection paraphernalia,^{30, 31} having high-risk sex partners,^{12, 32, 33} and engaging in sex trade.^{13, 34, 35} Therefore, it is also important to examine whether gender differences in HIV risk behavior are apparent among opioid-dependent youth.

Treatment for opioid dependence, including use of buprenorphine/naloxone (bup/nx), is an important component of HIV prevention.³⁶ Buprenorphine, a high-affinity μ -opioid partial agonist and κ -opioid antagonist, is a safe and effective medication for both opioid detoxification and longer-term agonist therapy in adults^{37–40} and adolescents.^{41, 42} To date, four studies have examined changes in HIV risk following bup/nx treatment. The first randomized 137 adults to bup/nx, methadone, or levomethadyl acetate.⁴³ Over 18 weeks of treatment, there were significant reductions in frequency of opioid injection and sharing of injection equipment, but no differences across conditions. Methadone was associated with decreased sexual activity, possibly due to its effect on libido,^{44, 45} but there was no change in number of sex partners in any condition. The second was an observational study of 166 adults initiating bup/nx treatment in a primary care clinic.¹⁰ Over 24 weeks of treatment, there were significant declines in IDU, sharing of injection equipment, and sex while intoxicated, but not in unprotected intercourse or sex with a new or secondary partner. The third study randomized 126 adults to 24 weeks of treatment with bup/nx, naltrexone, or placebo.⁴⁰ Among participants retained in treatment, there was a significant reduction in drug risk but not sex risk, and no differences across conditions. In the only published bup/nx treatment study that has examined HIV risk in youth, 36 adolescents were randomized to 4 weeks of detoxification with bup/nx or clonidine.⁴⁶ There was a significant decrease in drug risk during the first week, but no difference by gender or condition. This study did not report on sex risk and was limited by a small sample and short assessment period. Cumulatively, these studies suggest that pharmacotherapy for opioid dependence may promote reductions in HIV risk behavior, particularly related to drug risk. However, given the increasing prevalence of opioid abuse and HIV infection among adolescents and young adults, further

research is needed to understand if and how bup/nx may contribute to HIV risk reduction in opioid-dependent youth.

The current study, conducted within the NIDA Clinical Trials Network (CTN), was a multi-site randomized clinical trial of bup/nx treatment for opioid-dependent youth.⁴¹ As reported in the primary outcome paper, participants who received extended bup/nx therapy had significantly better treatment retention, greater reductions in opioid, marijuana, and cocaine use and drug injection, and less need for non-study addiction treatments (i.e., higher-level of care).⁴¹ The aims of this secondary analysis were to: (1) describe the prevalence of HIV drug and sex risk behaviors in the sample; (2) examine gender differences in HIV risk behaviors; and (3) determine whether HIV risk behavior changed over time as a function of treatment and gender.

METHODS

Participants and procedures

Full details of the methods have been reported previously.⁴¹ The study met all requirements of human subjects protection and was approved by the institutional review boards of participating institutions. Recruitment was open to patients aged 15–21 years who met DSM-IV criteria for opioid dependence with physiologic features and sought outpatient treatment at community-based clinics between July 2003 and December 2005. Eligible participants were randomly assigned to either (1) 14-day bup/nx detoxification (DETOX), or (2) 12-week extended bup/nx therapy (BUP). Randomization was stratified by gender (male/female), age (14–17/18–21), ethnicity (Caucasian/other), and injection status (injecting/non-injecting). Participants completed assessments at baseline (pre-treatment), weeks 4 and 8 (mid-treatment), and week 12 (post-treatment).

Of 154 randomized individuals, 4 were excluded from this analysis (2 did not enter treatment, 2 did not provide HIV risk data), leaving a final sample of 150. Of these participants, 74 were allocated BUP and 74 were allocated DETOX. Follow-up rates were 85%, 74%, and 67% at weeks 4, 8, and 12, respectively. There was a significant decline in follow-up over time (Wald $\chi^2(2) = 27.63, p < .001$), but no differences in overall follow-up by condition (Wald $\chi^2(1) = 1.91, p = .17$) or gender (Wald $\chi^2(1) = 0.66, p = .42$). Furthermore, participants who completed the 12 week follow-up did not differ from those who did not on any baseline characteristics.

Treatment

Consistent with standard bup/nx induction, patients were instructed to not use opioids for ≥ 6 hours and be in mild/moderate withdrawal before first dosing. Medication was administered under direct observation 5–7 days/week. BUP participants received a maximum dose of 24 mg/day of bup/nx for 9 weeks followed by a taper off the medication during weeks 10–12. DETOX participants were started on a maximum dose of up to 14 mg/day of bup/nx, followed immediately by a gradual taper off the medication during weeks 1–2, such that they were completed detoxified by day 14. In addition, participants in both groups were scheduled to receive standardized individual and group drug counseling weekly, with more frequent sessions if needed (www.nida.nih.gov/txmanuals/idca/idca1.html). Only 23 participants (15%) opted for extra sessions (12 had one, 6 had two, and 5 had three or more). The counseling focused on reducing drug use and relapse prevention (e.g., providing education about addiction, promoting positive relationships, learning how to tolerate stressful events, developing ways to avoid drug using situations), but they did not directly address HIV risk reduction.

Measures

At each visit, HIV drug and sex risk behavior in the past 30 days was assessed using the Risk Behavior Survey, an interviewer-administered survey used widely in CTN studies.⁴⁷ The following behaviors were modeled as binary outcomes (yes/no): self-report of any opioid use, IDU, injection risk (e.g., using dirty needles, sharing equipment, splitting drug solution), sexual activity (i.e., vaginal or anal intercourse), multiple partners (i.e., ≥ 2 sex partners), and unprotected intercourse (i.e., non-condom use). While opioid use and sexual activity are not necessarily HIV risk behaviors per se, they are a prerequisite for engaging in injection and sex risk behaviors associated with HIV infection. The CTN Baseline Demographics Form collected information about substance use in the past month (days) and lifetime (years).

Data analysis

First, prevalence of HIV risk behaviors was examined at baseline and weeks 4, 8, and 12, with gender comparisons made at each occasion using chi-square tests. Next, longitudinal changes in drug and sexual behaviors were examined over time using the generalized estimating equations (GEE) method. Specifically, a logistic regression model was fit to the repeated binary outcomes to examine predictors of change in these behaviors over the course of treatment. This model, fitted using the GEE approach, assumed a compound symmetry correlation structure to account for the correlation among repeated measures over time; the reported standard errors were based on the so-called “empirical” variance estimator. The GEE analysis allows for missing data, so all available data for each study visit were included. A sequence of models was fit in a hierarchical manner with variables entered in the following blocks: (1) condition, time, and gender; (2) condition by time interaction, and (3) condition by time by gender interaction. Other 2-way interactions were included in the models but are not reported. For opioid use, the GEE analysis modeled drug use at week 4, 8, and 12 follow-up visits; the pre-treatment visit could not be included because all participants had used opioids in the past month (i.e., there was no variation in the baseline measure). For all behaviors, the pre-treatment visit was included as the baseline comparison. For drug behaviors, data from weeks 4 and 8 were modeled as a single intermediate visit reflecting behavior at mid-treatment (i.e., average IDU and risk); the rationale for doing so was the sparseness of data at week 8 when stratified by gender (e.g., no females in BUP injected drugs). For sexual behaviors, all four visits were included. Analyses were conducted using SPSS 17.0.

RESULTS

Participant characteristics

The sample included 61 females and 89 males. Participants were 15–21 years old ($M = 19.15$, $SD = 1.49$) and primarily (72%) Caucasian. The mean education was 11.21 years ($SD = 1.60$), and approximately half (52%) were currently employed. As shown in Table 1, there were no gender differences on demographic variables. Females and males also had similar histories of substance use, except females had used cocaine for more years (1.08 vs. 0.47; $p < .05$). There were no significant gender differences in days of substance use in the month prior to baseline.

Baseline HIV risk behaviors

Table 2 shows the past-month prevalence of HIV risk behaviors by gender over time. At baseline, all participants had used opioids (as per inclusion criteria); nearly half reported IDU, with no difference by gender. Females were significantly more likely to report injection risk (77% vs. 35%). Specifically, they were more likely to share a cooker, cotton,

or rinse water (52% vs. 20%; $p=.005$) and to fix drugs and split the solution with someone else (52% vs. 20%; $p=.005$), but no more likely to use works after someone else (39% vs. 30%; $p=.44$).

At baseline, the majority of participants (77%) were sexually active in the past month. Among the sexually active, 20% had multiple partners and 66% engaged in unprotected intercourse. There were no significant gender differences in these sexual behaviors at any time point, except that significantly more males than females reported multiple sex partners at week 4 (31% vs. 8%) and week 8 (29% vs. 10%).

Changes in drug and sexual behaviors

Table 3 presents the results of the GEE analyses predicting changes in drug behaviors over the course of treatment. As reported in the main outcome paper⁴¹, there was a significant overall reduction in opioid use from 100% at baseline to 71%, 49%, and 57% at weeks 4, 8, and 12, respectively ($p<.001$). At each follow-up visit, BUP patients were less likely than DETOX patients to have used opioids ($p<.001$). For example, at week 8 (prior to the taper), 33% of BUP patients had used opioids compared to 67% of DETOX patients. There was no gender difference in opioid use over the course of treatment.

For IDU, there were significant effects for time, condition by time, and condition by time by gender. This means that, compared to baseline, the rate of IDU decreased over the course of treatment in both conditions ($p<.001$), but the decrease was significantly greater in BUP versus DETOX patients ($p=.03$). For example, at week 8, 9% of BUP patients compared to 30% of DETOX patients had injected drugs. Furthermore, the effect of condition differed by gender ($p=.02$). Figure 1 illustrates this 3-way interaction effect, suggesting that the benefit of BUP in terms of reducing the relative risk of IDU was greater for females than males. This effect was driven primarily by changes from baseline to mid-treatment (Wald $\chi^2(1) = 5.49$, $p=.02$) rather than changes from baseline to post-treatment when bup/nx had been tapered in both conditions (Wald $\chi^2(1) = 0.06$, $p=.81$). Among all injectors, however, there was no significant reduction in injection risk over time or by condition. For participants who continued to inject drugs, the odds of engaging in injection risk behavior did not decrease for either gender, but female injectors had a significantly higher overall rate of injection risk behavior compared to male injectors ($p=.001$).

Table 3 also shows the results of parallel GEE analyses predicting change in sexual behavior. Over the course of treatment, there was a significant reduction in sexual activity for all participants ($p=.008$). At baseline, 77% were sexually active, compared to 67%, 63%, and 73% at weeks 4, 8, and 12, respectively; there was no effect of condition or gender. Among sexually active participants, males were more likely than females to have multiple partners ($p=.006$). There was no change in the rate of multiple partners over time or by condition. Engagement in unprotected intercourse was unrelated to gender, time, or condition.

DISCUSSION

This is the first study to report on gender differences in HIV risk behaviors among opioid-dependent youth receiving outpatient bup/nx treatment, and to examine changes in drug and sexual behaviors as a function of gender and treatment. The majority of males and females in this sample reported one or more HIV risk behaviors in the month prior to treatment, including injection risk (e.g., sharing needles, splitting drug solution), unprotected intercourse, and multiple sex partners. These behaviors place opioid-dependent youth at high risk for infection with HIV and other diseases. This is especially alarming given that 18% of

this sample were seropositive for hepatitis C virus at baseline, and 4 of the 83 participants (5%) who were seronegative at baseline converted to seropositive status by week 12.⁴¹

Over the course of treatment, there were significant reductions in opioid use and IDU in both treatment conditions, but these reductions were greater for participants in BUP. By the end of treatment, only 15% of BUP participants were injecting compared to 35% of DETOX participants. This finding of decreased opioid use and IDU associated with bup/nx treatment is consistent with previous trials conducted among adults^{10, 40, 43} and adolescents.⁴⁶ Given that opioid-dependent youth are at such high risk for relapse to drug use,⁴⁸ these results provide further support that bup/nx treatment is a safe and effective option that should be considered. However, among participants who continued to inject drugs, there was no reduction in injection risk over time. This suggests that extended bup/nx therapy alone maybe insufficient for stopping behavioral risk among persistent drug injectors.

While males and females were equally likely to be injectors, females were significantly more likely to engage in injection risk behavior. For example, in the month prior to enrollment, 77% of female injectors compared to 35% of male injectors reported injection risk. Specifically, females were more likely to share a cooker, cotton, or rinse water and to fix drugs together and split the solution. These types of injection risk behaviors may be an increasingly important route of HIV and hepatitis C transmission.⁴⁹ While further research is needed to better understand this gender difference, studies with adult injectors suggest that intimate partnerships may play a role. Women are more likely than men to attribute their initiation to and continued use of heroin to social reasons, particularly the influence of an opioid-using partner.^{50–52} Furthermore, the primary reason injectors report sharing equipment is the use of drugs with sex partners.^{52–54} Qualitative and social network research with young drug users may help elucidate the interpersonal dynamic and context associated with injection risk in females.

Interestingly, females responded particularly well to extended bup/nx therapy. Females in BUP had larger declines in IDU compared to females in DETOX, whereas males in both groups showed only modest declines. The cause of this gender difference is unclear, but it cannot be explained by reductions in opioid use, as no gender difference in opioid use was observed. Due to social and economic disparities, women may be more likely to depend upon male partners to obtain drugs, perpetuating a pattern of using, injecting, and sharing drugs with them. Young females may be particularly vulnerable to such dependency. The physiological benefits of extended bup/nx therapy may, for some females, weaken this dependency, allowing them to change their patterns of drug use, including less IDU. Future research should test this and other potential explanations.

In terms of sexual risk, males and females were equally likely to be sexually active and not use condoms, but males were more likely to have multiple partners. This gender difference is consistent with nationally representative surveys of adolescents and young adults.^{55, 56} Having multiple sex partners, particularly concurrent partners, is associated with sexually transmitted infections,^{57, 58} and this subgroup of young males with multiple sex partners may play an important role in the transmission of HIV and other infections among drug-using youth.

As in other studies of treatment-seeking opioid abusers,^{43, 59} there was a slight drop in the rate of sexual activity during treatment. It is unclear from our results whether this was a deliberate attempt by participants to reduce their sexual activity or a reflection of their increased focus on recovery during the early part of treatment. This change was not sustained, however, as the rate of sexual activity returned to nearly baseline levels by the end of treatment. More importantly, there was no change in the rate of multiple partners or

unprotected intercourse. Many other studies have found that drug treatment is not associated with reductions in sex risk behaviors.^{10, 43, 46, 60} Thus, bup/nx and other treatments seem to have greater impact on drug risk and may be insufficient for promoting sex risk reduction.

While bup/nx therapy is an important component of HIV prevention for opioid-dependent youth, additional risk reduction counseling may be needed to promote greater decreases in both drug and sex risk behaviors. In one of the only randomized controlled trials to target drug using-adolescents, St. Lawrence and colleagues (2002) tested a 12-session group treatment based on the Information-Behavior-Motivation (IBM) Model among 161 adolescents in residential treatment.⁶¹ Compared to participants in the information only condition, participants who received both information and behavioral skills components had greater increases in condom use and sexual abstinence. Among adolescents in general, theoretically-based group interventions that are tailored to meet the needs of specific sub-groups of adolescents can effectively increase condom use and reduce number of sex partners.⁶²

To the best of our knowledge, no interventions have been developed to reduce injection drug risk among adolescents, though many trials have included adolescents and young adults. For example, early in the HIV epidemic, Des Jarlais and colleagues (1992) tested the effects of a 4-session intervention based on Social Learning Theory among heroin sniffers with a mean age of 27 years.⁶³ Participants who were randomized to the intervention rather than the control group were less likely to transition to IDU. Comprehensive reviews of the adult literature suggest that interventions can effectively reduce both drug and sex risk behaviors among IDUs.^{64–66} Successful programs have been theory-based and include the following components: HIV/AIDS education; assessment of personal risk and responsibility; behavioral skills training in safer sex and drug behaviors; development of intrapersonal skills (e.g., problem solving); reinforcement of positive changes; and discussion of practical and emotional issues related to HIV risk reduction.⁶⁴ Thus, youth receiving bup/nx or other pharmacological treatments for opioid dependence may benefit from multi-component HIV prevention services that include cognitive-affective-behavioral skills training delivered in group formats. Clinical trials are needed to test the effectiveness of such programs among youth receiving psychopharmacological treatment.

Several limitations of this study should be noted. First, the sample size was modest, although this is the largest randomized clinical trial of bup/nx treatment for adolescents conducted to date. It is also the first to examine gender differences in and the effects of bup/nx treatment on HIV risk among opioid-dependent youth. Second, the follow-up rate was 67%, but there was no difference by treatment condition or gender, and participants who completed the week 12 assessment did not differ from those who did not on baseline demographic characteristics, substance abuse, and HIV risk behavior. Third, no HIV testing was performed and self-reported HIV risk data are subject to response bias. However, the latter remains the standard assessment method for obtaining personal data, and other studies of substance abusers have documented test-retest reliability and predictive validity of self-reported sexual and drug use behaviors.^{67, 68} Finally, while the use of a convenience sample of treatment-seeking volunteers raises the possibility of selection bias, the multi-site design of this study attempts to improve generalizability. Yet, results may not generalize to adolescents who are not seeking addiction treatment or are unwilling to participate in a clinical trial.

Conclusion

Most opioid-dependent youth in this study engaged in one or more drug and sex risk behaviors that place them at heightened risk for HIV, hepatitis C, and other infectious

diseases. This is particularly concerning in light of the growing HIV epidemic among youth in the United States.⁵ Gender differences were evident, with females engaging in higher rates of injection risk behavior and males being more likely to have multiple sex partners. Extended bup/nx therapy was associated with significantly greater reductions in opioid use and IDU, and it should be considered as a safe and effective treatment option for opioid-dependent youth. However, it appears that adjunctive risk reduction counseling may be needed to promote greater decreases in injection and sex risk behaviors. Additional studies are warranted to determine the optimal length of bup/nx treatment and to identify effective behavioral risk reduction interventions for opioid-dependent youth that can be combined with extended bup/nx treatment and drug counseling.

Acknowledgments

Funding: This study was supported by the following grants from the National Institutes of Health: P30-AI064518 (Weinhold) and U10-DA013711 (Patkar) to Duke University, K05-DA017009 (Woody) and U10-DA13043 (Woody) to University of Pennsylvania, K24-DA022288 (Weiss) and U10-DA015831 (Weiss) to Harvard University, U10-013034 (Stitzer) and K12-DA000357 (Riggs) to Johns Hopkins University, and U10-DA015833 (Miller) to University of New Mexico.

We thank the staff and participants at the community treatment programs and regional research and training centers of the NIDA Clinical Trials Network for their involvement in this project, including Mountain Manor Treatment Program (Mid-Atlantic Node), Duke Addictions Program (North Carolina Node), Mercy Hospital (Northern New England Node), Addiction and Substance Abuse Program (Southwest Node), and Brandywine Treatment Center (Delaware Valley Node). We also thank the staff of the Clinical Coordinating Centers (Michele Strauss, NIDA CCC and Christopher Bertz, CCRA), the Data and Statistics Center (Duke Clinical Research Institute), and the Center for the Clinical Trials Network at NIDA for their work on this project. Finally, we acknowledge the Karen Dugosh, PhD and Kevin Lynch, PhD for their analysis of the primary outcomes.

References

1. Johnston, LD.; O'Malley, PM.; Bachman, JG.; Schulenberg, JE. National Results on Adolescent Drug Use: Overview of Key Findings, 2008. Bethesda, MD: National Institute on Drug Abuse; 2009.
2. SAMSHA. The NSDUH Report: Substance Use and Dependence Following Initiation of Alcohol or Illicit Drug Use. Rockville, MD: Substance Abuse and Mental Health Services, Office of Applied Studies; 2008.
3. Institute of Medicine. Dispelling the Myths About Addiction: Strategies to Increase Understanding and Strengthen Research. Washington, DC: National Academies Press; 1997.
4. Hser YI, Hoffman V, Grella CE, Anglin MD. A 33-year follow-up of narcotic addicts. Arch Gen Psychiatry. 2001; 58(5):503–508. [PubMed: 11343531]
5. Hall HI, Song R, Rhodes P, et al. Estimation of HIV incidence in the United States. JAMA. Aug 6; 2008 300(5):520–529. [PubMed: 18677024]
6. SAMSHA. The DASIS Report. Rockville, MD: Substance Abuse and Mental Health Services Administration, Office of Applied Studies; 2007. Heroin: Changes in how it is used, 1995 – 2005.
7. Gordon SM, Mulvaney F, Rowan A. Characteristics of adolescents in residential treatment for heroin dependence. Am J Drug Alcohol Abuse. Aug; 2004 30(3):593–603. [PubMed: 15540495]
8. Subramaniam GA, Stitzer MA. Clinical characteristics of treatment-seeking prescription opioid vs. heroin-using adolescents with opioid use disorder. Drug Alcohol Depend. Apr 1; 2009 101(1–2): 13–19. [PubMed: 19081205]
9. Hopfer CJ, Mikulich SK, Crowley TJ. Heroin use among adolescents in treatment for substance use disorders. J Am Acad Child Adolesc Psychiatry. Oct; 2000 39(10):1316–1323. [PubMed: 11026188]
10. Sullivan LE, Moore BA, Chawarski MC, et al. Buprenorphine/naloxone treatment in primary care is associated with decreased human immunodeficiency virus risk behaviors. J Subst Abuse Treat. 2008; 35(1):87–92. [PubMed: 17933486]

11. Chitwood DD, Griffin DK, Comerford M, et al. Risk factors for HIV-1 seroconversion among injection drug users: a case-control study. *Am J Public Health*. 1995; 85(11):1538–1542. [PubMed: 7485667]
12. Miller M, Neaigus A. Sex partner support, drug use and sex risk among HIV-negative non-injecting heroin users. *AIDS Care*. 2002; 14(6):801–813. [PubMed: 12511213]
13. Sánchez J, Comerford M, Chitwood DD, Fernandez MI, McCoy CB. High risk sexual behaviours among heroin sniffers who have no history of injection drug use: implications for HIV risk reduction. *AIDS Care*. 2002; 14(3):391–398. [PubMed: 12042084]
14. Meade CS, McDonald LJ, Weiss RD. HIV risk behavior in opioid dependent adults seeking detoxification treatment: an exploratory comparison of heroin and oxycodone users. *Am J Addict*. Jul–Aug; 2009 18(4):289–293. [PubMed: 19444732]
15. Cameron JL. Interrelationships between hormones, behavior, and affect during adolescence: understanding hormonal, physical, and brain changes occurring in association with pubertal activation of the reproductive axis. Introduction to part III. *Ann N Y Acad Sci*. Jun.2004 1021:110–123. [PubMed: 15251880]
16. Dahl RE. Adolescent brain development: a period of vulnerabilities and opportunities. Keynote address. *Ann N Y Acad Sci*. Jun.2004 1021:1–22. [PubMed: 15251869]
17. Collins WA, Welsh DP, Furman W. Adolescent romantic relationships. *Annu Rev Psychol*. 2009; 60:631–652. [PubMed: 19035830]
18. DiIorio C, Van Marter DF, Dudley WN. An assessment of the HIV/AIDS knowledge, attitudes, and behaviors of adolescents living in a large metropolitan area. *Ethn Dis*. Spring;2002 12(2):178–185. [PubMed: 12019925]
19. Romer D, Black M, Ricardo I, et al. Social influences on the sexual behavior of youth at risk for HIV exposure. *Am J Public Health*. 1994; 84(6):977–985. [PubMed: 8203696]
20. Prinstein MJ, Meade CS, Cohen GL. Adolescent oral sex, peer popularity, and perceptions of best friends' sexual behavior. *J Pediatr Psychol*. Jun; 2003 28(4):243–249. [PubMed: 12730281]
21. Arnett J. Contraceptive use, sensation seeking and adolescent egocentrism. *J Youth Adolesc*. Apr; 1990 19(2):171–180. [PubMed: 12288788]
22. Greene K, Krcmar M, Walters LH, Rubin DL, Jerold, Hale L. Targeting adolescent risk-taking behaviors: the contributions of egocentrism and sensation-seeking. *J Adolesc*. Aug; 2000 23(4): 439–461. [PubMed: 10936016]
23. Crone EA, Bullens L, van der Plas EA, Kijkuik EJ, Zelazo PD. Developmental changes and individual differences in risk and perspective taking in adolescence. *Dev Psychopathol*. Fall;2008 20(4):1213–1229. [PubMed: 18838039]
24. Raffaelli M, Crockett LJ. Sexual risk taking in adolescence: the role of self-regulation and attraction to risk. *Dev Psychol*. Nov; 2003 39(6):1036–1046. [PubMed: 14584983]
25. Curry LA, Youngblade LM. Negative affect, risk perception, and adolescent risk behavior. 2006; 27(5):468–485.
26. Ernst M, Nelson EE, Jazbec S, et al. Amygdala and nucleus accumbens in responses to receipt and omission of gains in adults and adolescents. *Neuroimage*. 2005; 25(4):1279–1291. [PubMed: 15850746]
27. Galvan A, Hare T, Voss H, Glover G, Casey BJ. Risk-taking and the adolescent brain: who is at risk? *Dev Sci*. Mar; 2007 10(2):F8–F14. [PubMed: 17286837]
28. Galvan A, Hare TA, Parra CE, et al. Earlier development of the accumbens relative to orbitofrontal cortex might underlie risk-taking behavior in adolescents. *J Neurosci*. Jun 21; 2006 26(25):6885–6892. [PubMed: 16793895]
29. CDC. HIV/AIDS Surveillance Report, 2006. Vol. 18. Atlanta: U.S. Department of Health and Human Services; 2008.
30. Bennett GA, Velleman RD, Barter G, Bradbury C. Gender differences in sharing injecting equipment by drug users in England. *AIDS Care*. Feb; 2000 12(1):77–87. [PubMed: 10716020]
31. Montgomery SB, Hyde J, De Rosa CJ, et al. Gender differences in HIV risk behaviors among young injectors and their social network members. *Am J Drug Alcohol Abuse*. 2002; 28(3):453–475. [PubMed: 12211360]

32. Evans JL, Hahn JA, Page-Shafer K, et al. Gender differences in sexual and injection risk behavior among active young injection drug users in San Francisco (the UFO Study). *J Urban Health*. Mar; 2003 80(1):137–146. [PubMed: 12612103]
33. Absalon J, Fuller CM, Ompad DC, et al. Gender differences in sexual behaviors, sexual partnerships, and HIV among drug users in New York City. *AIDS Behav*. Nov; 2006 10(6):707–715. [PubMed: 16676223]
34. Joe GW, Simpson DD. HIV risks, gender, and cocaine use among opiate users. *Drug Alcohol Depend*. 1995; 37(1):23–28. [PubMed: 7882870]
35. Tyndall MW, Patrick D, Spittal P, Li K, O'Shaughnessy MV, Schechter MT. Risky sexual behaviours among injection drugs users with high HIV prevalence: implications for STD control. *Sex Transm Infect*. Apr; 2002 78 (Suppl 1):i170–175. [PubMed: 12083439]
36. Metzger DS, Navaline H, Woody GE. Drug abuse treatment as AIDS prevention. *Public Health Rep*. Jun; 1998 113 (Suppl 1):97–106. [PubMed: 9722815]
37. Ling W, Amass L, Shoptaw S, et al. A multi-center randomized trial of buprenorphine-naloxone versus clonidine for opioid detoxification: findings from the National Institute on Drug Abuse Clinical Trials Network. *Addiction*. Aug; 2005 100(8):1090–1100. [PubMed: 16042639]
38. Johnson RE, Jaffe JH, Fudala PJ. A controlled trial of buprenorphine treatment for opioid dependence. *JAMA*. May 27; 1992 267(20):2750–2755. [PubMed: 1578593]
39. Fiellin DA, Pantalon MV, Chawarski MC, et al. Counseling plus buprenorphine-naloxone maintenance therapy for opioid dependence. *N Engl J Med*. Jul 27; 2006 355(4):365–374. [PubMed: 16870915]
40. Schottenfeld RS, Chawarski MC, Mazlan M. Maintenance treatment with buprenorphine and naltrexone for heroin dependence in Malaysia: a randomised, double-blind, placebo-controlled trial. *Lancet*. Jun 28; 2008 371(9631):2192–2200. [PubMed: 18586174]
41. Woody GE, Poole SA, Subramaniam G, et al. Extended vs short-term buprenorphine-naloxone for treatment of opioid-addicted youth: a randomized trial. *JAMA*. Nov 5; 2008 300(17):2003–2011. [PubMed: 18984887]
42. Marsch LA, Bickel WK, Badger GJ, Jacobs EA. Buprenorphine treatment for opioid dependence: the relative efficacy of daily, twice and thrice weekly dosing. *Drug Alcohol Depend*. Feb 14; 2005 77(2):195–204. [PubMed: 15664721]
43. Lott DC, Strain EC, Brooner RK, Bigelow GE, Johnson RE. HIV risk behaviors during pharmacologic treatment for opioid dependence: a comparison of levomethadyl acetate buprenorphine, and methadone. *J Subst Abuse Treat*. 2006; 31(2):187–194. [PubMed: 16919747]
44. Hallinan R, Byrne A, Agho K, McMahon C, Tynan P, Attia J. Erectile dysfunction in men receiving methadone and buprenorphine maintenance treatment. *J Sex Med*. Mar; 2008 5(3):684–692. [PubMed: 18093096]
45. Teusch L, Scherbaum N, Bohme H, Bender S, Eschmann-Mehl G, Gastpar M. Different patterns of sexual dysfunctions associated with psychiatric disorders and psychopharmacological treatment. Results of an investigation by semistructured interview of schizophrenic and neurotic patients and methadone-substituted opiate addicts. *Pharmacopsychiatry*. May; 1995 28(3):84–92. [PubMed: 7568370]
46. Marsch LA, Bickel WK, Badger GJ, et al. Comparison of pharmacological treatments for opioid-dependent adolescents: a randomized controlled trial. *Arch Gen Psychiatry*. Oct; 2005 62(10):1157–1164. [PubMed: 16203961]
47. NIDA. Risk Behavior Survey. 3. Rockville, MD: National Institute on Drug Abuse, Community Research Branch; 1993.
48. Fiellin DA. Treatment of adolescent opioid dependence: no quick fix. *JAMA*. Nov 5; 2008 300(17):2057–2059. [PubMed: 18984896]
49. Nelson KE, Galai N, Safaeian M, Strathdee SA, Celentano DD, Vlahov D. Temporal trends in the incidence of human immunodeficiency virus infection and risk behavior among injection drug users in Baltimore, Maryland, 1988–1998. *Am J Epidemiol*. Oct 1; 2002 156(7):641–653. [PubMed: 12244033]
50. Hser Y, Anglin MD, McGlothlin W. Sex differences in addict careers: initiation of use. *Am J Drug Alcohol Abuse*. 1987; 13:33–57. [PubMed: 3318399]

51. Rosenbaum M. Sex roles among deviants: the woman addict. *Int J Addict*. Jul; 1981 16(5):859–877. [PubMed: 7327770]
52. Dwyer R, Richardson D, Ross MW, Wodak A, Miller ME, Gold J. A comparison of HIV risk between women and men who inject drugs. *AIDS Educ Prev*. Oct; 1994 6(5):379–389. [PubMed: 7818974]
53. McKeganey N, Abel M, Taylor A, Frischer M, Goldberg D, Green S. The preparedness to share injecting equipment: an analysis using vignettes. *Addiction*. Sep; 1995 90(9):1253–1260. [PubMed: 7580822]
54. Sherman SG, Latkin CA, Gielen AC. Social factors related to syringe sharing among injecting partners: a focus on gender. *Subst Use Misuse*. Dec; 2001 36(14):2113–2136. [PubMed: 11794586]
55. Santelli JS, Lindberg LD, Abma J, McNeely CS, Resnick M. Adolescent sexual behavior: estimates and trends from four nationally representative surveys. *Fam Plann Perspect*. Jul–Aug; 2000 32(4):156–165. 194. [PubMed: 10942351]
56. Santelli JS, Brener ND, Lowry R, Bhatt A, Zabin LS. Multiple sexual partners among U.S. adolescents and young adults. *Fam Plann Perspect*. Nov–Dec; 1998 30(6):271–275. [PubMed: 9859017]
57. DiClemente RJ, Crosby RA, Wingood GM, Lang DL, Salazar LF, Broadwell SD. Reducing risk exposures to zero and not having multiple partners: findings that inform evidence-based practices designed to prevent STD acquisition. *Int J STD AIDS*. Dec; 2005 16(12):816–818. [PubMed: 16336765]
58. Crosby RA, DiClemente RJ, Wingood GM, et al. Associations between sexually transmitted disease diagnosis and subsequent sexual risk and sexually transmitted disease incidence among adolescents. *Sex Transm Dis*. Apr; 2004 31(4):205–208. [PubMed: 15028932]
59. Krupitsky EM, Zvartau EE, Masalov DV, et al. Naltrexone with or without fluoxetine for preventing relapse to heroin addiction in St. Petersburg, Russia. *J Subst Abuse Treat*. Dec; 2006 31(4):319–328. [PubMed: 17084785]
60. Gowing LR, Farrell M, Bornemann R, Sullivan LE, Ali RL. Methadone treatment of injecting opioid users for prevention of HIV infection. *J Gen Intern Med*. 2006; 21(2):193–195. [PubMed: 16336624]
61. St Lawrence JS, Crosby RA, Brasfield TL, O'Bannon RE III. Reducing STD and HIV risk behavior of substance-dependent adolescents: a randomized controlled trial. *J Consult Clin Psychol*. Aug; 2002 70(4):1010–1021. [PubMed: 12182264]
62. DiClemente RJ, Crittenden CP, Rose E, et al. Psychosocial predictors of HIV-associated sexual behaviors and the efficacy of prevention interventions in adolescents at-risk for HIV infection: what works and what doesn't work? *Psychosom Med*. Jun; 2008 70(5):598–605. [PubMed: 18541908]
63. Des Jarlais DC, Casriel C, Friedman SR, Rosenblum A. AIDS and the transition to illicit drug injection--results of a randomized trial prevention program. *Br J Addict*. Mar; 1992 87(3):493–498. [PubMed: 1559048]
64. Des Jarlais DC, Semaan S. HIV prevention for injecting drug users: the first 25 years and counting. *Psychosom Med*. 2008; 70(5):606–611. [PubMed: 18519886]
65. Copenhaver MM, Johnson BT, Lee IC, Harman JJ, Carey MP. Behavioral HIV risk reduction among people who inject drugs: meta-analytic evidence of efficacy. *J Subst Abuse Treat*. Sep; 2006 31(2):163–171. [PubMed: 16919744]
66. Semaan S, Des Jarlais DC, Sogolow E, et al. A meta-analysis of the effect of HIV prevention interventions on the sex behaviors of drug users in the United States. *J Acquir Immune Defic Syndr*. 2002; 30(Suppl 1):S73–93. [PubMed: 12107362]
67. Darke S. Self-report among injecting drug users: a review. *Drug Alcohol Depend*. Aug 1; 1998 51(3):253–263. [PubMed: 9787998]
68. Darke S, Hall W, Heather N, Ward J, Wodak A. The reliability and validity of a scale to measure HIV risk-taking among intravenous drug users. *AIDS*. 1991; 5(2):181–185. [PubMed: 2031690]

Figure 1.
Percent of participants who injected drugs in the past month by condition and gender over the course of treatment

Table 1

Participant characteristics at baseline by gender

	Females n = 61	Males n = 89	Test of difference
Age, M (SD)	19.18 (1.52)	19.13 (1.48)	t(148) = .183
Race, n (%)			$\chi^2(1) = .001$
White	44 (72.1%)	64 (71.9%)	
Other	17 (27.9%)	25 (28.1%)	
Years of education, M (SD)	11.30 (1.64)	11.15 (1.56)	t(148) = .563
Occupational status, n (%)			$\chi^2(2) = .352$
In school	11 (18.0%)	13 (14.6%)	
Working	44 (72.1%)	66 (74.2%)	
Unemployed	6 (9.8%)	10 (11.2%)	
Years of substance use, M (SD)			
Heroin	1.85 (1.73)	1.46 (1.52)	t(147) = 1.454
Methadone	0.07 (0.32)	0.24 (1.22)	t(147) = 1.055
Other opioid	1.48 (1.94)	1.49 (1.76)	t(147) = 0.036
Marijuana	3.55 (2.87)	4.10 (2.87)	t(147) = 1.150
Cocaine	1.08 (1.97)	0.47 (1.08)	t(147) = 2.440*
Alcohol	0.92 (1.70)	0.91 (1.83)	t(147) = 0.022
Days of substance use (past month), M (SD)			
Heroin	20.25 (12.75)	18.33 (13.44)	t(147) = 0.875
Methadone	1.67 (4.92)	1.45 (4.16)	t(147) = 0.290
Other opioid	9.05 (12.18)	10.62 (12.80)	t(147) = 0.752
Marijuana	8.90 (11.48)	10.15 (11.78)	t(147) = 0.640
Cocaine	3.16 (6.54)	1.70 (4.10)	t(147) = 1.690
Alcohol	0.60 (1.90)	0.83 (2.39)	t(147) = 0.627
Hepatitis C infection, n (%)	11 (18.0%)	16 (18.2%) ^I	$\chi^2(1) = 0.001$

*
p < .05^I
One person had missing data.

Table 3

GEE analyses predicting drug and sexual behaviors

	Opioid Use ¹		Injection drug use ²		Injection Risk ²		Sexual Activity ³		Multiple p partners ³		Non-condom use ³	
	Wald χ^2	p	Wald χ^2	p	Wald χ^2	p	Wald χ^2	p	Wald χ^2	p	Wald χ^2	p
<u>Step 1</u>												
Condition	16.74	<.001	3.56	.06	1.10	.30	0.34	.56	0.14	.71	1.87	.17
Time	25.76	<.001	34.30	<.001	5.05	.08	11.76	.008	0.90	.83	2.22	.53
Gender	1.23	.27	0.09	.77	10.60	.001	2.45	.12	7.61	.006	0.002	.97
<u>Step 2</u>												
Condition*Time	0.25	.88	6.83	.03	0.93	.63	6.49	.09	2.18	.54	3.88	.28
<u>Step 3</u>												
Condition*Time*Gender	0.19	.91	7.61	.02	2.90	.24	0.86	.84	7.24	.07	4.46	.22

¹Time points = week 4 (early/mid), week 8 (late/mid), and week 12 (post)

²Time points = week 0 (pre), week 4/8 (mid), and week 12 (post)

³Time points = week 0 (pre), week 4 (early/mid), week 8 (late/mid), and week 12 (post) Note: all 2-way interactions were included in the analysis but are not shown here.