CORRESPONDENCE

Physicians' Knowledge of and Compliance With Guidelines: An Exploratory Study in Cardiovascular Diseases

by Dr. rer. pol. Ute Karbach, Dr. rer. soc. Ingrid Schubert, Dr. med. Jens Hagemeister, Dr. rer. medic. Nicole Ernstmann, Prof. Dr. phil. Holger Pfaff, Prof. Dr. med. Hans-Wilhelm Höpp in volume 5/2011

REFERENCES

1. Karbach U, Schubert I, Hagemeister J, Ernstmann N, Pfaff H, Höpp HW: Physicians' knowledge of and compliance with guidelines: an exploratory study in cardiovascular diseases. *Dtsch Arztebl Int* 2011; 108(5): 61–9.

Klaus Lischka
Villingendorf, Germany
lilamed@gmx.de

Conflict of interest statement

The author declares that no conflict of interest exists.

Quality of Treatment Is Almost Identical

The authors sought to establish an association between (guideline related) knowledge of general practitioners and how competently they treated their patients. The conclusion, as befits the subject matter, could have been: insufficiently trained GPs (with inadequate knowledge of guidelines) achieve inadequate results with their treatment. The article, however, showed a different result: the quality of treatment was almost identical in both groups of physicians, whether a doctor had adequate or inadequate knowledge of the guidelines.

In spite of this finding, the conclusion was: “Increased efforts should be made—as has already happened in some integrated care projects—to implement guideline recommendations as in-process control variables in standardized software.”

The study was intended to provide proof that such software corridors are needed, but it didn't do so. And another, similarly inconsistent, result was not even included among the key messages: with respect to the four coronary heart disease (CHD)-related indicators (indicator-guided analysis), the doctors who had demonstrated inadequate knowledge in the test were actually superior in terms of the therapeutic criteria to their colleagues with better CME certifications. The most important indicator was patients' blood pressure, which was controlled better by doctors without adequate knowledge of guidelines than by those with certified knowledge. This is particularly notable because as far as the guideline test for blood pressure was concerned, only 11% of doctors had adequate knowledge of guidelines. The 89% of GPs whose knowledge of guidelines was inadequate had achieved better control of blood pressure in their patients.

The important components of the study's results were ignored and linked to the insistence on changing practices' IT systems. This cannot logically be deduced from the results, and the question is what the authors' motives might be.

DOI: 10.3238/arztebl.2011.0490a

Consultations not Informed by Mutual Trust

Guidelines are set out by experts and specialists who focus exclusively on their own specialties. Other disciplines are not included in the considerations. The result is that, for example, multimorbid patients with five conditions (not rare in this era of the International Classification of Diseases (ICD), and diagnosis related groups (DRG)), would have to be given a minimum of 15 medications. Drug interactions and adverse effects are unpredictable in such a scenario, and that does not even touch on lacking compliance. In geriatric patients, prioritization is usually required. Only few guidelines exist for patients older than 75. Medications have usually not been tested in this age group.

The authors say: “Guidelines are one of the means by which health care organizations try to improve health care and lower its cost.” The likely problem will be that for GPs, the doctor-patient relationship will be dominated by considerations of how to meet externally imposed targets (“quality indicators”) rather than the confidential and trusting consultation with the patient.

It is true that in the context of evidence based medicine, randomized controlled studies are an essential instrument for the purposes of evaluation. But significance is not always the same as clinical relevance. What may apply for a healthy 20-year-old may not apply for a multimorbid 85-year-old. In spite of our ever increasing knowledge of molecular and genetic associations in the human body, in spite of our insights into receptors, ion channels, or signal transduction, we still apply the instrument of (medication) treatment practically “blindly” to the complex networks of body and psyche, without in any way recognizing the possible long-term consequences of such therapies.

DOI: 10.3238/arztebl.2011.0490b

REFERENCES

1. Karbach U, Schubert I, Hagemeister J, Ernstmann N, Pfaff H, Höpp HW: Physicians' knowledge of and compliance with guidelines: an

exploratory study in cardiovascular diseases. *Dtsch Arztebl Int* 2011; 108(5): 61–9.

Dr. med. Robert Hector
Simmersfeld, Germany
roberthector@t-online.de

Conflict of interest statement

The author declares that no conflict of interest exists.

Barriers to Guideline Implementation

In order to investigate the high prevalence of cardiovascular risk factors in the Münster area that is still observed in spite of existing guidelines for secondary prevention of coronary heart disease (1) we conducted the COSIMA study from 2002–2004 at the Institute of Epidemiology and Social Medicine at the University of Münster (2). We asked 1023 practicing physicians organized in the administrative district of Münster of the Westphalia-Lippe Medical Association about the most important cardiovascular risk factors and about their knowledge and acceptance of guidelines in secondary coronary prevention. Our results confirm the findings described in the article (3)—namely, a higher acceptance of guideline-based medicine and a better knowledge and acceptance of guidelines among younger colleagues, internists, and doctors who take part in quality circles.

However, our results also indicate important barriers to guideline implementation. The majority of physicians primarily named organizational factors, such as a lack of reimbursement (84% agreement) and lack of time (51% agreed). Furthermore, poor patient compliance (70%) and contradictory recommendations in guidelines published by different medical societies (54%) were stated. The main criticisms of guidelines as an instrument of evidence-based medicine again focused on cost reduction and a perceived lack of flexibility in the treatment of patients.

In order to successfully implement guidelines in medical practice, a multi-level approach seems to be crucial. Local settings and circumstances, private physicians' values, and perceived barriers could be discussed by audits in the context of local quality circles in order to develop strategies for a better guideline implementation. The acceptance of guideline recommendations by treating physicians is of critical importance, and may be achieved by publishing guidelines of high scientific quality and transparency.

DOI: 10.3238/arztebl.2011.0491a

REFERENCES

1. Kotseva K, Wood D, De Backer G, De Bacquer D, Pyörälä K, Keil U: EUROASPIRE Study Group. Cardiovascular prevention guidelines in daily practice: a comparison of EUROASPIRE I, II, and III surveys in eight European countries. *Lancet* 2009; 373: 929–40.
2. Heidrich J, Behrens T, Raspe F, Keil U: Knowledge and perception of guidelines and secondary prevention of coronary heart disease among general practitioners and internists. Results from a physician

survey in Germany. *Eur J Cardiovasc Prev Rehabil* 2005; 12: 521–9.

3. Karbach U, Schubert I, Hagemeyer J, Ernstmann N, Pfaff H, Höpp HW: Physicians' knowledge of and compliance with guidelines: An exploratory study in cardiovascular diseases. *Dtsch Arztebl Int* 2011; 108(5): 61–9.

PD Dr. med. Thomas Behrens

Bremer Institut für Präventionsforschung und Sozialmedizin, Bremen, Germany
behrens@bips.uni-bremen.de

Prof. Dr. med. Ulrich Keil

Institut für Epidemiologie und Sozialmedizin, Münster, Germany

Dr. med. Jan Heidrich

Bremer Institut für Präventionsforschung und Sozialmedizin, Bremen, Germany

Conflict of interest statement

The authors declare that no conflict of interest exists.

Crucial Characteristics

Imparting guideline knowledge to doctors constitutes merely one of the building blocks for the practical therapeutic use of guideline knowledge. The path to clinical competence leads from the mere knowledge of facts to explaining the reasons/causes underlying those facts to the demonstration of skills to their independent application (1). The authors' conclusion—that physicians' knowledge of guidelines does not in itself lead to better guideline implementation—is self-evident rather than surprising from a didactic perspective.

The authors conclude that doctors' therapeutic decisions are shaped to a lesser extent by medical data and to a greater extent by organizational routines, financial framework conditions, and patient related aspects. This conclusion reflects important characteristics of general practitioners' work: including patients' preferences and collaborating specialist physicians into identifying a joint therapeutic decision (2). It cannot necessarily be assumed that these factors are equally distributed in the study practices of doctors with adequate or inadequate knowledge of guidelines. Regrettably, neither was the perspective of the patients elicited with regard to the presented guideline implementation nor were data collected on hard end points with relevance for patients (mortality, hospitalization).

To investigate the total process of different strategies for guideline implementation up to the effect on patients, cluster-randomized studies are appropriate. The different implementation strategies are randomly allocated to the practices, and the treatment of patients in the comparator groups needs to be standardized. Such studies exist for the treatment and secondary prophylaxis of cardiovascular disorders; there is a sufficiently large number of them and they are of sufficiently high quality (3). Before testing new implementation aids, existing knowledge should be systematically summarized and discussed, in manner that is appropriate to the target group and the indication.

DOI: 10.3238/arztebl.2011.0491b

REFERENCES

1. Miller GE. The assessment of clinical skills/competence/performance. *Acad Med* 1990; 65: 63–7.
2. Van Royen P, Beyer M, Chevallier P, et al.: The research agenda for general practice/family medicine and primary health care in Europe. Part 3. Results: person centred care, comprehensive and holistic approach. *Eur J Gen Pract* 2010; 16: 113–9.
3. Peters-Klimm F, Müller-Tasch T, Remppis A, Szecsenyi J, Schellberg D: Improved guideline adherence to pharmacotherapy of chronic systolic heart failure in general practice – results from a cluster-randomized controlled trial of implementation of a clinical practice guideline. *J Eval Clin Pract* 2008; 14: 823–9.
4. Karbach U, Schubert I, Hagemeister J, Ernstmann N, Pfaff H, Höpp HW: Physicians' knowledge of and compliance with guidelines: an exploratory study in cardiovascular diseases. *Dtsch Arztebl Int* 2011; 108(5): 61–9.

Dr. rer. nat. Susanne Unverzagt

Dr. med. Andreas Klement

Institut für Medizinische Epidemiologie, Biometrie und Informatik, Halle, Germany
susanne.unverzagt@medizin.uni-halle.de

Conflict of interest statement

The authors declare that no conflict of interest exists.

Incomprehensible Jargon

I have collaborated on some current guidelines (the German Program for Disease Management Guidelines on diabetic foot, diabetic nephropathy, etc). I have often asked myself why guidelines that are well intended in principle have a problem with their practical implementation. I regard the above article as a good example for the fact that a communication problem exists between those in charge of setting out guidelines and the doctor for whom they are intended. Might the lacking implementation of clinical guidelines be explained by the incomprehensible “sociolect” used in health services research or health economics, which is used in the general sphere of guideline business? On reading the article I notice how far the jargon in use (“indicator-guided analysis of patient data to determine physicians’ adherence to guidelines”) is removed from clinical practice. I think readers’ time is too limited to engage fully with an article that is written in such a complicated way. In my opinion the same holds true for many guidelines that provide no reading pleasure because the language and style they are written in is too removed from clinical practice.

Improvements could certainly be made in order to make guidelines more interesting for readers and to increase “the knowledge of guidelines as a valid surrogate parameter for the objective ‘guideline-conform clinical practice.’”

DOI: 10.3238/arztebl.2011.0492a

REFERENCES

1. Karbach U, Schubert I, Hagemeister J, Ernstmann N, Pfaff H, Höpp HW: Physicians' knowledge of and compliance with guidelines: an exploratory study in cardiovascular diseases. *Dtsch Arztebl Int* 2011; 108(5): 61–9.

Prof. Dr. med. Gerhard Rümenapf

Diakonissen-Stiftungs-Krankenhaus, Speyer, Germany
gerhard.rumenapf@diakonissen.de

Conflict of interest statement

Professor Rümenapf has received CME support from Jotext and honoraria from Mölnlycke.

Multimorbidity as a Dilemma

I completely agree with the authors’ conclusion, that therapeutic decisions are based on “other practice-relevant factors”; however, this is with regard to guidelines and NOT to “clinical data.”

The authors mention “internal organizational routines of the physician’s office” first, then “financial parameters,” with “patient-related aspects” mentioned last. The fact that they recommend “to implement guideline recommendations as in-process control variables in standardized software” is consistent with this.

From my perspective as a general practitioner the practice relevant factors are weighted in exactly the opposite order: patient-related aspects come first.

The most important aspect that hinders guideline-conform treatment is patients’ multimorbidities. These consist of many clinical data.

The second is people’s insistent adherence to their own habits, and the slight unease that most people develop when they have to take more than three tablets a day.

This is an obstacle to guideline-conform treatment of hypertension, which—as is well known— patients themselves do not notice .

I would ask to be spared further bureaucratization (“to implement guideline recommendations as in-process control variables in standardized software”). In my opinion it would make sense to set out overriding guidelines for how to deal with patients with multimorbidities. Geriatric medicine provides a good example in this context. Medical pr

actice based on geriatric principles often leads to minimalizing the instructions from other guidelines. For the individual physician faced with the dilemma of multimorbidity this provides substantial relief.

DOI: 10.3238/arztebl.2011.0492b

REFERENCES

1. Karbach U, Schubert I, Hagemeister J, Ernstmann N, Pfaff H, Höpp HW: Physicians' knowledge of and compliance with guidelines: an exploratory study in cardiovascular diseases. *Dtsch Arztebl Int* 2011; 108(5): 61–9.

Dr. med. Julia Beller

Weilheim, Germany
jubelokt@aol.com

Conflict of interest statement

The author declares that no conflict of interest exists.

In Reply:

We thank our correspondents for their interest in and critical comments on our study. Behrens and coauthors mostly confirm our results in their own study of guideline knowledge and implementation in secondary coronary prevention. In contrast to our data, however, they found that knowledge and acceptance of guidelines were better in younger physicians, specialists in internal medicine, and doctors who participated in quality circles. Although we suspected similar barriers to guideline implementation as Behrens et al, we were not able to investigate this question explicitly in our current project owing to scarce funding. We unreservedly agree with the comment that implementation of guidelines needs to happen at multiple levels and should acknowledge the known barriers in the sense of a complex intervention. In this sense we also understand the comment by Unverzagt and Klement—that cognitive communication of guideline recommendations is only one building block in the implementation chain and that the patient's perspective and the influence of the specialist should be considered in general practice. This is in principle identical to our own opinion. However, since in our study we attempted to prove the benefits neither of a guideline or therapeutic recommendation nor of a particular implementation strategy, we did not conduct a randomized controlled study with hard fixed end points.

Rümenapf names communication problems because of incomprehensible specialist jargon as a possible barrier to implementation. We agree in the sense that guideline authors should use language that reflects clinical practice and is concrete.

Beller and Hector mention the multimorbidity of elderly patients and the often associated limitations regarding the extent to which therapeutic recommendations can be implemented by doctors, but also by patients, as a crucial reason for why the treatment did not meet the expected guideline recommendations. Their criticism is that neither the guidelines nor our study paid enough attention to these patient related aspects. However, our study did not focus on implementation barriers from the patients' perspective. We did, however, take this aspect into consideration in that we did not include treatment data from patients older than 79 in our evaluations. Fundamentally we agree with Beller and Hector that the topic of multimorbidity has not received enough attention in guidelines—because of the

logic inherent in guideline development, which relies on high-level evidence from studies and therefore does not sufficiently include the general practice clientele. This problem is currently the subject of intense interdisciplinary debate among guideline authors—for example, in developing the National Disease Management Guidelines Program (“German DM-CPG Program”) or even in general practice guidelines (1). Whether multimorbidity is the crucial reason for deviating from guideline recommendations, however, remains to be investigated.

Our considerations on implementing guideline recommendations as in-process control variables in standardized software into to support doctors' decision making triggered fears of increasing bureaucratization of doctors' everyday clinical practice. Lischka furthermore questions the motives of our research team and suspects that the primary reasons for our study was to establish reasons for implementing “software corridors”. This was not our study objective but the conclusion we drew from our observation that both groups of doctors saw room for improvement in implementing the recommendations selected for our study. In any case, as Unverzagt and Klement say, retrievable guideline knowledge in itself cannot be used as a surrogate parameter for guideline-conform clinical practice. How to implement evidence based knowledge is a major challenge facing us all—as evidenced by the discussions at the 15th annual meeting of the German Agency for Quality in Medicine (*Ärztliches Zentrum für Qualität, AQuMed*).

DOI: 10.3238/arztebl.2011.0493

REFERENCES

1. Hausärztliche Leitliniengruppe Hessen. Leitlinie Geriatrie, Teil I Version 1.00 vom 17.11.2008 und Teil II, Version 1.02 vom 14.09.2009 [www.pmvforschungsguppe.de/pdf/03_publicationen/geriatrie1_II.pdf]
2. Karbach U, Schubert I, Hagemeister J, Ernstmann N, Pfaff H, Höpp HW: Physicians' knowledge of and compliance with guidelines: an exploratory study in cardiovascular diseases. *Dtsch Arztebl Int* 2011; 108(5): 61–9.

Dr. rer. pol. Ute Karbach

Institut für Medizinsoziologie, Versorgungsforschung und Rehabilitationswissenschaft der Humanwissenschaftlichen Fakultät und der Medizinischen Fakultät der Universität zu Köln
Ute.Karbach@uk-koeln.de

Conflict of interest statement

The author declares that no conflict of interest exists.