Published in final edited form as:

Org Lett. 2011 August 5; 13(15): 4144–4147. doi:10.1021/ol201704g.

Stereoselective Synthesis of the Eastern Quinolizidine Portion of Himeradine A

Nathan D. Collett and Rich G. Carter*

Department of Chemistry, Oregon State University, Corvallis, OR 97331

Abstract

The synthesis of the C₁₅-C₁₇/N₁'-C₁₁' quinolizidine portion of himeradine A is disclosed. An intramolecular, heteroatom Michael addition was employed to establish the C₆' stereogenic center with high diastereoselectivity. The quinolizidine ring was constructed using microwave-induced cyclization at the N₁'-C₂' position. The C₁₇ stereogenic center was introduced through a diastereoselective Overman rearrangement.

In 2003, Kobayashi and co-workers reported the isolation and structural determination of the alkaloid himeradine A (**1**) from the club moss *Lycopodium chinense* in small amounts (2 mg, 0.001%, Figure 1).¹ Polycycle **1** was assigned based on extensive 2D NMR techniques (COSY, HOHAHA, HMQC, HMBC and HMQC-HOHAHA); however, they were unable to establish the correlation between the eastern and western halves or the absolute stereochemistry of the molecule. Alkaloid **1** showed modest cytotoxicity against murine lymphoma L1210 cells (IC₅₀ 10 μg/mL), but a thorough biological screening of the compound has not been reported. Other members of the lycopodium family have shown intriguing biological activity.²

The heptacyclic himeradine A (**1**) possesses a challenging array of structural features: ten stereogenic centers including one all-carbon quaternary center, a densely packed pentacyclic western half, the C₈'-C₁₀'-*trans*-disubstituted quinolizidine and the challenging sigma linkage at C₁₁' (Figure 1). The carbon framework of the western portion is similar to that found in lycopodine (**2**);³ however, the additional challenge of the C₃-C₁₄ linkage increases the complexity of any synthetic endeavor. A related structural scaffold can be found in fastigiatine (**3**).⁴ The relative stereochemistries of the eastern quinolizidine ring systems are distinct from typical stereochemical combinations found in related natural products such as cermizine D (**4**).⁵ In particular, the axially disposed C₁₀' moiety creates added synthetic challenges. To date, no synthetic endeavors have been published towards himeradine A.⁶ Herein, we disclose the synthesis of the eastern portion of himeradine A and the

rich.carter@oregonstate.edu.

Supporting Information Available Complete experimental procedures are provided, including ¹H and ¹³C spectra, of all new compounds. X-ray crystallographic data (CIF) for compound **25** is also provided. This material is available free of charge via the Internet at <http://pubs.acs.org>.

development of a viable model coupling strategy for incorporation of the C₁₅-C₁₇ carbons including the C₁₇ stereogenic center.

Our retrosynthetic strategy is shown in Scheme 1. For the polycyclic western portion of himeradine, we intend to adapt our previously developed route to lycopodine for the construction of the carbon framework.^{3b, c} Compound **5** will be formed via a diastereoselective Overman rearrangement⁷ of the trichloroimidate derived from alcohol **6**. The required C₁₅ stereogenic center will be introduced through an organozinc addition to an α,β -unsaturated aldehyde. This aldehyde will be accessible in turn from a Julia-Kocienski-Blakemore olefination⁸ with the known 1-phenyl-1H-tetrazol-5-yl (PT) sulfone **8**⁹ and aldehyde **9** followed by acid-catalyzed deacetalization with *in situ* alkene migration. The quinolizidine ring system **9** will be formed via N₁-C₂ lactamization strategy. The piperidine ring system **10** will be constructed from a substrate-controlled, intramolecular heteroatom Michael addition of enal **11**.

Synthesis of cyclization precursor **11** is shown in Scheme 2. The key C_{10'} stereocenter was constructed from a Grignard addition of organomagnesium species **12**¹⁰ with the known Ellman imine **13**¹¹ in good levels of diastereoselectivity and chemical yield. Removal of the sulfoxamine under acid-catalyzed conditions followed by Cbz protection revealed the carbamate **15**. Cross metathesis of alkene **15** with crotonaldehyde (**16**) cleanly provided the enal **11**. We have previously shown that cross metathesis of monosubstituted alkenes with α,β -unsaturated carbonyl compounds proceeds in higher chemical yield when the unsaturated carbonyl compound contains a β -methyl substitution (e.g. **16**).^{3b-c,12}

The key Lewis acid-catalyzed intramolecular, heteroatom Michael addition is shown in Scheme 3. We had hypothesized that the C_{8'} and C_{10'} stereogenic centers would work in concert to direct to facial attack on an α,β -unsaturated oxonium ion, as shown in possible transition state model **17**. The planar carbamate at C_{10'} should force the CH₂OBn moiety into a pseudoaxial position in the transition state in order to minimize steric repulsions with the Cbz moiety.¹³ We were pleased to see that our hypothesis proved accurate as aldehyde **10** was isolated as the sole diastereomer (>20:1 dr) in excellent chemical yield (98%).

The next hurdle was the construction of the quinolizidine ring system (Scheme 4). Wittig olefination of aldehyde **10** provided the α,β -unsaturated ester **21** in excellent yield. We had anticipated that hydrogenation of **21** would not only reduce the alkene, but also induce Cbz deprotection and debenzoylation at C_{11'}. To our surprise, only two of the three predicted events occurred – delivering the benzyl ether **22** as the sole product in high yield. Lactam formation of **22** (or its corresponding carboxylic acid) under a variety of conditions proved unusually challenging. Fortunately, we ultimately identified that thermolysis under microwave irradiation cleanly induced lactam formation in high yield. The synthetic challenge with this bond formation may be due the required placement of the C_{10'} substituent in the axial orientation to facilitate C₂-N₁ bond formation. Interestingly, heating of amine **22** in refluxing xylenes (approximately 140 °C) using an oil bath provided the product **23** in significantly lower yield (40% conversion after 48 h). Microwave irradiation's ability to provide purely rotational energy transfer¹⁴ may facilitate this transformation more efficiently than standard thermal conditions – thereby inducing rotation of the sidearm into close proximity for the 2° amine to attack the C₂ ester. Next, hydrogenation of the benzyl ether **23** under essentially the same conditions as employed previously with compound **21** provided the free alcohol **24**. One possible explanation for the divergent reactivity (compounds **21** vs. **23**) may be the presence of the free amine in compound **22**, which may poison the catalyst. The stereochemistry of the newly formed quinolizidine ring system was conclusively established by x-ray crystallographic analysis on the thiolactam **25**.¹⁵ Oxidation of 1° alcohol **24** under buffered Dess-Martin conditions revealed the aldehyde **9**.

We had feared the aldehyde **9** might be prone to epimerization; however, **9** appeared to be configurationally stable at C_{10'} – likely due to the presence of the lactam carbonyl again disfavoring placement of the C_{10'} substituent in an equatorial position due to A^{1,3} strain.¹³

With the aldehyde **9** in hand, we shifted our attention to the incorporation of the C₁₅-C₁₇ carbon atoms and the C₁₇ stereogenic center (Scheme 5). Olefination of aldehyde **9** with the known PT sulfone **8**⁹ gave the desired β,γ-unsaturated acetal in good yield with an inconsequential 14:1 *E:Z* selectivity. Treatment of acetal **7** under aqueous acidic conditions induced both acetal deprotection and alkene isomerization to cleanly provide the desired α,β-unsaturated aldehyde **26** with excellent *E:Z* selectivity (>20:1). *N*-Methyl diphenylprolinol catalyzed addition of Et₂Zn to the aldehyde **26** revealed the allylic alcohol **27** in 10:1 dr.^{16,17} Formation of the trichloroimidate was accomplished using DBU as the base followed by thermolysis at 90°C in the presence of K₂CO₃ to cleanly generate the rearranged amide **5**. The presence of the carbonate base appeared to prevent decomposition under the reaction conditions.¹⁸

In summary, the synthesis of the C₁₅-C₁₇/N₁'-C₁₁' quinolizidine core of himeradine A has been accomplished. Key steps in the synthetic sequence include a diastereoselective Overman rearrangement, a substrate-controlled, intramolecular heteroatom Michael addition and a microwave-induced lactam formation. Further studies towards the total synthesis of himeradine A and other related *lycopodium* alkaloids will be reported in due course.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Financial support was provided by the National Institutes of Health (NIH) (GM63723). The National Science Foundation (CHE-0722319) and the Murdock Charitable Trust (2005265) are acknowledged for their support of the NMR facility. We thank Dr. Lev N. Zakharov (OSU and University of Oregon) for X-ray crystallographic analysis of compound **25**, Professor Max Deinzer and Dr. Jeff Morré (OSU) for mass spectra data. Finally, the authors are grateful to Professor James D. White (OSU) and Dr. Roger Hanselmann (Rib-X Pharmaceuticals) for their helpful discussions.

References

1. Morita H, Hirasawa Y, Kobayashi J. *J Org Chem.* 2003; 68:4563–4566. [PubMed: 12762772]
2. (a) Ma X, Gang DR. *Nat Prod Rep.* 2004; 21:752–772. [PubMed: 15565253] (b) Kobayashi J, Morita H. *Alkaloids.* 2005; 61:1–57. [PubMed: 16173399] (c) Hirasawa Y, Kobayashi J, Morita H. *Heterocycles.* 2009; 77:679–729.
3. Structure Determination: (a) Ayer WA, Iverach GG. *Tetrahedron Lett.* 1962; 3:87–92. Enantioselective total syntheses: (b) Yang H, Carter RG, Zakharov LN. *J Am Chem Soc.* 2008; 130:9238–9239. [PubMed: 18582046] (c) Yang H, Carter RG. *J Org Chem.* 2010; 75:4929–4938. [PubMed: 20586477] Racemic Total Syntheses: (d) Stork G, Kretschmer RA, Schlessinger RH. *J Am Chem Soc.* 1968; 90:1647–1648. [PubMed: 5636803] (e) Ayer WA, Bowman WR, Joseph TC, Smith P. *J Am Chem Soc.* 1968; 90:1648–1650. [PubMed: 5636804] (f) Kim S, Bando Y, Horii Z. *Tetrahedron Lett.* 1978:2293–2294. (g) Heathcock CH, Kleinman EF, Binkley ES. *J Am Chem Soc.* 1982; 104:1054–1068. (h) Schumann D, Mueller HJ, Naumann A. *Liebigs Ann Chem.* 1982:1700–1705. (i) Kraus GA, Hon YS. *Heterocycles.* 1987; 25:377–386. (j) Grieco PA, Dai Y. *J Am Chem Soc.* 1998; 120:5128–5129. Formal Syntheses: (k) Padwa A, Brodney MA, Marino JP Jr, Sheehan SM. *J Org Chem.* 1997; 62:78–87. [PubMed: 11671366] (l) Mori M, Hori K, Akashi M, Hori M, Sato Y, Nishida M. *Angew Chem Int Ed.* 1998; 37:637–638.
4. Structure Determination: (a) Gerard RV, MacLean DB, Fagianni R, Lock CJ. *Can J Chem.* 1986; 64:943–949. (b) Gerard RV, MacLean DB. *Phytochemistry.* 1986; 25:1143–1150. Total Synthesis: (c) Liao BB, Shair MD. *J Am Chem Soc.* 2010; 132:9594–9595. [PubMed: 20568766]

5. Structure Determination: (a) Morita H, Hiraswa Y, Shinzato T, Kobayashi J. *Tetrahedron*. 2004; 60:7015–7023. Total Synthesis: (b) Nishikawa Y, Kitajima M, Takayama H. *Org Lett*. 2008; 10:1987–1990. [PubMed: 18429621] (c) Nishikawa Y, Kitajima M, Kogure N, Takayama H. *Tetrahedron*. 2009; 65:1608–1617.
6. Collett, ND.; Carter, RG. 241st National American Chemical Society Meeting; Anaheim, CA. March 27–31, 2011; p. ORGN-400 Saha, M.; Carter, RG. 241st National American Chemical Society Meeting; Anaheim, CA. March 27–31, 2011; p. ORGN-263
7. (a) Overman LEJ. *Am Chem Soc*. 1974; 96:597–599. (b) Overman LE, Carpenter NE. *Org React*. 2005; 66:1–107. (c) Kitamoto K, Sampei M, Nakayama Y, Sato T, Chida N. *Org Lett*. 2010; 12:5756–5759. [PubMed: 21105709]
8. (a) Kocienski PJ, Bell A, Blakemore PR. *Synlett*. 2000:365–366. (b) Blakemore PR. *J Chem Soc, Perkin Trans*. 2002; 1:2563–2585.
9. Lear MJ, Hiram M. *Tetrahedron Lett*. 1999; 40:4897–900.
10. Comins DL, Libby AH, Al-awar RS, Foti CJ. *J Org Chem*. 1999; 64:2184–2185.
11. Tang TP, Volkman SK, Ellman JA. *J Org Chem*. 2001; 66:8772–8778. [PubMed: 11749605]
12. Carlson EK, Rathbone LK, Yang H, Collett ND, Carter RG. *J Org Chem*. 2008; 73:5155–5158. [PubMed: 18529081]
13. (a) Watson PS, Jiang B, Scott B. *Org Lett*. 2000; 2:3679–3681. [PubMed: 11073674] (b) Martin R, Murruzzu C, Pericas MA, Riera A. *J Org Chem*. 2005; 70:2325–2328. [PubMed: 15760222] (c) Coombs TC, Lushington GH, Douglas J, Aubé J. *Angew Chem Int Ed*. 2011; 50:2734–3737.
14. Loupy, A., editor. *Microwaves in Organic Synthesis*. Wiley-VCH Verlag GmbH & Co; Weinheim: 2002. p. 1–73.
15. See supporting information of crystallographic information.
16. (a) Soai K, Ookawa A, Kaba T, Ogawa K. *J Am Chem Soc*. 1987; 109:7111–7115. (b) Shibata T, Tabira H, Soai K. *J Chem Soc Perkin Trans*. 1998; 2:177–178. (c) Nishiyama T, Kusumoto Y, Okumura K, Hara K, Kusaba S, Hirata K, Kamiya Y, Isobe M, Nakano K, Kotsuki H, Ichikawa Y. *Chem Eur J*. 2010; 16:600–610.
17. Stereochemical assignment at C₁₅ was established via advanced Mosher ester analysis as detailed in the supporting information. Ohtani I, Kusumi T, Kashman Y, Kakisawa H. *J Am Chem Soc*. 1991; 113:4092–4096.
18. Nishikawa T, Asai M, Ohyabu N, Isobe M. *J Org Chem*. 1998; 63:188–192. [PubMed: 11674062]

Figure 1.
Himeradine A and Related *Lycopodium* Alkaloids.

Scheme 1.
Retrosynthesis.

Scheme 2.
Synthesis of Cyclization Precursor.

Scheme 3.
Intramolecular Heteroatom Michael Addition.

Scheme 4.
Synthesis of the Quinolizidine Core.

Scheme 5.
Incorporation of the C₁₅-C₁₇ Portion.