Published in final edited form as:

Ophthalmology. 2011 August ; 118(8): 1603–1611. doi:10.1016/j.opththa.2011.04.036.

CENTRAL AND HEMI-CENTRAL RETINAL VEIN OCCLUSION:

Role of Anti-platelet Aggregation Agents and Anticoagulant

Sohan Singh Hayreh, MD, PhD¹, Patricia A. Podhajsky, BSN¹, and M. Bridget Zimmerman, PhD²

¹Department of Ophthalmology and Visual Sciences, College of Medicine, University of Iowa, Iowa City, Iowa

²Department of Biostatistics, College of Public Health, University of Iowa, Iowa City, Iowa

Abstract

Objective—To investigate systematically the role of anti-platelet aggregating drugs or anticoagulant in central retinal vein occlusion (CRVO) and hemi-CRVO.

Design—Cohort study.

Participants—Consecutive 686 patients with CRVO (567 patients, 585 eyes) and non-ischemic hemi-CRVO (119 patients, 122 eyes).

Methods—At first visit, all patients had a detailed ophthalmic and medical history (including the use of anti-platelet aggregating drugs or anticoagulants), and comprehensive ophthalmic and retinal evaluation. Visual evaluation was done by recording visual acuity, using the Snellen visual acuity chart, and visual fields with a Goldmann perimeter. The same ophthalmic evaluation was performed at each follow-up visit. CRVO and hemi-CRVO were classified into non-ischemic and ischemic at initial visit.

Main Outcome Measures—Visual acuity, visual fields and severity of retinal hemorrhages.

Results—For all three type of CRVO, there was a significantly greater severity of retinal hemorrhages among aspirin users than non-users ($p < 0.001$). Initial visual acuity and visual fields were significantly worse in aspirin users than non-users in non-ischemic CRVO and hemi-CRVO, but did not differ for ischemic CRVO. Among the patients with non-ischemic CRVO that initially presented with 20/60 or better visual acuity, there was a significant association of aspirin use with visual acuity deterioration. The odds ratio of visual acuity deterioration, after adjusting for age, diabetes, ischemic heart disease, and hypertension, for aspirin users relative to non-users was 2.24 (95% CI (confidence interval): 1.14, 4.41; $p = 0.020$). Of those in whom macular edema resolved, over-all cumulative visual acuity outcome also suggested a higher percentage with deterioration among aspirin users, with the odds ratio for deterioration of 3.62 (95% CI: 0.97, 13.54; $p = 0.05$) for aspirin users relative to non-users. For the non-ischemic CRVO that had 20/70 or worse visual acuity at initial visit, after resolution of macular edema, improvement in visual acuity was less likely in the aspirin users than non-users (odds ratio 0.18; 95% CI: 0.04, 0.72; $p = 0.016$).

© 2011 American Academy of Ophthalmology, Inc. Published by Elsevier Inc. All rights reserved

Correspondence to: Dr. S.S. Hayreh, Department of Ophthalmology and Visual Sciences, University Hospitals & Clinics, 200 Hawkins Drive, Iowa City, Iowa 52242-1091 Telephone No. 319-356-2947. Fax No. 319-353-7996 sohan-hayreh@uiowa.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

The authors have no conflict of interest.

Conclusion—Findings of this study indicate that, for patients with CRVO and hemi-CRVO, the use of aspirin, other antiplatelet aggregating agents or anticoagulants was associated with a worse visual outcome, and no apparent benefit.

One of us (SSH), who has dealt with more than a thousand patients with central retinal vein occlusion (CRVO) during the past 45 years, has consistently found a fixed conviction among most ophthalmologists that aspirin or anticoagulants are beneficial in this. The genesis of this widespread belief in antiplatelet aggregation agents (most commonly aspirin) or anticoagulants for CRVO and other types of retinal vein occlusion comes from their proven effectiveness in major systemic venous thrombotic disorders (e.g., deep vein thrombosis); it has been assumed that they must be equally beneficial in various types of retinal vein occlusion.

Holmin and Ploman¹ in 1938 first used heparin in one case with CRVO and claimed that that improved the visual outcome. Following that, for almost 3 decades, a series of reports of the use of anticoagulants appeared in the literature^{2–20}, some claiming varying degrees of success while others believed it made no significant difference to the outcome as compared to untreated cases, or even was harmful. A review of these studies showed no definite proof that anticoagulants cured or significantly benefited CRVO. Moreover, in most of the studies, all types of RVOs were lumped together. In spite of the wide use of aspirin in CRVO, because of its use in systemic venous thrombotic disorders, there is actually little firm evidence for it in the literature. Thus, question is far from settled whether anticoagulants or anti-platelet aggregation agents in CRVO or hemi-CRVO are beneficial, have no effect or are harmful.

Since CRVO is a common visually disabling disease, with no definitely proven treatment so far, we conducted a comprehensive prospective study to find out the effect of this common mode of therapy on the clinical course and visual outcome in a large cohort of CRVO and hemi-CRVO.

Patients and Methods

We conducted a large, comprehensive, prospective study dealing with all aspects of CRVO and hemi-CRVO systematically in the Ocular Vascular Clinic at the University of Iowa Hospitals and Clinics, a tertiary care center, from 1973. This study was part of a prospective study on Ocular Vascular Occlusive Disorders funded by the National Institute of Health (RO1 grant), approved by the Institutional Review Board. Evaluation of various modes of treatments advocated for CRVO (including aspirin or anticoagulants) was part of the originally planned study and grant. This is because aspirin was advocated commonly as the treatment of choice for thrombosis, including retinal vein occlusion. In this segment of that study, we investigated the effect of aspirin on the clinical course and visual outcome in CRVO and hemi-CRVO. The patients were first seen in our clinic from 1973 to 2000. This study included 478 patients with non-ischemic CRVO, 89 with ischemic CRVO, and 119 with non-ischemic hemi-CRVO, who were either taking aspirin or anticoagulant treatment when they were first seen, or those that did not take aspirin or anti-coagulants. The anti-platelet aggregating drug primarily used by the patients in this study was aspirin and only a rare case took Dipyridamole during the very early stages of the study in early 1970s only. Table 1 gives the number of patients that took aspirin only and those that did not take aspirin or anti-anticoagulant (warfarin) that fulfilled our inclusion criteria for this study and their demographic characteristics. In addition, there were 24 patients who were taking anticoagulant warfarin, 14 with non-ischemic CRVO, 6 with non-ischemic hemi-CRVO, and 4 with ischemic CRVO. Ischemic hemi-CRVO was not analyzed due to small sample size limitation.

Inclusion Criteria—Only those patients for whom a definite diagnosis of CRVO or hemi-CRVO could be made were included. Both CRVO and hemi-CRVO consist of two distinct entities: non-ischemic and ischemic^{21–28}.

Distinction between ischemic and non-ischemic CRVO—The criteria to differentiate ischemic from non-ischemic CRVO were recently discussed in detail in our paper dealing with natural history of visual outcome in CRVO²⁹, and the subject is not discussed here to avoid repetition. Briefly, this was based on aggregated information from data acquired from six tests (*4 functional tests*: visual acuity, visual fields, relative afferent pupillary defect and electroretinography; and *2 morphological tests*: ophthalmoscopy and fluorescein fundus angiography), which we evaluated in a specifically designed study²⁸. Of the six tests, combined information from the relative afferent pupillary defect and electroretinography differentiated 97% of the cases. Thus, relative afferent pupillary defect and electroretinography often played the most important deciding role in determining the type of CRVO. Fluorescein fundus angiography (when it was of reliable quality) also provided reliable information in differentiation. Other tests often provided confirmatory evidence. No one test was decisive *per se*.

Distinction between ischemic and non-ischemic hemi-CRVO—Since only half of the retina or even less is involved in this clinical entity, in contrast to CRVO, visual acuity, relative afferent pupillary defect and electroretinography are not reliable tests to differentiate the two types of hemi-CRVO. Primary reliance was placed on the extensive retinal capillary non-perfusion in the involved segment. The extent of visual field loss also provided useful information.

Exclusion criteria—We excluded all other retinopathies mimicking CRVO or hemi-CRVO. All patients with inadequate information or doubtful diagnosis were excluded. Patients who had any retinal or optic nerve lesion or any other factor (e.g., cataract), including any treatment for CRVO or hemi-CRVO which could have influenced the visual status, were excluded. CRVO and hemi-CRVO patients with only background diabetic retinopathy were included, but those who had active neovascularization, vitreous hemorrhages, traction detachment or other complications influencing the visual acuity or fields were excluded. Patients who had a diagnosis of glaucoma and visual field loss were excluded; however, those with elevated intraocular pressure with documented normal visual fields prior to the onset of CRVO were included. Eyes with unreliable visual fields were excluded.

STUDIES PERFORMED

At the initial visit, all patients were seen by one of us (SSH) in the Ocular Vascular Clinic and had a detailed ocular and medical history, including all the medicines they were taking, especially use of anticoagulants (warfarin) or anti-platelet aggregating agent (Aspirin or Dipyridamole), as well as a comprehensive bilateral ophthalmic evaluation. This included, (i) careful testing of the best corrected visual acuity using the Snellen visual acuity chart, (ii) visual field plotting with a Goldmann perimeter (using I-2e, I-4e and V-4e targets regularly), (iii) intraocular pressure recording with a Goldmann applanation tonometer, (iv) relative afferent pupillary defect, (v) a thorough anterior segment examination, including slit lamp examination of the anterior segment, lens and vitreous, (vi) a meticulous fundus evaluation by direct and indirect ophthalmoscopy, and, if required, by contact lens, (vii) stereoscopic color fundus photography, (viii) stereoscopic fluorescein fundus angiography (only in the involved eye), and (ix) electroretinography. For this study, the data were collected prospectively and systematically on visual acuity, visual fields, and retinal changes, particularly retinal hemorrhages and macular edema.

In addition, the patients had a full systemic evaluation performed either by an internist at the University of Iowa Hospitals & Clinics, or by their local internist/physician.

At each follow-up visit, the same ophthalmic evaluation and stereoscopic color fundus photography were performed, except that fluorescein fundus angiography was performed only when considered essential. Electroretinography was usually performed only during the initial visit for CRVO; however, if a change in status of CRVO from non-ischemic to ischemic was suspected, it was repeated. Both eyes were examined at each visit.

Follow-up protocol for all patients—All patients were followed (by SSH) according to a protocol practiced in this Clinic for CRVO and hem-CRVO patients - at about 3 monthly intervals for 3 visits, then 6 monthly intervals for 4 visits, then annually. Both eyes were examined at each visit. No intervention of any kind whatsoever was undertaken in this cohort of patients.

VISUAL STATUS EVALUATION

Visual acuity—Best corrected visual acuity was tested using the Snellen visual acuity chart and under identical testing conditions, almost invariably by the same person (SSH), encouraging the patient to look around and take his/her own time in responding, to ensure that the testing provided the most reliable information about the visual acuity. The following steps of visual acuity were checked: 20/15, 20/20, 20/25, 20/30, 20/40, 20/50, 20/60, 20/70, 20/80, 20/100, 20/200, 20/400, counting fingers (CF), hand motion (HM), perception of light (PL), and no perception of light (NPL).

Visual fields—Throughout this study, we used manual kinetic perimetry. Automated perimetry did not exist when we started the study; moreover, the changing face of automated perimetry would make such long-term studies difficult - indeed, automated perimetry is still evolving. Both types of perimetry have their advantages and disadvantages, which are discussed elsewhere.^{30,31} Visual field plotting was attempted in all patients with a visual acuity of hand motion or better at all visits, with a Goldmann perimeter using I-2e, I-4e and V-4e targets regularly, although occasionally other targets were used if it was felt that that would provide additional information for evaluation of the visual status. In a few patients, we could not do the visual fields at the initial visit for various logistic reasons. The method of testing visual fields is described in detail elsewhere.^{30,31}

The central visual field was also tested by using the Amsler grid chart, which provided very useful information about the central defect, particularly metamorphopsia, which is usually missed on perimetry.

Evaluation of changes in the visual acuity and visual field defects—Each was evaluated separately in a masked fashion, i.e., changes in visual acuity and visual fields were evaluated independently of each other, as well as independent of assessment of the severity of retinal hemorrhages, and presence or absence of macular edema, so that the severity of one did not influence the evaluation of the other.

Visual acuity evaluation—A change of at least 3 lines in the Snellen visual acuity chart was considered a significant change, which is equivalent to a logMAR (logarithm of the minimum angle of resolution) change of at least 0.30. We divided visual acuity into two categories for evaluation purposes. (i) Normal visual acuity was defined as 20/30 or better, because that category cannot show an improvement of 3 lines to achieve 20/20. (ii) As in our natural history study on visual outcome in non-arteritic anterior ischemic optic neuropathy³² and CRVO²⁹, and in the Ischemic Optic Neuropathy Decompression Trial study³³, poor

visual acuity was defined as 20/70 or worse, because that is markedly disabling and persons with that degree of disability are not considered fit to drive a vehicle.

Visual field evaluation—This is discussed in detail elsewhere.²⁹ Visual fields were graded into minimal, mild, moderate, marked and severe loss. These grades are best described by examples given elsewhere.³⁴

EVALUATION OF RETINAL HEMORRHAGES

Final evaluation was always done from stereoscopic color fundus photographs (30° and 60° photographs).and in some based on the indirect and direct ophthalmoscopic evaluation. For each visit, the retinal hemorrhages at various locations in the retina were evaluated (comparing with the standard photos for various grades), by a single investigator (SSH), using the following criteria:

Subjective grading of retinal hemorrhages—1= no hemorrhages; 2 = minimal/small/ isolated hemorrhages; 3 = medium amount of hemorrhages; 4 = extensive hemorrhages; 5 = uncertain/questionable hemorrhages; 9 =no data.

Regions of the retina—The entire fundus was divided into the following 5 regions: (1) foveal center, (2) perifoveal region, (3) rest of the macular region, (4) rest of posterior pole (in the region of the arcades and around that), and (5) peripheral region. This is because different areas of the retina have different effects on visual function.

EVALUATION OF MACULAR EDEMA

Poor visual acuity in CRVO and hemi-CRVO, particularly in non-ischemic CRVO and hemi-CRVO, is due primarily to macular edema. Therefore, macular edema was evaluated critically in each eye at each visit. This was done by using: (i) direct ophthalmoscopy, (ii) Hruby non-contact lens, (iii) stereoscopic fundus photographs of the macular region, and (iv) frequent stereoscopic fluorescein fundus angiography of the macular region. OCT or other newer technologies to evaluate macular edema were not available during the period of this study.

Statistical Analysis

The demographic and clinical characteristics of patients that used aspirin and those that did not was compared using either the two-sample t-test for continuous variables or the Pearson Chi-square test for categorical variables. The Cochran-Mantel-Haenszel statistic was used to compare severity of hemorrhage, initial visual acuity, and visual defect grade at initial visit controlling for age, presence of systemic conditions and smoking status. The effect of aspirin use on the outcomes of visual acuity and visual field defect over time were examined using logistic regression fitted by the method of generalized estimating equations to account for correlation of outcomes over time from the same eye. The outcome of interest was visual deterioration for those that initially presented with 20/60 or better visual acuity, or minimal to mild visual field defect grade. For those that initially presented with 20/70 or worse visual acuity, or marked to severe visual field defect grade, the outcome was visual improvement. To account for differences in age and systemic conditions between those that used aspirin and those that did not, age, diabetes, hypertension, and stroke were included as covariates in the model.

Results

The demographic and clinical characteristics of patients by use of aspirin

These are shown in Table 1. Among the non-ischemic CRVO, those who were using aspirin were significantly older ($p=0.0002$), and had a higher prevalence of arterial hypertension ($p=0.014$), ischemic heart disease ($p<0.0001$), diabetes mellitus ($p=0.014$), and transient ischemic attack/cerebrovascular accident (TIA/CVA) ($p=0.024$) compared to the non-aspirin users. There was also a higher prevalence that ever smoked among aspirin users compared to non-aspirin users ($p=0.055$). In the non-ischemic hemi-CRVO patients there was a significantly greater prevalence of ischemic heart disease ($p=0.028$) in aspirin users compared to the non-aspirin users. For the ischemic CRVO patients, aspirin and aspirin/anticoagulant users differed in the prevalence of ischemic heart disease ($p=0.032$).

The demographic and clinical characteristics of patients by use of anticoagulant

There were 24 patients who were taking anticoagulant (warfarin): 14 with non-ischemic CRVO, 6 with non-ischemic hemi-CRVO, and 4 with ischemic CRVO. Among the 14 non-ischemic CRVO patients, there were 8 (57%) male and 4 female (43%), with mean age of 63.7 years (range 23–86). All patients had only one eye involved (7 right eye and 7 left eye). Of the relevant systemic diseases, 8 (57%) had arterial hypertension, 4 (29%) had ischemic heart disease, and 1 (7%) had TIA/CVA. Seven (50%) either currently or had smoked in the past. Of the 6 patients with non-ischemic hemi-CRVO, 5 (83%) were male, mean age was 74 (range 66 to 81) years; 1 (17%) had arterial hypertension, 2 (33%) had ischemic heart disease, and 2 (33%) had TIA/CVA. Of the 4 patients with ischemic CRVO, 2 were male, mean age was 69 (range 61 to 78) years. 2 had diabetes, 1 had arterial hypertension, and 1 had TIA/CVA.

At the initial visit, aspirin and non-aspirin/anticoagulant users were compared with respect to degree of fundus hemorrhage, visual acuity, and visual field defect. For all three type of CRVO, those that were taking aspirin had significantly greater severity of hemorrhage compared to those that were not taking aspirin (see 2 (available at <http://aaojournal.org>). This finding was seen for all regions examined except in the optic disc in non-ischemic hemi-CRVO and optic disc and peripapillary region in ischemic CRVO. In the case of visual acuity and visual field defect, aspirin users had significantly worst visual findings compared to non-aspirin users in non-ischemic CRVO (see Table 3). This was also observed for visual field defect in those with hemi-CRVO. No significant difference in initial visual acuity and visual field was found between aspirin users and non-users among the ischemic CRVO patients.

In the non-ischemic CRVO who were on anticoagulant, the degree of fundus hemorrhage at the initial visit was also of greater severity compared to the non-aspirin/anticoagulant group. This was observed for all regions examined ($p<0.003$ for fovea, perifovea, post pole, and peripheral; $p=0.018$ for macular; $p=0.056$ for peripapillary), except in the optic disc ($p=0.76$). Severe hemorrhage was present in 14% in the fovea, perifovea, and macular regions; 21% in the peripheral, 29% in the post pole, and 36% in the peripapillary region. For the 6 eyes with non-ischemic hemi-CRVO on anticoagulant, 4 of the 6 eyes showed moderate to severe fundus hemorrhage in all regions, except the optic disc. All 4 eyes with ischemic CRVO on anticoagulant had moderate to severe hemorrhage in the macular, post-pole, and peripapillary regions.

For visual assessment at the initial visit among those with non-ischemic CRVO on anticoagulant, 5 (36%) had 20/30 or better and 6 (43%) had 20/200 or worse visual acuity. Visual field defect was minimal in 9 (64%), mild in 4 (29%), and marked in 1 (7%). In the non-ischemic hemi-CRVO on anticoagulant, 5 of the 6 eyes had initial visual acuity worse

than 20/60, with 2 having moderate visual field defect. All 4 with ischemic CRVO on anticoagulant had initial visual acuity of CF; 2 had moderate and 2 had marked visual field defect.

The change in visual outcome was compared between aspirin users and no aspirin/anticoagulant users. Table 4 (available at <http://aaojournal.org>) and Table 5 (available at <http://aaojournal.org>) shows the proportion, at various follow-up times, that improved or worsened among those with 20/60 or better visual acuity (or minimal to mild visual field defect) and those with 20/70 or worse visual acuity (or moderate to severe visual field defect) at the initial visit. For eyes with non-ischemic CRVO that initially presented with 20/60 or better visual acuity, the effect of aspirin consistently showed a greater proportion with visual acuity deterioration in those that used aspirin compared to those that did not use aspirin over the follow-up duration (aspirin follow-up time interaction $p=0.268$). Adjusting for the effect of age, diabetes, ischemic heart disease, and hypertension, the odds ratio of visual acuity deterioration in aspirin users relative to non-users was 2.24 (95% CI (confidence interval): 1.14, 4.41; $p=0.020$). Of those that had macular edema resolved, overall cumulative visual acuity outcome also suggested an effect of aspirin use on deterioration, with the odds ratio for deterioration of 3.62 (95% CI: 0.97, 13.54; $p=0.055$) for aspirin users relative to non-users. For the non-ischemic CRVO that had 20/70 or worse visual acuity at initial visit, a significant difference on improvement in visual acuity was only found at the 2–5 year follow-up, with those that used aspirin were less likely to show improvement in visual acuity compared to non-users (odds ratio: 0.32; 95% CI: 0.11, 0.89; $p=0.029$). Among those where macular edema had resolved, over-all cumulative visual acuity outcome for improvement showed similar results, with the odds ratio for improvement of 0.18 (0.04, 0.72; $p=0.016$) for those that used aspirin relative to those that did not use aspirin.

Since there was large number of patients that did not have a 2–5 year follow-up, we examined how this possibly may have biased the results in the reported visual acuity change. This was done by comparing those with follow-up at 2–5 years and those with shorter follow-up with respect to visual acuity change. For the non-ischemic CRVO, in eyes with initial visual acuity of 20/60 or better, the distribution of visual acuity change for those with 2–5 year follow-up ($n=137$) was 12% improved, 20% worsened, and 68% with no change. For those not followed at 2–5 years ($n=108$), their observed change at the last follow-up also showed the same percentage of improved, worsened, and no change as those with the 2–5 year follow-up. Similarly, for those with initial visual acuity of 20/70 or worse, those with 2–5 year follow-up ($n=68$) had 47% improved, 24% worsened, and 29% no change, with the same distribution observed at the last follow-up for those not followed at 2–5 years ($n=61$). This suggests that those who had shorter follow-up had similar visual acuity change than those with the longer follow-up.

For changes in visual field defect among the non-ischemic CRVO eyes that initially presented with minimal to mild defect, significant differences in the proportion with visual field worsening due to aspirin use was seen at the 6 and 9 months follow-up ($p=0.003$ and $p=0.019$, respectively), but was not observed at the later follow-up (15 months $p=0.540$; 2–5 years $p=0.724$). In those where macular edema had resolved, over-all cumulative visual field outcome did not show a significant effect of aspirin use on visual field deterioration, with the odds ratio for deterioration of 2.40 (95% CI: 0.76, 7.64; $p=0.138$) for aspirin users relative to non-users. Among the non-ischemic CRVO eyes with moderate to severe initial visual field defect, there was no significant difference between aspirin users and non-users with respect to improvement in visual field.

Among those with non-ischemic hemi-CRVO and those with ischemic CRVO, there was no significant effect of aspirin use in changes in visual acuity and visual field.

For the non-ischemic CRVO eyes, there was no significant effect of aspirin use on conversion to the ischemic type ($p=0.827$). Within 6 months of onset of non-ischemic CRVO, there was conversion to the ischemic type in 9.1% of aspirin users and 9.2% of non-users. By 12 months after onset, 11.9% of patients that used aspirin and 12.7% of those that did not use aspirin have converted to ischemic CRVO.

Aspirin use did not have a significant effect on time to resolution of macular edema ($p=0.632$) in eyes with non-ischemic CRVO. The median (25th–75th percentile) time to macular edema resolution was 33.4 (11.5–58.6) months in those who used aspirin and 23.8 (8.5–71.3) months in those who did not use aspirin. The proportion(SE) of eyes that had resolved within 6, 12, and 24 months, was 13.5%(2.8%), 25.1%(3.7%), and 43.1%(4.5%), respectively, for aspirin users, and 17.4%(2.7%), 31.6%(3.5%), and 50.7%(4.0%), respectively, for those that did not use aspirin.

Discussion

This study comprised 686 patients (707 eyes) who either were on aspirin only or on anticoagulant (warfarin), or were not taking aspirin/anticoagulant treatment when they were first seen. Of these, 24 patients were on warfarin (14 with non-ischemic CRVO, 6 with non-ischemic hemi-CRVO, and 4 with ischemic CRVO) when first seen in the clinic. The study shows that among CRVO and hemi-CRVO, patients who took aspirin versus those who did not, there were significantly more marked retinal hemorrhages (Table 2 and Figs 1, 2 [available at <http://aaojournal.org>]) worse visual acuity and more marked visual field loss (Table 3), and on follow-up their final visual outcome was worse [Tables 4,5 (available at <http://aaojournal.org>)]. These findings show that aspirin has a deleterious effect on visual outcome in non-ischemic, as well as in non-ischemic hemi-CRVO. However, aspirin use did not have a significant effect ($p=0.63$) on the resolution of macular edema and conversion to ischemic CRVO ($p=0.83$) in those with non-ischemic CRVO. In our experience of over a thousand cases of CRVO and hemi-CRVO eyes, we have found that the common description of the appearance of the fundus in CRVO and hemi-CRVO as “blood and thunder” or “tomato ketchup” seems to represent an iatrogenic phenomenon caused by the use of aspirin (Figs. 1,2, available at <http://aaojournal.org>) or anticoagulants (Fig. 3 available at <http://aaojournal.org>) rather than the actual appearance in the vast majority of CRVO eyes, without the use of aspirin or anticoagulants. We have had a similar experience in dealing with branch retinal vein occlusion. In this study, among the patients who were on aspirin/anticoagulants when first seen, 60% of non-ischemic CRVO, 62% of ischemic CRVO and 77% of non-ischemic hemi-CRVO were already on aspirin (Fig. 2 available at <http://aaojournal.org>) or anticoagulant for other reasons at the time of development of CRVO or hemi-CRVO (Table 1). These findings are also reported by other studies^{35,36}.

Sedney³⁷, on a review of all the reports published until then, concluded that the vast amount of information on the use of anticoagulant therapy in retinal vein occlusion has failed to give convincing evidence of the value of this therapy. In his small series of CRVO cases treated with anticoagulants, he found that among the non-ischemic CRVO eyes, “the clinical course was disappointing”, with visual deterioration in 92% and no change in 8%. Among the ischemic CRVO cases in his study, anticoagulant therapy did not help. As discussed in our previous study²⁹ on the natural history of visual outcome in CRVO, the primary cause of poor visual outcome in ischemic CRVO is permanent ischemic damage to retinal ganglion cells in the macular region, resulting in permanent central scotoma, so that there is little chance of visual acuity improvement on follow-up. That explains why the ischemic CRVO cases in the series of Sedney³⁷ showed no change. All this contradicts the findings of Ageno et al.³⁸, who claimed that low molecular weight heparin appeared to be more effective than aspirin in preventing functional worsening in patients with retinal vein occlusion.

Koizumi et al.³⁹, in a case control study of 144 patients with CRVO, on multivariate logistic regression stated that the independent predictors for CRVO were glaucoma, aspirin use, and warfarin use. They identified both aspirin and warfarin use as independent risk factors for CRVO. Our studies⁴⁰ on CRVO do not support the concept that use of aspirin or anticoagulants are “independent risk factors for CRVO” *per se*, but the present study showed that they simply aggravate the retinal hemorrhages and worsen the visual outcome when CRVO or hemi-CRVO develops. There is no scientifically valid reason to believe that use of aspirin or anticoagulants can act as independent risk factors for CRVO. However, our previous study⁴¹, like that of Koizumi et al.³⁹, did show a significant association between CRVO and hemi-CRVO and glaucoma and ocular hypertension.

The genesis for the conventional wisdom of prescribing anticoagulants or aspirin for CRVO and other types of retinal vein occlusion comes from their proven effectiveness in major systemic venous thrombotic disorders (e.g., deep vein thrombosis); it has been assumed that they must also be equally beneficial in various types of retinal vein occlusion. But most important fact is that the treatment of a disease must be guided by its pathophysiology and pathogenesis. Although both deep vein thrombosis and CRVO are due to “thrombosis”, the similarity almost stops there. After all, the eye is not a leg anatomically or in any other characteristic. The anatomy of the veins and the morphology of the tissues in the leg differ fundamentally from those of the central retinal vein and morphology of the retina. Moreover, the pathogenesis of deep vein thrombosis is very different from that of CRVO and hemi-CRVO⁴². An increase in retinal hemorrhages (Figs. 1–3, available at <http://aaojournal.org>) can be devastating, destroying the delicate retinal neural tissue and resulting in visual deterioration, as shown in this study; however, similar hemorrhages in the leg tissue cause no such morphological and functional damage. Thus, to equate the two conditions and their management is a fundamental mistake.

The argument invariably put forward for treating CRVO and hemi-CRVO patients with aspirin or anticoagulants is that hematologic abnormalities are responsible for their development and that treatment of those abnormalities with aspirin or anticoagulants has protective or beneficial effect in such cases. A brief discussion of the pathogenesis of CRVO, hemi-CRVO and branch retinal vein occlusion is essential here to explain the primary cause of the occlusion. In the center of the optic nerve, the central retinal artery and central retinal vein lie side by side in close apposition, both enclosed by a common fibrous tissue envelope (Fig. 4, available at <http://aaojournal.org>). With aging and in glaucoma⁴³, there is thickening and sclerotic change in the fibrous tissue envelope as well as in the central retinal artery (due to atherosclerosis/arteriosclerosis). The sclerotic changes in the fibrous tissue envelope reduce the space in it; furthermore, the thickened central retinal artery compresses the adjacent thin-walled central retinal vein (Fig. 4 available at <http://aaojournal.org>). That results in marked narrowing of the lumen of the central retinal vein, causing stagnant flow and hemodynamic changes in the central retinal vein. According to Virchow's triad, blood flow stasis promotes thrombosis. A similar mechanism most probably works in branch retinal vein occlusion because invariably the site of occlusion in it is at the arteriovenous crossing. Thus, pathogenetically, the primary factor in the development of CRVO and branch retinal vein occlusion is almost invariably the change in the adjacent arteries, and not systemic hematologic abnormalities. This is very different from deep vein thrombosis and other systemic venous occlusive disorders where hematologic abnormalities, and not arterial disease, play the primary role. Thus, pathogenetically there is a fundamental difference between retinal vein occlusion and deep vein occlusion and other systemic venous occlusive disorders.

Hayreh et al.³⁵ critically reviewed the literature about the prevalence of various hematologic abnormalities in CRVO and other types of retinal vein occlusion. That review showed that

the negative findings about the association between those hematologic abnormalities and development of retinal vein occlusion outweighed the positive findings, with flaws in the latter studies. The conclusion was that, contrary to many claims, there is no definite evidence of a cause-and-effect relationship between the various hematologic abnormalities and the development of various types of retinal vein occlusion in the vast majority of the retinal vein occlusion patients; however, there is always the possibility of a purely chance occurrence of some of these hematologic abnormalities in an occasional retinal vein occlusion cases or the possibility of the findings due to unrelated associated systemic disease cannot be ruled out. The misconception of equating retinal vein occlusion with systemic venous occlusive disorders is responsible for blaming hematologic abnormalities responsible for retinal vein occlusion.

Therefore, what is the practical implication of all this from the clinical point of view? Ingerslev⁴⁴ very well summarized the state of our knowledge on the subject when he stated that: "Most well-characterised (hematologic) risk factors for general venous thrombosis occur sporadically only in RVO, and it seems these have no major importance in the pathophysiology of RVO." In view of that, he rightly recommended, "there seems to be no particular reason that this should include a complete haemostasiological investigation like that offered to patients with spontaneous major venous thromboembolism." The study by Hayreh et al.³⁵ in 645 patients supported that view.

Interestingly, relevant to our study is a recent report⁴⁵ of the effect of daily use of antiplatelet or anticoagulant "(AP/AC)" medication in patients with neovascular age-related macular degeneration. In that study there were 195 patients, 96 (49.2%) of them were on daily AP/AC. They found that of the patients taking AP/AC, 63.5% developed hemorrhages compared with 29.2% of patients not taking (odds ratio = 4.21; 95% confidence interval = 1.42–8.46; $P < 0.001$). They concluded that use of AP/AC: (i) was significantly associated with an increased risk of development of intraocular hemorrhages in patients with neovascular AMD, (ii) was also an independent risk factor for the development of intraocular hemorrhages in these patients, and (iii) "presence of bilateral neovascular AMD was a significant association in those taking daily A/AC." The findings of this study lend support to those of our study.

In conclusion, findings of this study contradict the conventional wisdom of prescribing aspirin or other antiplatelet aggregating agents or anticoagulants to treat patients with CRVO and hemi-CRVO, because the use of these drugs was associated with a worse visual outcome, and no apparent benefit.

In the light of these findings, we give the following advice to our patients with CRVO and hemi-CRVO:

1. Frequently, the question arises about what to advise these patients who have been taking aspirin or anticoagulants for various systemic diseases, e.g., myocardial infarction, coronary artery disease, stroke, carotid artery disease and other systemic disorders, where continuation of aspirin or anticoagulants is essential to prevent life threatening problems. We always advise them that they should continue with those treatments because on a risk/benefit ratio, the eye takes a second place to the heart or brain; and that they should discuss the whole matter with their internists.
2. Patients who develop CRVO or hemi-CRVO, have never taken aspirin and have no indication to use it for any systemic reason are advised not to take aspirin or anticoagulants.

3. Patients who have been started on aspirin or anticoagulants solely to treat CRVO or hemi-CRVO, or are taking ASA on their own just because they think it is beneficial (without any specific medical indication), are advised to stop taking that.

Precis

Findings of this study indicate that, for patients with CRVO and hemi-CRVO, the use of aspirin, other antiplatelet aggregating agents or anticoagulants was associated with a worse visual outcome, and no apparent benefit.

Acknowledgments

Supported by grant EY-1151 from the National Institutes of Health.

References

1. Holmin N, Ploman KG. Thrombosis of central vein of retina treated with heparin. *Lancet*. 1938; 231:664–5.
2. Huggett AS, Juler FA. Heparin in thrombosis of the central vein of the retina. *Trans Ophthalmol Soc U K*. 1942; 62:123–34.
3. Odqvist B. Prognose vid stan - och gromboser i nathinnans vener utan behandling med heparin. *Nord Med*. 1942; 16:3349–50.
4. Rosengren B, Stenstrom S. 30 Falle von Venenthrombose der Netzhaut, behandelt mit Heparin. *Acta Ophthalmol (Copenh)*. 1942; 20:145–52.
5. Klien BA. Anticoagulant therapy of occlusion of central vein of retina: in relation to pathogenesis and differential diagnosis. *Arch Ophthalmol*. 1943; 29:699–710.
6. Blum JD, Jeanneret H. Traitement des thromboses des veines retiennes par un nouvel anticoagulant: la dicoumarine. *Ophthalmologica*. 1944; 108:129–36.
7. Ploman KG. Treatment of thrombosis of the veins of the retina with heparin. *Acta Ophthalmol (Copenh)*. 1943; 21:190–203.
8. Maclean AL, Brambel CE. Dicumarol and rutin in vascular disease. *Trans Am Ophthalmol Soc*. 1946; 44:194–213.
9. Banks-Smith RG. A case of partial thrombosis of central retinal vein treated by dicoumarin. *Trans Ophthalmol Soc Aust*. 1947; 7:103–5.
10. Rosengren B. The value of anti-coagulators in the treatment of retinal thrombosis. *Acta Ophthalmol (Copenh)*. 1948; 26:275–80. [PubMed: 18878069]
11. Benedict WL. The clinical significance of closure of the retinal vessels. *J Med Assoc Ga*. 1949; 38:423–33. [PubMed: 18141837]
12. Braendstrup P. Central retinal vein thrombosis and hemorrhagic glaucoma. *Acta Ophthalmol Suppl*. 1950; 35:1–162. [PubMed: 14782975]
13. Larsson S, Nord B. Some remarks on retinal vein thrombosis and its treatment with anticoagulants. *Acta Ophthalmol*. 1950; 28:187–201. [PubMed: 14777290]
14. Duff IF, Falls HF, Linman JW. Anticoagulant therapy in occlusive vascular disease of the retina. *Arch Ophthalmol*. 1951; 46:601–17.
15. Cassady JV. Central retinal vein thrombosis. *Am J Ophthalmol*. 1953; 36:331–5. [PubMed: 13030633]
16. Klien BA, Olwin JH. A survey of the pathogenesis of retinal venous occlusion: emphasis upon choice of therapy and an analysis of the therapeutic results in fifty-three patients. *Arch Ophthalmol*. 1956; 56:207–47.
17. Vannas S, Orma H. Anticoagulant treatment of retinal venous occlusion with anticoagulant and antisclerosis therapy. *AMA Arch Ophthalmol*. 1957; 58:812–28. [PubMed: 13478227]
18. van Loon J. The causes and therapy of thrombosis of the retinal veins. *Ophthalmologica*. 1961; 141:467–75. [PubMed: 13763407]

19. Meythaler H. On the occlusions of retinal vessels [in German]. *Klin Monbl Augenheilkd*. 1966; 149:32–41. [PubMed: 5966438]
20. Vannas S, Raitta C. Anticoagulant treatment of retinal venous occlusion. *Am J Ophthalmol*. 1966; 62:874–84. [PubMed: 5928837]
21. Hayreh SS. Occlusion of the central retinal vessels. *Br J Ophthalmol*. 1965; 49:626–45. [PubMed: 4954984]
22. Hayreh SS. So-called “central retinal vein occlusion”: 1. Pathogenesis, terminology, clinical features. *Ophthalmologica*. 1976; 172:1–13. [PubMed: 1250571]
23. Hayreh SS. Pathogenesis of occlusion of the central retinal vessels. *Am J Ophthalmol*. 1971; 72:998–1011. [PubMed: 5315812]
24. Hayreh SS, van Heuven WA, Hayreh MS. Experimental retinal vascular occlusion. I. Pathogenesis of central retinal vein occlusion. *Arch Ophthalmol*. 1978; 96:311–23. [PubMed: 415709]
25. Hayreh SS, Hayreh MS. Hemi-central retinal vein occlusion: pathogenesis, clinical features, and natural history. *Arch Ophthalmol*. 1980; 98:1600–9. [PubMed: 7425921]
26. Hayreh SS. Classification of central retinal vein occlusion. *Ophthalmology*. 1983; 90:458–74. [PubMed: 6877778]
27. Hayreh SS, Rojas P, Podhajsky P, et al. Ocular neovascularization with retinal vascular occlusion: III. Incidence of ocular neovascularization with retinal vein occlusion. *Ophthalmology*. 1983; 90:488–506. [PubMed: 6192376]
28. Hayreh SS, Klugman MR, Beri M, et al. Differentiation of ischemic from non-ischemic central retinal vein occlusion during the early acute phase. *Graefes Arch Klin Exp Ophthalmol*. 1990; 228:201–17.
29. Hayreh SS, Podhajsky PA, Zimmerman MB. Natural history of visual outcome in central retinal vein occlusion. *Ophthalmology*. 2011; 118:119–33. [PubMed: 20723991]
30. Hayreh SS, Zimmerman B. Visual field abnormalities in nonarteritic anterior ischemic optic neuropathy: their pattern and prevalence at initial presentation. *Arch Ophthalmol*. 2005; 123:1554–62. [PubMed: 16286618]
31. Hayreh SS. Prevalent misconceptions about acute retinal vascular occlusive disorders. *Prog Retin Eye Res*. 2005; 24:493–519. [PubMed: 15845346]
32. Hayreh SS, Zimmerman MB. Nonarteritic anterior ischemic optic neuropathy: natural history of visual outcome. *Ophthalmology*. 2008; 115:298–305. [PubMed: 17698200]
33. Ischemic Optic Neuropathy Decompression Trial Research Group. Optic nerve decompression surgery for nonarteritic anterior ischemic optic neuropathy (NAION) is not effective and may be harmful. *JAMA*. 1995; 273:625–32. [PubMed: 7844872]
34. Hayreh SS. Posterior ischaemic optic neuropathy: clinical features, pathogenesis, and management. *Eye (Lond)*. 2004; 18:1188–206. [PubMed: 15534605]
35. Hayreh SS, Zimmerman MB, Podhajsky PA. Hematologic abnormalities associated with various types of retinal vein occlusion. *Graefes Arch Clin Exp Ophthalmol*. 2002; 240:180–96. [PubMed: 11935275]
36. Browning DJ, Fraser CM. Retinal vein occlusions in patients taking warfarin. *Ophthalmology*. 2004; 111:1196–200. [PubMed: 15177971]
37. Sedney, SC. The Hague. Vol. 21–7. W Junk; The Netherlands: 1976. Photocoagulation in Retinal Vein Occlusion [dissertation]; p. 80-100.
38. Ageno W, Cattaneo R, Manfredi E, et al. Parnaparin versus aspirin in the treatment of retinal vein occlusion: a randomized, double blind, controlled study. *Thromb Res*. 2010; 125:137–41. [PubMed: 19477488]
39. Koizumi H, Ferrara DC, Bruè C, Spaide RF. Central retinal vein occlusion case-control study. *Am J Ophthalmol*. 2007; 144:858–63. [PubMed: 17916319]
40. Hayreh SS, Zimmerman B, McCarthy MJ, Podhajsky P. Systemic diseases associated with various types of retinal vein occlusion. *Am J Ophthalmol*. 2001; 131:61–77. [PubMed: 11162981]
41. Hayreh SS, Zimmerman MB, Beri M, Podhajsky P. Intraocular pressure abnormalities associated with central and hemicentral retinal vein occlusion. *Ophthalmology*. 2004; 111:133–41. [PubMed: 14711725]

42. Hayreh SS. Central retinal vein occlusion. *Ophthalmol Clin North Am.* 1998; 11:559–90.
43. Hayreh SS, Pe'er J, Zimmerman MB. Morphologic changes in chronic high-pressure experimental glaucoma in rhesus monkeys. *J Glaucoma.* 1999; 8:56–71. [PubMed: 10084276]
44. Ingerslev J. Thrombophilia: a feature of importance in retinal vein thrombosis? *Acta Ophthalmol Scand.* 1999; 77:619–21. [PubMed: 10634549]
45. Kiernan DE, Hariprasad SM, Rusu IM, et al. Epidemiology of the association between anticoagulants and intraocular hemorrhage in patients with neovascular age-related macular degeneration. *Retina.* 2010; 30:1573–8. [PubMed: 21060269]

Figure 1.

A 61-year old man with non-ischemic CRVO, when first seen by his local ophthalmologist, had visual acuity of 20/30, and was started on aspirin. When seen in the Ocular Vascular Clinic at Iowa City 2 weeks later, the fundus showed extensive retinal hemorrhages and visual acuity in 2 months deteriorated to 20/400 and developed a huge centrocecal scotoma.

Figure 2.

An 83-year old woman with superior hemi-non-ischemic CRVO, when seen 2½ months after its onset, showed extensive retinal hemorrhages and visual acuity of 20/200, which deteriorated to counting fingers. She was on aspirin for angina and coronary artery bypass graft prior to the onset venous occlusion.

Figure 3.

A 35-year-old man with non-ischemic CRVO, when seen by his local ophthalmologist had visual acuity of 20/15, and was started on Coumadin. When seen at the Ocular Vascular Clinic at Iowa City 3 months after the onset, fundus showed extensive retinal hemorrhages, especially marked in the peripheral retina. His visual acuity deteriorated to 20/80, with a marked loss of peripheral visual field (with only an island field in the inferior nasal region with I-2e isopter).

Figure 4. Transverse sections through the anterior part of the optic nerve show the central retinal vessels and the surrounding fibrous tissue envelope (FTE) in the center of the optic nerve. Upper picture shows the normal pattern. The lower picture shows markedly thickened perivascular fibrous tissue envelope (FTE), marked thickening of the wall of the central retinal artery (CRA), and marked stenosis of the central retinal vein (CRV).

Table 1

Demographic and Clinical characteristics

Demographic/Clinical Variable	Non-ischemic CRVO		Non-ischemic Hemi-CRVO		Ischemic CRVO as first diagnosis	
	On Aspirin (n=189 patients*, n=197 eyes)	Not on Aspirin (n=277 patients*, n=284 eyes)	On Aspirin (n=57 patients, n=58 eyes)	Not on Aspirin (n=56 patients, n=58 eyes)	On Aspirin (n=38 patients, n=38 eyes)	Not on Aspirin (n=47 patients, n=48 eyes)
Gender (Male)	104 (55%)	142 (51%)	32 (56%)	26 (46%)	22 (58%)	22 (47%)
Age at initial visit						
Mean±SD	63±14	58±17	67±12	63±14	70±12	68±16
Range	20–90	18–89	39–90	21–84	30–87	32–89
Distribution by age (count, %)						
<45	17 (9%)	65 (23%)	2 (4%)	9 (16%)	1 (3%)	7 (15%)
45–<65	73 (398%)	96 (35%)	18 (32%)	16 (29%)	7 (18%)	7 (15%)
65 and older	99 (52%)	116 (42%)	37 (65%)	31 (55%)	30 (79%)	33 (70%)
Eye involvement						
OD	98 (52%)	129 (47%)	28 (49%)	21 (38%)	15 (39%)	19 (40%)
OS	83 (44%)	141 (51%)	28 (49%)	33 (59%)	23 (61%)	27 (57%)
OU	8 (4%)	7 (3%)	1 (2%)	2 (4%)	0 (0%)	1 (2%)
Follow-up @	(n=148 eyes)	(n=205 eyes)	(n=49 eyes)	(n=47 eyes)	(n=28 eyes)	(n=42 eyes)
Median (25 th –75 th percentile)	3.1 (1.0–6.3) yrs	2.8 (0.8–6.1) yrs	4.0 (2.1–7.3) yrs	1.9 (0.9–6.1) yrs	1.9 (0.9–5.1) yrs	2.9 (1.1–6.6) yrs
Minimum-Maximum	3.2 mos-23.6 yrs	3.1 mos-28.2 yrs	3.0 mos-20.9 yrs	3.0 mos-17.4 yrs	3.0 mos-15.3 yrs	3.0 mos-14.1 yrs
Systemic Conditions						
Arterial hypertension	93 (49%)	104 (38%)	27 (47%)	20 (36%)	18 (45%)	24 (52%)
Ischemic heart disease	35 (19%)	17 (6%)	9 (16%)	2 (4%)	11 (29%)	5 (11%)
Diabetes mellitus	26 (14%)	19 (7%)	5 (9%)	4 (7%)	8 (21%)	7 (15%)
TIA/CVA	14 (7%)	8 (3%)	4 (7%)	0 (0%)	1 (3%)	1 (2%)
Smoked current/past	(n=184) 92 (50%)	(n=262) 107 (41%)	(n=56) 24 (33%)	(n=50) 21 (42%)	(n=36) 17 (47%)	(n=40) 14 (35%)
Aspirin started after onset of CRVO	80 (42%)	–	15 (26%)	–	16 (38%)	–

CRVO=Central retinal Vein Occlusion

SD=Standard deviation

CVA = Cerebrovascular accident;

TIA = Transient ischemic attack

* Total patients with non-ischemic CRVO is 464, with 2 patients having 1 eye in each aspirin use group

@ Of those with follow-up of at least 3 months.

Table 2

Hemorrhage at initial visit by Aspirin use

Location of Hemorrhage	On Aspirin at initial visit				Not on Aspirin/Anticoagulant				p-value*		
	n	Hemorrhage			n	Hemorrhage					
		None	Mild	Moderate		Severe	None	Mild		Moderate	Severe
Non-ischemic CRVO											
Fovea (center)	191	70 (37%)	64 (34%)	34 (18%)	23 (12%)	277	177 (64%)	71 (26%)	21 (8%)	8 (3%)	<0.0001
Perifovea	191	36 (19%)	66 (35%)	66 (35%)	23 (12%)	277	79 (29%)	157 (57%)	34 (12%)	7 (3%)	<0.0001
Macular	190	19 (10%)	58 (31%)	83 (44%)	30 (16%)	275	47 (17%)	166 (60%)	56 (20%)	6 (2%)	<0.0001
Post Pole	191	15 (8%)	55 (29%)	82 (43%)	39 (20%)	276	33 (12%)	178 (65%)	57 (21%)	8 (3%)	<0.0001
Peripheral	190	2 (1%)	96 (51%)	70 (37%)	22 (12%)	275	6 (2%)	229 (83%)	36 (13%)	4 (1%)	<0.0001
Disc	190	98 (52%)	55 (29%)	33 (17%)	4 (2%)	277	183 (66%)	64 (23%)	24 (9%)	6 (2%)	0.0006
Peripapillary	190	35 (18%)	44 (23%)	56 (29%)	55 (29%)	277	98 (35%)	108 (39%)	50 (18%)	21 (8%)	<0.0001
Non-ischemic Hemi-CRVO											
Fovea (center)	58	15 (26%)	13 (22%)	16 (28%)	14 (24%)	58	28 (48%)	23 (40%)	3 (5%)	4 (7%)	0.002
Perifovea	58	5 (9%)	15 (26%)	22 (38%)	16 (28%)	58	13 (22%)	35 (60%)	9 (16%)	1 (2%)	<0.0001
Macular	58	4 (7%)	6 (10%)	31 (53%)	17 (29%)	58	9 (16%)	31 (53%)	18 (31%)	0 (0%)	<0.0001
Post Pole	58	1 (2%)	13 (22%)	30 (52%)	14 (24%)	58	5 (9%)	33 (57%)	20 (34%)	0 (0%)	<0.0001
Peripheral	58	1 (2%)	15 (26%)	31 (53%)	11 (19%)	58	0 (0%)	44 (76%)	14 (24%)	0 (0%)	<0.0001
Disc	58	33 (57%)	15 (26%)	9 (16%)	1 (2%)	58	35 (60%)	15 (26%)	7 (12%)	1 (2%)	0.851
Peripapillary	58	5 (9%)	16 (28%)	20 (35%)	16 (28%)	58	14 (24%)	26 (45%)	17 (29%)	1 (2%)	0.0008
Ischemic CRVO at first diagnosis											
Fovea (center)	38	4 (11%)	5 (13%)	6 (16%)	23 (61%)	46	5 (11%)	16 (35%)	14 (30%)	11 (24%)	0.038
Perifovea	38	1 (3%)	6 (16%)	7 (18%)	24 (63%)	46	2 (4%)	8 (17%)	24 (52%)	12 (26%)	0.015
Macular	38	0 (0%)	4 (11%)	10 (26%)	24 (63%)	46	0 (0%)	10 (22%)	22 (48%)	14 (30%)	0.002
Post Pole	37	1 (3%)	4 (11%)	8 (22%)	24 (65%)	46	0 (0%)	8 (17%)	24 (52%)	14 (30%)	0.028
Peripheral	38	0 (0%)	5 (13%)	10 (26%)	23 (61%)	46	0 (0%)	10 (22%)	24 (52%)	12 (26%)	0.003
Disc	36	20 (56%)	7 (19%)	7 (19%)	2 (6%)	45	25 (56%)	11 (24%)	3 (7%)	6 (13%)	0.894
Peripapillary	38	2 (5%)	4 (11%)	9 (24%)	23 (61%)	46	2 (4%)	10 (22%)	17 (37%)	17 (37%)	0.124

CRVO = Central retina vein occlusion

* Cochran-Mantel-Haenszel test comparing On Aspirin vs. Not On Aspirin controlling for age and presence of systemic conditions and smoking status

Table 3

Visual acuity and visual field at initial visit

Visual assessment	On Aspirin	Not on Aspirin/Anticoagulant	p-value*
Non-ischemic CRVO			
Visual acuity	(n=197 eyes)	(n=284 eyes)	0.017
20/15–20/20	30 (15%)	84 (30%)	
20/25–20/30	34 (17%)	48 (17%)	
20/40–20/60	48 (24%)	71 (25%)	
20/70–20/100	27 (14%)	35 (12%)	
20/200–400	50 (25%)	43 (15%)	
CF or worse	8 (4%)	3 (1%)	
Visual field defect	(n=193)	(n=274)	0.016
Minimal	107 (55%)	207 (76%)	
Mild	64 (33%)	50 (18%)	
Moderate	21 (11%)	17 (6%)	
Marked	1 (1%)	0 (0%)	
Severe	0 (0%)	0 (0%)	
Non-ischemic Hemi-CRVO			
Visual acuity	(n=57 eyes)	(n=57 eyes)	0.233
20/15–20/20	10 (18%)	14 (25%)	
20/25–20/30	9 (16%)	16 (28%)	
20/40–20/60	12 (21%)	12 (21%)	
20/70–20/100	8 (14%)	10 (18%)	
20/200–400	14 (25%)	4 (7%)	
CF or worse	4 (7%)	1 (2%)	
Visual field defect	(n=54)	(n=52)	0.003
Minimal	29 (54%)	45 (87%)	
Mild	14 (26%)	5 (10%)	
Moderate	11 (20%)	2 (4%)	
Marked	0 (0%)	0 (0%)	
Severe	0 (0%)	0 (0%)	
Ischemic CRVO at first diagnosis			
Visual acuity	(n=38 eyes)	(n=45 eyes)	0.905
Better than 20/70	0 (0%)	0 (0%)	
20/70–20/100	1 (3%)	0 (0%)	
20/200–400	7 (18%)	11 (24%)	
CF or worse	30 (79%)	34 (76%)	
Visual field defect	(n=36)	(n=42)	0.667

Visual assessment	On Aspirin	Not on Aspirin/Anticoagulant	p-value*
Minimal	0 (0%)	0 (0%)	
Mild	4 (11%)	2 (5%)	
Moderate	16 (44%)	21 (50%)	
Marked	14 (44%)	16 (38%)	
Severe	0 (0%)	3 (7%)	

CF = Counting fingers

CRVO = Central retinal vein occlusion

* Cochran-Mantel-Haenszel test comparing On aspirin vs. Not on aspirin controlling for age and presence of systemic conditions and smoking status

Table 4

Visual acuity change from initial visit to 3 months, 6 months, 9 months, 15 months, and 2 to 5 years follow-up, and overall cumulative change after resolution of macular edema in those first seen within 3 months from onset, by aspirin use at initial visit

Follow-up/Initial Visual Acuity	On Aspirin		Not on Aspirin/Anticoagulant			
	Number (%) of eyes		Number (%) of eyes			
	n		n			
	Improved	Worsened	Improved	Worsened		
Non-ischemic CRVO						
<u>3 months</u> *						
20/60 or better	66	7 (11%)	15 (23%)	133	8 (6%)	16 (12%)
20/70 or worse	58	16 (28%)	21 (36%)	51	18 (35%)	8 (16%)
<u>6 months</u> *						
20/60 or better	59	8 (14%)	18 (31%)	93	10 (11%)	13 (14%)
20/70 or worse	53	16 (30%)	20 (38%)	45	19 (42%)	8 (18%)
<u>9 months</u> *						
20/60 or better	47	2 (4%)	15 (32%)	97	10 (10%)	16 (16%)
20/70 or worse	41	13 (32%)	15 (37%)	34	15 (44%)	7 (21%)
<u>15 months</u> *						
20/60 or better	45	2 (4%)	14 (31%)	69	10 (14%)	7 (10%)
20/70 or worse	41	11 (27%)	17 (41%)	29	13 (45%)	6 (21%)
<u>2 to 5 years</u> *						
20/60 or better	48	3 (6%)	14 (29%)	85	12 (14%)	12 (14%)
20/70 or worse	38	12 (32%)	13 (34%)	30	20 (67%)	3 (10%)
<u>Cumulative after macular edema resolved</u> @						
20/60 or better	57	8 (14%)	9 (16%)	112	15 (13%)	4 (4%)
20/70 or worse	34	17 (50%)	6 (18%)	27	22 (81%)	2 (7%)
Non-ischemic Hemi-CRVO						

Follow-up/Initial Visual Acuity	On Aspirin				Not on Aspirin/Anticoagulant			
	n		Number (%) of eyes		n		Number (%) of eyes	
			Improved	Worsened			Improved	Worsened
<u>3 months</u> *								
20/60 or better	23		2 (9%)	1 (4%)	31		4 (13%)	3 (10%)
20/70 or worse	19		8 (42%)	4 (21%)	10		4 (40%)	2 (20%)
<u>6 months</u> *								
20/60 or better	20		1 (5%)	2 (10%)	19		6 (32%)	3 (16%)
20/70 or worse	15		7 (47%)	2 (13%)	5		0 (0%)	1 (20%)
<u>9 months</u> *								
20/60 or better	19		3 (16%)	3 (16%)	23		7 (30%)	3 (13%)
20/70 or worse	11		7 (64%)	2 (18%)	7		1 (14%)	2 (29%)
<u>15 months</u> *								
20/60 or better	20		4 (20%)	3 (15%)	18		5 (28%)	2 (11%)
20/70 or worse	12		8 (67%)	2 (17%)	7		2 (29%)	2 (29%)
<u>2 to 5 years</u> *								
20/60 or better	22		4 (18%)	2 (9%)	18		3 (17%)	1 (6%)
20/70 or worse	14		4 (29%)	2 (14%)	5		0 (0%)	0 (0%)
Cumulative after macular edema resolved @								
20/60 or better	23		3 (13%)	0 (0%)	33		6 (18%)	2 (6%)
20/70 or worse	13		8 (62%)	1 (8%)	8		4 (50%)	2 (25%)
Ischemic CRVO at first diagnosis								
<u>3 months</u> *								
20/70 or worse	26		2 (8%)	5 (19%)	32		4 (12%)	4 (12%)
<u>6 months</u> *								
20/70 or worse	22		3 (14%)	4 (18%)	29		5 (17%)	6 (21%)

Follow-up/Initial Visual Acuity	On Aspirin		Not on Aspirin/Anticoagulant	
	Number (%) of eyes		Number (%) of eyes	
	n		n	
9 months [*]				
20/70 or worse	18	3 (17%)	32	5 (16%)
				8 (25%)
15 months [*]				
20/70 or worse	16	5 (31%)	21	5 (24%)
				5 (24%)
2 to 5 years [*]				
20/70 or worse	9	2 (22%)	17	4 (24%)
				5 (29%)
Cumulative after macular edema resolved [@]				
20/70 or worse	11	3 (27%)	21	9 (43%)
				4 (19%)

CRVO = Central retina vein occlusion

^{*} ±6 weeks for 3, 6, and 9 months; ±12 weeks for 15 months

[@] Change after macular edema has resolved (from initial visit to visit following resolution for eyes where macular edema resolved)

Table 5

Visual field change from initial visit to 3 months, 6 months, 9 months, 15 months, and 2 to 5 years follow-up, and overall cumulative change after resolution of macular edema in those first seen within 3 months from onset, by aspirin use at initial visit

Follow-up/Initial Visual Fields	On Aspirin		Not on Aspirin/Anticoagulant				
	n	Number (%) of eyes		Number (%) of eyes			
		Improved	Worsened	Improved	Worsened		
Non-ischemic CRVO							
3 months *							
	Minimal-mild	105	20 (19%)	23 (22%)	166	33 (20%)	21 (13%)
	Moderate-severe	15	4 (27%)	5 (33%)	11	7 (64%)	1 (9%)
6 months *							
	Minimal-mild	99	23 (23%)	29 (29%)	121	33 (27%)	13 (11%)
	Moderate-severe	10	4 (40%)	3 (30%)	11	5 (45%)	1 (9%)
9 months *							
	Minimal-mild	76	20 (26%)	22 (29%)	115	28 (24%)	13 (11%)
	Moderate-severe	10	6 (60%)	3 (30%)	12	7 (58%)	1 (8%)
15 months *							
	Minimal-mild	73	18 (25%)	14 (19%)	88	29 (33%)	12 (14%)
	Moderate-severe	12	8 (67%)	3 (25%)	6	4 (67%)	1 (17%)
2 to 5 years *							
	Minimal-mild	76	21 (28%)	13 (17%)	100	40 (40%)	12 (12%)
	Moderate-severe	9	7 (78%)	2 (22%)	10	8 (80%)	1 (10%)
@							
Cumulative after macular edema resolved							
Minimal-mild	82	26 (32%)	8 (10%)		128	43 (34%)	6 (5%)
Moderate-severe	8	6 (75%)	2 (25%)		8	8 (100%)	0 (0%)
Non-ischemic Hemi-CRVO							

Follow-up/Initial Visual Fields	On Aspirin				Not on Aspirin/Anticoagulant			
	n	Number (%) of eyes		n	Number (%) of eyes		n	Worsened
		Improved	Worsened		Improved	Worsened		
<u>3 months</u> *								
Minimal-mild	32	7 (22%)	4 (12%)	37	3 (8%)	1 (3%)		
<u>6 months</u> *								
Minimal-mild	30	13 (43%)	3 (10%)	21	4 (19%)	1 (5%)		
<u>9 months</u> *								
Minimal-mild	25	11 (44%)	3 (12%)	28	5 (18%)	2 (7%)		
<u>15 months</u> *								
Minimal-mild	25	10 (40%)	4 (16%)	23	4 (17%)	3 (13%)		
<u>2 to 5 years</u> *								
Minimal-mild	30	7 (23%)	2 (7%)	19	2 (11%)	1 (5%)		
Cumulative after macular edema resolved @								
Minimal-mild	30	13 (43%)	2 (7%)	37	5 (14%)	2 (5%)		
Ischemic CRVO at initial diagnosis								
<u>3 months</u> *								
Moderate-Severe	23	4 (17%)	11 (48%)	33	4 (12%)	12 (36%)		
<u>6 months</u> *								
Moderate-Severe	21	6 (29%)	10 (48%)	29	4 (14%)	13 (44%)		
<u>9 months</u> *								
Moderate-Severe	15	4 (27%)	8 (53%)	29	3 (10%)	14 (48%)		
<u>15 months</u> *								
Moderate-Severe	15	5 (33%)	8 (53%)	22	4 (18%)	12 (55%)		

Follow-up/Initial Visual Fields	On Aspirin			Not on Aspirin/Anticoagulant		
	n	Number (%) of eyes		n	Number (%) of eyes	
		Improved	Worsened		Improved	Worsened
2 to 5 years [*] Moderate-Severe	11	4 (36%)	7 (64%)	13	5 (24%)	11 (52%)
Cumulative after macular edema resolved [@] Moderate-Severe	11	3 (27%)	2 (18%)	16	6 (38%)	6 (38%)

CRVO = Central retina vein occlusion

^{*} ± 6 weeks for 3, 6, and 9 months; ±12 weeks for 15 months

[@] Change after macular edema has resolved (from initial visit to visit following resolution for eyes where macular edema resolved)