Published in final edited form as:

J Child Psychol Psychiatry. 2003 October ; 44(7): 988–996.

Genetic epidemiology of self-reported lifetime DSM-IV major depressive disorder in a population-based twin sample of female adolescents

Anne L. Glowinski, Pamela A.F. Madden, Kathleen K. Bucholz, Michael T. Lynskey, and Andrew C. Heath

Washington University School of Medicine, St. Louis, USA

Abstract

Background: In adults, about 40% of the variance in risk of Major Depressive Disorder (MDD) is due to genetic factors, but little data exist on the heritability of youth MDD. The goal of this study was the genetic analysis of MDD in an epidemiologically and genetically representative sample of adolescent female twins.

Methods: A sample of 3416 female adolescent twins systematically ascertained from birth records was assessed using a structured telephone interview that included a comprehensive DSM-IV-based section for the diagnostic assessment of MDD. Mean subject age at time of assessment was 15.5 and participation rate exceeded 85%. Genetic modeling was conducted taking into consideration the problem of censoring, i.e., that younger adolescents were not through their period of risk for adolescent onset of MDD.

Results: Lifetime self-reported MDD prevalence ranged from 1% under age 12 to 17.4% at age 19 and older. The genetic variance in risk of MDD was 40.4% (95% confidence interval (CI): 23.9–55.1), with the remaining variance explained by non-shared environmental effects 59.6% (95% CI: 44.9–76.1). Shared environmental effects were not significant. A significant recall bias was observed with older respondents on average reporting later onsets for their first episode of MDD.

Conclusions: The genetic and environmental contributions to risk of MDD in this representative sample of female adolescent twins are remarkably analogous to findings from adult samples. These results are congruent with a conceptualization of adolescent MDD and adult MDD as having very similar etiologic determinants.

Keywords

Major Depressive Disorder; genetic epidemiology; adolescent twin studies

‘True, I have cried too much; I am heartsick at dawn. The moon is bitter and the sun is sour ...’

Arthur Rimbaud, 1870 (Age 16)

Previous findings

In a recent meta-analysis of adult twin and adoption studies, the liability to Major Depressive Disorder (MDD) in adults was estimated to be partitioned into approximately

Correspondence to Anne L. Glowinski, Washington University School of Medicine, Department of Psychiatry, Division of Child and Adolescent Psychiatry, 40 N. Kingshighway, Suite One, St. Louis, MO 63108, USA, Box 8134; Email: glowinsa@matlock.wustl.edu.

40% due to additive genetic factors and 60% due to non-shared environmental factors, with no evidence of shared environmental factors (Sullivan, Neale, & Kendler, 2000). In contrast, the genetic epidemiology of youth MDD remains largely unelucidated.

Reports on the epidemiology of youth MDD have multiplied in the past 15 years. The prevalence of this disorder is low before puberty and increases steadily during adolescence (e.g., Birmaher et al., 1996; Fleming & Offord, 1990; Kessler, Avenevoli, & Ries Merikangas, 2001). In epidemiological samples, the well-described phenomenon of greater prevalence of MDD in females seen in adult samples (e.g., Kessler, McGonagle, Swartz, Blazer, & Nelson, 1993) does not start until adolescence (e.g., Angold & Worthman, 1993). Evidence is emerging strongly suggesting a continuity of MDD from adolescence to adulthood in young women (e.g., Rao, Hammen, & Daley, 1999), in both genders (e.g., Lewinsohn, Rohde, Klein, & Seeley, 1999) and even when controlling for comorbidity patterns (e.g., Fombonne, Wostear, Cooper, Harrington, & Rutter, 2001).

Family studies of MDD have shown familial aggregation of adolescent MDD in community samples (e.g., Klein, Lewinsohn, Seeley, & Rohde, 2001), but the nature of this familial transmission remains unclear. Thus far, twin studies of youth depression (comprehensively reviewed recently by Rice, Harold, & Thapar, 2002) have not estimated the heritability of youth MDD as they have mostly focused on continuous measures of depression such as scores obtained from non-diagnostic instruments, for instance the parent-reported Achenbach (1991) Child Behavioral Checklist (e.g., Edelbrock, Rende, Plomin, & Thompson, 1995; Hewitt et al., 1997), the child-reported Kovacs Children Depression Inventory (e.g., Rende, Plomin, Reiss, & Hetherington, 1993), or Angold, Costello, Pickles, and Winder's (1987) Mood and Feelings Questionnaire (e.g., Thapar & McGuffin, 1994). Notably, the design of the Virginia Twin Study of Adolescent Behavioral Development (VTSABD) included both parent- and child-reported depressive symptoms. Reports from the VTSABD have suggested significant heritability of depressive symptoms in male and female youth aged 8 to 16 when parental reports are considered but much less when youth reports are considered (Eaves et al., 1997). However, when stratifying the VTSABD sample by age and gender and using a different measure of depression based on the Child and Adolescent Psychiatry Assessment (CAPA) (Angold et al., 1995), greater heritability was found for the 314 twin pairs of adolescent girls (Silberg et al., 1999). Although the CAPA allows for clinical MDD diagnosis, the measurement of depression used in this latter analysis was a subscale score composed of the 9 child-rated symptoms included in the CAPA. For this non-diagnostic measure of depression, the portion of the variance attributed to additive genetic effects was reported to be 28%, with 0% attributed to shared environmental effects and 72% to unique environmental effects.

Methodological issues

Rater effects.

The poor agreement between child- and parent-reported psychopathology when using either diagnostic interviews (Edelbrock, Costello, Dulcan, Conover, & Kala, 1986) or scales (e.g., Achenbach, McConaughy, & Howell, 1987) has been well documented. Notably however, child-reported depressive symptoms have been reported to be more concordant with professional clinical diagnoses than other forms of self-reported child psychopathology (Rubio-Stipec et al., 1994).

One obvious advantage of using adolescent-reported MDD as a phenotype is that it provides the most meaningful comparison to adult studies, which typically use individual reported phenotypes.

Categorical construct of depression.

The cross-sectional categorical construct of DSM-IV (American Psychiatric Association, 1994) Major Depressive Disorder, or previously DSM-III-R (American Psychiatric Association, 1991) Major Depression, has been shown to poorly encapsulate the reality of impairment associated with a continuum of depressive symptoms (as reported for a large population-based sample of adolescents and adults by Lewinsohn, Solomon, Seeley, and Zeiss (2000)) or in adults, of chronic lifetime depression often characterized by an alternation of MDD and subsyndromal depressive disturbances both phenotypically (e.g., Judd et al., 1998) and genetically (e.g., Kendler & Gardner, 1998). Studies which attempt to record lifetime rather than just recent MDD bypass this problem to some degree, although other issues must be considered, notably the issue of recall bias.

Recall bias.

In adult samples, the reliability of a lifetime history of Major Depression has been shown to be only fair (e.g., Foley, Neale, & Kendler, 1998, who reported a kappa of .43 for reliability over 5 years). This is obviously an important issue to consider when examining cohort effects (e.g., Giuffra & Risch, 1994), which are also affected by the problem of recall bias for age at onset (e.g., Stassen, Ragaz, & Reich, 1997). Potential recall bias should be considered when interpreting genetic epidemiological data based on only one cross-sectional assessment.

Data censoring in youth samples.

The issue of data incompleteness or data censoring whereby not all subjects in a sample have passed the age of risk for developing a disorder is crucial in youth samples, particularly for a disorder such as MDD known to steadily increase in prevalence throughout adolescence. Pragmatically this means that data from an 'unaffected' youth should be treated statistically as censored.

Our goals were to investigate the genetic epidemiology of youth MDD in an epidemiologically and genetically informative sample and to compare the fit of models with and without recall bias.

Materials and methods

The data were collected through the Missouri Adolescent Female Twin Study (MOAFTS), a prospective twin-family study of alcohol use, alcohol problems and psychiatric disorders in young adolescent females. The study subjects were ascertained through the Missouri Twin Register, which originated from a systematic review of all Missouri birth records from 1968 and onward. The MOAFTS methodology has been described elsewhere (Heath et al., 1999 and 2002) and is summarized briefly: 95% of all female like-sex twin pairs identified from birth records were located. Twins adopted at birth were excluded from the study. The subjects were 13, 15, 17 and 19 years old at initial enrollment into the study. New cohorts of twins aged 13 were added each year to the study, using a cohort sequential design. Twin zygosity was determined based on parental (usually and whenever feasible the biological mother's) answers to standard interview questions (Loehlin & Nichols, 1975), which is a method shown to yield 95% accuracy when compared to genotyping (Eaves, Eysenck, & Martin, 1989). A total of 3416 female adolescent twins (consisting of about 55% monozygotic (MZ) and 45% dizygotic (DZ)) were interviewed between 1995 and 2000, in the first MOAFTS assessment wave. The participation rate was over 85%. The age at assessment was distributed as follows: 12 (.5%), 13 (34%), 14 (8.4%), 15 (13%), 16 (7.5%), 17 (12.3%), 18 (5.8%), 19 (14.1%), 20 (3.1%), 21 (.9%), 22 (.2%) and 23 (.1%) with an average age of 15.5 (SD: 2.4). The sample included 13% minority subjects, almost

exclusively African American, congruently with the minority composition of the Missouri general population. The general representativeness of this sample was recently investigated by Heath and others (2002a), using 1990 US census block group data for family residence at times of twins' birth, and sampling biases were found to be minimal.

Assessment

The MOAFTS adolescent interview was a telephone adaptation of the Child Semi-Structured Assessment for the Genetics of Alcoholism (C-SSAGA), a polydiagnostic instrument specifically developed for psychiatric genetics research and based on the DSM-IV. The C-SSAGA is adapted from the Diagnostic Interview for Children and Adolescents (DICA; Reich, 1996). This instrument yields lifetime diagnoses. Good validity and reliability data have been published on the DICA (Reich, 2000).

The diagnosis of lifetime DSM-IV MDD was made with a 'Depression' section shown to have excellent inter-rater reliability (Bucholz et al., 1994). This section includes 30 comprehensive question categories, including history of DSM-IV MDD symptoms, dating of onset and recency of symptoms, time, duration and clustering of symptoms for 'worst episode', and functional impairment history during MDD episodes. This section also contains DSM-IV-based exclusion questions (e.g., episode reported to be secondary to a medical illness, bereavement, or substance use). Congruently with the DSM-IV criteria for MDD, respondents were skipped out of the depression section if they reported that they had never had a lifetime period of at least 2 weeks of depressed mood, irritable mood (if before age 18), or loss of interest that represented a change from previous functioning and lasted most of the day almost every day. The age at onset of MDD was derived also from this section, which recorded both age at 'worst episode' but also age at onset of other episodes, if more than one was reported. The MDD diagnoses and ages at onset were derived by computer algorithms for self-reported MDD using the SAS software (SAS Institute, 1990). The MDD diagnosis was based on DSM-IV criteria including the requirement for clinically significant functional impairment.

Statistical analyses

Descriptive analyses.—Frequencies or means were computed using SAS to summarize MDD prevalences, average symptom count, average age at onset, episodes' durations, average number of episodes, and trends in health service use. Measures of twin–cotwin resemblances for MDD were computed based on the SAS outputs, including probandwise concordances (defined as the proportions of cotwins with MDD among the cotwins of twins with MDD) and the tetrachoric correlations (defined as the twin–cotwin correlations for the liability to MDD).

Survival analyses.—Given the wide age spread of this youth sample, survival analysis (where failure to survive is defined as reporting onset of MDD) provides a useful way to visualize how subjects are differentially affected by lifetime MDD as a function of age. To capitalize on the unique information provided by a twin sample, one can also use survival analysis to visualize the development of MDD in subjects grouped by putative genetic risk for MDD. For that purpose, a multilevel dummy variable was created to simultaneously specify zygosity and history of MDD of the respondent's cotwin (level 1 indicated an MZ cotwin with a lifetime history of MDD, level 2 a DZ cotwin with lifetime MDD, level 3 a DZ cotwin without lifetime MDD and level 4 a MZ cotwin without lifetime MDD). Using the Kaplan–Meier method, the survival analysis curves for lifetime MDD were obtained for the whole sample and for the sample stratified by levels of the dummy variable, leading to stratification by degree of genetic risk for MDD: twins with MZ cotwins affected by MDD (highest risk), twins with DZ cotwins affected by MDD (high risk), twins with DZ cotwins

unaffected by MDD (low risk) and twins with MZ cotwins unaffected by MDD (lowest risk).

Genetic analyses

Given this adolescent sample's age distribution, many subjects without lifetime MDD remained at risk for developing adolescent MDD. When estimating twin pairs' concordance or discordance for MDD, simply counting subjects without lifetime MDD as 'non-cases' would ignore this censoring issue. Genetic analyses were conducted in Mx (Neale & Cardon, 1992), using the method of Pickles and others (1994) for the analysis of censored age-of-onset data subject to recall bias. We constructed 5 sets of bivariate contingency tables for MZ and DZ twin pairs corresponding to ages at assessment of 12–14, 15–16, 17–18, and 19 and older. These sets included: a first set for the youngest age group in the sample (twins 12 to 14) with 3 categories (onset of MDD before age 12, onset of MDD between 12 and 14, and 'censored'); a second set for the second age group (twins 15 to 16) with 4 categories (onset of MDD before age 12, onset of MDD between 12 and 14, onset of MDD between 15 and 16, and 'censored'), and similarly a third set for the third age group (twins 17 to 18) with 5 categories and a fourth set for the fourth age group (twins 19 and older) with 6 categories. We then fitted normal liability threshold models to the data, by the method of maximum-likelihood, using Mx (Neale, 1999). We tested models where the phenotypic variance for MDD liability was decomposed into three components: additive genetic factors (A), shared environmental factors (C) (such as attending the same school during childhood or being reared by the same parents) and unique or non-shared environmental factors (E) (individual environmental influences not shared by the twins). We also tested models including only AE or CE and models including dominant genetic factors (D), in addition to A and E factors. The models were compared for goodness of fit by likelihood-ratio chi square and Akaike criteria. Prevalence estimates were derived from the estimated threshold values, using a table of the cumulative normal frequency distribution (Snedecor & Cochran, 1980).

Recall bias analyses

Tests of possible recall bias were conducted by determining whether thresholds (i.e., prevalences) corresponding to a given age-at-onset could be constrained across age groups. Thus, under the most general model, we would estimate 5 thresholds corresponding to the category of onset of depression before age 12 and to the four age groups classified by age at assessment: 12–14, 15–16, 17–18 and 19 and older. With substantial forgetting of an episode of MDD before age 12 by older respondents, or a tendency to recall the onset as having occurred later than in fact was the case, then we would expect to observe significant differences in threshold between the four age-at-assessment groups. If there were no recall bias, it should be possible to constrain those thresholds to be the same for the four age-at-assessment groups, without a significant worsening of fit. For the categories of onset of depression corresponding to ages 12–14 and 15–16, in the absence of recall bias effects, we would expect to be able to constrain thresholds, respectively, across only the three oldest, or two oldest, age-at-assessment groups since a proportion of the younger unaffected respondents in the 12–14 (or 15–16) age at assessment groups would still be expected to have onset of depression during this age window. To examine the impact of recall bias on the estimation of genetic and environmental contribution to risk of MDD, we compared the parameter estimates under the model assuming no recall bias and under models that freed corresponding thresholds across age-at-assessment groups as necessary.

Results

Out of the 3364 subjects who answered the questions in the depression section, 8.9% ($N = 299$) met full criteria for DSM-IV MDD by self-report. The average age of onset of MDD was 14.4 (STD = 2.9). The mean number of episodes per subject reporting lifetime MDD was 1.6 (SD = 1.5). The average duration of the MDD episode reported to be 'the worst' was 14 weeks (STD = 31.2). Only 100 subjects (33.4%) met criteria for MDD without self-reported comorbid psychiatric or substance use disorders (i.e., neither social phobia, conduct disorder, oppositional defiant disorder, nicotine dependence, alcohol dependence, nor significantly impairing non-time-clustered anxiety symptoms), consistent with known patterns of comorbidity in adolescent depression (e.g., Angold, Costello, & Erkanli, 1999). Of the subjects meeting criteria for MDD, 37.9% ($N = 113$) reported seeking help with either a psychiatrist, a psychologist, another doctor, a social worker, a counselor, a clergyman or another professional. Even fewer of the subjects meeting MDD criteria were prescribed medications during their 'worst episode' (17.4% ($N = 52$)), and notably all but one of those subjects who reported being prescribed antidepressants met criteria for MDD in this interview. A small number of subjects ($N = 26$ or 8.7% of the respondents meeting MDD criteria) were hospitalized in a psychiatric facility during their 'worst episode'.

Survival analyses

The Kaplan–Meier age-at-onset curves for MDD are presented in Figure 1 for the whole sample and for the sample stratified by cotwin zygosity and lifetime history of MDD. Genetic influences on liability to MDD are strongly suggested by these curves with earlier onset of MDD and a significantly steeper increase in prevalence for cotwins of MZ twins reporting lifetime MDD, followed by cotwins of DZ twins with MDD, in contrast to the cotwins of DZ twins without MDD who approximate the trajectory of the whole sample and the cotwins of MZ twins without MDD who show a trend for slightly later onset of MDD and a slightly lower prevalence for MDD at all ages after 12.

Genetic analyses

Resemblance measures for MZ versus DZ twin pairs in each age interval are summarized in Table 1. Notably, the prevalences for MDD are not significantly different in MZ versus DZ twins, congruently with our assumption that the phenotypes of MZ twins, DZ twins or other individuals are affected by genetic and environmental influences in a similar way. The finding that the DZ tetrachoric correlation is a little less than half of the MZ is compatible with the contribution of both genetic dominant and additive effects on individual differences in liability to MDD.

Results of genetic model-fitting are summarized in Table 2. The most parsimonious model fitting the data is an AE model, where A and E are set to be equal across all the sample's age groups, and some of the thresholds for age at onset of MDD are allowed to differ for the various age groups. Under this model, MDD is partly heritable (40.4% additive genetic effects, 95%CI: 23.9–55.1), but an even larger portion of the variance is accounted for by non-shared environmental influences (59.6%, 95%CI: 44.9–76.1). Even in the less well-fitting ACE models, the point estimates for C are 0, but the 95% confidence interval for C is broad (0–30%). Models of no familial resemblance (E models) are not shown but provided a very poor fit to the data. Table 3 illustrates the MDD prevalence estimates stratified by age group for each age category derived from the best-fitting AE genetic model and from the AE model where thresholds are fixed (except in the age groups where censoring is an issue). There is evidence for a significant recall bias, with onset of MDD more likely to be reported as having occurred between 12 and 14 by those respondents who had just passed this age and were between 15 and 16 than by those who were age 17 or older at assessment (6.9%

versus 4.7%). Likewise, onset of MDD in the 15–16 interval is more likely to be reported by respondents aged 17–18 than by those 19 and older (11.1% versus 9.4%).

Discussion

Continuity of MDD from adolescence to adulthood

We believe that this report constitutes the first youth twin study genetic modeling of DSM-IV-defined MDD. We find that the genetic architecture of MDD in these female adolescents is strikingly similar to the genetic architecture of MDD in adult samples (e.g., Sullivan et al., 2000). It is characterized by modest but significant heritability of around 40%, with the remaining variance attributable to non-shared environmental effects, which include measurement error. We were able to constrain the estimated additive genetic variance to be the same across age groups ranging from 12–14 to 19 and older. This supports the hypothesis of continuity between adolescent and adult MDD. The finding in some family studies that familial loading for adolescent MDD (defined by presence of parental MDD) is not particularly associated with continuity of MDD into adulthood (e.g., Wickramaratne, Greenwald, & Weissman, 2000) is illustrated by the results in DZ twins of our survival analysis of MDD in this youth sample. By the beginning of adulthood the DZ cotwins of respondents with MDD had only a modestly elevated prevalence of MDD when compared to the whole sample (20.75% versus 16.5% by age 18). It also illustrates dramatically the power of twin studies for deriving information about genetic risk: when familial loading is now defined by having an MZ cotwin with MDD, the prevalence rate by age 18 of MDD is 36%. This group of girls at highest genetic risk also show substantially earlier onset of MDD than the rest of the sample. Notably, under the method we used to conduct these genetic analyses, the heritability estimates extend at the same time to MDD and to the age at onset of MDD.

The shared environment issue

As in many previous studies of adult samples, we obtained a zero estimate for the shared environmental variance. As discussed by Rutter, Pickles, Murray, and Eaves (2001), lack of shared environmental variance must be interpreted intelligently. Firstly, measurement error, which as mentioned earlier does not occur uncommonly in the cross-sectional diagnosis of MDD, will inflate the findings of non-shared environmental effects, notably at the expense of shared environmental effects. Secondly, if we analyze for instance a phenotype consisting of reporting either 2 weeks of substantial sadness, anhedonia or irritability (with the same method we used to analyze the categorical diagnosis of MDD), we cannot discard the influence of shared environment on the liability to such a phenotype; finding the best-fitting model allows for 24% additive genetic effects, 30% shared environmental effects and 45% non-shared environmental effects. This suggests that there might be family environments where young people are at high risk of experiencing the same stress and demoralization, but that the actual development of a disorder such as adolescent MDD is more dependent on individual factors (genetic or others). This is recapitulated by Rutter and others (2001): ‘Thus if a familywide influence such as discord or poverty tends to impinge differentially on individual children, the effects will be largely non-shared (even though the influence, as measured, is family wide or may appear as a genetic effect; ...)’. Another third crucial point to consider when interpreting the consistency of this pattern of zero shared environmental variance for MDD across multiple samples, is that there probably are non-additive genetic effects (either dominance or epistasis) as well as additive genetic effects on risk of MDD (Heath et al., 2002b). In the twin design, we decompose the phenotypic variance into additive genetic factors (A), shared environmental factors (C), non-shared environmental factors (E) and dominant genetic factors (D). In the absence of separated (at birth) twin data, only three of these components can be estimated in the same model: A, C and E or A, D and

E, since we have three independent statistics (MZ correlation, DZ correlation and total variance standardized to unity, from which E is estimated). Thus it is not possible to model the contribution of all these components at once. E can never be dropped from our model fitting (as that would predict perfect MZ twin correlations), so we can evaluate the fit of ADE models or ACE models but not of models containing A, D and C at the same time. Given that the DZ tetrachoric correlation is slightly less than half the MZ correlation, and that the ADE models provide reasonable fits to the data (though inferior to AE models), there is a weak suggestion of genetic dominance in these data, but too weak an effect to be confirmed with model fitting with these sample sizes. It is significant enough, though, to potentially lead to the inflation of additive genetic influences estimates and the masking of the shared environment component in an ACE model, when the greater phenotypic concordance in MZ twins is actually partially due to non-additive genetic factors (i.e., dominant factors).

Recall bias

Our findings suggest that about $100\% - 4.7\% / 6.9\% = 31.9\%$ of the subjects 17 and older and $100\% - 9.4\% / 11.1\% = 15.3\%$ of subjects 19 and older could be either forgetting their first episode of MDD or overestimating its date to displace it in the age category immediately preceding their own. (As illustrated in Table 3, the MDD prevalence between the ages of 12 and 14, for instance, is 4.7% when derived from the reports of subjects 17 and older, in contrast to an MDD prevalence of 6.9% between the ages of 12 and 14 when examining the reports of subjects between 15 and 16.) Follow-up data will be required to determine whether forgetting, or recall of an episode as having occurred more recently than in fact was the case, has occurred. This latter phenomenon has also been described by Pickles and others (1994) and is called by some a ‘bo-peep’ effect. If forgetting is responsible for our observed findings, then this effect does not appear to increase with increasing age. Prevalences of MDD before age 12 and in the 12–14 age group were the same as in the 19 and older and 17–18 age groups, i.e., we did not observe any further increase in the 19 and older age group. In either case, the very similar estimates obtained in models adjusted or unadjusted for recall bias provide little evidence that recall problems are leading to biased estimates of genetic parameters.

Limitation and future directions

Despite the evidence that DSM-IV MDD is genetically correlated with subsyndromal depressive disturbances, we did not include subthreshold syndromes in our analyses. In the MOAFTS interview, subjects who do not meet initial criteria for MDD (i.e., at least 4 DSM-IV symptoms which cluster in time continually for at least 2 weeks with either 2 weeks or more of depression, irritability (before age 18) or anhedonia) are not asked questions about functional impairment. We therefore elected to limit our analysis to a clinically meaningful phenotype, which at least allows a valid comparison to adult findings. We also limited our analyses to self-reported DSM-IV MDD and chose not to incorporate parental reports into our analyses. Notably, we initially explored the nature of parental reports and found a predictably low agreement between youth and parental reports of MDD ($\kappa = .23$). We also find that parents with MDD appear to over-report MDD in their children with relative risk of parental MDD being 6.0 for parent-reported youth MDD versus 1.7 for youth-reported MDD.

It is important to comment that while our model fitting takes into account both MDD and its age of onset, it does not address the issue of prepubertal versus postpubertal onset of MDD given the age of our sample and the very small numbers of recalled prepubertal onset MDD. This does not allow an independent genetic analysis of prepubertal onset MDD (a total of 2 MZ pairs and 0 DZ pairs report concordance for prepubertal onset MDD). Also, we cannot

currently contribute to the literature on early onset recurrent MDD, defined as at least 2 major depressive episodes and onset before age 25 (e.g., Zubenko, Zubenko, Spiker, Giles, & Kaplan, 2001; Zubenko, Hughes, Stiffler, Zubenko, & Kaplan, 2002): The numbers for recurrence are too small for separate genetic analyses with 5 MZ pairs endorsing concordance for recurrent MDD in contrast to no DZ pairs endorsing concordance for recurrent MDD. The analysis of recurrence will become interesting once the sample has aged. This is a cross-sectional assessment only and one would expect that the heritability of depression could change once several MOAFTS waves are analyzed together. Finally, our design would be more informative with the inclusion of adolescent male twin pairs.

In conclusion, a lifetime DSM-IV MDD phenotype is common in girls by the end of adolescence. Girls with a high genetic risk of MDD are at a significantly higher risk than other girls for both MDD and earlier onset of the disorder; however, that genetic risk cannot easily be evaluated from a family history of MDD as is often done in clinical practice. The fact that a significant portion of a liability to youth MDD in adolescent girls appears due to non-shared environmental factors makes a creative search for these environmental etiological factors particularly important (as suggested by Rutter, Silberg, O'Connor, & Simonoff, 1999). Explorations of environmental factors' effects in children at high genetic risk are likely to be particularly informative.

Acknowledgments

This research was supported by National Institutes of Health (NIH) grants AA09022, AA07728 and AA11998 to Andrew Heath, NIH grant DA12540 to Pam Madden, NIH grant MH64134 to Anne Glowinski, the Klingenstein Third Generation Foundation (KTGF) Award in adolescent depression and the National Alliance for Research on Schizophrenia and Depression (NARSAD) Young Investigator Award (2001 Callaghan Award) to Anne Glowinski.

References

- Achenbach, T.M. (1991). *Manual for the Child Behavior Checklist/4-18 and 1991 Profile* Burlington: University of Vermont Department of Psychiatry.
- Achenbach TM, McConaughy SH, Howell CT. Child/adolescent behavioral and emotional problems: Implications of cross-informant correlations for situational specificity. *Psychological Bulletin*. 1987; 101:213–232. [PubMed: 3562706]
- American Psychiatric Association. (1991). *Diagnostic and statistical manual of mental disorders, revised 3rd edition (DSM-III-R)* Washington, DC: American Psychiatric Association.
- American Psychiatric Association. (1994). *Diagnostic and statistical manual of mental disorders, 4th edition (DSM-IV)* Washington, DC: American Psychiatric Association.
- Angold, A., Costello, E.J., Pickles, A., & Winder, F. (1987). *The development of a questionnaire for use in epidemiological studies of depression in children and adolescents* London: Medical Research Council Child Psychiatry Unit.
- Angold A, Worthman C. Puberty onset of gender differences in rates of depression: A developmental, epidemiological, and neuroendocrine perspective. *Journal of Affective Disorders*. 1993; 29:145–158. [PubMed: 8300975]
- Angold A, Prendergast A, Cox R, Harrington E, Simonoff E, Rutter M. The Child and Adolescent Psychiatric Assessment (CAPA). *Psychological Medicine*. 1995; 25:739–753. [PubMed: 7480451]
- Angold A, Costello EJ, Erkanli A. Comorbidity. *Journal of Child Psychology and Psychiatry*. 1999; 40:57–87. [PubMed: 10102726]
- Birmaher B, Ryan ND, Williamson DE, Brent DA, Kaufman J, Dahl RE, Perel J, Nelson B. Childhood and adolescent depression: A review of the past 10 years. Part I *Journal of the American Academy of Child and Adolescent Psychiatry*. 1996; 35:1427–1439.
- Bucholz KK, Cadoret R, Cloninger CR, Dinwiddie SH, Hesselbrock VM, Nurnberger JI Jr, Reich T, Schmidt I, Schuckit MA. A new, semistructured psychiatric interview for use in genetic linkage

- studies: A report on the reliability of the SSAGA. *Journal of Studies on Alcohol*. 1994; 55:149–158. [PubMed: 8189735]
- Eaves, L.J., Eysenck, H.J., & Martin, N.G. (1989). *Genes, culture and personality: An empirical approach* London: Academic Press.
- Eaves LJ, Silberg JL, Meyer JM, Maes HH, Simonoff E, Pickles A, Rutter M, Neale MC, Reynolds CA, Erikson MT, Heath AC, Loeber R, Truett KR, Hewitt JK. Genetics and developmental psychopathology: 2. The main effects of genes and environment on behavioral problems in the Virginia Twin Study of Adolescent Behavioral Development. *Journal of Child Psychology and Psychiatry*. 1997; 38:965–980. [PubMed: 9413795]
- Edelbrock C, Costello AJ, Dulcan MK, Conover NC, Kala R. Parent–child agreement on child psychiatric symptoms assessed via structured interview. *Journal of Child Psychology and Psychiatry*. 1986; 27:181–190. [PubMed: 3958075]
- Edelbrock C, Rende R, Plomin R, Thompson LA. A twin study of competence and problem behavior in childhood and early adolescence. *Journal of Child Psychology and Psychiatry*. 1995; 36:775–785. [PubMed: 7559844]
- Fleming JE, Offord DR. Epidemiology of childhood depressive disorders: A critical review. *Journal of the American Academy of Child and Adolescent Psychiatry*. 1990; 29:571–580. [PubMed: 2201675]
- Fombonne E, Wostear G, Cooper V, Harrington R, Rutter M. The Maudsley long-term followup of child and adolescent depression. 1. Psychiatric outcomes in adulthood. *British Journal of Psychiatry*. 2001; 179:210–217. [PubMed: 11532797]
- Foley DL, Neale MC, Kendler KS. Reliability of a lifetime history of major depression: Implications for heritability and co-morbidity. *Psychological Medicine*. 1998; 28:857–870. [PubMed: 9723141]
- Giuffra LA, Risch N. Diminished recall and the cohort effect of major depression: A simulation study. *Psychological Medicine*. 1994; 24:375–383. [PubMed: 8084933]
- Heath AC, Madden PAF, Grant JD, McLaughlin TL, Todorov AA, Bucholz KK. Resiliency factors protecting against teenage alcohol use and smoking: Influences of religion, religious involvement and values, and ethnicity in the Missouri Adolescent Female Twin Study. *Twin Research*. 1999; 2:145–155. [PubMed: 10480749]
- Heath AC, Howells W, Bucholz KK, Glowinski AL, Nelson EC, Madden PAF. Ascertainment of a Mid-Western US female adolescent twin cohort for alcohol studies: Assessment of sample representativeness using birth record data. *Twin Research*. 2002a; 5:107–112. [PubMed: 11931688]
- Heath, A.C., Madden, P.A.F., Bucholz, K.K., Nelson, E.C., Todorov, A.A., Price, R.K., Whitfield, J.B., & Martin, N.G. (2002b). Genetic and environmental risks of dependence on alcohol, tobacco and other drugs. In R. Plomin, I. Craig, J. DeFries, & P. McGuffin (Eds.), *Behavioral genetics in the postgenomic era* Washington, DC: American Psychological Association.
- Hewitt JK, Silberg JL, Rutter M, Simonoff E, Meyer JM, Maes H, Pickles A, Neale MC, Loeber R, Erickson MT, Kendler KS, Heath AC, Truett KR, Reynold CA, Eaves LJ. Genetics and developmental psychopathology: 1. Phenotypic assessment in the Virginia Twin Study of Adolescent Behavioral Development. *Journal of Child Psychology and Psychiatry*. 1997; 38:943–963. [PubMed: 9413794]
- Judd LL, Akiskal HS, Maser JD, Zeller PJ, Endicott J, Coryell W, Paulus MP, Kunovac JL, Leon AC, Mueller TI, Rice JA, Keller MB. A prospective 12-year study of subsyndromal and syndromal depressive symptoms in unipolar major depressive disorders. *Archives of General Psychiatry*. 1998; 55:694–700. [PubMed: 9707379]
- Kendler KS, Gardner CO Jr. Boundaries of major depression: An evaluation of DSM-IV criteria. *American Journal of Psychiatry*. 1998; 155:172–177. [PubMed: 9464194]
- Kessler RC, McGonagle KA, Swartz M, Blazer DG, Nelson CB. Sex and depression in the National Comorbidity Survey I: Lifetime prevalence, chronicity and recurrence. *Journal of Affective Disorders*. 1993; 29:85–96. [PubMed: 8300981]
- Kessler RC, Avenevoli S, Ries Merikangas K. Mood disorders in children and adolescents: An epidemiologic perspective. *Biological Psychiatry*. 2001; 49:1002–1014. [PubMed: 11430842]

- Klein DN, Lewinsohn PM, Seeley JR, Rohde P. A family study of major depressive disorder in a community sample of adolescents. *Archives of General Psychiatry*. 2001; 52:13–20. [PubMed: 11146753]
- Lewinsohn PM, Rohde P, Klein DN, Seeley JR. Natural course of adolescent major depressive disorder: I. Continuity into young adulthood. *Journal of the American Academy of Child and Adolescent Psychiatry*. 1999; 38:56–63. [PubMed: 9893417]
- Lewinsohn PM, Solomon A, Seeley JR, Zeiss A. Clinical implications of ‘subthreshold’ depressive symptoms. *Journal of Abnormal Psychology*. 2000; 109:345–351. [PubMed: 10895574]
- Loehlin, J.C., & Nichols, R.C. (1975). *Heredity, environment and personality* Austin, TX: University of Texas Press.
- Neale, M.C., & Cardon, L.R. (1992). *Methodology for genetic studies of twins and families, NATO ASI series* Dordrecht, The Netherlands: Kluwer Academic Publishers.
- Neale, M.C. (1999). *Mx: Statistical modeling* Richmond, Virginia: Department of Psychiatry, Virginia Commonwealth University.
- Pickles A, Neale M, Simonoff E, Rutter M, Hewitt J, Meyer J, Crouchley R, Silberg J, Eaves L. A simple method for censored age-of-onset data subject to recall bias: Mothers’ reports of age of puberty in male twins. *Behavior Genetics*. 1994; 24:457–468. [PubMed: 7993322]
- Rao U, Hammen C, Daley SE. Continuity of depression during the transition to adulthood: A 5-year longitudinal study of young women. *Journal of the American Academy of Child and Adolescent Psychiatry*. 1999; 38:908–915. [PubMed: 10405510]
- Reich, W. (1996). *Diagnostic Interview for Children and Adolescents (DICA)* St. Louis, MO: Washington University Division of Child Psychiatry.
- Reich W. Diagnostic Interview for Children and Adolescents (DICA). *Journal of the American Academy of Child and Adolescent Psychiatry*. 2000; 39:59–66. [PubMed: 10638068]
- Rende RD, Plomin R, Reiss D, Hetherington EM. Genetic and environmental influences on depressive symptomatology in adolescence: Individual differences and extreme scores. *Journal of Child Psychology and Psychiatry*. 1993; 34:1387–1398. [PubMed: 8294525]
- Rice F, Harold G, Thapar A. The genetic aetiology of childhood depression: A review. *Journal of Child Psychology and Psychiatry*. 2002; 43:65–79. [PubMed: 11848337]
- Rimbaud, A. (1870). ‘Le bateau ivre’ [‘The drunken boat’]. In *Arthur Rimbaud Complete Works* (trans Paul Schmidt) New York: Harper & Row Publishers, 1976
- Rubio-Stipek M, Canino GJ, Shrout P, Dulcan M, Freeman D, Bravo M. Psychometric properties of parents and children as informants in child psychiatry epidemiology with the Spanish Diagnostic Interview Schedule for Children (DISC. 2). *Journal of Abnormal Child Psychology*. 1994; 22:703–720. [PubMed: 7876458]
- Rutter M, Silberg J, O’Connor T, Simonoff E. Genetics and child psychiatry: II Empirical research findings. *Journal of Child Psychology and Psychiatry*. 1999; 40:19–55. [PubMed: 10102725]
- Rutter M, Pickles A, Murray R, Eaves L. Testing hypotheses on specific environmental causal effects on behavior. *Psychological Bulletin*. 2001; 127:291–324. [PubMed: 11393298]
- SAS Institute. (1990). *SAS/STAT User’s Guide, Version 6* (4th edn). Cary, NC: SAS Institute, Inc.
- Silberg J, Pickles A, Rutter M, Hewitt J, Simonoff E, Maes H, Carbonneau R, Murrelle L, Foley D, Eaves L. The influence of genetic factors and life stress on depression among adolescent girls. *Archives of General Psychiatry*. 1999; 56:225–232. [PubMed: 10078499]
- Snedecor, G.W., & Cochran, W.G. (1980). *Statistical methods* (7th edn). Ames, IA: The Iowa State University Press.
- Stassen HH, Ragaz M, Reich T. Age-of-onset or age-cohort changes in the lifetime occurrence of depression? *Psychiatric Genetics*. 1997; 7:27–34. [PubMed: 9264135]
- Sullivan PF, Neale MC, Kendler KS. Genetic epidemiology of major depression: Review and meta-analysis. *American Journal of Psychiatry*. 2000; 157:1552–1562. [PubMed: 11007705]
- Thapar A, McGuffin P. A twin study of depressive symptoms in childhood. *British Journal of Psychiatry*. 1994; 165:259–265. [PubMed: 7953041]

- Wickramaratne PJ, Greenwald S, Weissmen MM. Psychiatric disorders in the relatives of probands with prepubertal-onset or adolescent-onset major depression. *Journal of the American Academy of Child and Adolescent Psychiatry*. 2000; 36:1396–1405. [PubMed: 11068895]
- Zubenko GS, Zubenko WN, Spiker DG, Giles DE, Kaplan BB. Malignancy of recurrent, early-onset major depression: A family study. *American Journal of Medical Genetics*. 2001; 105:690–699. [PubMed: 11803516]
- Zubenko GS, Hughes HB, Stiffler JS, Zubenko WN, Kaplan BB. Genome survey for susceptibility loci for recurrent, early-onset major depression: Results at 10cM resolution. *American Journal of Medical Genetics*. 2002; 114:413–422. [PubMed: 11992563]

Figure 1. Age at onset of MDD in whole sample and in sample stratified by zygosity and cotwin history of lifetime MDD

Table 1

Measures of twin–cotwin resemblance

Age group	MZ pairs (855 pairs)			DZ pairs (661 pairs)		
	<i>N</i> pairs	MDD prevalence	Probandwise concordance	<i>N</i> pairs	MDD prevalence	Probandwise concordance
12–14	388	.025	.1	305	.025	0
15–16	183	.08	.21	146	.10	.27
17–18	149	.15	.33	105	.13	.17
Over 19	135	.17	.45	105	.16	.13
Tetrachoric correlations and 95% confidence intervals						
			.49 (.32–.64)			.22 (0–.42)

Table 2

Comparison of best models of familial resemblance for lifetime MDD

	A 95%CI	C 95%CI	D 95%CI	E 95%CI	χ^2	df	AIC	p
ACE model: family resemblance due to additive genetic and shared environmental effects. Age at onset thresholds constrained	42(4–57)	0(0–30)	–	58(43–75)	142.6	154	–165.4	.73
CE model: family resemblance due only to shared environmental effects. Age at onset thresholds constrained	–	31(17–44)	–	69(56–83)	147.2	155	–162.8	.66
ADE model: family resemblance due to additive and dominant genetic effects. Age at onset thresholds constrained	6(0–55)	–	38(0–59)	56(41–73)	142.0	154	–166.0	.75
AE model: family resemblance only due to additive genetic effects. Age at onset thresholds constrained	42(25–57)	–	–	58(43–75)	142.6	155	–167.4	.75
ACE model Age at onset thresholds relaxed	41(5–56)	0(0–28)	–	59(44–76)	134.2	148	–161.8	.78
ADE model Age at onset thresholds relaxed	0(0–54)	–	44(0–58)	59(44–76)	133.3	148	–162.7	.80
AE model Age at onset thresholds relaxed	41(24–56)	–	–	59(44–76)	134.2	149	–163.8	.80
ACE model Some age at onset thresholds relaxed	40(4–55)	0(0–29)	–	60(45–76)	135.0	152	–169.0	.84
ADE model Some age at onset thresholds relaxed	.7(0–54)	–	42(0–58)	57(0–58)	134.2	152	–170.0	.85
AE model Some age at onset thresholds relaxed	40(24–55)	–	–	60(45–76)	135.0	153	–171.0	.85

A: additive genetic variance, C: shared environmental variance, D: dominant genetic variance, E: non-shared environmental variance, AIC: Aikake information criterion, df: degrees of freedom.

Table 3

Model-based estimates of the lifetime prevalence of MDD as a function of age, without versus with adjustment for recall bias

		Lifetime prevalence by age (%)				
		Before 12	12–14	15–16	17–18	19 and over
<i>Unadjusted:</i>						
Age group:	12–14	1.0%	2.6%	–	–	–
	15–16	1.0%	6.9%	9.3%	–	–
	17–18	1.0%	4.7%	11.1%	13.8%	–
	19 and over	1.0%	4.7%	9.4%	15.4%	17.4%
<i>Adjusted:</i>						
Age group:	12–14	1.0%	2.6%	–	–	–
	15–16	1.0%	5.8%	8.0%	–	–
	17–18	1.0%	5.8%	10.8%	13.4%	–
	19 and over	1.0%	5.8%	10.8%	17.4%	19.5%