Published in final edited form as:

Int Psychogeriatr. 2011 August ; 23(6): 923–929. doi:10.1017/S1041610210002322.

Chronic Pain Conditions and Depression in the Ibadan Study of Ageing

Bibilola D. Oladeji [(Dr)], Victor A. Makanjuola [(Dr)], Oluyomi B. Esan [(Dr)], and Oye Gureje [(Prof)]

Department of Psychiatry, University College Hospital, College of Medicine, University of Ibadan, Ibadan

Abstract

Background—Chronic pain is quite common in the elderly and is often associated with co-morbid depression, limitation of functioning and reduced quality of life. The aim of this study was to ascertain whether there is a differential risk of depression among persons with pain in different anatomical sites and to determine which pain conditions are independent risk factors for depression.

Methods—Data is from the Ibadan Study of Ageing (ISA), a community based longitudinal survey of persons aged 65 years and older from eight contiguous Yoruba Speaking states in Nigeria (n = 2152). Data was collected in face-to-face interviews; depression was assessed using the World Mental Health initiative version of the Composite International Diagnostic Interview (CIDI) while chronic pain was assessed by self-report (response rate = 74%).

Results—Estimates of persistent pain (lasting more than 6 months), in different anatomical sites range from 1.3% to 12.8%, with the commonest being joint pains (12.8%), neck or back (spinal) pain (7.6%) and chest pain (3.0%). Significantly more pain conditions were reported by females and by respondents who were aged over 80 years. The risk for depression was higher in respondents with spinal, joint and chest pain. However, only chest pain was independently associated with depression after adjustments were made for pains at other sites and for functional disability.

Conclusion—Our data suggests that, among elderly persons, there is a differential association of depression with chronic pain that is related to the anatomical site of the pain.

Keywords

Chronic pain; Depression; Disability; Elderly; Epidemiology

INTRODUCTION

Pain conditions are common in the elderly. Reported estimates of chronic pain range between 20% and 50% in elderly populations (Chou, 2007; Mossey and Gallagher, 2004; Reyes-Gibby *et al.*, 2002; Zancocci *et al.*, 2008). Many authors believe the documented

Corresponding Author: Bibilola Oladeji, Department of Psychiatry, College of Medicine, University of Ibadan, Ibadan, bibideji@yahoo.com, Telephone: 234 802 332 4989, Fax: 234-2-2413545.

Description of Author Roles

O. Gureje designed the study and supervised data collection. B. Oladeji and O Gureje interpreted the data and wrote the first draft. V. Makanjuola and O. Esan assisted in reviewing the manuscript. All authors approved the final draft for submission.

Conflicts of interest

None

prevalence of pain in elderly populations may be an underestimate because older adults are less likely to complain of pain as it is seen as a normal, expected consequence of ageing (Gagliese and Melzack, 1997; Melding, 2004). The under reporting of pain in elderly populations are reportedly attributed to the presence of other medical problems, cognitive impairment, presence of depression and the definition of what constitutes chronic pain (Gagliese and Melzack, 1997; Mantyselka *et al.*, 2004). General population studies of the elderly report that pain tends to be persistent over time (Geerlings *et al.*, 2002) and has significant adverse impact on health outcomes, functioning, and quality of life (Reyes-Gibby *et al.*, 2002; Zanolchi *et al.*, 2008).

Pain conditions frequently co-exist with depression in community dwelling elderly (Bair *et al.*, 2003; Onder *et al.*, 2005). The comorbidity of pain and depression is associated with worse outcomes than either condition alone (Arnow *et al.*, 2006). Comorbidity is also more likely to be associated with insomnia, longer duration of depressive symptoms, more intense pain, disability and reduced functioning (Leong *et al.*, 2007; Mossey and Gallagher, 2004; Ohayon, 2004). Individuals with pain and depression are less likely to seek mental health care, to incur higher total health care costs and to use more complementary and alternative treatments (Bao *et al.*, 2003; Emptage *et al.*, 2005).

Epidemiological studies also show that chronic pains commonly co-occur and that the occurrence of multiple pains increases the risk of comorbid mental health conditions. Despite these well established relationships between pain and depression, little is known about the differential risk of depression with pain at different anatomic sites, or whether the association of depression with a particular pain condition is partly or fully accounted for by the presence of other pain conditions or associated disability. It is of course plausible to expect that the presence of functional limitation may influence the association of pain with depression. Also, there is some evidence that the association of pain with depression may be partly influenced by the anatomic site of the pain and the presence or absence of associated functional limitation. For example, Blay *et al.* (Blay *et al.*, 2007) found increased psychiatric morbidity in elderly persons with chest, head or joint pain compared with those with abdominal and back pains while Ohayon and Schatzberg (Ohayon and Schatzberg, 2003), reporting on a general adult population in some European countries, found higher odds for depression among persons with backache or headaches compared with persons with gastrointestinal and limb pains. There is therefore a need to consider anatomical sites as well as functional limitation in exploring the association of pain with depression.

The developing world is currently witnessing a demographic transition with the rapid growth in elderly populations. According to the WHO, of the current population of 580 million for persons aged 60 years and over worldwide, 355 million live in developing countries. By 2020, the projected population of 1 billion elderly persons worldwide will consist of 700 million living in developing countries. There is need for empirical data on mental and physical health needs of the elderly to enable countries prioritize the allocation of health service resources for the elderly. Chronic pain, with its associated disability, will be an important health issue for this growing elderly population and about which information is required for health planning for the population. In this report, we present prevalence estimates of chronic pain in different anatomic sites, examine the relationship of the pain conditions to major depressive disorder, and determine the mediating role of functional impairment in the association between pain conditions and depression.

METHODS

This report is based on data from the Ibadan Study of Aging (ISA). The ISA is a community-based, longitudinal survey of the profile and determinants of successful aging.

The ISA examines the mental and physical health status as well as functioning and disability in elderly persons aged 65 years and over. The details of the methods of the survey have been described in full in earlier reports (Bekibele and Gureje, 2008; Gureje *et al.*, 2007) and only a summary is provided here.

Sample

The survey is being conducted in the Yoruba-speaking areas of Nigeria, comprising of eight contiguous states, six in the south-west (Lagos, Ogun, Osun, Ondo, Oyo and Ekiti) and two in the north-central (Kogi and Kwara) regions of the country. These states account for approximately 22% of the Nigerian population. Nigeria has a population of approximately 130 million people; persons aged 65 years and older constitute about 4% of the population.

A four-stage area probability sampling of households was used to select respondents aged 65 years and over. In households with more than one eligible person (≥ 65 and fluent in Yoruba, the language of the study), the Kish table selection method was used to select one respondent. The baseline survey was fielded between November 2003 and August 2004. On the basis of the selection procedure, face-to-face interviews were conducted on a total of 2152 respondents who provided consent, representing a response rate of 74.2%. Non-response was due to non-availability after repeated visits (14%), interviewers' inability to trace the original address (4%), death (3%), physical incapacitation (2%) and refusal (2%).

The survey was approved by the University of Ibadan/University College Hospital Joint Ethical Review Board.

Measures

Chronic Pain was assessed by self-report. Respondents were asked whether they had any chronic pain, defined as persistent pains present most of the time for a period of 6 months or more during the previous year that interfered with daily activities or caused significant worry. The checklist ascertained the presence of pain at different sites; back or neck, chest, stomach or abdomen, joint, headache and a residual category of pain in any other body parts.

Socio-demographic characteristics included the variables age, sex, marital status, years of education, area of residence and economic status. Economic status was assessed by taking an inventory of a list of 24 household and personal items such as buckets, radio, chairs, clocks, stoves or cookers, telephones, cars or motorbikes present in the household. This is a standard and validated method for estimating economic wealth of elderly persons in low-income settings (Ferguson *et al.*, 2003). The economic status of respondents was categorized by relating each person's possessions to the median number of possessions of the entire sample. Thus the economic status was rated low when its ratio to the median was 0.5 or less, low average if the ratio was 0.5-1, high average if it was 1.0-2.0 and high if it was greater than 2.0.

Depression was assessed using the World Health Organization Composite International Diagnostic Interview (CIDI) Version 3, a fully structured, lay-administered diagnostic interview. The diagnosis of major depression was based on meeting the diagnostic criteria of the Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (DSM-IV) in the preceding 12 months. DSM-IV organic exclusion rules were imposed in making the diagnosis of depression. Judgments about which organic factors could explain a depressive syndrome (organic conditions such as anemia, infections *e.t.c.*) were made by a psychiatrist after clinical review of all questionnaires that had endorsements of depressive features. Respondents with probable dementia (diagnosed using a previously validated cognitive screen) were excluded from the analysis.

Functional disability was rated in two domains. Activities of daily living (ADL) was rated using the Katz index of independence in ADLs, in which the activities assessed included bathing, dressing, toileting, rising and transferring, continence, and feeding. Instrumental activities of daily living (IADLs) were rated on a scale of seven items adapted from the Nagi Physical Performance scale and the Health Assessment questionnaire; items rated included walking around the house, climbing a flight of stairs, reaching above the head to carry something as heavy as 10 pounds, stooping, gripping small objects with the hands, shopping, and activities such as sweeping the floor with a broom or pruning grass around the yard.

The interviews were conducted by 24 trained interviewers, all of whom had at least a high school education. Many of them had been involved in previous field surveys and had experience of face-to-face interviews. Interviewers had two weeks of training prior to the commencement of the survey which consisted of initial 6-day training by OG which included item by item description of the questionnaires and role play. This was followed by a further two days of debriefing and review after each interviewer had done two practice interviews in the field.

Statistical Analysis

To take into account the stratified multistage sampling procedure with the associated clustering, weights were derived and applied to the rates presented in this report (Heeringa *et al.*, 2008). In addition, adjustments were made for the differences between the sample and the total Nigerian population (according to the 2000 United Nations projections) by post-stratification to the target sex and age range. The derived weights were adjusted to normal values to reset the sum of weights back to the original sample size of 2152. Individuals who had a diagnosis of probable dementia (255 respondents) were excluded leaving 1897 records for analysis.

The complex sample design and weighting were accounted for in the analysis by making use of the jackknife replication method implemented with the STATA version 7.0 to estimate standard errors for proportion. Demographic correlates were explored with logistic regression analysis and estimates of standard errors of odds ratios were made with STATA. All the reported confidence intervals are adjusted for design effects.

RESULTS

Prevalence of Pain

The sample consisted of 52.5% male and had a mean age of 75.0 (s.d. 9.2) years. Chronic pain was reported by 315 subjects (16.6%). The most common chronic pain conditions were joint pain (12.8%), neck or back pain (7.6%) and chest pain (3.0%).

Associations of Chronic Pain and Demographic Variables

Table 1 shows the association of chronic pain conditions with age, sex and economic status. Females reported more pain conditions than males (20.1% and 14.1% respectively; $p=0.01$) Headache and neck or back pain were significantly more common in women than men ($p=0.003$ and 0.014 respectively).

Increasing age was associated with increasing pain prevalence with the trend being significant for neck or back pain, joint pain, chest pain as well as the presence of pain in any anatomical site. In general, respondents in the oldest age group (80 years) had higher rates of pain at all the different anatomic sites except for stomach or abdominal pain.

Even though the prevalence of pain was lower in respondents from the highest economic groups, the trend was only significant for stomach or abdominal pain ($p=0.015$).

Pain and Depression

The 12-month prevalence estimate for major depressive disorder in those reporting a chronic pain condition is 11.9% whereas it is 7.1% for the entire sample and 6.7% in those with no pain. In logistic regression analyses, adjusting for age and sex, there was a two-fold increased likelihood of comorbid depression in persons with joint pain and neck or back pain and a three-fold increased likelihood among those with chest pain (Table 2). When the presence of other pain conditions were taken into consideration in the analysis, only chest pain remained significantly associated with major depression (table 3).

To assess whether the presence of disability mediates the association between pain and depression, multivariate analyses were carried out that controlled for the occurrence of ADL and IADL disability. The risk for depression among individuals with any chronic pain remained unchanged when controls were introduced for both ADL (OR(95%CI)-1.8 [1.1-3.0]) and IADL (OR(95%CI)- 1.9[1.1-3.1]) disability.

As shown in table 4, while the presence of ADL disability slightly attenuated the relationship of chest pain to depression, the risk remains unchanged when the presence of IADL disability was taken into consideration.

DISCUSSION

To our knowledge, this is the first large scale study to consider the prevalence and correlates of pain in a representative sample of community-dwelling elderly persons in Sub-Saharan Africa. Close to one in five elderly persons reported pain that was persistent for at least six months and interfered with their daily activities. The demographic profile of respondents who are more likely to experience chronic pain include being female, older age and being socio-economically disadvantaged.

Comparison of chronic pain prevalence across studies is somewhat difficult as different studies have used different definitions for chronic pain. However, our results are comparable to those from studies which incorporated interference with daily activities into their definition of pain. For example Reyes-Gibby et al (Reyes-Gibby *et al.*, 2002) in a population based survey of the elderly in the USA reported a prevalence of 20% for pain resulting in significant activity limitation lasting up to twelve months and Mossey and Gallagher (Mossey and Gallagher, 2004) in a longitudinal study of residents in a continuing care retirement community, reported chronic activity limiting pain in 37% of the population who reported pain at three or more assessments six months apart.

In keeping with many studies from both general adult (Gureje *et al.*, 1998; Tsang *et al.*, 2008) and elderly populations (Brochet *et al.*, 1998; Chou, 2007; Reyes-Gibby *et al.*, 2002), persistent pain was more often reported by females in this sample. The reason for this increased burden of pain in females is unknown. Only a few studies have attempted to explain the reasons for this gender difference in the experience of pain. Brochet et al (Brochet *et al.*, 1998) in their study attributed the gender difference to the difference in the type of pain experienced by men and women. In their study, women more often experienced persistent pain while men tend to more frequently report episodic pain. There is a possibility that this could be a factor in this study as our definition of pain focused on persistent pain hence there might be the possibility that more men with pain were screened out. Other authors have suggested that the higher prevalence of pain in females might be due to greater sensitization to pain and its reporting possibly arising from hormonal and/or psychosocial

factors specific to the female gender (Breivik *et al.*, 2006). The aetiological basis for the increased prevalence of pain in women warrants further investigation.

Most epidemiologic studies of pain complaints in the community have found that pain peaks in the middle ages and declines thereafter (Gallagher *et al.*, 2000; Thomas *et al.*, 2004). However the prevalence of persistent pain, joint pain and pain interfering with daily activities increases with increasing age (Ohayon and Schatzberg, 2003; Sternbach, 1986; Thomas *et al.*, 2007). This age related increase in the occurrence of persistent activity-limiting pain was also demonstrated in this population of elderly persons from a developing country which was more pronounced for joint and chest pain where there was a stepwise increase in the reporting of pain with increasing age. Given the effect that joint pain may have for mobility and the central role of mobility to full functional independence, our observation raises the possibility of the likelihood of increased disability related to joint pains in the elderly. In the context of the projected increase in the growth of elderly populations in developing nations, this observation has serious public health implications. Facilities and resources to cater for the health needs of the elderly are not widely available in most low-income nations. In addition to this, the burden of care for elderly people in Nigeria currently falls on family members. With the rapid rate of urbanization and migration resulting from the social and economic changes being witnessed in many African societies, it will become increasingly difficult for elderly people requiring assistance in daily activities to get family members who are able or willing to provide such care. As shown by an earlier report from the Ibadan Study of Ageing, up to 19% of elderly persons with disability requiring assistance were unable to access such help (Gureje *et al.*, 2006).

Consistent with many previous studies on pain and depression, there was an increased risk of depression amongst individuals with chronic pain. Interestingly, chest pain stood out as being a significant predictor for major depression in this study even after adjusting for the co-occurrence of pain at other sites. Blay *et al.* (Blay *et al.*, 2007), in a survey of community-dwelling elderly persons in Brazil found chest, head and joint pains to be a significant predictor of psychiatric morbidity. This differential risk for depression amongst individuals with chest pain is difficult to explain. However, different locations of pain refer to different bodily systems with different impact on the individual's every day functioning and their interpretation of the significance of the pain. The occurrence of chest pain might also be an indicator of underlying cardiac, respiratory or other physiological dysfunction. For example, studies have shown that up to 20% of coronary heart disease patient develop depression (Rybakowski, 2003). Unfortunately, our study is not designed to explore possible etiological factors for the pains. This association of chest pain to depression and the possible role of underlying physical dysfunction might be worth exploring in future studies

In general, the presence of disability had no major impact on the relationship between chronic pain and depression. However, when considering association of depression to different pain sites, while performance on IADL did not alter the strength of the association between chest pain and depression, that on ADL attenuated the relationship but not to a large extent. It is not surprising that the effect of ADL is more pronounced than that of IADL given that disability in ADLs constitutes a more severe form of disability than that in IADL. Our findings support those of earlier studies that have demonstrated that functional disability is not a mediating factor in the pain-depression relationship (Geerlings *et al.*, 2002; Parmelee *et al.*, 1991).

The results of this study need to be interpreted in the context of its limitations. The assessment of pain, disability and depression were based on self-report. However, previous studies have demonstrated the validity of self-report in the assessment of depression, pain and physical disability (Myers *et al.*, 1993). In addition to this, interference with daily

function was incorporated in our definition for pain, this might have affected our analysis of the role of disability in the association between pain and depression. Another limitation is the cross-sectional nature of the data presented here which does not permit any inference about direction of causality. While we have demonstrated associations between pain and depression, we cannot aver that one causes the other. Nevertheless the results presented here are of value in considering the import of the well-known relationship between pain and depression. At least in the elderly, our findings suggest that in considering this relationship, due attention should always be given to the anatomical sites of pain.

Acknowledgments

Funding for The Ibadan study of Ageing was provided by The Wellcome Trust. The Wellcome Trust was not involved in the design, methods, subject recruitment, data collection, analysis or preparation of this manuscript.

REFERENCES

- Arnow BA, et al. Comorbid depression, chronic pain, and disability in primary care. *Psychosom Med*. 2006; 68:262–268. [PubMed: 16554392]
- Bair MJ, Robinson RL, Katon W, Kroenke K. Depression and pain comorbidity: a literature review. *Arch Intern Med*. 2003; 163:2433–2445. [PubMed: 14609780]
- Bao Y, Sturm R, Croghan TW. A National Study of the Effect of Chronic Pain on the Use of Health Care by Depressed Persons. *Psychiatr Serv*. 2003; 54:693–697. [PubMed: 12719500]
- Bekibele CO, Gureje O. Self-Reported Visual Impairment and Impact on Vision-Related Activities in an Elderly Nigerian Population: Report from the Ibadan Study of Ageing. *Ophthalmic Epidemiology*. 2008; 15:250–256. [PubMed: 18780258]
- Blay SL, Andreoli SB, Dewey ME, Gastal FL. Co-occurrence of chronic physical pain and psychiatric morbidity in a community sample of older people. *International Journal of Geriatric Psychiatry*. 2007; 22:902–908. [PubMed: 17252525]
- Breivik H, Collett B, Ventafridda V, Cohen R, Gallacher D. Survey of chronic pain in Europe: prevalence, impact on daily life, and treatment. *Eur J Pain*. 2006; 10:287–333. [PubMed: 16095934]
- Brochet B, Michel P, Barberger-Gateau P, Dartigues J. Population-based study of pain in elderly people: a descriptive survey. *Age Ageing*. 1998; 27:279–284.
- Chou KL. Reciprocal relationship between pain and depression in older adults: evidence from the English Longitudinal Study of Ageing. *J Affect Disord*. 2007; 102:115–123. [PubMed: 17240455]
- Emptage NP, Sturm R, Robinson RL. Depression and comorbid pain as predictors of disability, employment, insurance status, and health care costs. *Psychiatr Serv*. 2005; 56:468–474. [PubMed: 15812099]
- Ferguson, BD.; Tandon, A.; Gakidou, E., et al. Estimating permanent income using indicator variables. In: Murray, CJL.; Evans, DB., editors. *Health Systems Performance Assessment: Debates, Methods and Emiricism*. World Health Organization; Geneva, Switzerland: 2003.
- Gagliese L, Melzack R. Chronic pain in elderly people. *Pain*. 1997; 70:3–14. [PubMed: 9106804]
- Gallagher RM, Verma S, Mossey J. Chronic pain. Sources of late-life pain and risk factors for disability. *Geriatrics*. 2000; 55:40–44. 47. [PubMed: 10997125]
- Geerlings SW, Twisk JW, Beekman AT, Deeg DJ, van Tilburg W. Longitudinal relationship between pain and depression in older adults: sex, age and physical disability. *Soc Psychiatry Psychiatr Epidemiol*. 2002; 37:23–30. [PubMed: 11926200]
- Gureje O, Kola L, Afolabi E. Epidemiology of major depressive disorder in elderly Nigerians in the Ibadan Study of Ageing: a community-based survey. *The Lancet*. 2007; 370:957–964.
- Gureje O, Ogunniyi A, Kola L, Afolabi E. Functional disability in elderly Nigerians: Results from the Ibadan Study of Aging. *J Am Geriatr Soc*. 2006; 54:1784–1789. [PubMed: 17087709]
- Gureje O, Von Korff M, Simon GE, Gater R. Persistent pain and well-being: a World Health Organization Study in Primary Care. *JAMA*. 1998; 280:147–151. [PubMed: 9669787]

- Heeringa, S., et al. Sample Designs and Sampling Procedures. In: Kessler, RC.; Ustun, TB., editors. The WHO World Mental Health Surveys: Global Perspectives on the Epidemiology of Mental Disorders. Cambridge University Press; New York: 2008.
- Leong IY, Farrell MJ, Helme RD, Gibson SJ. The relationship between medical comorbidity and self-rated pain, mood disturbance, and function in older people with chronic pain. *J Gerontol A Biol Sci Med Sci*. 2007; 62:550–555. [PubMed: 17522361]
- Mantyselka P, Hartikainen S, Louhivuori-Laako K, Sulkava R. Effects of dementia on perceived daily pain in home-dwelling elderly people: a population-based study. *Age Ageing*. 2004; 33:496–499. [PubMed: 15271639]
- Melding PS. Who is losing the language? Persistent pain in home-dwelling elders. *Age Ageing*. 2004; 33:432–434. [PubMed: 15315914]
- Mossey J, Gallagher RM. The longitudinal occurrence and impact of comorbid chronic pain and chronic depression over two years in continuing care retirement community residents. *Pain Med*. 2004; 5:335–348. [PubMed: 15563319]
- Myers AM, Holliday PJ, Harvey KA, Hutchinson KS. Functional performance measures: are they superior to self assessments? *J Gerontol*. 1993; 48:M196–206. [PubMed: 8366262]
- Ohayon MM. Specific characteristics of the pain/depression association in the general population. *J Clin Psychiatry*. 2004; 65:5–9. [PubMed: 15315471]
- Ohayon MM, Schatzberg AF. Using Chronic Pain to Predict Depressive Morbidity in the General Population. *Arch Gen Psychiatry*. 2003; 60:39–47. [PubMed: 12511171]
- Onder G, et al. Association between pain and depression among older adults in Europe: results from the Aged in Home Care (AdHOC) project: a cross-sectional study. *J Clin Psychiatry*. 2005; 66:982–988. [PubMed: 16086612]
- Parmelee PA, Katz IR, Lawton MP. The relation of pain to depression among institutionalized aged. *J Gerontol*. 1991; 46:15–21.
- Reyes-Gibby CC, Aday L, Cleeland C. Impact of pain on self-rated health in the community-dwelling older adults. *Pain*. 2002; 95:75–82. [PubMed: 11790469]
- Rybakowski JK. Depression and cardiovascular disease. *WPA Bulletin on Depression*. 2003; 26:12–14.
- Sternbach RA. Survey of pain in the United States: the Nuprin pain report. *Clin J Pain*. 1986; 2:49–53.
- Thomas E, Mottram S, Peat G, Wilkie R, Croft PR. The effect of age on the onset of pain interference in a general population of older adults: Prospective findings from the North Staffordshire Osteoarthritis Project (NorStOP). *Pain*. 2007; 129:21–27. [PubMed: 17084980]
- Thomas E, Peat G, Harris L, Wilkie R, Croft PR. The prevalence of pain and pain interference in a general population of older adults: cross-sectional findings from the North Staffordshire Osteoarthritis Project (NorStOP). *Pain*. 2004; 110:361–368. [PubMed: 15275787]
- Tsang A, et al. Common chronic pain conditions in developed and developing countries: gender and age differences and comorbidity with depression-anxiety disorders. *J Pain*. 2008; 9:883–891. [PubMed: 18602869]
- Zanocchi M, et al. Chronic pain in a sample of nursing home residents: prevalence, characteristics, influence on quality of life (QoL). *Arch Gerontol Geriatr*. 2008; 47:121–128. [PubMed: 18006088]

Prevalence of chronic pain and association with sex, age and economic status

Table 1

Pain site	Sex (%)			Age (%)					Economic status (%)				
	Total n (%)	Male	Female	P value	65-69 (n=598)	70-74 (n=425)	75-79 (n=250)	80 (n=514)	P value	Low (n=520)	Low average (n=640)	High average (n=430)	High (n=197)
Neck or Back	136(7.6)	5.4	8.0	0.014 [*]	5.1	4.5	7.1	12.8	0.004 [*]	7.8	6.9	5.8	4.7
Stomach or abdomen	34(1.9)	1.3	2.8	0.072	1.3	1.6	3.6	2.0	0.268	3.5	1.0	2.6	0.2
Joints	228(12.8)	10.9	15.0	0.062	9.4	12.3	13.7	19.9	0.012 [*]	10.8	15.7	11.7	10.0
Face or jaw	48(2.7)	1.8	2.8	0.280	2.3	2.2	1.4	3.4	0.470	4.0	2.2	1.5	1.2
Chest	53(3.0)	1.7	3.3	0.083	1.6	1.7	2.5	5.5	0.030 [*]	3.7	1.8	2.9	0.6
Headache	40(2.2)	0.8	3.6	0.003 [*]	1.8	1.5	2.1	3.2	0.496	2.6	2.6	1.4	0.8
Pain at other sites	23(1.3)	0.8	1.3	0.406	0.8	0.4	1.8	2.0	0.141	0.8	1.7	0.8	0.2
Any Chronic pain	315(16.6)	14.1	20.1	0.010 [*]	12.1	15.7	19.9	26.2	0.001 [*]	16.5	18.8	16.6	11.6

^{*} Significant at 0.05 level

Table 2Association of **12-month major depressive disorder with persistent pain (controlling for age and sex)**

Pain site	Prevalence of Depression (%)	OR (95% CI)⁺
Neck or Back	13.3	2.0 (1.1-3.7) *
Stomach or abdomen	19.5	2.8 (0.8-9.5)
Joints	11.7	1.8 (1.0-3.1) *
Face or jaw	12.0	1.7 (0.7-3.9)
Chest	19.7	3.3 (1.5-7.1) *
Headache	14.4	1.8 (0.5-6.3)
Pain at other sites	6.2	0.8 (0.2-3.3)
Any Chronic pain	11.9	1.9 (1.2-3.2) *
No Pain	6.7	0.4 (0.3-0.8) ⁺

OR- Odds ratio, CI- Confidence Interval

* Significant

⁺ Values adjusted for age and sex

Table 3

Association of **12-month major depressive disorder with persistent pain (Controlling for age, sex, other pain conditions)**

Pain site	OR (95% CI)⁺⁺
Neck or Back	1.2 (0.5-3.2)
Stomach or abdomen	2.0 (0.6-6.6)
Joints	1.5 (0.7-3.5)
Face or jaw	1.2 (0.4-3.8)
Chest	2.1 (1.0-4.4) *
Headache	1.1 (0.3-4.5)
Pain at other sites	0.3 (0.1-2.2)

OR- Odds ratio, CI- Confidence Interval

* Significant

⁺⁺ Values adjusted for age, sex and other pain conditions

Table 4Association of 12-month major **depressive disorder with persistent pain at different anatomic sites**

Pain site	OR (95%CI)⁺ (Controlling for ADL Disability)	OR (95% CI)⁺ (Controlling for IADL Disability)
Neck or Back	1.2 (0.5-3.2)	1.2 (0.5-3.3)
Stomach or abdomen	2.0 (0.6-6.9)	1.8 (0.5-6.5)
Joints	1.4 (0.6-3.2)	1.5 (0.6-4.4)
Face or jaw	1.2 (0.4-3.8)	1.2 (0.4-3.8)
Chest	2.0 (0.9-4.4)	2.1 (1.0-4.5) *
Headache	1.0 (0.2-4.1)	1.0 (0.2-4.7)
Pain at other sites	0.3 (0.1-2.1)	0.3 (0.1-2.1)

ADL-Activities of daily living, IADL- Instrumental activities of daily living

OR- Odds ratio, CI- Confidence Interval

* Significant

⁺ Values adjusted for age, sex and other pain conditions