Published in final edited form as:

Tetrahedron Lett. 2011 August 10; 52(32): 4182–4185. doi:10.1016/j.tetlet.2011.06.006.

Preparation of phenalenes and hydronaphthacenes through tandem alkyne Fischer-carbene complex coupling and inter- or intramolecular Diels-Alder reactions

Rajesh Kumar Patti, Shaofeng Duan, Zhipeng Wang, and James W. Herndon*

Department of Chemistry & Biochemistry, New Mexico State University, Las Cruces, NM, 88003, USA

Abstract

The rapid construction of phenalenes through the reaction of 8-alkynyltetralones with Fischer carbene complexes followed by either inter- or an intramolecular Diels-Alder reaction is presented. As a showcase of the synthetic utility of this process, the rapid construction of polycyclic ring systems containing the tetracycline core has been demonstrated through an iterative application of this reaction sequence.

Keywords

carbene complexes; isobenzofurans; cycloaddition; phenalenes; alkynes; naphthacenes

Introduction

The phenalene ring system comprises the core of several natural products, including the pseudopterisins and helioporins, compounds which possess anti-inflammatory¹ and antitumor² activity. Non-natural phenalenes have been evaluated as dopamine receptors.³ Higher annulated aromatic species (*e.g.* pyrenes and benzo analogs) form the core structures several antibiotics⁴ and of various fluorophores useful as biological probes.⁵ Phenalenes are substructures of benzo[a]pyrene, which is one of the most persistent environmental carcinogens and synthetic benzo[a]pyrene derivatives are frequently required to assess mechanisms of carcinogenesis.⁶ The phenalene ring system is most commonly accessed through intramolecular electrophilic aromatic substitution reactions⁷ and occasionally through Diels-Alder processes.⁸ The phenalene ring system is potentially accessible in a single reaction event through a recently-discovered sequence involving tandem carbene complex alkyne coupling⁹ / isobenzofuran formation¹⁰ / and Diels-Alder reaction (Scheme 1).¹¹ Operationally this is a very simple process involving only thermolysis and avoids a multi-step series of reactions required to perform similar transformations without chromium.¹² Formation of phenalenes requires the use of alkynylbenzenes fused to cyclic ketones (**A**), which are likely ideal substrates for isobenzofuran formation since the carbonyl group is locked in the orientation required for the key C-O bond formation step. In this manuscript the synthesis of 8-alkynyl-2-tetralone derivatives and their subsequent reaction with Fischer carbene complexes is presented. An iterative version of this process leads to

compounds possessing the hydronaphthacene skeleton, a ring system that is potentially useful for in the development of organic semiconductors.¹³

Results and Discussion

The preparation of the *o*-alkynylbenzocycloalkanones (**1**, **3**, **5**) used in this study from commercially-available starting materials is depicted in Scheme 2. Alkynyltetralone **1** was prepared from 1-amino-5,6,7,8-tetrahydronaphthalene via the known transformation to 8-bromo-1-tetralone¹⁴ followed by Sonogashira coupling. This method, though lengthy, is superior to the alternative preparation of 8-bromo-1-tetralone from α -tetralol,¹⁵ since the ortho-metallation process would never go to completion and separation of tetralol and bromotetralol was quite difficult. The five-membered ring analog **3** was prepared from 2-bromobenzoyl chloride using the Nazarov cyclization as a key step.¹⁴ The bis(alkyne) derivative **5** was synthesized from 1,3-dibromobenzene through regioselective lithiation,¹⁶ followed by formylation and double Sonogashira coupling.

Initial studies focused on the systems depicted in Scheme 3, involving the three-component coupling of alkynyltetralone **1**, methylcarbene complex **7**, and either dimethyl acetylenedicarboxylate (DMAD) (**9**) or dimethyl fumarate. A complex reaction mixture resulted when all three components were present at the onset of the reaction, presumably due to multiple components present in solution that are reactive to carbene complexes.¹⁷ If a sequential addition involving (1) reaction of the carbene complex and alkyne at 100 °C for 1 h and (2) addition of DMAD was employed, a surprisingly excellent yield of phenalene-containing product **11** was observed after silica gel purification. Initially, an efficient process was not anticipated under these conditions since isobenzofurans bearing α -hydrogens are unstable with respect to conversion to alkylidenephthalans,¹⁸ and success thus requires this critical intermediate to survive intact for more than an hour at 100 °C. Naphthalene **11** can in theory be obtained using the *N,N*-dimethylhydrazone derivative of ketone **1**,¹⁹ however this process was considerably less efficient. A problem with this alternate strategy is that hydrazone formation from **1** would never go to completion, and the reaction was initiated using a 2:3 mixture of the ketone and hydrazone as starting material.

Ketone **1** was also tested as a substrates for the net [5+5]-cycloaddition process employing the butenylcarbene complex **12a** (see Scheme 4). In this case the reaction was highly efficient leading to the tetracycle **15**. The anticipated product is simple dehydration product **14**,¹¹ which undergoes an unanticipated oxidation under the reaction conditions. Interestingly, the product containing the phenanthrene ring was obtained when the reaction was conducted in anhydrous dioxane, however high yields of alcohol **16** were obtained when the reaction was conducted in dioxane/water mixtures.²⁰ This latter process was completely diastereoselective. The stereochemistry of compound **16** was assigned based on the known exo preference for intramolecular six-membered ring-forming reactions involving isobenzofurans,²¹ which would afford the depicted stereoisomer after ring opening.

Unfortunately, reactions involving the related five-membered ring systems were unsuccessful (Scheme 5). Coupling of alkynylindanone **3** and carbene complex **12a** afforded a complex reaction mixture where the pattern for a monosubstituted alkene is a predominant feature of the crude ¹H NMR spectrum. This observation suggests failure in the formation of the isobenzofuran intermediate,²⁰ which is likely due to ring strain.²² An examination of the isodesmic reactions in Scheme 5 (based on DFT calculations employing the B3LYP method and 6-31G(d) basis sets) reveals that formation of the five-membered fused ring system **20** is unfavorable relative to the six-membered fused ring-fused system **18**. The energy for the isodesmic reaction is 26 kcal/mol greater in the five-membered ring case. A simple MM2 calculation revealed the angle strain in the five-membered fused ring system to be 34.1 kcal/

mol while the strain in the six-membered ring case was considerable less at only 15.5 kcal/mol.

This novel [5+5]-cycloaddition reaction was also examined iteratively as a synthetic route to the hydronaphthacene ring system. The initial product (**21**, Scheme 6) of net [5+5]-cycloaddition employing bis(alkyne) derivative **5** and butenylcarbene complex **12a** is an incipient alkynyltetralone system, and thus the iterative [5+5]-cycloaddition can provide rapid access to this medicinally-important ring system. The crude product from cycloaddition of bis(alkyne) **5** and carbene complex **12a** was subjected to oxidation with PCC, affording alkyne-dione **22**, followed by an additional [5+5]-cycloaddition sequence. This reaction sequence resulted in a 3:2 mixture of diastereomers (out of four possible) favoring the symmetrical isomer **25a**. The stereoisomers arise through exo selective intramolecular Diels-Alder reaction, which proceeds with relative asymmetric induction from the only stereocenter present in **22** to afford either of intermediates **23** or **24**. The stereochemistry of the major isomer (**25a**) was assigned based on the appearance of only 13 signals in the ¹³C NMR spectrum, compared with 21 signals in the unsymmetrical minor isomer (**26a**). This chromium(0-6-0)-based synthesis of the tetracyclines was also demonstrated for a system employing two different carbene complexes. Reaction of alkynone **22** with methylbutenylcarbene complex **12b** afforded a 5:1 mixture of diastereomers. The structure of the minor isomer was determined by X-ray crystallography and was determined to be stereoisomer **25b** (see Supporting Information). The crystal consisted of two independent enantiomeric molecules where one of molecule of the pair shows a disorder at the chiral centers and the other member of the pair is well-resolved. The reactions of alkyne-dione **22** with the γ,δ -unsaturated carbene complexes **12a** and **12b** proceeded with the opposite stereoselectivity in the [5+5]-cycloaddition event. The carbene complex featuring a 1,1-disubstituted alkene group likely affords an alkene-isobenzofuran that undergoes a slower and more thermodynamically selective Diels-Alder reaction due to the more electron rich nature of the dienophile.

In summary, tandem isobenzofuran generation / Diels-Alder processes that employ 8-alkynyltetralone systems are quite efficient. In these systems one can employ ketones as substrates in sequential addition processes due to the enhanced lifetime of the isobenzofuran intermediate, despite the presence of hydrogens poised for rearrangement to alkylidenephthalans. Use of dialkylbenzaldehydes allows for a rapid gain in molecular complexity through an iterative process leading to selective formation of two out of the possible four diastereomers in the resulting complex hydronaphthacene ring systems in three reaction events.

Experimental²³

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by the SCORE program of NIH (SC1GM083693). The Bruker X8 X-ray diffractometer was purchased via an NSF CRIF:MU award to The University of New Mexico, CHE-0443580. We thank Dr. Eileen Duesler for X-ray data collection. We thank Prof. William Maio for helpful comments.

References

1. Potts BCM, Faulkner DJ, Jacobs RS. J. Nat. Prod. 1992; 55:1701–1717. [PubMed: 1294693]
2. Tanaka J, Ogawa N, Liang J, Higa T, Garcia Gravalos D. Tetrahedron. 1993; 49:811–822.

3. Tang AH, Franklin SR, Code RA, Althaus JS, VonVoigtlander PF, Darlington WH, Szmuszkovicz J. *Drug Dev. Res.* 1990; 21:53–62.
4. Inokoshi J, Shiomi K, Masuma R, Tanaka H, Yamada H, Omura S. *J. Antibiotics.* 1999; 52:1095–1100. [PubMed: 10695672] b Huang X, He J, Niu X, Menzel KD, Dahse HM, Grabley S, Fiedler HP, Sattler I, Hertweck C. *Angew. Chem. Int. Ed.* 2008; 47:3995–3998.
5. a Vullev VI, Jiang H, Jones G II. *Topics in Fluorescence Spectroscopy.* 2005; 10:211–239. b Pereira G, Abreu AS, Castanheira EMS, Coutinho PJG, Ferreira PMT, Queiroz MJRP. *Eur. J. Org. Chem.* 2009:3906–3916.
6. a Xu D, Penning TM, Blair IA, Harvey RG. *J. Org. Chem.* 2009; 74:597–604. [PubMed: 19132942] b Ran C, Xu D, Penning TM, Blair IA, Harvey RG. *Tetrahedron Lett.* 2008; 49:4531–4533. c Ran C, Dai Q, Ruan Q, Penning TM, Blair IA, Harvey RG. *J. Org. Chem.* 2008; 73:992–1003. [PubMed: 18181642] d Harvey RG. *Curr. Org. Chem.* 2004; 8:303–323.
7. a Kim SH, Kim YM, Lee HS, Kim JN. *Tetrahedron Lett.* 2010; 51:1592–1595. b Nanclares J, Gil J, Rojano B, Saez J, Schneider B. Otávaro, F. *Tetrahedron Lett.* 2008; 49:3844–3847. c Barluenga J, Trincado M, Rubio E, González JM. *Angew. Chem. Int. Ed.* 2006; 45:3140–3143. d Benoit-Marquie F, Csaky AG, Esteban G, Martínez ME, Plumer J. *Tetrahedron Lett.* 2000; 41:3355–3358. e Wichmann J, Adam G, Rover S, Hennig M, Scalone M, Cesura AM, Dautzenberg FM, Jenck F. *Eur. J. Med. Chem.* 2000; 35:839–851. [PubMed: 11006485] f Lazerwith SE, Johnson TW, Corey EJ. *Org. Lett.* 2000; 2:2389–2392. [PubMed: 10930291]
8. a Enders D, Httl MRM, Raabe G, Bats JW. *Adv. Synth. Catal.* 2008; 350:267–279. b Brummond KM, You L. *Tetrahedron.* 2005; 61:6180–6185. c Nicolaou KC, Sugita K, Baran PS, Zhong YL. *J. Am. Chem. Soc.* 2002; 124:2221–2232. [PubMed: 11878976] d Piers E, Friesen RW, Kao P, Rettig SJ, Trotter J. *Can. J. Chem.* 1993; 74:1463–1483.
9. For detailed mechanistically-based experimental studies, see: A Bos ME, Wulff WD, Miller RA, Chamberlin S, Brandvold TA. *J. Am. Chem. Soc.* 1991; 113:9293–9319. b. For the seminal theoretical paper on carbene complex-alkyne coupling, see: Hoffman P, Hammerle M, Unfried G. *New J. Chem.* 1991; 15:769–789.
10. Herndon JW, Wang H. *J. Org. Chem.* 1998; 63:4564–4565.
11. For the most recent example of this reaction, see Camacho-Davila A, Gamage LSR, Wang Z, Herndon JW. *Tetrahedron.* 2010; 66:4954–4960. [PubMed: 20711418]
12. Duan S, Cress K, Waynant K, Ramos-Miranda E, Herndon JW. *Tetrahedron.* 2007; 63:2959–2965. [PubMed: 18382602]
13. For synthetic studies, see: a Seri T, Qu H, Zhou L, Kanno K, Takahashi T. *Chem. Asian J.* 2008; 3:388–392. [PubMed: 18219642] . b. For applications in materials science, see: Sundar VC, Zaumseil J, Podzorov V, Menard E, Willet RL, Someya T, Gershenson ME, Rogers JA. *Science.* 2004; 303:1644–1647. [PubMed: 15016993]
14. Nguyen P, Corpuz E, Heidelbaugh TM, Chow K, Garst ME. *J. Org. Chem.* 2003; 68:10195–10198. [PubMed: 14682726]
15. Lebedev AY, Khartulyari AS, Voskoboinikov AZ. *J. Org. Chem.* 2005; 70:596–602. [PubMed: 15651807]
16. Lulinski S, Serwatowski J. *J. Org. Chem.* 2003; 68:5384–5387. [PubMed: 12816504]
17. Jiang D, Herndon JW. *Org. Lett.* 2000; 2:1267–1269. [PubMed: 10810724]
18. a Friedrichsen W. *Struct. Chem.* 1999; 10:47–52. b Chacko E, Sardella DJ, Bornstein J. *Tetrahedron Lett.* 1976:2507–2510.
19. Duan S, Sinha-Mahapatra DK, Herndon JW. *Org. Lett.* 2008; 10:1541–1544. [PubMed: 18351767]
20. Li R, Zhang L, Camacho-Davila A, Herndon JW. *Tetrahedron Lett.* 2005; 46:5117–5120.
21. For a review of the chemistry of isobenzofurans, see: Friedrichsen W. *Adv. Heterocycl. Chem.* 1999; 73:1–96.
22. Early investigators of 6,5,5-tricyclic ring systems declared this ring system to be “unsynthesizable” until the first successful synthesis. Rapoport H, Pasky JZ. *J. Am. Chem. Soc.* 1956; 78:3788–3792.
23. For a General Experimental, see the Supporting Information.

Scheme 1.

Scheme 2.

Scheme 3.

Scheme 4.

Scheme 5.

Scheme 6.