The Alberta Health Technology Assessment (HTA) Ambassador Program: The Development of a Contextually Relevant, Multidisciplinary Clinical Practice Guideline for Non-specific Low Back Pain: A Review

Greg Cutforth, Aaron Peter, Paul Taenzer

ABSTRACT

Purpose: To describe the development of a contextually relevant multidisciplinary clinical practice guideline (CPG) for non-specific low back pain (LBP) and to discuss its value to the management of LBP and the practice of physiotherapy.

Method: To mitigate an identified knowledge gap for Alberta primary-care practitioners in the management of non-specific LBP, a collaborative process was developed to engage multidisciplinary health care providers in designing a primary-care CPG for non-specific LBP. A comprehensive review of published LBP guidelines identified the seven highest-quality CPGs; these were used to inform a multidisciplinary guideline development group (GDG) as they developed the CPG.

Results: The GDG constructed a CPG for non-specific LBP along with point-of-care decision-support and patient-education tools.

Conclusions: The Ambassador Program on Low Back Pain worked with front-line clinicians from across Alberta to review the best available evidence in developing a CPG responsive to the Alberta context. This CPG is intervention specific and provides a wide range of primary-care practitioners with the best available evidence to inform their clinical decisions in managing non-specific LBP.

Key Words: evidence-based practice, low back pain, multidisciplinary communication, practice guidelines, rehabilitation

Cutforth G, Peter A, Taenzer P. The Alberta Health Technology Assessment (HTA) Ambassador Program: the development of a contextually relevant, multidisciplinary clinical practice guideline for non-specific low back pain: a review. *Physiother Can.* 2011; 63(3):278–286; doi:10.3138/ptc.2009-39P

RÉSUMÉ

Objectif : Décrire l'élaboration d'une ligne directrice multidisciplinaire de pratique clinique pertinente au contexte pour la lombalgie non spécifique et discuter de sa valeur pour la gestion de la lombalgie et la pratique de la physiothérapie.

Méthode : Dans le but d'atténuer les lacunes dans les connaissances des praticiens en soins primaires de l'Alberta dans la gestion de la lombalgie non spécifique, un mode de collaboration a été mis sur pied pour que les fournisseurs de soins de santé de multiples disciplines participent à l'élaboration d'une ligne directrice clinique en soins primaires pour la lombalgie non spécifique. Une analyse approfondie des lignes directrices pour la lombalgie a permis de répertorier sept lignes directrices de la plus grande qualité; elles ont été utilisées pour inspirer le groupe multidisciplinaire dans la création d'une ligne directrice.

Résultats : Le groupe multidisciplinaire a élaboré une ligne directrice pour la lombalgie non spécifique de même qu'une aide à la décision sur les lieux de soins et des outils éducatifs pour le patient.

Conclusions : L'équipe de l'*Ambassador Program on Low Back Pain* a travaillé avec des cliniciens de première ligne de partout en Alberta pour analyser les meilleures preuves scientifiques disponibles pour l'élaboration d'une ligne directrice clinique convenant au contexte de l'Alberta. Cette ligne directrice est une intervention précise et offre à un large éventail de praticiens en soins primaires les meilleures preuves scientifiques disponibles pour les aider dans leurs décisions cliniques pour la gestion de la lombalgie non spécifique.

Mots clés : communication multidisciplinaire, lignes directrices de pratique, lombalgie, pratique fondée sur l'expérience clinique, réadaptation

Christa Harstall, MHSA; Carmen Moga, MD, MSc; Alayne Lobay-Ness, BScPT; Rachelle Slavik, BASW; Douglas P. Gross, PhD; Todd Wolansky, BScPT; Cathie Scott, PhD; Lynne Dunsmuir, BScPT, MSc; and Audrey Lowe, PT, Msc PT, BscPT, BPE, all contributed to the editing process for this manuscript.

The authors have no conflicts of interest to declare.

Greg Cutforth, PT, BScPT: Regional Manager, Allied Health Services (Jasper to Barrhead), Alberta Health Services, Hinton, Alberta.

Aaron Peter, BScPT, CAFCI: Physical Therapist, Medicine Hat Therapy Services and Healthy Living Program, Alberta Health Services, Medicine Hat, Alberta.

Paul Taenzer, PhD, RPsych: Specialist/Clinical Psychologist, Calgary Pain Program, Alberta Health Services, Calgary, Alberta.

Address correspondence to *Greg Cutforth*, 1280a Switzer Drive, Hinton, AB T7V 1T5 Canada; Tel.: 780-817-4121; Fax: 780-865-3727;

E-mail: greg.cutforth@albertahealthservices.ca; *Aaron A. Peter*, 715 Sprague Way SE, Medicine Hat, AB T1B 4P9 Canada; Tel.: 403-548-0084;

E-mail: aaann.peter@gmail.com; *Paul Taenzer*, 100, 2310 – 2nd Street SW, Calgary, AB T2S 3C4 Canada; E-mail: paul.taenzer@albertahealthservices.ca.

DOI:10.3138/ptc.2009-39P

INTRODUCTION

The Alberta Health Technology Assessment (HTA) Ambassador Program was established in 2004 and was based on the original knowledge-translation model developed in Sweden.^{1–4} The initial goal of the Ambassador Program was to develop and test a case-based interactive multidisciplinary workshop as a model of knowledge translation. In following up this initial work, the programme team next identified knowledge gaps in the assessment and management of low back pain (LBP) among primary-care practitioners through a province-wide survey. Based on this identified knowledge gap, the most recent objective of the Ambassador Program was to apply the knowledge-translation model to the development of a contextually relevant clinical practice guideline (CPG) for LBP for use by primary-care practitioners in Alberta (see Figure 1). This article describes the development process and discusses the relevance of the CPG to physiotherapy (PT) and the management of LBP.

Low Back Pain

Estimates of prevalence and incidence of LBP vary widely, depending on how LBP is defined and on the populations studied. The available data suggests that between 49% and 90% of people in developed countries will experience at least one episode of LBP during their lifetime.^{5–9} As back pain is often more recurrent than acute and self-limiting, its management can be complex and costly for the individual, the health care system, and society. In the United States, back problems are associated with nearly one-quarter of all lost work days;⁸ back pain is now the second leading cause of work absenteeism, and approximately 2% of American adults are either temporarily or chronically disabled by LBP.⁸ Canadian data^{10,11} indicates that the prevalence of and clinical course for LBP are similar in Canada as in other developed countries. The significant economic burden to society consists largely of the indirect costs associated with employee absenteeism and disablement.^{6,12}

In Alberta and Saskatchewan, close to 40% of patients with back pain seek help from a health care provider.⁹ Primary-care physicians undertake the initial evaluation in 65% of LBP cases and are often the sole care provider for these patients.^{10,13} It has been estimated that 2% of patients with acute LBP initially seek PT services;¹⁴ the

same study found that those initially seeking PT services for acute LBP typically present with higher baseline disability scores.¹⁴ A recent telephone survey by MPT students at the University of Alberta asked participants who had experienced LBP about their care-seeking behaviour. Only 38% of participants had sought medical attention for their last episode of LBP;¹⁵ PT was reportedly used by 3% of those surveyed, while 31% sought care from a medical doctor and 28% sought care from a chiropractor.¹⁵ In 2008 the province of Alberta began working to amalgamate the nine existing health regions into one provincial health region, now called Alberta Health Services. This amalgamation officially came into effect on June 1, 2009. Data from one of the former rural Alberta health regions indicate that LBP was the primary complaint for approximately 15% of patients seen in publicly funded outpatient PT departments.¹⁶ Data provided by a former urban health region in Alberta suggest that LBP was the chief complaint for between 20% and 31% of the population receiving publicly funded musculoskeletal PT services.¹⁷

The prevalence of LBP, the burden it places on people and their families, and the varied strategies used to manage this condition all underscore the importance of informing primary-care practitioners about the best evidence in the management of LBP.¹⁸

Knowledge Gap—Survey and Literature Results

Given the human and economic impact of LBP, it is crucial that health care professionals be well trained and knowledgeable about current treatment strategies. In following up on its initial work on knowledge translation, the Ambassador Program Advisory Committee recommended that a detailed knowledge assessment be conducted. The primary purpose of the assessment was to determine the level of knowledge of specific health care providers (physicians, nurses, physiotherapists, and occupational therapists) as it related to the management (i.e., diagnosis and treatment) of LBP and headache. A secondary objective was to elicit these practitioners' views on barriers to effective management. The survey sought to identify respondents' knowledge of specific aspects of care based on published clinical practice guidelines (CPGs) related to LBP and headache. A total of 105 respondents, including 12 physiotherapists, completed the survey.¹⁹

The average score for all professions on the LBP section of the assessment was 55%. Both medical doctors (MDs) and physiotherapists attained an average knowledge score of 64% on the LBP section; other non-MDs, including nurses, pharmacists, and occupational therapists, achieved an average score of 52%. Respondents typically knew more about diagnosis and non-pharmacological care than about surgical and pharmacological treatments.¹⁹

Figure 1 The Ambassador Program

Although this knowledge assessment had a number of important methodological limitations that may have implications for its generalizability, it did suggest that Alberta health care providers have significant knowledge gaps related to the diagnosis and treatment of LBP. Respondents indicated that lack of access to CPGs is one of the major barriers to effective management of chronic pain. The survey results are consistent with a review of the available published literature on primary-care practitioners' knowledge gaps undertaken by the Ambassador Program research team.²⁰

METHODS

The Guideline-Development Process

The development of the CPG was directed through the organizational structure of the Ambassador Program (see Figure 2).

The Guideline Development Group (GDG) was given the task of developing the CPG for non-specific LBP. A call for volunteers was sent to previous participants in the original Ambassador workshop to invite their further involvement in the GDG; at the same time, the programme undertook targeted recruitment of professionals and specialists with a special interest in LBP from across Alberta. Ultimately, this group was composed of family physicians, specialist physicians, physiotherapists, occupational therapists, nurses, pharmacists, researchers, psychologists, an administrator, and a knowledge-translation specialist. The group included both urban and rural clinical practitioners and achieved broad geographical representation of the province. One result of the selec-

tion process, however, was a lack of participation from the chiropractic profession. This was mitigated by a face-to-face follow-up with the Alberta College and Association of Chiropractors, during which the content of the CPG and resources were reviewed and this group of professionals was given an opportunity to provide feedback.

The GDG was supported by the programme's Steering Committee, which consisted of the HTA research team, an experienced guideline-development consultant, a communications and knowledge-translation expert, the project co-chairs, and a project assistant. The Steering Committee guided the overall project and provided research, administrative, and communications support.

An Advisory Committee directed the Steering Committee on issues related to the design of the overall project and the dissemination of the completed guideline, and its members served as project champions in their respective organizations. The members of the Advisory Committee were representatives of stakeholder agencies, including the project host, the Institute of Health Economics, the provincial Ministry of Health, several of the former Alberta health regions, a patient-advocacy group, professional colleges, and the provincial physician guideline organization, Toward Optimized Practice (TOP).

Within this organizational structure, a multistage process was initiated that began with a search for and appraisal of existing guidelines for the management of non-specific LBP. This work was undertaken by the research team from the Steering Committee and generated a review and synopsis of the recommendations from the highest-quality existing guidelines ("seed guidelines"), which the GDG used to develop the new tool.

The research team conducted an extensive search for LBP guidelines published between January 1996 and February 2008. The search identified 58 potentially relevant guidelines published in countries with developed economies. Based on the inclusion criteria, nine guidelines were selected. A more detailed description of the inclusion and exclusion process and criteria can be found in the supporting background document.²¹ Two of the nine guidelines focused on prevention, five on acute LBP, and two on chronic LBP. These nine guidelines were appraised for quality using a modification of the Appraisal of Guidelines for Research and Evaluation (AGREE) instrument, an internationally developed generic tool that is validated, transparent, and widely accepted, with satisfactory reliability for most domains.²²⁻²⁴ The research team modified the AGREE instrument to reduce the ambiguity and subjectivity associated with item scoring and to enable the differentiation of good- from poor-quality guidelines.²⁵ A sub-committee of clinical experts excluded two lower-quality guidelines that focused only on acute LBP; seven seed guidelines were ultimately chosen to be used in the adaptation process.

Figure 2 Structure of the Ambassador Program

Tables of convergent and divergent recommendations and supporting evidence were produced to inform the GDG.

The GDG reviewed the evidence compiled by the research team from the seven seed guidelines and considered each recommendation in the context of the Alberta health care environment. Using videoconferencing technology, the working group evolved a process to align the seed guideline recommendations with contextual and clinical input from the GDG. Decisions were made via consensus. Dialogue between and input from GDG members was key to ensuring that the agreed-upon recommendations were meaningful, clinically relevant, and feasible within Alberta's working health care environment. To maintain a level of scientific rigour, the committee retained the original wording for each specific recommendation from the seed guidelines whenever possible. Where a change in wording was necessary to ensure that a recommendation reflected and/or was clinically relevant in the Alberta context, the group used the Guideline Implementability Appraisal (GLIA) tool to structure the modified recommendations.²⁶ The majority of the GDG's group work was accomplished through a series of 10 half-day videoconferences and a single full-day meeting. Varying amounts of work were completed by GDG members between sessions, typically in preparation for each group meeting.

To present the evidence and make recommendations, the GDG used a framework consistent with that generally observed in the seven seed CPGs. The recommendations were listed in the categories *prevention*, *acute/subacute*, and *chronic*. While the *prevention* category listed evidence on interventions only, the two remaining headings listed recommendations first on diagnostics and assessment, then on interventions.

Items for which the evidence was conflicting, limited, or equivocal were further investigated by sub-groups composed of GDG members with relevant expertise. In an effort to ensure a consistent approach within these sub-groups, all sub-groups also included the chair of the GDG. The sub-groups were supported by an HTA researcher, who assembled the research evidence for consideration prior to discussion. Once each sub-group had reached consensus, its recommendation was brought back to the GDG for further discussion and ratification. One example of this process that involved PT was the recommendation on physical exercise and therapeutic exercise for chronic LBP. The equivocal nature of the evidence in this area made the wording of the recommendation challenging; in light of the limited evidence, there was a strong push from the physiotherapists on the GDG to ensure that the importance of exercise be upheld. In the end, by means of group discussion and follow-up through electronic correspondence, the sub-group agreed to wording that was accurate but not dismissive of the intervention. Ultimately, physiotherapists were involved in 16 of these sub-groups during the

development process, including those that addressed recommendations on spinal manipulation, physical activity, advice/reassurance, and return to work.

Given the range of LBP interventions available and the number of health care disciplines working with LBP, it would be naïve, if not impossible, to assume that one CPG could specify detailed best practice for all disciplines. The intent in developing an overarching guideline for multiple disciplines was to achieve consistent messaging that would be relevant across the disciplines. With this understanding, the GDG made every effort to present the evidence as it applied to the intervention in question rather than focusing on the profession or the individual practitioner providing the intervention or service. For certain restricted activities, such as spinal manipulation, references were provided to inform the user which professions are able to practise the technique safely. This approach respects the autonomy of the individual practitioner and is in keeping with the spirit of health professions legislation in Alberta. Developing guidelines with this principle in mind avoids pitting professions against one another, which can only work against the spirit of integrated services for patients.

RESULTS

As the evidence was compiled and organized, it became evident that if the guideline were to be both relevant and useable for a wide range of users, a number of resources would need to be developed.

The Steering Committee decided to focus on the following evidence-based resources:

1. Guideline summary (see Figure 3)²⁷
2. Full clinical practice guideline²⁸
3. Background document that outlines the guideline-development process in more detail²¹

Additional resources were also compiled, including information sheets for patients and a document developed by the New Zealand acute LBP Guideline Group²⁹ that describes how to assess and manage psychosocial yellow flags.

All of the resources developed and used to support the clinicians in the management of LBP, including the CPG and the background document, are posted on the TOP Web site.³⁰ TOP is a joint venture of the College of Physicians and Surgeons of Alberta, the Alberta Medical Association, and Alberta Health and Wellness. This provincial initiative develops, maintains, and distributes Alberta CPGs for physicians and the teams they work with. TOP will retain the intellectual property rights of the Ambassador Program guideline and, in partnership with the IHE, will be responsible for ongoing maintenance and updating of the guideline and associated resources. This update process is scheduled to occur every 2 years, the first of which began in January of 2011.

A Summary of the Guideline for the Evidence-Informed Primary Care Management of Low Back Pain

This evidence-informed guideline is for non-specific, non-malignant low back pain in adults only

For complete guideline refer to the TOP Website: www.topalbertadoctors.org

①

Low Back Pain

Key Messages

- Do a full clinical assessment; rule out red flags
- In the absence of red flags, reassure the patient there is no reason to suspect a serious cause
- Reinforce that pain typically resolves in a few weeks without intervention
- Encourage patient to keep active
- Consider evidence-based management as per the guideline
- Recommend physical activity and/or exercise to prevent recurrence
- If pain continues beyond 6 weeks, reassess and consider additional treatment and referrals
- The goal of chronic pain management is improved quality of life
- Encourage and support pain self-management
- Monitor patient for relative benefit versus side effects

Contraindications

Evidence indicates these actions are ineffective or harmful

- Lab tests and diagnostic imaging in the absence of red flags
- Prolonged bed rest
- Tradition
- Oral steroids
- Epidural steroid injections in the absence of radicular pain
- TENS for acute pain
- Back schools for acute pain
- Massage therapy for acute pain

Medication Table¹

Pain Type	Medication	Dosage range
Acute and sub-acute low back pain or flare-up of chronic low back/spinal pain	1st line Acetaminophen	Up to 1000 mg QID (max of 4000 mg/day)
	2nd line NSAIDs	Up to 800 mg TID (max of 800 mg QID)
	Diclofenac	Up to 50 mg TID
	Add: Cyclobenzaprine for prominent muscle spasm	10 to 30 mg/day; Greatest benefit seen within one week; therapy up to 2 weeks may be justified
Chronic low back/spinal pain	If taking controlled release opioids: add a short-acting opioid or increase controlled release opioid by 20 to 25%	See opioids below
	1st and 2nd lines	See acute pain, above
	3rd line Weak Opioids	Codeine 30 to 60 mg every 3 to 4 hours
	Controlled release codeine	50 to 200 mg Q8h, may also be given Q12h
TRICYCLICS	Tricyclics (TCAs)	Amiripryline 10 to 100 mg HS
	Nortriptyline	10 to 100 mg HS
	fewer adverse effects	
	4th line Tramadol (not currently covered by Alberta Blue Cross)	Slow titration up to max of 400 mg/day; short acting form is only available in combination with acetaminophen. Monitor for total combined daily acetaminophen dose.
5th line Strong Opioids (controlled release)	Morphine sulfate	15 to 100 mg BID
	Hydromorphone HCl	3 to 24 mg BID
	Oxycodone HCl	10 to 40 mg BID-TID
	Fentanyl patch	25 to 50 µg Q3days

¹ Adapted from the Calgary Regional Pain Program, September 19, 2006

- This guideline was written to provide healthcare providers and patients with guidance about appropriate prevention, assessment and intervention strategies
- It was developed by a multidisciplinary team of Alberta clinicians and researchers
- This guideline is for adults 18 years of age or older with low back pain and is not applicable to pregnant women
- It is recognized that not all recommended treatment options are available in all communities
- For further details on the recommendations, see the guideline and background document

②

Figure 3 The LBP Guideline Summary (intellectual property of Toward Optimized Practice; reproduced under Creative Commons Attribution-NonCommercial-ShareAlike 2.5 Canada licence (CC BY-NC-SA 2.5))

The guideline summary²⁷ (see Figure 3) is a two-page document printed on both sides of one sheet that is intended to act as a quick reference for clinicians. It includes only information and recommendations for which the evidence was unequivocal and either clearly supported or did not support a particular intervention. The information is presented both as a visual algorithm and in narrative point form. A list of red and yellow flags is also included, along with a medication table.

The full CPG²⁸ provides a much more detailed review of the evidence and some description of the process used to arrive at the specific recommendations. A more detailed review of red and yellow flags for clinicians is also included. The recommendations are presented in table format and classified according to the available evidence of what a practitioner should either "Do" (recommended) or "Do Not Do" (not recommended). Where the evidence reviewed was equivocal, the recommendations are listed as "Do Not Know" (uncertain).

The background document²¹ is a comprehensive review of the evidence, the process used to select the seed guidelines, and the development of the recommendations themselves. The target audience for this detailed resource is academic researchers and guideline developers or appraisers.

Once the general content of the CPG had been finalized by the GDG, the information was presented in draft form to the provincial medical colleges and associations. Through a 1-day face-to-face meeting with these groups and a meeting with the Alberta College and Association of Chiropractors, feedback and commentary were compiled that helped to further refine the CPG and provided further suggestions and opportunities for dissemination.

A dissemination plan³¹ was developed with input from the Advisory Committee and from communication experts and consultants. It was agreed that the guiding principle for the dissemination plan would be to link where possible with existing initiatives and resources, such as TOP. The supporting background document²¹ provides a more detailed description of this work and the planning involved.

DISCUSSION

It is a great challenge to develop a multidisciplinary CPG that is structured to integrate knowledge into practice for multiple health professions. If this is to be done effectively, information must be presented in a way that is clinically relevant and can be readily implemented by multiple clinicians. The multidisciplinary membership of

the GDG, its broad geographical representation, and the use of videoconferencing technology to communicate were meaningful steps toward ensuring that the CPG would be more easily adopted by Alberta primary-care practitioners.

The level of detail that practitioners require in a CPG will be influenced by the experience and backgrounds of individual practitioners as well as by the environments in which they work. The issue is further complicated when the target audience is multidisciplinary in nature. To address some of these challenges, the Ambassador Program chose to focus its work on a first layer of evidence for LBP. The information included in the Ambassador Program CPG is intended to be applicable, at least in some way, to all relevant health care providers. Other care pathways and CPGs exist that are more service or discipline specific, and these perhaps constitute a second layer of information that can more specifically direct the management of LBP for smaller target audiences of health care practitioners. The Clinic on Low Back Pain in Interdisciplinary Practice (CLIP) guideline,³² developed in the province of Quebec, is a good example of a potential second layer of evidence. The CLIP guideline can better direct the work of rehabilitation practitioners by providing a relevant review of evidence and outcome measures more specific to the rehabilitation disciplines, but it is less applicable for other health care disciplines, such as primary-care physicians.³³ Another example of more discipline-specific layers of evidence might be the National Guidelines for Physical Therapy and Manual Therapy in patients with LBP.^{34–35} These two CPGs, published in the Netherlands, potentially represent a second and third layer of evidence respectively. The first guideline is focused more directly on the management of LBP through a broader understanding of PT practice;³⁴ the second narrows its focus one step further and looks specifically at manual therapy in the management of LBP.³⁵

If information already exists to direct the management of LBP more specifically, what is the purpose or value of this first layer of evidence? We believe that its value lies in its potential to align the efforts of the many health care disciplines and services that are working to help people living with LBP. Providing consistent and evidence-informed knowledge to health care professionals creates an opportunity to enhance the integration of health care services. This type of integration can lead to more consistent messaging for people living with LBP and for the general public. The current inconsistencies in messaging between and within various health professions are a challenge for both patients and professionals trying to manage LBP, and this situation is further complicated by the wide array of conflicting sources of information available to the public. Proper integration of this first layer of evidence has the potential to minimize confusion for patients seeking credible information. Broad, credible, and consistent messaging

can also serve to support those professionals who are already disseminating advice and providing treatment in alignment with the current best evidence by increasing the credibility of the services and advice these professionals are providing to patients and other health care providers. Used in conjunction with tools more specific to the direct patient care provided by physiotherapists for non-specific LBP, this multidisciplinary CPG can further support the work of physiotherapists at a level consistent with the practice of other evidence-informed health care professionals. It is therefore conceivable that both broad and targeted uses of this CPG have the potential to contribute improving quality of care and reducing the economic burden of LBP.

Effective integration of guidelines into practice is a challenge. The Ambassador Program team has worked closely with its partners to develop dissemination strategies that go beyond traditional thinking in this area. The Ambassador concept is an excellent example of how the health care community can rethink strategies to integrate knowledge and evidence more effectively into front-line care. There is a consensus within the Ambassador Program team that the dissemination strategy will need to establish long-term objectives that will work toward the gradual integration of the guideline. Traditional strategies for innovation and practice change have typically adopted more of a “big bang” approach, which can seem threatening to the prospective audience and may lead to a loss of impact once the initial effect wears off. The Ambassador Program has identified opportunities for the CPG to link to potential and existing initiatives; in this way the programme can “market” its work as a knowledge-transfer tool that will enhance service, not redefine it.

As noted above, the Ambassador Program has partnered with TOP. The goal of this partnership is to ensure that the LBP guideline is disseminated and made available to Alberta physicians and other health care professionals; this relationship also provides a structure that will ensure that the guideline is regularly updated to reflect the most recent evidence. Given TOP’s status among Alberta physicians, it is anticipated that the Ambassador/TOP partnership will enhance credibility for the guideline as a source of evidence for clinicians.

Integrating guidelines into practice can be challenging for front-line practitioners. All health care providers work within the context of their experience, their education, the service environment, and the situations that their patients present to them. Introducing CPGs and evidence-based practice can be daunting. In the absence of strong supporting evidence, clinicians may be left with a sense that what they are doing is incorrect or of no value; this is especially true when practice and experience do not align well with the available evidence. This quandary of clinical decision making for health care providers introduces some interesting points of discussion.

First, a lack of evidence of effectiveness does not necessarily mean that there is evidence of a lack of effectiveness; in the case of LBP, there are significant gaps in the knowledge base around the effectiveness of many of our traditional interventions. This lack of evidence does not indicate that a particular intervention is ineffective; rather, it means only that there is a lack of evidence to support its efficacy. Clinicians can apply that understanding to their clinical decision making.

The second point relates to the application of evidence in clinical decision making. Rycroft-Malone et al.³⁶ suggested a four-pillared model of evidence to support clinical decision making. The four pillars are research, clinical experience, patient experience, and local context. Research evidence is thus only one component of clinical decision making; in the absence of strong evidence, practitioners must rely more heavily on clinical experience, patient experience, and local context. Likewise, in the absence of clinical experience—as is the case for less experienced practitioners—effective clinical decisions rely more heavily on local context, patient experience, and the available evidence. In developing the CPG for LBP, the Ambassador Program has tried to provide practitioners from all disciplines with an updated and accessible research pillar to support their clinical decision making.

Finally, individual practitioners have an obligation to make balanced clinical decisions that help patients work toward their individual goals for care. A recent study by Rutten et al. found that increased adherence by physiotherapists to the Dutch physical therapy and manual therapy guidelines lead to greater improvements in physical functioning and lower utilization of care.³⁷ Despite this emerging evidence supporting the use of CPG's in the management of LBP, a recent paper by Harman et al. documented the prevalence of interventions poorly supported by evidence in the management of subacute LBP by physiotherapists working with injured workers.³⁸ Using CPGs or the best available evidence in clinical decision making is not compulsory, but the onus is on health care providers to integrate evidence into their clinical decisions whenever possible. Guidelines such as those developed by the Ambassador Program can be a useful tool to facilitate this process.

LIMITATIONS

Several important limitations should be considered when interpreting the results of the work described in this article. First, as stated previously, a number of methodological limitations existed that limit the generalizability of the data obtained in the knowledge-gap analysis, most notably the very low response rate of the original survey. These limitations were mitigated to some degree by the consistency of the findings with a

review of the published literature on knowledge gaps and primary-care practitioners.

A second limitation of the study was the current lack of research investigating the general impact of the CPG on clinical practice. First, it is unclear whether the model used by the Ambassador Program to develop the CPG for LBP improved the integration of research evidence into the clinical practice of primary-care practitioners. Second, more research is needed to evaluate what effect enhanced integration of the CPG into clinical practice will have on the lives of people living with LBP.

CONCLUSION

The Ambassador Program identified a knowledge gap among Alberta primary-care practitioners in the management of LBP. Given the prevalence of LBP and the burden it imposes on individuals, families, and society, the need for an evidence-based tool to inform clinical decision making by primary-care practitioners was evident. The Ambassador Program team worked with front-line clinicians from across the province to review the best available evidence to develop a CPG responsive to the Alberta context. This CPG is intervention specific and provides a wide range of primary-care practitioners with the best available evidence to inform their clinical decisions in the management of LBP.

KEY MESSAGES

What Is Already Known on This Topic

The burden of LBP on individuals, families, and society is significant. Health care providers and users have a common interest in minimizing the functional impact of LBP by aligning practice with the current best evidence for the management of LBP. Currently there are multiple barriers to the adoption of evidence into clinical practice.

What This Study Adds

This paper describes the strategies and processes used by the Health Technology Assessment Ambassador Program in the province of Alberta to identify a knowledge gap in the management of LBP and to develop tools to inform front-line health care providers with the current best evidence in the management of LBP. The tools described were developed with consideration of the health care environment in Alberta. The evidence-based recommendations within the tools are applicable across multiple health care disciplines. This paper describes and provides access to the resources developed. It also describes the relevance and applicability of these resources to the practice of physiotherapy.

REFERENCES

- Rashiq S, Barton P, Harstall C, Schopflocher D, Taenzer P. The Alberta Ambassador Program: delivering Health Technology Assessment results to rural practitioners. *BMC Med Educ* [Internet]. 2006 Mar 31 [cited 2011 Jan 10];6(21):6 p.]. Available from: <http://www.biomedcentral.com/1472-6920/6/21>
- Scott NA, Moga C, Barton P, Rashiq S, Schopflocher D, Harstall C, et al. Creating clinically relevant knowledge from systematic reviews: the challenges of knowledge translation. *J Eval Clin Pract*. 2007;13:681–8. doi:10.1111/j.1365-2753.2007.00830.x
- Taenzer P, Schopflocher D, Rashiq S, Harstall C. The Alberta chronic pain ambassador program. In: Rashiq S, Schopflocher D, Taenzer P, Jonsson E, editors. *Chronic pain: a health policy perspective*. Weinheim (Germany): Wiley Blackwell; 2008. p. 255–72.
- The Alberta Ambassador Program [Internet]. Edmonton: Institute of Health Economics; c2010 [cited 2009 Jun 26]. Available from: <http://www.ihe.ca/research/ambassador-program/>
- Woolf AD, Pfleger B. Burden of major musculoskeletal conditions. *Bull World Health Organ*. 2003;81:646–56.
- van Tulder M, Koes B, Bombardier C. Low back pain. *Best Pract Res Clin Rh*. 2002;16:761–75.
- Margarido MS, Kowalski SC, Natour J, Ferraz MB. Acute low back pain: diagnostic and therapeutic practices reported by Brazilian rheumatologists. *Spine*. 2005;30:567–71.
- Devereaux MW. Low back pain. *Primary Care: Clin Off Pract*. 2004;31(1):33–51.
- Nyiendo J, Haas M, Goldberg B, Sexton G. Pain, disability, and satisfaction outcomes and predictors of outcomes: a practice-based study of chronic low back pain patients attending primary care and chiropractic physicians. *J Manip Physiol Ther*. 2001;24:433–9.
- Moulin DE, Clark AJ, Speechley M, Morley-Forster PK. Chronic pain in Canada—prevalence, treatment, impact and the role of opioid analgesia. *Pain Res Manage*. 2002;7(4):179–84.
- Schopflocher D. Chronic pain in Alberta: a portrait from the 1996 National Population Health Survey and the 2001 Canadian Community Health Survey [Internet]. Edmonton: Health Surveillance, Alberta Health and Wellness; 2003 [cited 2011 Jan 10]. Available from: <http://www.health.alberta.ca/documents/Chronic-Pain-Survey-2003.pdf>
- Coyte P, Asche C, Croxford R, Chan B. The economic cost of musculoskeletal disorders in Canada. *Arthritis Care Res*. 1998;11: 315–25. doi:10.1002/art.1790110503
- Bishop PB, Wing PC. Compliance with clinical practice guidelines in family physicians managing worker's compensation board patients with acute lower back pain. *Spine J*. 2003;3:442–50. doi:10.1016/S1529-9430(03)00152-9
- Mielenz TJ, Carey TS, Dyrek DA, Harris BA, Garrett JM, Darter JD. Physical therapy utilization by patients with acute low back pain. *Phys Ther*. 1997;77:1040–51.
- Brunke KM, Huls CR, LaRoy JN, Reagan KS, Wilton JA, Gross DP. Descriptive study of care-seeking behaviour of people with back pain [Internet]. Paper presented at the Reaching Forward Physical Therapy Conference; 2008 Oct 24–26 [cited 2011 Jan 18]; Edmonton, AB. Available from: http://journals.lww.com/spinejournalabstracts/Citation/2010/00001/Care_Seeking_Behaviour_and_Back_Pain_Are_Beliefs.22.aspx.
- van der Nest, Dalique (Rehabilitation Coordinator, Aspen Regional Health). E-mail correspondence and phone conversation with: Greg Cutforth (Physical Therapy Clinical Lead, Aspen Regional Health). 2009 Jun 12.
- Pratt, Doug (Program Coordinator, Community Physiotherapy Program, Calgary Health Region). E-mail correspondence and phone conversation with: Greg Cutforth (Physical Therapy Clinical Lead, Aspen Regional Health). 2009 Feb 20.
- Rossignol M, Abenhaim L, Bonvalot Y, Gobeille D, Shrier I. Should the gap be filled between guidelines and actual practice for management of low back pain in primary care? the Quebec experience. *Spine*. 1996;21:2893–8.
- Lopatka H. Primary care practitioner knowledge assessment for low back pain and headache management: summary report. Edmonton: Institute of Health Economics; 2007 Sep [cited 2009 Jun 8]. Available from: <http://www.ihe.ca/documents/Knowledge-Assessment-Survey.pdf>
- Scott NA, Moga C, Harstall C. Managing low back pain in the primary care setting: the know–do gap. *Pain Res Manage*. 2010;15:392–400.
- Institute of Health Economics. Ambassador Program guideline for the evidence-informed primary care management of low back pain: background document [Internet]. Edmonton (AB): The Institute; 2009 [cited 2011 Jan 10]. Available from <http://www.ihe.ca/research/ambassador-program/-low-back-pain/low-back-pain-guideline/guideline-documents/>
- Appraisal of Guidelines for Research and Evaluation (AGREE) instrument [Internet]. London: AGREE Collaboration; c2001 [updated 2001 Sep; cited 2009 Jun 26]; [22p]. Available from: <http://www.agreecollaboration.org/pdf/agreecollaborationfinal.pdf>
- The AGREE Collaboration. Development and validation of an international appraisal instrument for assessing the quality of clinical practice guidelines: the AGREE project. *Qual Saf Health Care*. 2003;12(1):18–23. doi:10.1136/qhc.12.1.18
- MacDermid JC, Brooks D, Solway S, Switzer-McIntyre S, Brosseau L, Graham ID. Reliability and validity of the AGREE instrument used by physical therapists in assessment of clinical practice guidelines. *BMC Health Serv Res*. 2005;5(18). doi:10.1186/1472-6963-5-18
- Scott NA, Moga C, Harstall C. Making the AGREE tool more user-friendly: the feasibility of a user guide based on Boolean operators. *J Eval Clin Prac*. 2009;15(6):1061–73. doi:10.1111/j.1365-2753.2009.01265.x
- Shiffman RN, Dixon J, Brandt C, Essaihi A, Hsiao A, Michel G, et al. The GuideLine Implementability Appraisal (GLIA): development of an instrument to identify obstacles to guideline implementation. *BMC Med Inform Decis Mak*. 2005;5(23). doi:10.1186/1472-6947-5-23
- A summary of the guideline for the evidence-informed primary care management of low back pain [Internet]. Edmonton: Toward Optimized Practice Alberta; 2009 Mar [cited 2010 May 31]. Available from: http://www.topalbertadoctors.org/cpgs/back_pain.html.
- Guideline for the evidence-informed primary care management of low back pain [Internet]. Edmonton: Toward Optimized Practice Alberta; 2009 Mar [cited 2010 May 31]. Available from: http://www.topalbertadoctors.org/cpgs/back_pain.html.
- New Zealand acute low back pain guide, incorporating the guide to assessing psychosocial yellow flags in acute low back pain [Internet]. Wellington: Accident Compensation Corporation; c2004 [cited 2010 May 31] Available from: http://www.nzgg.org.nz/guidelines/dsp_guideline_popup.cfm?guidelineID=72&guideline_popup_mode=index
- Management of low back pain [Internet]. Edmonton: Toward Optimized Practice Alberta; 2009 Mar [cited 2010 May 31]. Available from: http://www.topalbertadoctors.org/informed_practice/cpgs/low_back_pain.html.
- Angus D. Draft of the dissemination and implementation of evidence-informed primary care management of low back pain guideline [Internet]. Edmonton: Alberta HTA Ambassador Program; 2009 Mar [cited 2010 May 31]. Available from: <http://www.ihe.ca/documents/DisseminationPlan1.pdf>
- Rossignol M, Arseneault B, Dionne C, Poitras S, Tousignant M, Truchon M, et al. Clinic on low back pain in interdisciplinary practice (CLIP) guidelines [Internet]. Montreal: Direction de santé publi-

- que, Agence de la santé et des services Sociaux de Montréal; 2007 [cited 2009 Jun 8]. Available from: <http://www.santepub-mtl.qc.ca/Publication/pdftravail/CLIPenglish.pdf>.
33. Côté AM, Durand MJ, Tousignant M, Poitras S. Physiotherapists and use of low back pain guidelines: a qualitative study of the barriers and facilitators. *J Occup Rehabil.* 2009;19(1):94–105. doi:10.1007/s10926-009-9167-2
 34. Bekkering GE, Hendriks HJM, Koes BW, Oostendorp RAB, Ostelo RWJG, Thomassen JMC, et al. KNGF-guidelines for physical therapy in patients with low back pain [Internet]. Royal Dutch Association of Physical Therapy; 2003 [cited 2009 Jun 8]. Available from: https://www.kngfrichtlijnen.nl/downloads/Low_Back_Pain_Guideline.pdf
 35. Heijmans WFGJ, Hendriks HJM, van der Esch M, Pool-Goudzwaard A, Scholten-Peeters GGM, van Tulder MW, et al. KNGF-guideline: manual therapy on low back pain [Internet]. Royal Dutch Association of Physical Therapy; 2003 [cited 2009 Jun 8]. Available from: https://www.kngfrichtlijnen.nl/downloads/Manual_Therapy_in_case_of_Low_Back_Pain_Guideline.pdf
 36. Rycroft-Malone J, Seers K, Titcher A, Harvey G, Kitson A, McCormack B. What counts as evidence in evidence-based practice? *J Adv Nurs.* 2004;47(1):81–90. doi:10.1111/j.1365-2648.2004.03068.x
 37. Rutten GM, Degen S, Hendriks EJ, Braspenning JC, Harting J, Oostendorp RA. Adherence to clinical practice guidelines for low back pain in physical therapy: do patients benefit? *Phys Ther.* 2010;90(8):1111–22.
 38. Harman K, Fenety A, Hoens A, Crouse J, Padfield B. Physiotherapy and low back pain in the injured worker: an examination of current practice during the subacute phase of healing. *Physiother Can.* 2009;61:88–106. doi:10.3138/physio.61.2.88