Oxidation and Assimilation of Carbohydrates by *Micrococcus sodonensis*¹

JEROME J. PERRY AND JAMES B. EVANS

Department of Microbiology, North Carolina State University, Raleigh, North Carolina

Received for publication 7 August 1965

ABSTRACT

PERRY, JEROME J. (North Carolina State University, Raleigh), AND JAMES B. EVANS. Oxidation and assimilation of carbohydrates by *Micrococcus sodonensis*. J. Bacteriol. 91:33-38. 1966.—*Micrococcus sodonensis* is a biotin-requiring strict aerobe that cannot utilize carbohydrates as sole sources of carbon and energy. However, addition of mannose, glucose, sucrose, or maltose to a medium on which the organism can grow resulted in an increase in total growth. *M. sodonensis* oxidized these sugars without induction, thus indicating the presence of constitutive enzymes for their transport, activation, and metabolism. Under appropriate nonproliferating cell conditions, glucose was readily incorporated into essential constituents of the cell. When glucose-1-C¹⁴ and glucose-6-C¹⁴ were oxidized by nonproliferating cells, the label was found in both the protein and nucleic acid fractions of the cell. The respiratory quotients of cells oxidizing glucose in saline and in phosphate buffer indicated assimilation of sugar carbon in buffer and virtually no assimilation in saline. The ability of *M. sodonensis* to completely oxidize glucose and to grow on intermediates of glucose oxidation but not on glucose suggests that glucose may suppress or repress some reaction(s) necessary for growth, and that growth substrates either derepress or circumvent this block.

Micrococcus sodonensis is an obligate aerobe that can grow on a biotin-containing synthetic medium with lactate, succinate, pyruvate, or acetate as the carbon and energy source. In the original description of the organism, the inability to produce acid or gas from carbohydrates was noted (1). Subsequent research (14) demonstrated that *M. sodonensis* does not grow when glucose or other carbohydrates are the sole source of carbon and energy.

Recently, we observed, however, that the addition of glucose to a medium containing lactate as substrate increased the rate and amount of growth of *M. sodonensis* (Perry and Evans, Bacteriol. Proc., p. 77, 1965). This increase in rate of growth and total cell yield also occurred with other substrates. Experiments with nonproliferating cells indicated that glucose is readily oxidized by this organism without prior induction. This observation has been extended,

and the oxidative and assimilatory capacity of the organism has been investigated.

MATERIALS AND METHODS

M. sodonensis ATCC 11880 was employed for this study. The basal medium was the same as the L-salts medium of Leadbetter and Foster (9) with NH₄Cl replacing NaNO₃. The medium was supplemented with 10 µg of biotin per liter. Substrates (lactate or succinate) were added at 0.5%, and glycine was routinely added at 0.1% to enhance growth.

Manometric techniques were those described in Umbreit et al. (18).

Cells were fractionated into the cold trichloroacetic acid-soluble fraction, hot trichloroacetic acid-soluble fraction, and protein, according to the procedure of Roberts et al. (16).

Glucose was quantified by the Nelson (13) modification of the Somogyi test.

Radioactive chemicals were obtained from the Volk Radiochemical Corp., Chicago, Ill.

The uptake of radioactivity by cells was measured by filtering suitable dilutions onto membrane filters (Schleicher and Schuell Co., Keene, N.H.), followed by thorough washing of the collected cells with water. The filter and cells were placed in a counting vial and dried in a hot-air oven. Scintillation fluid was

¹ Contribution from Microbiology, North Carolina Agricultural Experiment Station, Raleigh. Published with the approval of the Director of Research as Paper No. 2035 of the Journal Series.

added, and radioactivity was measured in a Packard Tri-carb liquid scintillation spectrometer.

RESULTS

The number of compounds capable of supporting the growth of *M. sodonensis* is severely limited. Lactate, pyruvate, and acetate, as well as some of the intermediates in the tricarboxylic acid cycle, are the sole growth substrates for this organism (14, 15). Glucose and other carbohydrates added as the sole carbon source will not support growth, whether added to media before autoclaving, autoclaved separately, or filter-sterilized. However, inoculation of cells into media (lactate or succinate) supplemented with glucose results in a significant and reproducible increase in rate of growth and dry weight of cells produced. The effect of added sugars on total growth is shown in the first column of Table 1. Growth was markedly increased by incorporation of maltose, glucose, sucrose, or mannose in the medium. At conventional levels, the enhancement of growth by carbohydrates was proportional to the amount added.

Nonproliferating cells harvested from the succinate medium were tested for the ability to

oxidize various carbohydrates (Table 1). The sugars that are effective growth stimulants (sucrose, glucose, mannose, and maltose) were readily oxidized by nonproliferating cells. The addition of 50 μ g of chloramphenicol per ml to cells in the presence of the oxidizable sugars had no effect on the initial rate of oxidation. The rapid initial oxidation rate and lack of interference by chloramphenicol indicate that enzyme induction is not required for the oxidation of sugars.

As shown in Fig. 1, fructose is oxidized very slowly by this organism, at a rate that is proportional to the substrate concentration. Glucose and sucrose are oxidized at a much higher rate than is fructose, and the rate is essentially independent of substrate concentration. It also can be seen from Fig. 1 that sucrose oxidation continues at the same rate after the comparable level of glucose is used up (curves B and D) and is equivalent to twice the comparable level of glucose (curves B and C). Thus, it appears that the fructose moiety of sucrose is oxidized at a rate and to an extent that is equivalent to glu-

TABLE 1. Effect of various carbohydrates on the growth of *Micrococcus sodonensis* in a succinate medium, and the ability of resting cells to oxidize these compounds

Carbohydrate added	Total growth*	Resting-cell† oxidation (O ₂ uptake)
	mg/ml	μ liters
Maltose.....	1.20	477
Glucose.....	1.04	320
Sorbitol.....	0.56	100
Fructose.....	0.56	48
Xylose.....	0.40	47
Lactose.....	0.52	34
Sucrose.....	1.02	709
Mannose.....	1.02	219
Arabinose.....	0.52	8
Ribose.....	0.52	8
No addition....	0.45	—

* Culture conditions: 10-ml amounts of media in 50-ml Erlenmeyer flasks were placed on a rotary shaker at 30 C. The substrate (succinate) and carbohydrate were added at 0.5% each. Growth was estimated after 72 hr by measuring the optical density of a culture at 620 m μ , and is expressed as dry weight.

† Respirometer temperature, 30.5 C. Each vessel contained 10 μ moles of substrate. Succinate-grown cells were diluted in 0.067 M phosphate buffer at pH 7.0 to 12 mg per vessel (dry weight); 0.2 ml of 20% KOH in center well; total volume, 2.4 ml. All data corrected for endogenous metabolism. Time, 300 min.

FIG. 1. Oxidation of various carbohydrates by succinate-grown *Micrococcus sodonensis*. A 20-mg amount (dry weight) of cells was added to a total volume of 2.4 ml; 0.2 ml of 20% KOH in the center well. Temperature, 30.5 C. All data corrected for endogenous.

cose, and that the poor oxidation of the mono-saccharide fructose results from deficiencies in the transport mechanism for fructose. Other experiments, not shown here, indicate that maltose is oxidized at a rate similar to sucrose, and that mannose and sorbitol are oxidized at rates that are proportional to their concentration.

During respirometer studies, an apparent anomaly was encountered in that the diluent for resting-cell experiments had a striking effect on the total oxidation of glucose. Dilution of cells in 0.067 M phosphate buffer (pH 7.0) resulted in a total O_2 uptake of less than 50% of that observed when cells were diluted in physiological saline (Fig. 2). Monitoring the pH during oxidation confirmed that there was no change. Quantification of glucose remaining in the supernatant fraction indicated that under both conditions over 90% disappeared within 360 min.

Carbon dioxide production was measured during oxidation of glucose in buffer and in saline. Table 2 gives data on the relative level of CO_2 - O_2 during oxidation of glucose and sucrose. The respiratory quotient (RQ = micromoles of CO_2 /micromoles of O_2) for oxidation is 1.0 in saline and about 0.86 in buffer. Since glucose disappears at equal rates under either condition, more glucose is assimilated with phosphate buffer as diluent than with saline. The addition of 10^{-4} M 2,4-dinitrophenol (DNP) alleviated the reduced total oxidation in buffer. There was also some increase with saline as diluent, but the total O_2 uptake in the presence of DNP with either buffer or saline was equivalent.

FIG. 2. Oxidation of glucose with physiological saline and with 0.067 M phosphate buffer (pH 7.0) as diluents. Temperature, 30.5 C. Each vessel contained 20 mg (dry weight) of succinate-grown cells. All data corrected for endogenous.

TABLE 2. Respiratory quotient of succinate-grown cells during oxidation of glucose and sucrose*

Diluent	Substrate	Time	O_2 uptake	CO_2	RQ, CO_2/O_2
		min	μ liters	μ liters	
Saline	Glucose	120	350	352	1.0
	Glucose	300	833	840	1.0
	Glucose	300	465	465	1.0
	Sucrose	180	603	608	1.0
Buffer	Glucose	120	264	234	0.89
	Glucose	300	336	286	0.85
	Glucose	300	301	265	0.88

* The amount of substrate in each vessel was 10 μ moles.

TABLE 3. Fate of uniformly labeled glucose added to resting cells of *Micrococcus sodonensis**

Fraction	Total counts per min
CO_2	3,480,384
Supernatant fluid.....	520,920
Cold trichloroacetic acid (transient intermediates).....	629,463
Alcohol (lipid protein).....	784,588
Ether alcohol.....	140,743
Hot trichloroacetic acid (nucleic acid).....	42,810
Protein.....	92,046

* Succinate-grown cells (20 mg, dry weight) diluted in saline; 10 μ moles of carrier glucose added. Time of experiment, 135 min. Reaction was stopped with 0.2 ml of 2 N H_2SO_4 , and cell fractions were analyzed.

To measure the fate of glucose, 10 μ c of the uniformly labeled compound were added to resting cells, with carrier glucose, in a respirometer vessel. After oxidation for 135 min, the reaction was stopped by addition of 0.2 ml of 2 N H_2SO_4 , and the contents of the vessel were removed for analysis. The amount of oxygen taken up by the cells was two-thirds of theoretical for complete oxidation of glucose (Table 3); 62% of the glucose was oxidized to CO_2 , 9% remained in the supernatant fraction, and 29% was assimilated into cell material. Most of the assimilated material was found in the cold trichloroacetic acid (transient intermediates) fraction and the alcohol-soluble (lipid) fraction of the cells. No great loss of glycolytic intermediates from *M. sodonensis* was observed on addition of N_2SO_4 , in contrast to the findings of Clifton (2). Comparable counts of activity were found in cells whether vessels were cooled immediately or when acid was added.

TABLE 4. *Distribution of radioactivity from respirometer studies of Micrococcus sodonensis oxidizing variously labeled glucose**

Glucose	Count/min			O ₂ uptake μmoles
	CO ₂	Super-natant fluid	Cells	
Saline				
Glucose-1-C ¹⁴	475,335	56,293	9,600	48
Glucose-6-C ¹⁴	376,528	111,321	28,710	45
Buffer				
Glucose-1-C ¹⁴	168,139	106,032	96,608	18
Glucose-6-C ¹⁴	86,028	207,205	147,502	17

* Time, 240 min. KOH in center well; 600,000 count/min of appropriately labeled glucose added to saline, and 400,000 count/min added to buffer with 10 μmoles of carrier glucose per 20 mg (dry weight) of succinate-grown cells.

Table 4 shows the conditions and results of studies on the fate of glucose-1-C¹⁴ and glucose-6-C¹⁴ during oxidation in buffer and saline. The reaction was stopped after 240 min by addition of acid from the side arm, and the distribution of radioactivity was ascertained. During oxidation in saline, 88% of the label in glucose-1-C¹⁴ and 73% of the label in glucose-6-C¹⁴ appeared as CO₂. When buffer was the diluent, 45% of the glucose-1-C¹⁴ and 20% of the glucose-6-C¹⁴ label appeared as CO₂. In saline, 2 and 6%, respectively, of the added label were fixed into the cells, whereas more significant levels (26 and 33%) were incorporated during oxidation in phosphate buffer.

The cells from the previous experiment were fractionated according to the procedure of Roberts et al. (16) to obtain the glycolytic intermediates and the protein and nucleic acid fractions (Table 5). Cells suspended in saline assimilated glucose into the glycolytic (cold trichloroacetic acid) fraction, whereas cells in buffer assimilated more into the protein and nucleic acid portions of the cell.

Since glucose was assimilated into the protein and nucleic acid fractions of the cell in phosphate buffer, an experiment was run to determine whether the organism might reproduce when inoculated at high levels into a complete medium with glucose as carbon source. To the mineral salts plus biotin medium with 0.5% glucose as substrate, 10⁷ cells were added. The cells were counted in a Petroff-Hausser counter at the start of the experiment, and after 18 hr the material was removed from the shaker and counted again. Virtually the same number of cells were present as at the start of the experiment. No reproduction

TABLE 5. *Distribution of radioactivity in cells after oxidation of labeled glucose**

Glucose	Per cent of total count		
	Cold tri-chloroacetic acid and lipid	Hot tri-chloroacetic acid	Protein
Saline			
Glucose-1-C ¹⁴	62	18	9
Glucose-6-C ¹⁴	64	11	10
Buffer			
Glucose-1-C ¹⁴	35	37	26
Glucose-6-C ¹⁴	36	31	28

* Cells from experiment presented in Table 3.

TABLE 6. *Growth of Micrococcus sodonensis on lowered levels of lactate in the presence of constant levels of glucose**

Substrate (mg/10 ml)		Cells (mg/10 ml)
Lactate	Glucose	
20	0	6.3
20	50	10.8
10	50	8.5
5	50	7.3
1	50	2.9
0.1	50	1.4
0	50	0

* Growth for 72 hr without added glycine.

occurred with glucose as the sole carbon and energy source.

To ascertain whether the organism would ferment glucose, uniformly labeled glucose (1 μc), with 10 μmoles of carrier glucose, was placed in the side arm of a respirometer vessel, with cells in the main compartment. The flask was flushed with nitrogen. At 2 hr after addition of the substrate, the reaction was stopped by the addition of acid, and the center well paper was checked for radioactive CO₂. No significant count was present. The supernatant fraction was also quantified for glucose, and it was found that none was utilized. Insignificant amounts of radioactive glucose were taken up by the cells.

Oxidative and fermentative products of glucose often have a marked effect on the synthesis of enzymes in bacteria. Enzymes concerned with oxidation through the tricarboxylic acid cycle are repressed by the presence of glucose. An experiment was designed to test whether glucose present at disproportionately high levels would suppress growth on low levels of lactate (Table 6). Glucose was present at a constant level and lactate at decreasing levels. After 72 hr, the flasks

were removed from the shaker and dry weights were determined. Since lactate, which is obligatory for growth, disappeared from the medium at the lower levels, it does not appear that glucose exerts its effect on the oxidation of lactate. In a parallel experiment without added glucose, the 1 and 0.1 mg per 10 ml levels of lactate were insufficient for growth to visible turbidity.

DISCUSSION

The inability of an organism to grow on a particular substrate can be attributed to a number of factors. Major conditions that presuppose an organism's capacity to grow on a specific compound are permeability and the heritable enzyme composition. The enzyme make-up must enable the cell to gain requisite energy from the breakdown of the substrate and to convert the products to essential cell constituents.

The ability of *M. sodonensis* to oxidize glucose and some disaccharides containing glucose to CO_2 and water indicates that these substrates enter the cells and are extensively metabolized. Furthermore, abundant growth of this bacterium occurs on the intermediates of total glucose oxidation, e.g., pyruvate, succinate, and α -ketoglutarate. The capacity for glucose oxidation is not affected by the growth medium but is constitutive. Since glucose is readily oxidized by nonproliferating whole cells, the existence of a permeability barrier is essentially eliminated.

Glucose can serve as a source of major cell constituents, as shown by the incorporation of labeled glucose into cells, particularly in the presence of phosphate buffer. Products of glucose oxidation are distributed (Table 4) in the lipid, nucleic acid, and protein fractions. The level in each fraction is similar to the pattern in *Pseudomonas aeruginosa* grown on glucose (4). Phosphate buffer has a marked effect on the oxidation of glucose by nonproliferating cells. R Q values in the presence and absence of buffer indicate that assimilation of products of glucose metabolism is much greater when a source of inorganic phosphate is available. This is confirmed (Table 4) by use of glucose labeled in the 1 and 6 position. There is a 10-fold increase in label uptake from glucose-1- C^{14} and a 5-fold increase from glucose-6- C^{14} in the presence of phosphate buffer as contrasted with saline. Analogous results have been obtained with resting cells of *Escherichia coli* (8), where increased glucose assimilation occurs in phosphate buffer over that obtained in distilled water.

The capacity of cells to oxidize the fructose moiety of sucrose at the same rapid rate as glucose when it enters the cell as a constituent of a

disaccharide and at a lower concentration-dependent rate when added as the monosaccharide has also been observed in *Pseudomonas saccharophila* (3). These results suggest that a permease exists for sucrose but that fructose enters by diffusion, or that the affinity of fructose-oxidizing enzymes for substrate is affected by the mode of entrance of the compound into the cell. The concentration dependence would indicate that the former explanation is more plausible.

Sugars that are readily oxidized by *M. sodonensis* stimulate the growth of the organism, and those not metabolized have little effect on growth. Using *Clostridium tetani*, reported to be non-saccharolytic, Martinez and Rittenberg (11) demonstrated that sugars added to the growth medium increased the total growth, although they were utilized only after an initial lag. During this lag, significant cell growth occurred. This is in contrast to *M. sodonensis* where the initial growth rate is increased and lag time is decreased by the addition of glucose.

Glucose repression of enzyme synthesis is an established phenomenon in microorganisms (5, 10, 12). Strasters and Winkler (17) found that growth of *Staphylococcus aureus* on glucose enhanced the glycolytic enzymes and suppressed the tricarboxylic acid cycle enzymes. This repression of the tricarboxylic acid cycle has been attributed to the accumulation of lactic acid in *S. aureus* by Gershanovitch et al. (6). Wittenberg and Haaf (19) found that the lactate-degrading system in *Butyribacterium rettgeri* is repressed by the presence of glucose. The addition of high levels of glucose to cells growing on diminishing levels of lactate did not interfere with the growth of *M. sodonensis*. When glucose was present, in the external environment, at 500 times the concentration of lactate, there was no evident suppression of growth. The lack of cell division on a medium containing glucose as sole carbon source suggests that the effect of glucose may be an absolute repression of some reaction(s) necessary for growth.

There was no evidence for mutation of this organism to glucose utilization, as was observed by Klein and Doudoroff (7) in *Pseudomonas putrefaciens*. Repeated experiments where high levels (10^7 to 10^8 cells per milliliter) of cells were inoculated into media with glucose as the sole carbon source consistently failed to show growth.

ACKNOWLEDGMENTS

This investigation was supported by Public Health Service research grant AI-03565 from the National Institute of Allergy and Infectious Diseases.

The technical assistance of Edith J. Dobbins is gratefully acknowledged.

LITERATURE CITED

1. AARONSON, S. 1955. Biotin assay with a coccus, *Micrococcus sodonensis*, nov. sp. *J. Bacteriol.* **69**:67-69.
2. CLIFTON, C. E. 1963. Oxidative assimilation by *Bacillus megaterium*. *J. Bacteriol.* **85**:1365-1370.
3. DOUDOROFF, M. 1940. The oxidative assimilation of sugars and related substances by *Pseudomonas saccharophila* with a contribution to the problem of the direct respiration of di- and polysaccharides. *Enzymologia* **9**:59-72.
4. DUNCAN, M. G., AND J. J. R. CAMPBELL. 1962. Oxidative assimilation of glucose by *Pseudomonas aeruginosa*. *J. Bacteriol.* **84**:784-792.
5. EPPS, H. M. R., AND E. F. GALE. 1942. The influence of the presence of glucose during growth on the enzymic activities of *Escherichia coli*: comparison of the effect with that produced by fermentation acids. *Biochem. J.* **36**:619-623.
6. GERSHANOVITCH, V. N., N. A. PALKINA, AND G. I. BURD. 1963. The role of lactate in regulation of the enzymatic system participating in the acetate oxidation of *Staphylococcus aureus*. *Biochem. Biophys. Res. Commun.* **13**:12-19.
7. KLEIN, H. P., AND M. DOUDOROFF. 1950. The mutation of *Pseudomonas putrefaciens* to glucose utilization and its enzymatic basis. *J. Bacteriol.* **59**:739-750.
8. KOPPER, P. H. 1954. The assimilation of glucose by resting cells of *Escherichia coli*. *J. Bacteriol.* **67**:507-510.
9. LEADBETTER, E. R., AND J. W. FOSTER. 1958. Studies on some methane-utilizing bacteria. *Arch. Mikrobiol.* **30**:91-118.
10. MCQUILLAN, A. M., AND H. O. HALVORSON. 1962. Physiological changes occurring in yeast undergoing glucose repression. *J. Bacteriol.* **84**:31-36.
11. MARTINEZ, R. J., AND S. C. RITTENBERG. 1959. Glucose dissimilation by *Clostridium tetani*. *J. Bacteriol.* **77**:156-163.
12. NEIDHARDT, F. C., AND B. MAGASANIK. 1956. The effect of glucose on the induced biosynthesis of bacterial enzymes in the presence and absence of inducing agents. *Biochim. Biophys. Acta* **21**:324-334.
13. NELSON, N. A. 1944. A photometric adaptation of the Somogyi method for the determination of glucose. *J. Biol. Chem.* **153**:375-380.
14. PERRY, J. J., AND J. B. EVANS. 1960. Oxidative metabolism of lactate and acetate by *Micrococcus sodonensis*. *J. Bacteriol.* **79**:113-118.
15. PERRY, J. J., AND J. B. EVANS. 1961. Role of potassium in the oxidative metabolism of *Micrococcus sodonensis*. *J. Bacteriol.* **82**:551-555.
16. ROBERTS, R. B., D. B. COWIE, P. H. ABELSON, E. T. BOLTON, AND R. J. BRITTEN. 1955. Studies on biosynthesis in *Escherichia coli*. *Carnegie Inst. Wash. Publ.* **607**, p. 13-30.
17. STRASTERS, K. C., AND K. C. WINKLER. 1963. Carbohydrate metabolism of *Staphylococcus aureus*. *J. Gen. Microbiol.* **33**:213-229.
18. UMBREIT, W. W., R. H. BURRIS, AND J. F. STAUFER. 1964. *Manometric techniques*. Burgess Publishing Co., Minneapolis.
19. WITTENBERG, C. L., AND A. S. HAAF. 1964. Lactate-degrading system in *Butyribacterium rettgeri* subject to glucose repression. *J. Bacteriol.* **88**:896-903.