Published in final edited form as:

J Am Chem Soc. 2011 August 31; 133(34): 13311–13313. doi:10.1021/ja205655w.

Discovery and Assembly Line Biosynthesis of the Lymphostin Pyrroloquinoline Alkaloid Family of mTOR Inhibitors in *Salinispora* Bacteria

Akimasa Miyanaga^{†,Δ}, Jeffrey E. Janso^{‡,Δ}, Leonard McDonald[‡], Min He[‡], Hongbo Liu[‡], Laurel Barbieri[‡], Alessandra S. Eustáquio[‡], Elisha N. Fielding[‡], Guy T. Carter[‡], Paul R. Jensen[‡], Xidong Feng[‡], Margaret Leighton[‡], Frank E. Koehn^{*,‡}, and Bradley S. Moore^{†,§,*}

[†]Center for Marine Biotechnology and Biomedicine, Scripps Institution of Oceanography, University of California at San Diego, La Jolla, California 92093-0204

[‡]Natural Products Laboratory, Worldwide Medicinal Chemistry, Pfizer Worldwide Research and Development, 558 Eastern Point Road, Groton, Connecticut 06340

[§]Skaggs School of Pharmacy and Pharmaceutical Sciences, University of California at San Diego, La Jolla, California 92093

Abstract

The pyrroloquinoline alkaloid family of natural products that includes the immunosuppressant lymphostin has long been postulated to arise from tryptophan. We now report the molecular basis of lymphostin biosynthesis in three marine *Salinispora* species that maintain conserved biosynthetic gene clusters harboring a hybrid nonribosomal peptide synthetase-polyketide synthase central to lymphostin assembly. Through a series of experiments involving gene mutations, stable isotope profiling, and natural product discovery, we report the assembly line biosynthesis of lymphostin and nine new analogues that exhibit potent mTOR inhibitory activity.

Since the discovery of discorhabdin C in 1986,¹ a group of cytotoxic natural products has emerged that contain a pyrroloquinoline core (Figure 1). Most members of this diverse family have been isolated from marine sponges² and include the batzellines, damirones, and makaluvamines.³ More recently, compounds of this class have been obtained from terrestrial organisms including makaluvamine A from a myxomycete⁴ and the sanguinones and mycenarubins from mushrooms of the genus *Mycena*.⁵ Variations in the structural complexity of these compounds arise from substitutions about the pyrrolo[4,3,2-*de*]quinoline core, which is thought to be derived from tryptophan. However, despite the large number of compounds currently known, biosynthetic studies have been hampered due to the uncultivability and genetic intractability of the producer organisms.

To date, two natural products containing a pyrrolo[4,3,2-*de*]quinoline core have been isolated from bacteria, the myosin-targeting ammosamides from *Streptomyces* strain CNR-698⁶ and the immunosuppressant lymphostin (**1**) that was first isolated from *Streptomyces* sp. KY11783.⁷ Lymphostin has been shown to inhibit both lymphocyte kinase (IC₅₀ 0.05 μM) and phosphatidylinositol 3-kinase (IC₅₀ 0.001 μM).⁸ This biological activity, along with lymphostin's tricyclic pyrroloquinoline skeleton, inspired a biomimetic

frank.koehn@pfizer.com; bsmoore@ucsd.edu.

^ΔThese authors contributed equally

Supporting Information Available: Figures S1–S4, Tables S1–S6, experimental methods, HRMS data, NMR spectra. This material is available free of charge via the Internet at <http://pubs.acs.org>.

total synthesis starting from tryptophan in 2004.⁹ We report here the molecular basis for lymphostin (**1**) biosynthesis in *Salinispora* spp. which involves a uniquely organized modular synthetase that through gene targeting provided the novel analogue lymphostinol (**2**). We also report eight new *N*-acyl derivatives including the potent mTOR inhibitor neolymphostin A (**3**).

Genome sequencing of the actinomycetes *Salinispora tropica* strain CNB-440 and *Salinispora arenicola* strain CNS-205 revealed that these marine bacteria have the genetic capacity to produce a wide variety of structurally and biologically diverse natural products.¹⁰ While the majority of their secondary metabolism genes are relatively unique, they share at least a dozen natural product gene clusters. Amongst these is the hybrid nonribosomal peptide synthetase-polyketide synthase (NRPS-PKS) locus *lym* (Stro3051–3057, Sare3277–3283) tentatively assigned to encode the biosynthesis of **1** (Figure 2A).¹⁰ The *lym* cluster is maintained in a third species, “*Salinispora pacifica*”, including strain DPJ-0019 where its gene organization is conserved (Table S1, Supporting Information [SI]).

Bioinformatic analysis of the syntenic *lym* loci in *S. tropica*, *S. arenicola* and “*S. pacifica*” established the approximate physical boundaries of the gene cluster that includes 7 genes covering 14.8-kb (Figure 2A). The central *lymA* gene encodes a heptadomain synthetase that putatively functions to malonate extend a tryptophan-derived peptidyl carrier protein (PCP)-bound residue and reductively release an *N*-acetylated diketide product as an aldehyde from an acyl carrier protein (ACP)-bound intermediate (Figure 2B). Trans methylation by the LymB methyltransferase (MT) would preserve the oxidation state of the released aldehyde as a methoxy enolate characteristic of **1** as a product of an atypical hybrid NRPS-PKS. Further sequence analysis of the LymA adenylation (A) domain revealed that the invariant Asp residue, which is required to stabilize the α -amino group of an amino acid substrate,¹¹ is replaced by Asn 226 that may activate an aryl acid substrate which may already contain the pyrrolo[4.3.2-*de*] core.

In order to explore this biosynthetic model of **1** assembly, we first established lymphostin production in *Salinispora* independently at UCSD and Pfizer. Standard growth conditions in A1 media that yielded products such as the salinosporamide proteasome inhibitors in *S. tropica*¹² and the cyclomarin peptides in *S. arenicola*,¹³ however, did not yield **1**. At UCSD, we explored different fermentation methods leading to a lymphostin production media that ultimately gave **1** in *S. tropica* and *S. arenicola* at ~0.4 mg/L and ~3 mg/L, respectively. We additionally observed a new derivative in *S. arenicola* with the molecular composition C₁₅H₁₄N₄O₃ as established by high-resolution electrospray ionization mass spectrometry (m/z [$M+H$]⁺ 299.1141 obs, 0.7 ppm error). Analysis of the ¹H and ¹³C NMR spectra, with the aid of gradient-enhanced HMBC data, clearly indicated that lymphostinol (**2**) was a novel lymphostin derivative with a modified C-4 side chain in which the β -methoxy enone moiety of **1** was replaced with a β -hydroxy ketone residue. Key HMBC signals connected the C-10/C-11 ethyl alcohol residue to the C-9 carbonyl at 200.2 ppm, which was supported by the difference in MS data of a single carbon atom between **1** and **2**.

In the Pfizer laboratory, **1** production conditions were first established with *S. arenicola* strain MOSO-0003 in 12 different media that ultimately gave a titer of ~20 mg/L in the medium M48-9. Subsequently, five of 11 different marine-invertebrate-derived strains of “*S. pacifica*” were found to produce **1**. Although most strains, such as DPJ-0024, produced **1** alone, strain DPJ-0019 was found to produce a mixture of lymphostin analogues. To improve the fermentation profile and facilitate downstream processing of material, a simpler production medium was developed containing only soluble starch and yeast extract (YESS). For comparative HPLC analysis of **1** production in strains from all three species, *S. tropica*

CNB-440, *S. arenicola* CNS-205, “*S. pacifica*” DPJ-0024 and DPJ-0019 were fermented in medium YESS at various scales, processed and analyzed in parallel (Figure 3).

Analysis of the comparative *lym* chemistry of the three strains revealed that *S. tropica* CNB-440 produced **1** alone, *S. arenicola* CNS-205 and “*S. pacifica*” DPJ-0024 produced **1**, **2**, **4**, and **8**, and “*S. pacifica*” DPJ-0019 uniquely produced neolymphostins **3**, **5**, and **6** and the corresponding neolymphostinols **7**, **9**, and **10** (Figure 3, Table S6). In the neolymphostins and neolymphostinols, the C-13 *N*-acetyl found in **1** and **2** is replaced by larger acyl groups, such as isobutyryl, propyl, sec-pentyryl, and isopentyryl (Figure 2B). All eight derivatives were isolated and fully characterized by HRMS and NMR analysis (Table S4). The most abundant member was neolymphostin A (**3**), whose ¹H and ¹³C NMR spectra showed the presence of a 2-propyl amide moiety (δ3.06(1H, m)/δ34.7 and δ1.22(6H,d)/δ18.9) in place of the acetamide in **1**. Lymphostin and the neolymphostins showed potent mTOR inhibition and cytotoxicity (Table 1), whereas the neolymphostinols as a whole were 1000-fold less active than their neolymphostin counterparts (Table S5).

Buoyed by the characterization of lymphostin chemistry, we next interrogated the *lym* clusters by PCR-targeted gene replacement recently optimized for *Salinispora* genetics.^{13,14} Inactivation of the conserved *lymA* gene resulted in *S. tropica* and *S. arenicola* mutants devoid of **1** and **2** (Figure S1), thereby confirming the central role of the LymA synthetase. To further establish the biosynthetic relationship between **1** and **2**, we inactivated the *S. arenicola* *lymB* MT. On the basis of biosynthetic precedence,¹⁵ we anticipated that the resulting mutant would accumulate an aldehyde intermediate that would lead to further carbonyl reduction by the LymA reductase in the absence of LymB (Figure 2B). Chemical analysis of the *lymB::apr^R* mutant revealed the complete loss of **1** and increased production of **2**, thereby confirming their biosynthetic linkage (Figure S1).

While the biosynthetic endgame is understood, we have yet to clarify the early stages of **1** biosynthesis that includes the formation of the pyrrolo[4,3,2-*de*]quinoline core structure. We established its biogenesis in “*S. pacifica*” strain DPJ-0019 from tryptophan through isotope experiments in which both nitrogen atoms were retained in the lymphostin products at N-1 and N-5 (Figure 2B). FTMS/MS analysis of ¹⁵N₂-L-tryptophan-derived **3** revealed both mono (21%) and di (41%) ¹⁵N-labeling. The remaining nitrogen atoms in **3** were likewise shown to derive from glutamine, providing a possible role of the conserved amidotransferase encoded by Strop_3057 in adding the aryl amine groups.

The molecular basis for the larger diversity of analogues produced by “*S. pacifica*” strain DPJ-0019 likely relies on the different substrate discrimination of the unusual LymA *N*-acetyltransferase (NAT) domain. As expected in the case of **3**, we confirmed that the *N*-isobutyryl group is derived from isobutyrate through ¹³C isotope labeling (Figure 2B). While NATs are common biosynthetic enzymes, its inclusion in a modular synthetase is to the best of our knowledge unprecedented. We are presently studying the molecular basis behind this difference in substrate specificity that offers an exciting opportunity to explore acyl group discrimination and to biosynthetically engineer new lymphostin and other pyrroloquinoline alkaloids.

In summary, we established the molecular basis of lymphostin biosynthesis in three *Salinispora* species by an unusual hybrid nonribosomal peptide-polyketide synthetase pathway that for the first time sheds light on the formation of pyrrolo[4,3,2-*de*]quinoline group alkaloids. Our work also resulted in the discovery of nine new lymphostin derivatives and the preliminary description of their potent mTOR inhibitory activity.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We gratefully acknowledge Dr. Yongxuan Su of the UCSD Mass Spectrometry Facility for MS assistance, Jan Kieleczawa and Tony Li at Pfizer's BioCore facility (Cambridge, MA) for DNA sequencing of *lym* genes in DPJ-0019, Dr. Nobuhiro Fusetani at Hokkaido University for providing *D. proliferum* tissue from which DPJ-0019 was isolated, and Dr. Ker Yu for biological assays. This work was generously supported in part by a research grant from the NIH (GM085770) to P.R.J. and B.S.M., and a postdoctoral fellowship from the JSPS to A.M.

References

1. Perry NB, Blunt JW, McCombs JD, Munro MHG. J. Org. Chem. 1986; 51:5476–5478.
2. Antunes EM, Copp BR, Davies-Coleman MT, Samaai T. Nat. Prod. Rep. 2005; 22:62–72. [PubMed: 15692617]
3. Sakemi S, Sun HH, Jefford CW, Bernardinelli G. Tetrahedron Lett. 1989; 30:2517–2520. Chang LC, Otero-Quintero S, Hooper JNA, Bewley CA. J. Nat. Prod. 2002; 65:776–778. [PubMed: 12027767] Stierle DB, Faulkner DJ. J. Nat. Prod. 1991; 54:1131–1133. Schmidt EW, Harper MK, Faulkner DJ. J. Nat. Prod. 1995; 58:1861–1867. [PubMed: 8691207] Radisky DC, Radisky ES, Barrows LR, Copp BR, Kramer RA, Ireland CM. J. Am. Chem. Soc. 1993; 115:1632–1638.
4. Ishibashi M, Iwasaki T, Imai S, Sakamoto S, Yamaguchi K, Ito A. J. Nat. Prod. 2000; 64:108–110. [PubMed: 11170681]
5. Peters S, Spiteller P. J. Nat. Prod. 2007; 70:1274–1277. [PubMed: 17658856] Peters S, Spiteller P. Eur. J. Org. Chem. 2007:1571–1576.
6. Hughes CC, MacMillan JB, Gaudêncio SP, Fenical W, La Clair JJ. Angew. Chem. Int. Ed. 2009; 48:728–732. Hughes CC, MacMillan JB, Gaudêncio SP, Jensen PR, Fenical W. Angew. Chem. Int. Ed. 2009; 48:725–727.
7. Nagata H, Ochiai Y, Aotani Y, Ando K, Yoshida M, Takahashi I, Tamaoki T. J. Antibiot. 1997; 50:537–542. [PubMed: 9711243] Aotani Y, Nagata H, Yoshida M. J. Antibiot. 1997; 50:543–545. [PubMed: 9711244]
8. Nagata H, Yano H, Sasaki K, Sato S, Nakanishi S, Takahashi I, Tamaoki T. Biosci. Biotechnol. Biochem. 2002; 66:501–507. [PubMed: 12005041]
9. Tatsuta K, Imamura K, Itoh S, Kasai S. Tetrahedron Lett. 2004; 45:2847–2850.
10. Udway DW, Zeigler L, Asolkar RN, Singan V, Lapidus A, Fenical W, Jensen PR, Moore BS. Proc. Natl. Acad. Sci. U.S.A. 2007; 104:10376–10381. [PubMed: 17563368] Penn K, Jenkins C, Nett M, Udway DW, Gontang EA, McGlinchey RP, Foster B, Lapidus A, Podell S, Allen EE, Moore BS, Jensen PR. ISME J. 2009; 3:1193–1203. [PubMed: 19474814] Nett M, Ikeda H, Moore BS. Nat. Prod. Rep. 2009; 26:1362–1384. [PubMed: 19844637]
11. May JJ, Kessler N, Marahiel MA, Stubbs MT. Proc. Natl. Acad. Sci. U.S.A. 2002; 99:12120–12125. [PubMed: 12221282]
12. Gulder TAM, Moore BS. Angew. Chem. Int. Ed. 2010; 49:9346–9367.
13. Schultz AW, Oh D-C, Carney JR, Williamson RT, Udway DW, Jensen PR, Gould SJ, Fenical W, Moore BS. J. Am. Chem. Soc. 2008; 130:4507–4516. [PubMed: 18331040]
14. Eustáquio AS, Pojer F, Noel JP, Moore BS. Nat. Chem. Biol. 2008; 4:69–74. [PubMed: 18059261]
15. Du L, Lou L. Nat. Prod. Rep. 2010; 27:255–278. [PubMed: 20111804]

Figure 1.
Structures of pyrrolo[4,3,2-*de*]quinoline natural products

Figure 2.

Organization of the *lym* gene cluster and proposed biosynthesis of lymphostin (**1**), lymphostinol (**2**), and related *N*-acyl derivatives **3–10**. **A**. The putative *lym* gene cluster is colored blue (including genes *lymA* and *lymB* that were inactivated in this study) with adjacent open reading frames (ORFs) in gray that are associated with ribosomal peptide biosynthesis. Stop gene numbers are shown flanking the cluster. Each arrow represents the direction of transcription of an ORF. See the SI for the deduced functions of the ORFs and a comparison of the *lym* gene products (Tables S2 and S3, respectively). **B**. Neolymphostin A (**3**) biosynthesis was probed in “*S. pacifica*” strain DPJ-0019 by isotope labeling with ¹⁵N₂-L-tryptophan, ¹⁵N-L-glutamine and ¹³C₄-isobutyrate as shown. Abbreviations: A,

adenylation; ACP, acyl carrier protein; AT, acyltransferase; KS, ketosynthase; MT, methyltransferase; NAT, *N*-acyltransferase; PCP, peptidyl carrier protein; R, reductase.

Figure 3.

Comparative HPLC analysis (467 nm) of crude extracts from “*S. pacifica*” DPJ-0019, *S. arenicola* CNS-205, “*S. pacifica*” DPJ-0024, and *S. tropica* CNB-440 fermentations. Lymphostin (1), lymphostinol (2), neolymphostins A (3), B (4), C (5) & D (6), neolymphostinols A (7) B (8) C (9) & D (10) are noted. See Table S6 for titers.

Table 1

mTor Kinase inhibition and cytotoxicity of lymphostin and neolymphostins

Compound	mTOR (nM)	LNCap (nM)	MDA-468 (nM)
1	1.7	38	14
3	0.8	22	58
4	1.5	48	85
5	1.8	230	700
6	1.8	230	700