Published in final edited form as:

J Am Chem Soc. 2011 August 31; 133(34): 13268–13271. doi:10.1021/ja204989p.

Reaction of O₂ with [(–)-Sparteine]Pd(H)Cl: Evidence for an Intramolecular [H–L]⁺ “Reductive Elimination” Pathway

Nattawan Decharin, Brian V. Popp, and Shannon S. Stahl*

Department of Chemistry, University of Wisconsin - Madison, 1101 University Avenue, Madison, Wisconsin 53706

Abstract

(Sp)PdCl₂ [Sp = (–)-sparteine] catalyzes a number of different aerobic oxidation reactions, and reaction of O₂ with a Pd^{II}–hydride intermediate, (Sp)Pd(H)Cl (**1**), is a key step in the proposed catalytic mechanism. Previous computational studies suggest that O₂ inserts into the Pd^{II}–H bond, initiated by abstraction of the hydrogen atom by O₂. Experimental and computational results obtained in the present study challenge this conclusion. Oxygenation of in-situ-generated (Sp)Pd(H)Cl exhibits a zero-order dependence on [O₂]. This result is inconsistent with a bimolecular H-atom-abstraction pathway, and DFT computational studies identify a novel “reductive elimination” mechanism, in which the chelating nitrogen ligand undergoes intramolecular deprotonation of the Pd^{II}–hydride. The relevance of this mechanism to other Pd^{II} oxidation catalysts with chelating nitrogen ligands is evaluated.

Pd-catalyzed aerobic oxidation reactions have expanded significantly over the past decade, and these advances are closely linked to the identification of oxidatively stable ancillary ligands that promote these reactions.¹ These ligands play a key role in stabilizing the reduced Pd catalyst formed upon substrate oxidation and promoting its reaction with molecular oxygen. The mechanism of catalyst oxidation by O₂ has been the focus of considerable recent study by us² and others.³ For catalytic reactions that proceed via Pd^{II}–hydride intermediates, two mechanistic pathways appear to be viable (Scheme 1A): (1) direct reaction of O₂ with the Pd–H bond via a hydrogen-atom-abstraction pathway (HAA, Path A) and a pathway initiated by H–X reductive elimination from Pd^{II}, followed by a reaction of O₂ with Pd⁰ (HXRE, Path B). Insights from recent experimental and computational studies suggest that all catalyst systems with labile monodentate ligands (e.g., pyridine, Et₃N, DMSO) strongly favor the HXRE pathway.⁴ Many catalysts feature chelating nitrogen ligands, however (Scheme 1B), and previous computational studies of the reaction of O₂ with (Sp)Pd(H)Cl (**1**) [Sp = (–)-sparteine] support an HAA mechanism.^{3a,d,5} Here, we present experimental data that are incompatible with an HAA mechanism, together with DFT computational studies that reveal a novel “reductive elimination” pathway for aerobic oxidation of **1**. These and additional results presented herein suggest that no existing Pd catalysts undergo aerobic oxidation via the HAA pathway.

The reaction (Sp)Pd(H)Cl with O₂ has been the focus of two previous computational studies by Goddard and coworkers.^{3a,d} The first study identified the unprecedented HAA pathway as a possible mechanism for O₂ “insertion” into a Pd–H bond (Scheme 2A).^{3a} The second

* stahl@chem.wisc.edu .

Supporting Information Available: Experimental procedures for (Sp)Pd(H)Cl synthesis and in situ oxygenation. Computational methods, gas-phase and toluene-solvated total energies with thermodynamic corrections, geometry optimization details, frequencies, Cartesian coordinates for optimized complexes. Detailed reaction coordinate for the oxygenation/protonolysis from complex **3** to form **9** and HLRE from (N–N)Pd(H)Cl complexes. These materials is available free of charge via the Internet at <http://pubs.acs.org>.

study showed that an alternative mechanism, initiated by deprotonation of the Pd–H, has a significantly higher activation energy relative to the HAA pathway (Scheme 2B; $\Delta\Delta G^\ddagger = 7.5$ kcal/mol).^{3d,6} These results have not been confirmed experimentally. In principle, these two mechanisms could be distinguished on the basis of their kinetic dependence on [O₂]: the HAA pathway predicts a first-order dependence, whereas the deprotonation mechanism predicts a zero-order dependence. This possibility prompted us to pursue the synthesis of (Sp)Pd(H)Cl (**1**).

We examined a number of methods for the preparation of **1**, including the reaction of (Sp)PdCl₂ with various Si-, B- and Sn-hydride reagents,⁷ β -hydride elimination from in-situ-generated Pd–formate and alkoxide derivatives,⁸ and addition of HCl to Pd⁰ sources in the presence of (–)-sparteine.⁹ With many of these methods, a common major Pd–H species was evident by ¹H NMR spectroscopy at reduced temperatures (≤ -35 °C); however, the complex was thermally unstable, and efforts to isolate the complex were unsuccessful. The instability of this Pd–H species is not entirely surprising. No mononuclear, nitrogen-chelated Pd–H complexes have been reported in the literature, and **1** has never been observed under catalytic conditions.

Among the synthetic routes evaluated, the most promising result was obtained from the reaction of excess (EtO)₃SiH (10 equiv) with (Sp)PdCl₂ in CD₂Cl₂ at –10 °C (Scheme 2). Over 25 min, the reaction turned from yellow to dark brown and a single Pd–hydride complex was formed in 40% yield by ¹H NMR spectroscopy ($\delta_{\text{Pd-H}} = -25.97$ ppm).¹⁰ The Pd–H complex decomposes if it is maintained at the reaction temperature, and it appears to have at least two accessible decomposition pathways (Scheme 3): (1) formation of H₂, evident by the emergence of a peak in the ¹H NMR spectrum at 4.59 ppm,¹¹ and (2) formation of the protonated sparteine ligand (Sp–H⁺) and Pd black, formally arising from N_{Sp}–H “reductive elimination” and aggregation of the Pd⁰ byproduct. The former decomposition pathway is especially evident when hydride reagents are used, and it may proceed via a Pd^{II}-dihydride intermediate (top pathway, Scheme 3). As discussed below, the latter pathway has important implications for the reaction of **1** with O₂.

The difficulties in isolating **1** prompted us to investigate the O₂ reactivity of the Pd^{II}–H complex in situ. When the reaction mixture, containing ~40% yield of Pd^{II}–H, was exposed to an atmosphere of O₂, the solution changed color from dark brown to yellow, and a ¹H NMR spectrum of the product solution revealed (Sp)PdCl₂ as the predominant product (Figure S1). A plausible explanation for this observation is that the Pd^{II}–H reacts with O₂ to afford a Pdhydroperoxide that undergoes rapid reaction with (EtO)₃SiCl present in the reaction mixture to afford (Sp)PdCl₂ and (EtO)₃SiOOH (Figure 1A). It was possible to monitor kinetics of the O₂ reactivity by ¹H NMR spectroscopy at –30 °C ($t_{1/2} \sim 20$ min),¹² and increasing the O₂ pressure from 1–5 atm exhibited no effect on the reaction rate (Figure 1B).

Interpretation of this result is subject to a number of caveats associated with the complexity of the reaction mixture and lack of full characterization of **1** or the Pd–hydroperoxide intermediate. Conservatively, however, the data indicate that a sparteine-coordinated Pd–hydride species undergoes a reaction that is induced by the presence of O₂, but is independent of the [O₂]. While this conclusion lacks mechanistic certainty, the data are not consistent with an HAA mechanism, which should exhibit a first-order dependence on [O₂].

Previous studies of the reactions between O₂ and well-defined (albeit catalytically unreactive) Pd–hydride complexes demonstrate close agreement between experimental and computational results.^{2b,c,e,3b,c} Consequently, the discrepancy between the experimental and computational results noted above suggest a mechanism different from those shown in

Scheme 2 might be involved. In order to examine this possibility, we decided to reinvestigate the reaction of **1** with O₂ using DFT computation methods.¹³

Analysis of the ground-state structure of (Sp)Pd(H)Cl **1** reveals that the two Pd–N bonds differ in length by 0.15 Å, with the longer one *trans* to the hydride ligand (Figure 2). This observation, together with the formation of Sp–H⁺ observed experimentally (cf. bottom pathway, Scheme 3), raised the possibility that sparteine could serve as an internal base upon dissociation of the amine ligand *trans* to the hydride ligand. This mechanism would be analogous to the deprotonation pathway noted in Scheme 2B, but it would avoid the entropic cost associated with a bimolecular reaction.^{3d}

A transition state for dissociation of the N^B amine of sparteine was identified computationally (**2**^{TS}, Figure 3). The imaginary frequency associated with this step includes both Pd–N^B bond lengthening and rotation of the sparteine ligand about the Pd–N^A bond. The rotation enables the basic N^B atom to approach the hydride ligand, and deprotonation of the hydride ligand by the N^B amine occurs, yielding the zwitterionic-Pd⁰ product **3** (Figure 3). The calculated free-energy barrier for this concerted process is 18.6 kcal/mol (Figure 3A). The Pd–H bond distance in **3** is 1.72 Å, and the Pd–H–N interaction is best described as a hydrogen bond between a Lewis-basic Pd⁰ center and an acidic H–N_{Sp} fragment.¹⁴ Natural charge analysis suggests the deprotonation can be described formally as an N–H “reductive elimination” reaction in which the Pd center starts with a +0.36 charge and ends with a –0.15 charge. The H atom simultaneously shifts from having slight hydridic character (–0.054 charge) in **1** to protic character, (+0.32 charge) in **3**. To summarize, the amine-dissociation and deprotonation steps occur as a concerted process. By analogy to the HXRE nomenclature employed when the reductive elimination step involves an anionic X-type ligand, we assign the name HLRE to the present reaction, reflecting the involvement of a neutral L-type ligand in the “reductive elimination” reaction.

Aerobic oxidation of the zwitterionic intermediate **3** proceeds via an oxygenation/protonolysis sequence in which all of the calculated intermediates and transition states are lower in energy than **2**^{TS} (Figures 3A and S3). These results indicate that the HLRE step is rate-limiting, and oxygenation of **1** by this mechanism should exhibit a zero-order dependence on [O₂]. Reinvestigation of the HAA mechanism, using the same computation package and methods employed for the HLRE study, revealed that the HLRE mechanism is more favorable than the HAA pathway: $\Delta G^\ddagger_{\text{HLRE}} - \Delta G^\ddagger_{\text{HAA}} = -3.5$ kcal/mol.¹⁵ The free energy of activation derived from the experiments in Figure 1 ($\Delta G^\ddagger_{\text{expt}} \sim 18$ kcal/mol) and the calculated barrier for the HLRE pathway are in good agreement,¹⁶ and we propose that aerobic oxidation of (Sp)Pd(H)Cl proceeds via this novel HLRE mechanism.

In order to probe the generality of the HLRE pathway, we performed analogous calculations with three other commonly used bidentate nitrogen ligands: phenanthroline (phen), bipyridine (bpy), and pyridine-oxazoline (py-ox). A slight difference observed in HLRE mechanism with these three ligands relative to (–)-sparteine was the presence of a stepwise, rather than concerted, ligand-dissociation/deprotonation pathway. The highest barrier for the pathway was associated with ligand dissociation.¹⁷ A comparison between the rate-limiting transition-state energies for the HLRE and HAA mechanisms with each of these ligands reveals that the HLRE mechanism is always lower in energy (Figure 4), although the difference is relatively small with the phenanthroline and bpy ligands ($\Delta\Delta G^\ddagger \sim 1$ –2 kcal/mol). Despite the relative small energy differences, however, these observations reveal the potential for these “ancillary” ligands to participate directly in reactions at the Pd center.

The mechanisms depicted in Figure 4 feature Pd–hydride complexes with chloride as the second anionic ligand; however, the (Sp)PdCl₂ catalyst system is unusual in its use of

chloride. The vast majority of nitrogen-ligated Pd catalysts for aerobic oxidation reactions feature carboxylates as the anionic ligands. We have previously highlighted the ability of acetate and other carboxylate ligands to undergo intramolecular deprotonation of Pd–hydrides, formally H–O₂CR "reductive elimination" from the Pd^{II} center.^{2b–e} Calculation of the rate-limiting transition states for this HXRE pathway for (N–N)Pd(H)(OAc) complexes (**10^{N–N}**) reveals that this mechanism is strongly favored¹⁸ relative to the HAA mechanism ($\Delta\Delta G^\ddagger = 8.8\text{--}11.9$ kcal/mol) (Figure 5). This mechanism is also significantly favored relative to the HLRE mechanism with Pd–chloride complexes by ($\Delta\Delta G^\ddagger = 6.4\text{--}10.1$ kcal/mol). The carboxylate HXRE pathway benefits energetically from the ability of the carboxylate ligand to retain partial bonding to the Pd center as it participates in the deprotonation step. In contrast, the bidentate nitrogen ligands must fully dissociate one of the nitrogen atoms before it can interact with the hydride ligand.

These observations have clear implications for Pd-catalyzed aerobic oxidation reactions. Pd^{II}–hydride intermediates in Pd-catalyzed aerobic oxidation reactions are elusive, and insights into their reactivity are limited by the lack of direct investigations. Nevertheless, results from experimental and computational studies of well-defined model systems suggested that nearly all catalyst systems undergo oxidation by O₂ via a Pd⁰ intermediate. The (Sp)PdCl₂ catalyst system appeared to be the sole exception. The present work, however, provides evidence for a previously unrecognized mechanism in which the chelating nitrogen ligand participates in intramolecular deprotonation of the hydride ligand, resulting in the formation of a Pd⁰ species that reacts with O₂. This mechanism exhibits many features in common with the previously characterized HX–reductive elimination pathway. In both pathways, one portion of the ligand remains coordinated to the Pd center while a remote site deprotonates the Pd–H. The intramolecular nature of this process avoids the unfavorable entropy associated with a bimolecular reaction. The hydrogen-atom-abstraction pathway remains an intriguing pathway for O₂ insertion into a Pd–H bond, but we are not aware of any aerobic oxidation reactions in which it represents a viable pathway for catalyst oxidation.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We are grateful to NIH for financial support of this work (R01 GM67163). Computational resources were supported by the NSF (CHE-0840494 and TG-CHE-070040) NMR spectroscopy facilities were partially supported by the NSF (CHE-9629688).

References

- (1). (a) Stahl SS. *Angew. Chem., Int. Ed.* 2004; 43:3400–3420. (b) Gligorich KM, Sigman MS. *Chem. Commun.* 2009:3854–3867.
- (2). (a) Konnick MM, Gandhi BA, Guzei IA, Stahl SS. *Angew. Chem., Int. Ed.* 2006; 45:2904–2907. (b) Popp BV, Stahl SS. *J. Am. Chem. Soc.* 2007; 129:4410–4422. [PubMed: 17371024] (c) Konnick MM, Stahl SS. *J. Am. Chem. Soc.* 2008; 130:5753–5762. [PubMed: 18393426] (d) Popp BV, Stahl SS. *Chem. Eur. J.* 2009; 15:2915–2922. (e) Konnick MM, Decharin N, Popp BV, Stahl SS. *Chem. Sci.* 2011:326–330.
- (3). For leading references, see: (a) Keith JM, Nielsen RJ, Oxgaard J, Goddard WA III. *J. Am. Chem. Soc.* 2005; 127:13172–13179. [PubMed: 16173744] ; (b) Denney MC, Smythe NA, Cetto KL, Kemp RA, Goldberg KI. *J. Am. Chem. Soc.* 2006; 128:2508–2509. [PubMed: 16492014] ; (c) Keith JM, Muller RP, Kemp RA, Goldberg KI, Goddard WA III, Oxgaard J. *Inorg. Chem.* 2006; 45:9631–9633. [PubMed: 17112254] ; (d) Keith JM, Goddard WA III, Oxgaard J. *J. Am. Chem.*

- Soc. 2007; 129:10361–10369. [PubMed: 17676735] ; (e) Keith JM, Goddard WA III. J. Am. Chem. Soc. 2009; 131:1416–1425. [PubMed: 19133788]
- (4). For justification of this claim, see refs. 2c and 2d and discussion therein
- (5). Application of (–)-sparteine as an ancillary ligand for Pd-catalyzed aerobic oxidation reactions was first reported by the groups of Sigman and Stoltz in the context of alcohol oxidation: (a) Jensen DR, Pugsley JS, Sigman MS. J. Am. Chem. Soc. 2001; 123:7475–7476. [PubMed: 11472200] ; (b) Ferreira EM, Stoltz BM. J. Am. Chem. Soc. 2001; 123:7725–7726. [PubMed: 11481006]
- (6). Direct H–Cl reductive elimination was determined to be highly unfavorable, so an external amine base was invoked to catalyze the transformation. See ref 3d for details
- (7). Reaction of L_2PdCl_2 with hydride donor reagents: (a) Brooks EH, Glocklin F. Chem. Commun. 1965:510.; (b) Moulton CJ, Shaw BL. J. Chem. Soc., Dalton Trans. 1976:1020–1024.; (c) Saito T, Munakata H, Imoto H. Inorg. Synth. 1977; Vol. 17:83–88.; (d) Imoto H, Moriyama H, Saito T, Sasaki Y. J. Organomet. Chem. 1976; 120:453–460.; (e) Moriyama H, Saito T, Sasaki Y. Chem. Lett. 1976:175–176.; (f) Reger DL, Garza DG. Organometallics. 1993; 12:554–558.; (g) Fryzuk MD, Lloyd BR, Clentsmith GKB, Rettig SJ. J. Am. Chem. Soc. 1991; 113:4332–4334.; (h) ref. 3b; (i) Nguyen DH, Laurenczy G, Urrutigoñy M, Kalck P. Eur. J. Inorg. Chem. 2005; 2005:4215–4225.
- (8). β -hydride elimination from Pd-alkyl, -alkoxide and -formate complexes: (a) Hosokawa T, Maitlis PM. J. Am. Chem. Soc. 1972; 94:3238–3240.; (b) Goel AB, Goel S. Inorg. Chim. Acta. 1980; 45:L85–L86.; (c) Clegg W, Eastham GR, Elsegood MRJ, Heaton BT, Iggo JA, Tooze RP, Whyman R, Zacchini S. J. Chem. Soc., Dalton Trans. 2002:3300–3308.; (d) Johansson R, Wendt OF. Organometallics. 2007; 26:2426–2430.
- (9). Oxidative addition of an HM^+ source to a L_2Pd^0 : (a) Ozawa F, Ito T, Nakamura Y, Yamamoto A. J. Organomet. Chem. 1979; 168:375–391.; (b) Seligson AL, Cowan RL, Troglor WC. Inorg. Chem. 1991; 30:3371–3381.
- (10). The large upfield signal of Pd–H resonances is common: Grushin VV. Chem. Rev. 1996; 96:2011–2034. [PubMed: 11848819]
- (11). Fulmer GR, Miller AJM, Sherden NH, Gottlieb HE, Nudelman A, Stoltz BM, Bercaw JE, Goldberg KI. Organometallics. 2010; 29:2176–2179.
- (12). Background decomposition of (sp)Pd(H)Cl at this temperature is > 7-fold slower than the O_2 -induced reaction, and the decomposition reaction does not afford (Sp)PdCl₂ as the product. These consideration suggest that the background process should has little effect on the observed oxygenation kinetics
- (13). Computational methods: All computations were performed with the *Gaussian 03* program. Geometry optimizations were performed using the B3LYP functional with the Stuttgart RSC 1997 ECP basis set for Pd and 6–31+G(d) for all other atoms. At the calculated stationary points, solvation-corrected single-point-energy calculations (toluene solvent at 298 K) were carried out with the Pd basis detailed above and the 6–311+G(d,p) basis for all other atoms. See Supporting Information for details
- (14). Falvello LR. Angew. Chem., Int. Ed. 2010; 49:10045–10047.
- (15). The calculated HAA transition state, employing our DFT methods, is 22.1 kcal/mol relative to (Sp)Pd(H)Cl
- (16). The extent of agreement between the experimental and calculated barriers is almost certainly fortuitous, considering the typical errors associated with DFT calculations (~3–5 kcal/mole). See, for example: Siegbahn PEM. J. Bio. Inorg. Chem. 2006; 11:695–701. [PubMed: 16830147]
- (17). See Supporting Information for the energy diagrams
- (18). The reductive-elimination mechanism from (Sp)Pd(H)OAc system has been previously studied: Keith JM. Palladium Mediated Activation of Molecular Oxygen. Ph.D. Thesis. 2008 California Institute of Technology

Figure 1.

(A) Proposed pathway for oxygenation of in-situ-generated (Sp)Pd(H)Cl. (B) Oxygen-pressure effects on the oxygenation reactions, demonstrating zero-order kinetics. Reaction conditions: [Pd-H] = 0.53 – 0.92 mM, p_{O_2} = 1.0 – 5.0 atm (9.0 – 45 mM of dissolved O₂), CD₂Cl₂, –30 °C, 600 μ L. Error bars reflect standard deviations based on independently prepared samples of the Pd^{II}-H.

Figure 2.
Ball-and-stick illustrations of (Sp)Pd(H)Cl with key metrics

Figure 3. Computed reaction coordinate for oxygenation of **1** initiated by H-ligand-reductive-elimination (HLRE). (A) Reaction profile for the HLRE, oxygenation, and protonolysis steps in toluene solvent. See Supporting Information for complete reaction coordinate. (B) Ball-and-stick illustrations with key metrics for the complexes **2^{TS}** and **3**.

Figure 4. Comparison of the rate-limiting transition-state energies for the HLRE and HAA mechanisms for O_2 insertion into a Pd–H bond with four different bidentate nitrogen ligands. Full reaction coordinates are provided in the Supporting Information.

Figure 5. Comparison of the rate-limiting transition-state energies for the HXRE (X = OAc) and HAA mechanisms for O₂ insertion into a Pd–H bond with four different bidentate nitrogen ligands.

Scheme 1.
Possible Mechanisms for the Oxidation of Pd-H Species by Dioxygen and Representative Catalyst Systems with *N*-Chelating Ligands

(A) H-Atom-Abstraction (HAA) Pathway **(B) External Base Deprotonation Pathway** **Scheme 2.**

Two Proposed Mechanistic Pathways for O₂ Insertion into Pd–Hydride Bond of (Sp)Pd(H)Cl Complex.^{3a,d}

Scheme 3.

Synthesis of (Sp)Pd(H)Cl and Proposed Decomposition Pathways under the Reaction Conditions.