Published in final edited form as:

Org Lett. 2011 September 2; 13(17): 4684–4687. doi:10.1021/ol201900d.

Copper-Catalyzed β -Boration of α,β -Unsaturated Carbonyl Compounds with Tetrahydroxydiborane

Gary A. Molander* and Silas A. McKee

Roy and Diana Vagelos Laboratories, Department of Chemistry, University of Pennsylvania, Philadelphia, Pennsylvania 19104-6323, United States

Abstract

The copper-catalyzed β -boration of α,β -unsaturated carbonyl compounds with tetrahydroxydiborane has been developed. This diboron reagent allows direct, efficient access to boronic acids and their derivatives. Primary, secondary and tertiary α,β -unsaturated amides are converted to the corresponding β -trifluoroboratoamides in good to excellent yields. The β -boration of a variety of α,β -unsaturated esters and ketones is also reported.

Boron homoenolates are versatile compounds that have been developed as synthetic intermediates for a variety of transformations,¹ including Suzuki coupling reactions.² Significantly, secondary β -trifluoroboratoamides have been demonstrated to react stereospecifically with inversion of configuration in palladium catalyzed cross-couplings.^{2b} These developments have greatly expanded the utility of trifluoroborato homoenolates in organic synthesis.

A particularly appealing approach to the synthesis of these compounds is the transition metal-catalyzed β -boration of readily available α,β -unsaturated carbonyl compounds using diboron reagents. The β -boration of α,β -unsaturated carbonyl compounds with bis(pinacolato)diboron is well established.³ Rhodium,⁴ platinum,⁵ palladium,⁶ nickel⁷ and copper^{6a–b,8} catalyzed methods have been previously reported. Of these, the copper catalyzed systems have received the most attention. Yun and coworkers were instrumental in expanding the substrate scope from the more traditional unsaturated ketones and esters to include the more challenging α,β -unsaturated amides.⁹

Although previously reported catalytic systems provide efficient access to pinacol boronate esters, they are less convenient for the syntheses of other boronate derivatives. Conversion of the pinacol esters to the corresponding trifluoroborate can be problematic,¹⁰ as pinacol can be difficult to remove from the crude reaction mixtures, requiring tedious purification protocols. Additionally, in the interest of atom economy, use of a borylating reagent that does not require the intermediacy of the pinacol ester would prove beneficial.

Despite the interest in copper catalyzed beta-boration, there have been only a few publications examining alternative diboron reagents,¹¹ each of which provides access to a

gmolandr@sas.upenn.edu.

Supporting Information Available. Experimental procedures and spectral data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

single boronate ester. Our group has recently established tetrahydroxydiborane (bisboronic acid, BBA)¹² as an effective replacement for bis(pinacolato)diboron in palladium catalyzed Miyaura borylation reactions.¹³ This reagent offers direct access to boronic acids, simplifying purification and providing access to a variety of boronate derivatives through a single intermediate. Herein we report a method for the β -boration of α,β -unsaturated carbonyl compounds using BBA as an effective and atom economical source of boron. To our knowledge this is the first example of the β -boration of α,β -unsaturated carbonyl compounds with BBA, as well as the first copper-catalyzed reaction of this reagent.

Reaction conditions for the β -borylation were optimized on (*E*)-*N*-cyclohexylbut-2-enamide. For ease of analysis, the initially formed boronic acid was converted in situ to the corresponding alcohol by oxidation with NaOH/H₂O₂. A screen of various solvents, alkoxide bases, copper(I) sources and phosphine ligands established EtOH, NaOt-Bu, CuCl, and 2-(dicyclohexylphosphino)biphenyl (CyJohnPhos) as the preferred conditions. A 1:1 ratio of ligand to copper and 1.2 equivalents of BBA was effective for >90% conversion after 4 hours.

We next sought to reduce the catalyst loading. Interestingly, initial attempts to reduce the catalyst load resulted in incomplete conversion even at long reaction times (Table 1, entries 1–3), despite the relatively short reaction times at high catalyst load. Consequently, attention was focused on the mechanism of the process and specifically the role of the base in these transformations (Scheme 1).^{9,14}

Previous studies have established that the base is critical in the formation of an sp²–sp³ species (**I**), creating a more nucleophilic intermediate that facilitates transmetalation to the active borylcopper species **II**.^{11c} Insertion of the α,β -unsaturated carbonyl compound into the copper boron bond of **II** gives a copper enolate (**III**), which accepts a proton from EtOH to give the borylated product **IV**, regenerating a copper alkoxide catalyst. We hypothesized that the formation of a borate ester byproduct **V** sequestered the NaOt-Bu necessary for catalyst regeneration as the reaction proceeded toward completion. Therefore, the incomplete conversion at low catalyst loads could be due to insufficient quantities of base rather than of catalyst.

Encouragingly, we found that keeping 30 mol % of NaOt-Bu allowed a reduction in the amount of CuCl and CyJohnPhos to 1 mol % while retaining efficient conversion to the product (entries 4–6). Reactions run with 2 mol % of CuCl and CyJohnPhos were found to be more reproducible, and thus 2 mol % of both catalyst and ligand were used as the general conditions for subsequent reactions. Increasing the amount of NaOt-Bu beyond 30 mol % provided no advantage.

With optimized conditions in hand, we examined the substrate scope of this reaction (Table 2). We first varied the substitution on a series on *N*-cyclohexyl amides. Methyl substitution at the alpha position was well tolerated, as was mono-alkyl or phenyl substitution at the beta position. Branching at the gamma position was tolerated, but required a longer reaction time, increased catalyst loading, and 2 equiv of BBA for efficient conversion (Table 2, entry 5). Effective conversion could not be obtained, however, for compounds containing multiple substituents at the beta position, presumably for steric reasons (Table 2, entry 6). Next we examined the effect of the amine portion of the amide on the borylation reaction. Primary, secondary and tertiary amides were all tolerated, although primary amides (Table 2, entry 12) required longer reaction times. The reaction could be run on gram scale with only a slight reduction in yield (Table 2, entry 1). Unfortunately, attempts to isolate the boronic acid intermediate synthesized from **1a** yielded an inseparable mixture of boronic acid and what appears to be boroxine, along with ligand and traces of starting material.

We were gratified to find that α,β -unsaturated esters and ketones were also efficiently borylated under the standard reaction conditions (Table 3). A variety of substitution patterns were effective for ketones, with cyclic and acyclic substrates proving to be good substrates for the reaction. Even sterically hindered ketones were borylated in good yield, albeit with a longer reaction time (Table 3, entry 3). Esters were equally effective coupling partners (Table 3, entries 5–8), and notably lactone **5h** was a competent substrate.

In summary, we have developed an efficient copper catalyzed method for the synthesis of β -trifluoroborato carbonyl compounds using bisboronic acid as a source of boron – the first copper-catalyzed process for this atom economical reagent. This method uses EtOH as an environmentally benign solvent, avoids purification difficulties encountered in the conversion of pinacol esters to the corresponding trifluoroborates, and provides efficient access to β -amido trifluoroborates. Efforts toward an enantioselective variant are ongoing in our laboratory.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This research was supported by the NIGMS (R01 GM-035249). Dr. Rakesh Kohli (University of Pennsylvania) is acknowledged for obtaining HRMS data. We thank BASF for a generous donation of tetrahydroxydiborane.

References

1. Bonet A, Sole C, Gulyas H, Fernandéz E. *Curr Org Chem*. 2010; 14:2531.
2. (a) Molander GA, Petrillo DE. *Org Lett*. 2008; 10:1795. [PubMed: 18393522] (b) Sandrock DL, Jean-Gérard L, Cheng-yi C, Dreher SD, Molander GA. *J Am Chem Soc*. 2010; 132:17108.(c) Molander GA, Jean-Gérard L. *J Org Chem*. 2009; 74:1297. [PubMed: 19133770] (d) Molander GA, Jean-Gérard L. *J Org Chem*. 2009; 74:5446. [PubMed: 19572522]
3. For a current review see: Schifner JA, Mütter K, Oestreich. *Angew Chem Int Ed*. 2010; 49:1194.
4. Kabalka GW, Das BC, Das S. *Tetrahedron Lett*. 2002; 43:2323.(b) Shiomi T, Adachi T, Toribatake K, Zhou L, Nishiyama H. *Chem Commun*. 2009:5987.
5. (a) Lawson Y, Lesly G, Marder T, Norman N, Rice C. *Chem Commun*. 1997:2051.(b) Bell J, Cox A, Cameron N, Evans J, Marder T, Duin M, Elsevier C, Baucherel X, Tulloch A, Tooze R. *Chem Commun*. 2004:1854.
6. (a) Lillo V, Geier MJ, Westcott SA, Fernández E. *Org Biomol Chem*. 2009; 7:4674. [PubMed: 19865703] (b) Bonet A, Gulyás H, Koshevoy IO, Estevan F, Sanaú M, Úbeda MA, Fernandéz E. *Chem Eur J*. 2010; 16:6382.
7. Hirano K, Yorimitsu H, Oshima K. *Org Lett*. 2007; 9:5031. [PubMed: 17973483]
8. (a) Takahashi K, Ishiyama T, Miyaura N. *Chem Lett*. 2000:982.(b) Ito H, Yamanaka H, Tateiwa J, Hosomi A. *Tetrahedron Lett*. 2000; 41:6821.(c) Chea H, Sim HS, Yun J. *Adv Synth Catal*. 2009; 351:855.
9. Mun S, Lee J, Yun J. *Org Lett*. 2006; 8:4887. [PubMed: 17020328]
10. Bagutski V, Ros A, Aggarwal V. *Tetrahedron*. 2009; 65:9956.
11. (a) Gao M, Thorpe SB, Santos WL. *Org Lett*. 2009; 11:1899. [PubMed: 19331360] (b) Thorpe SB, Guo X, Santos WL. *Chem Commun*. 2011; 47:424.(c) Gao M, Thorpe SB, Kleeberk C, Slebodnick C, Marder TB, Santos WL. *J Org Chem*. 2011; 76:3997. [PubMed: 21491953]
12. BBA reactions in the literature: (a) Sebelius S, Olsson V, Szabo K. *J Am Chem Soc*. 2005; 127:10478. [PubMed: 16045321] (b) Olsson V, Sebelius S, Selander N, Szabo K. *J Am Chem Soc*. 2006; 128:4588. [PubMed: 16594692] (c) Selander N, Kipke A, Sebelius S, Szabo K. *J Am Chem Soc*. 2007; 129:13723. [PubMed: 17929926] (d) Selander N, Szabo K. *Chem Commun*. 2008:3420.(e) Pilarski LT, Szabo KJ. *Angew Chem Int Ed*. 10.1002/anie.201102384

13. Molander GA, Trice SLJ, Dreher SD. J Am Chem Soc. 2010; 132:17701. [PubMed: 21105666]
14. Dang, Li; Lin, Z.; Marder, T. Organometallics. 2008; 27:4443.

Scheme 1.
Proposed Mechanism

Table 1

Catalyst Load Optimization^a

1) CyJohnPhos, CuCl
NaOt-Bu, Bz₂(OH)₄ (1.2 equiv),
EtOH, rt
2) NaOH, H₂O₂

entry	CuCl (mol %)	CyJohnPhos (mol %)	NaOtBu (mol %)	NaOH (mol %)	reaction time (h)	P:SM ^b
1	10	10	30		18	>20:1
2	5	5	15		18	1.4:1
3	3	3	9		18	0.4:1
4	5	5	30		18	>20:1
5	3	3	30		18	>20:1
6	1	1	30		18	>20:1
7	1	1	30		1	>18:1
8	1	1	30		2	>19:1
9	1	1	30		3	>19:1

^a All reactions were carried out on a 0.3 mmol scale in 3 mL of EtOH with 1.2 equiv of BBA.

^b Ratio of product to starting material calculated by ¹H NMR.

Table 2

 β -Boration of α,β -Unsaturated Amides with Bisboronic Acid^a

$ \begin{array}{c} \text{R}^1_2\text{N}-\text{C}(=\text{O})-\text{CH}=\text{CH}-\text{R}^3 \\ \text{1a-l} \end{array} \xrightarrow[2) \text{KHF}_2, \text{MeOH}, \text{H}_2\text{O}]{1) \text{CyJohnPhos (2 mol \%), CuCl (2 mol \%), NaOt-Bu (30 mol \%), B}_2(\text{OH})_4 \text{ (1.2 equiv), EtOH, rt}} \begin{array}{c} \text{R}^1_2\text{N}-\text{C}(=\text{O})-\text{CH}(\text{BF}_3\text{K})-\text{CH}(\text{R}^3)-\text{R}^3 \\ \text{3a-l} \end{array} $				
entry	time (h)	product	yield ^b (%)	
1	2		3a	90 (82) ^c
2	2		3b	89
3	2		3c	98
4	2		3d	86
5	18		3e	75 ^d
6	18		3f	— ^d
7	24		3g	98
8	2		3h	96
9	2		3i	90
10	2		3j	90
11	2		3k	88
12	18		3l	78

^a General conditions: Amide (1.0 equiv), BBA (1.2 equiv), CuCl (2 mol %), CyJohnPhos (2 mol %), NaOt-Bu (30 mol %), EtOH (0.1 M), rt. After filtration and solvent removal, MeOH, KHF₂ (8.2 equiv), 0 °C to rt, 20 min.

^b Yield is given for isolated products.

^c Gram scale.

^d Used 2.0 equiv BBA, CuCl (10 mol %), NaOt-Bu (50 mol %) CyJohnPhos (10 mol %).

Table 3 β -Boration of α,β -Unsaturated Carbonyl Compounds with Bisboronic Acid^a

entry	time (h)	product	yield ^b (%)
1	2		71
2	2		93
3	24		77
4	2		76
5	2		93
6	2		84
7	2		82
8	2		83

^a General conditions: Carbonyl substrate (1.0 equiv), BBA (1.2 equiv), CuCl (2 mol %), CyJohnPhos (2 mol %), NaOt-Bu (30 mol %), EtOH (0.1 M), rt. After filtration and solvent removal, MeOH, KHF₂ (8.2 equiv), 0 °C to rt, 20 min.

^b Yield is given for isolated products.