Published in final edited form as:

J Nat Prod. 2011 June 24; 74(6): 1535–1538. doi:10.1021/np200362q.

The Biosynthetic Origin of the 3-Amino-2,5,7,8-tetrahydroxy-10-methylundecanoic Acid Moiety and Absolute Configuration of Pahayokolides A and B

Li Liu[†], Daniel W. Bearden[‡], and Kathleen S. Rein^{*†}

[†]Department of Chemistry and Biochemistry, Florida International University, Miami, FL 33199, USA

[‡]Analytical Chemistry Division, National Institute of Standards and Technology, Hollings Marine Laboratory, Charleston, South Carolina 29412

Abstract

Pahayokolides A (**1**) and B (**2**) are cyclic undecapeptides that were isolated from the cyanobacterium *Lyngbya* sp. They contain the unusual α -hydroxy- β -amino acid, 3-amino-2,5,7,8-tetrahydroxy-10-methylundecanoic acid (Athmu). The absolute configurations of the amino acids of the pahayokolides, except for the four oxygen bearing stereocenters of Athmu, have been determined by Marfey's method. Incorporation of labeled leucine and acetate precursors into the pahayokolides have established that Athmu is derived from a leucine or α -keto isocaproic acid starter unit which is further extended with three acetate units.

Cyanobacteria have proven to be a rich source of biologically active secondary metabolites.^{1, 2} Natural product classes isolated from cyanobacteria include ribosomal and non-ribosomal peptides, mixed non-ribosomal peptides, polyketides and alkaloids.^{3, 4} The Florida Everglades is an oligotrophic marsh and an abundant source of diverse species of cyanobacteria. As a potential source of secondary metabolites however, the Florida Everglades remains largely unexplored. We have previously reported studies on the planar structure⁵, isolation⁶ and cytotoxicity⁷ of two related cyclic peptides, pahayokolides A (**1**) and B (**2**) which are produced by a *Lyngbya* sp., isolated from the Florida Everglades. Pahayokolides A and B contain the same cyclic undecapeptide core and pahayokolide A contains a pendant *N*-acetyl-*N*-methylleucine moiety which is absent in pahayokolide B. The pahayokolides are remarkably similar in structure to tychonamides A and B⁸ (**4** and **5**), schizotrin A⁹ (**3**), portoamides A and B¹⁰ (**6**) and lyngbyazothrins¹¹ (Figure 1), cyclic peptides isolated from *Tychonema* sp., *Schizothrix* sp. and *Oscillatoria* sp., respectively. The pahayokolides, schizotrin A and the portoamides and lyngbyazothrins¹¹ have an unusual β -amino acid, 3-amino-2,5,7,8-tetrahydroxy-10-methylundecanoic acid (Athmu). The tychonamides possess a similar β -amino acid, 3-amino-2,5,7-trihydroxy-8-phenyloctanoic acid (Atpoa). Like pahayokolide A, tychonamides A and B and portoamide contain a pendant *N*-acetyl-*N*-methylleucine connected via an ester linkage to the 5-hydroxy group of the β -amino acid whereas the appendage on schizotrin A is an *N*-butyryl-*N*-methylalanine. Herein we report the absolute configuration of the amino acids of pahayokolide A (**1**) and

*To whom correspondence should be addressed. Tel.: +1-305-348-6682; Fax: +1-305-348-3772. reink@fiu.edu.

Supporting Information Available

Marfey's analysis of pahayokolide A and ¹³C NMR spectra for natural abundance pahayokolide A and B, ¹³C enriched pahayokolide A [1-¹³C]leucine, [1, 2-¹³C]leucine, and ¹³C enriched pahayokolide B, [1-¹³C]acetate. This material is available free of charge via the Internet at <http://pubs.acs.org>.

the results of stable isotope incorporation experiments to determine the precursors of the Athmu moiety.

The absolute configurations of the common amino acids and homophenylalanine were determined by Marfey's HPLC method¹² and the advanced Marfey's method¹³ by comparison to authentic D and L standards derivatized with L-FDLA (1-fluoro-2,4-dinitrophenyl-5-L-leucinamide). The assignments are as follows: L-proline (X2), L-phenylalanine, L-serine, L-threonine, D-homophenylalanine, D-glutamine and *N*-methyl-L-leucine. In the case of *N*-methyl-leucine only the L isomer was commercially available. Comparison to the diastereomeric derivatives of *N*-methyl-L-leucine prepared with both L-FDLA and D-FDLA by the advanced Marfey's method showed the presence of *N*-methyl-(D/L)-leucine. However, the D/L ratio was lower under milder hydrolysis conditions and we conclude that the D isomer arose from partial racemization of *N*-methyl-L-leucine. A similar observation was made for the tychonamides.⁸ The configuration at C-54 is tentatively assigned as *R* (or D). This assignment is based on relative retention times of the diastereomeric Athmu derivatives, assuming that D-FLDA-D-Athmu (which is equivalent to L-FLDA-L-Athmu) < L-FDLA-D-Athmu.¹³

Assignments of ^{13}C NMR signals and the results of isotope incorporations into the Athmu and *N*-methyl-*N*-acetyl leucine subunits of pahayokolides A (**1**) and B (**2**) are presented in Table 1. The first five carbons in Table 1 are those which we anticipated would be enriched by one or more of the following feeding experiments. Because there is significant spectral overlap and the presence of several, slowly-interconverting conformers of pahayokolide A (**1**), leading to multiple overlapping peaks in the ^{13}C -NMR spectrum, we chose for comparison only those ^{13}C signals which are well resolved, always present as a single peak and which we did not expect to be enriched. These are the last eight carbons in Table 1. The ^{13}C NMR spectrum ($\text{DMSO}-d_6/\text{D}_2\text{O}$) of **1** derived from $[1-^{13}\text{C}]$ leucine fed cultures showed significant enrichment at C-58 and C-64¹⁴ while doubly labeled $[1, 2-^{13}\text{C}]$ leucine showed enrichment at C-58, C-59, C-64 and C-65 as well as the anticipated ^{13}C - ^{13}C coupling. Because of the presence of multiple conformers of **1**, the *N*-methyl-*N*-acetyl leucine of $[1-^{13}\text{C}]$ acetate labeled **1** was cleaved by treatment with dilute base to provide pahayokolide B (**2**) as previously described.⁵ The ^{13}C NMR spectrum ($\text{CD}_3\text{OD}/\text{D}_2\text{O}$) derived from $[1-^{13}\text{C}]$ acetate fed cultures showed enrichment at C-6, C-51, C-52, C-54, C-56 and C-58. Not shown in Table 1 are the results of feeding experiments with $[1-^{13}\text{C}]$ valine which failed to yield enrichment at any position.

β -Amino acids are not uncommon in cyanobacterial peptides and include Adda (3-amino-9-methoxy-2,6,8-trimethyl-10-phenyl-4,6-decadienoic acid) in the microcystins,¹⁶ and nodularin,¹⁷ Ahoa (3-amino-2,5-dihydroxy-8-phenyloctanoic acid) in nostophycin,¹⁸ and Ahda (3-amino-2,5,9-trihydroxy-10-phenyldecanoic acid) in scytonemin A.¹⁹ The Adda moiety of the microcystins has been shown to be of mixed peptide and polyketide origin. Stable isotope feeding experiments demonstrated that the biosynthesis of the Adda sidechain begins with a phenylalanine starter unit which is extended by four rounds of condensation

with malonate.²⁰ Thus four PKS (polyketide synthase) modules are involved in the biosynthesis of the polyketide amino acid Adda.²¹

Inspection of the Athmu residue of the pahayokolides suggests a mixed peptide and polyketide origin as well. We anticipated that the starter unit for the biosynthesis of the Athmu sidechain could be valine, leucine or the corresponding α -keto acids: α -ketoisovaleric acid, or α -ketoisocaproic acid. Leucine or α -ketoisocaproic acid seemed the more likely starter unit, as this could be further extended by three rounds of condensation with malonate units (Figure 2). Mehner, et al., proposed a similar polyketide pathway for the α -hydroxy- β -amino acids Ahda, Ahoa and Atpoa⁸ and Leao, et al.,¹⁰ proposed a similar pathway for Athmu. Alternatively, Leao, et al.,¹⁰ suggested that the starter unit might be acetate which is then doubly methylated at C-2. Our data are consistent with the first hypothesis. The anticipated enrichment of C-58 upon feeding of [1-¹³C] leucine as well as enrichment at C-58 and C-59 and ¹³C-¹³C coupling upon feeding of [1,2-¹³C] leucine was observed. Enrichment at C-52, C-54 and C-56 of **2** from [1-¹³C] acetate fed culture demonstrates the extension of leucine with three acetate units. The enrichment of C-58 in **2** from [1-¹³C] acetate fed cultures as well as other sites of enrichment (C-51 and C-6) can be easily rationalized. The incorporation of [1-¹³C] acetate at C-58 occurs via the condensation of acetate with α -ketoisovalerate during the biosynthesis of leucine. Enrichment at C-51 occurs via the condensation of [1-¹³C] acetate with oxaloacetate during the biosynthesis of α -ketoglutarate to produce glutamate. Through a similar mechanism, incorporation at C-6 results from the biosynthesis of homophenylalanine from phenylalanine.^{22,23} Clearly the Athmu moiety is derived from leucine and three units of acetate.

Experimental Section

General Experimental Procedures

One dimensional high resolution ¹³C NMR data for pahayokolide A (**1**) were acquired in 30:70 DMSO-*d*₆: D₂O using a Bruker AVANCE 400 MHz spectrometer (Bruker Biospin, Inc.) operating at 100 MHz for ¹³C. One dimensional high resolution ¹³C NMR spectra for pahayokolide B (**2**) were acquired in 50:50 CD₃OD/D₂O on a Bruker Avance II 700 MHz spectrometer equipped with a TCI cryoprobe operating at 176 MHz for ¹³C. All data were collected at a temperature of 298K. Chemical shifts were referenced to the deuterium CD₃ lock reference at 3.30 ppm using an absolute referencing scheme based on nuclear gyromagnetic constant ratios; this gives the center peak of the ¹³C CD₃OD septet a chemical shift of 47.603 ppm. The NMR data were processed with Topspin software. All mass spectra were acquired on single quadrupole mass spectrometer (ThermoQuest Finnigan Navigator) in ESI negative ion mode. High performance liquid chromatography (HPLC) was performed using a Thermo Finnigan Spectra SYSTEM HPLC (model P4000 pump; model AS3000 autosampler; model UV6000LP PDA UV detector) with an Apollo C18 (250×4.6 mm i.d., 5 μ , Alltech) column. Stable isotope precursors were purchased from Cambridge Isotope Laboratories.

Culture of *Lyngbya* sp

Lyngbya sp. strain 15-2 was maintained in 2 L cultures in BG11 medium, buffered with 2-(N-morpholino)ethanesulfonic acid buffer (2.6 mM) at pH 7.2. Cultures were supplemented with 150 mg of ¹³C labeled L-leucine (50 mg each on days 30, 34 and 38) or 1.3 g of ¹³C labeled sodium acetate (430 mg each on days 30, 34 and 38). Cultures were harvested after six weeks of growth. Pahayokolide A (**1**) and B (**2**) were isolated from the biomass as previously described.⁵

Preparation of FDLA derivatives

To 40 μ L of a 2 μ M solution of each amino acid standard was added 20 μ L of 1 M sodium bicarbonate and 80 μ L of 1% (w/v) L- or D-FDLA in acetone as previously described.^{11, 12} After incubating at 40 °C for 60 min the reactions were quenched by the addition of 10 μ L of 2 M HCl and stored at 4 °C. As *N*-Me-D-Leu was not available, the D-FDLA derivative of *N*-Me- L-Leu was used as a standard in lieu of its enantiomer the L-FDLA- *N*-Me-D-Leu derivative. One hundred μ g of pahayokolide A was hydrolyzed at 110 °C for 14 h with 500 μ L of 4M HCl. This solution was divided into two portions and dried under a stream of N₂. Each portion was derivatized with either L- or D-FDLA as described above and stored at 4 °C.

HPLC/PDA Conditions

The mobile phase used for the separation of the L- and D, L-FDLA derivatives of pahayokolide A was CH₃CN: 0.01 M TFA, step gradient [4:6 for 24 min; ramp to 1:1 from 24 to 34 min; 7:3 after 34 min] at a flow rate of 0.4 mL/min. The FDLA derivatives were detected with a PDA UV detector at 340 nm. The retention times for the L-FDLA derivatives of *N*-Me- L-Leu and L-homoPhe were very close at 47.3 and 47.7 min respectively. In separate experiments, the L-FDLA derivatized pahayokolide A hydrolysate was spiked with the L-FDLA derivatives of *N*-Me- L-Leu or L-homoPhe, confirming the presence of *N*-Me- L-Leu and the absence of L-homoPhe. Similarly, the retention times for the L-FDLA derivatives of D-Phe and *N*-Me- D-Leu were very close at 50.0 and 50.2 min respectively. In separate experiments, the L-FDLA derivatized pahayokolide A hydrolysate was spiked with the L-FDLA- D-Phe derivative and the D-FDLA-*N*-Me-L-Leu derivative confirming the presence of *N*-Me- D-Leu and the absence of D-Phe. When pahayokolide A was hydrolyzed in 6M HCl, the L:D ratio of *N*-Me-Leu was 2.75:1. However, hydrolysis of pahayokolide A in 4 M HCl, resulted in a L:D ratio of 5.5:1. We conclude that the *N*-Me- D-Leu arose from epimerization of *N*-Me-L-Leu during hydrolysis. A similar observation was made for the tychonamides.⁸ The FDLA derivative of Athmu was observed only when the hydrolysis was performed in 4 M HCl. When the hydrolysis was carried out in 6 M HCl a derivative having a molecular ion at *m/z* 668 was observed, corresponding to a loss of water from the FDLA-Athmu derivative. Either one of the alcohols is dehydrated to an alkene or the lactone is formed under these conditions.

ESI LC/MS Conditions

The sample probe was set at 400 °C and 4 kV with an entrance cone voltage of 10V. Chromatographic conditions were as described for the HPLC-PDA analysis with a flow rate of 0.5 ml/min.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by the National Institute of Environmental Health Sciences (NIEHS) Grant S11 ES11181. We acknowledge the support of the Hollings Marine Laboratory NMR Facility. Commercial equipment or materials are identified in this paper to specify adequately the experimental procedure. Such identification does not imply recommendation or endorsement by NIST, nor does it imply that the materials or equipment identified are necessarily the best available for the purpose.

References and notes

1. Gademann K, Portmann C. *Curr Org Chem*. 2008; 12:326–341.

2. Sivonen, K.; Börner, T. The Cyanobacteria: Molecular Biology, Genomics and Evolution. Herrero, A.; Flores, E., editors. Vol. 7. Caister Academic Press; 2008. p. 159-197.
3. Tan LT. Phytochemistry. 2007; 68:954–979. [PubMed: 17336349]
4. Banker R, Teltsch B, Sukenik A, Carmeli S. J Nat Prod. 2000; 63:387–389. [PubMed: 10757726]
5. An T, Kumar TKS, Wang M, Liu L, Lay JO Jr, Liyanage R, Berry J, Gantar M, Marks V, Gawley RE, Rein KS. J Nat Prod. 2007; 70:730–735. [PubMed: 17432902]
6. Berry, JP.; Gantar, M.; Gawley, RE.; Rein, KS. Harmful Algae 2002, Xth International Conference. Steidinger, KA.; Landsberg, JH.; Tomas, CR.; Vargo, GA., editors. Vol. 1. Florida Fish and Wildlife Conservation Commission, Florida Institute of Oceanography, Intergovernmental Oceanographic Commission of UNESCO; St. Petersburg, FL: 2004. p. 192-194.
7. Berry JP, Gantar M, Gawley RE, Wang M, Rein KS. Comp Biochem Physiol C. Toxicol Pharmacol. 2004; 139C:231–238.
8. Mehner C, Mueller D, Krick A, Kehraus S, Löeser R, Güetschow M, Maier A, Fiebig HF, Brun R, Köenig GM. Eur J Org Chem. 2008; 10:1732–1739.
9. Pergament I, Carmeli S. Tetrahedron Lett. 1994; 35:8473–8476.
10. Leão PN, Pereira AR, Liu WT, Ng J, Pevzner PA, Dorrestein PC, König GM, Vasconcelos VM, Gerwick WH. Proc Natl Acad Sci U S A. 2010; 107:11183–11188. [PubMed: 20534563]
11. Zainuddin EN, Jansen R, Nimtz M, Wray V, Preisitsch M, Lalk M, Mundt S. J Nat Prod. 2009; 72:1373–1378. [PubMed: 19621877]
12. Fujii K, Ikai Y, Oka H, Suzuki M, Harada K. Anal Chem. 1997; 69:5146–5151.
13. Fujii K, Shimoya T, Ikai Y, Oka H, Harada K. Tetrahedron Lett. 1998; 39:2579–2582.
14. As a result of experiments reported here we have revised our original assignments of C-31 and C-64. The correct assignments are C-31; 172.6 and C-64; 172.4.
15. Sitachitta N, Marquez BL, Thomas Williamson R, Rossi J, Ann Roberts M, Gerwick WH, Nguyen V-A, Willis CL. Tetrahedron. 2000; 56:9103–9113.
16. Carmichael WW. Sci Am. 1994; 270:64–72.
17. Annala A, Lehtimäki J, Mattila K, Eriksson JE, Sivonen K, Rantala TT, Drakenberg T. J Biol Chem. 1996; 271:16695–16702. [PubMed: 8663277]
18. Fujii K, Sivonen K, Kashiwagi T, Hirayama K, Harada K. J Org Chem. 1999; 64:5777–5782.
19. Helms GL, Moore RE, Niemczura WP, Patterson GML, Tomer KB, Gross ML. J Org Chem. 1988; 53:1296–1307.
20. Moore RE, Chen JL, Moore BS, Patterson GML, Carmichael W. J Am Chem Soc. 1991; 113:5083–5084.
21. Tillett D, Dittmann E, Erhard M, von Döhren H, Börner T, Neilan BA. Chem Biol. 2000; 7:753–764. [PubMed: 11033079]
22. Dewick, PM. Medicinal Natural Products: A Biosynthetic Approach. 2. John Wiley & Sons; New York: 2001. p. 459
23. Underhill, E.; Wetter, LR.; Chisholm, MD. Nitrogen Metabolism in Plants. In: Goodwin, TW.; Smelhe, RMS., editors. Biochemistry Society Symposium. Vol. 38. The Biochemical Society; 1973. p. 303-326.

1	Pro	Hphe	Thr	Dhb	Ser	Dhb	Phe	Pro	Gly	Gln	Athmu	[N-Ac-N-Me-Leu]
3	Pro	O-MeHty	Ser	Dhb	Ser	Val	Phe	Pro	Gly	Gln	Athmu	[N-Ac-N-Me-Leu]
4	Pro	O-MeHty	Thr	Dhb	Ser	Ile	Pro	Pro	Gly	Gln	Atpoa	[N-Ac-N-Me-Leu]
5	Pro	Hphe	Thr	Dhb	Ser	Ile	Pro	Pro	Gly	Gln	Atpoa	[N-Ac-N-Me-Leu]
6	Pro	O-MeHty	Thr	Dhb	Ser	Ile	Pro	Pro	Gly	Gln	Athmu	[N-Ac-N-Me-Tyr]

Figure 1.

Alignments of the core amino acids of the pahayokolide A (**1**), schizotrin A (**3**), tychonamides A and B (**4** and **5**), and portoamides (**6**). (The portoamides and the lyngbyazothrins are identical compounds isolated from different sources.)

Figure 2.
Proposed biogenic pathway for the β-amino acid Athmu.

Table 1

Isotopic enrichment based on ^{13}C NMR for pahayokolides A (**1**) and **B** (**2**)^a

position	δ_c (ppm) Pah A (1)	% enrichment Pah A (1) ^b	^{13}C leucine Pah A (1) ^b	% enrichment Pah A (1) ^b	^{13}C leucine Pah B (2) ^c	δ_c (ppm) Pah B (2) ^c	% enrichment Pah B (2) ^c	^{13}C acetate Pah B (2)
52	173.0	125			90	173.0		235
53	72.9	112			133	73.1		84
54	48.8	104			73	49.5		193
56	70.1	67			61	64.7		211
57	36.9	108			94	36.3		99
58	71.4	332			410	71.9		197
59	72.4	100			260	72.6		96
60	41.1	94			86	39.2		98
64	172.4	436			311	-		-
65	55.4	90			379	-		-
71	174.4	105			52	-		-
2	60.9	78			118	60.7		85
6	171.6	109			95	171.4		193
13	126.7	111			99	126.7		116
18	67.4	112			125	67.4		126
24	171.3	103			91	170.8		103
44	47.0	88			129	46.5		141
51	176.7	136			110	176.7		220
70	32.8	98			97	-		-

^a Carbon resonances were integrated relative to the signal for C-24. Enriched signals are shown in bold.

^b DMSO- d_6 /D $_2$ O.

^c CD $_3$ OD/D $_2$ O. Enrichment was calculated using the integral values for carbons shown on this table and applying the method described by Sitachitta.¹⁵ Average isotopic enrichment for carbons 2–70 above (excluding signals in bold) = 101 ± 20 .