Published in final edited form as:

Mol Ecol. 2011 September ; 20(17): 3503–3512. doi:10.1111/j.1365-294X.2011.05166.x.

Connecting the Sun to Flowering in Sunflower Adaptation

Benjamin K. Blackman^{1,2,4}, Scott D. Michaels¹, and Loren H. Rieseberg^{1,3}

¹Department of Biology, Indiana University, 1001 East Third Street, Bloomington, IN 47405, USA

²Department of Biology, Duke University, Box 90338, Durham, NC 27701, USA

³Department of Botany, University of British Columbia, 3529-6270 University Boulevard, Vancouver, BC V6T 1Z4, Canada

Abstract

Species living in seasonal environments often adaptively time their reproduction in response to photoperiod cues. We characterized the expression of genes in the flowering-time regulatory network across wild populations of the common sunflower, *Helianthus annuus*, that we found to be adaptively differentiated for photoperiod response. The observed clinal variation was associated with changes at multiple hierarchical levels in multiple pathways. Paralog-specific changes in *FT* homolog expression and tissue-specific changes in *SOC1* homolog expression were associated with loss and reversal of plasticity respectively, suggesting that redundancy and modularity are gene network characteristics easily exploited by natural selection to produce evolutionary innovation. Distinct genetic mechanisms contribute to convergent evolution of photoperiod responses within sunflower, suggesting regulatory network architecture does not impose strong constraints on the evolution of phenotypic plasticity.

Keywords

phenotypic plasticity; flowering time; sunflower; *Helianthus annuus*; photoperiod response; gene network evolution

Introduction

The timing and course of development can be highly plastic. Many traits change in response to environmental cues, and these plastic responses may provide important adaptations to changing environmental conditions (Schlichting & Pigliucci 1998). As species expand their ranges, colonize novel habitats, or face anthropogenic changes, phenotypic plasticity may be gained, lost, or otherwise altered by natural selection such that phenotypes adapted to these environments continue to be produced under locally predictive combinations of cues (Bradshaw & Holzapfel 2008; Moczek & Nijhout 2003). Complex gene regulatory networks sense and integrate various environmental signals to produce appropriate phenotypic responses. However, few studies have leveraged knowledge of these pathways to explore the mechanisms underlying evolutionary changes in plasticity in natural populations (Abouheif & Wray 2002).

⁴Author for correspondence: Benjamin K. Blackman, Department of Biology, Duke University, Box 90338, Durham, NC 27701. Phone: 919.660.7223. Fax: 919.660.7293. bkb7@duke.edu.

Data Accessibility

DNA sequences: GenBank accessions HM163571-HM163573

Phenotypic data for sunflowers and *A. thaliana* transformants, microsatellite data, and gene expression data: DRYAD entry doi: 10.5061/dryad.49sk0

In plants, a complex gene network regulates phenotypic plasticity in reproductive timing by integrating information from diverse cues including photoperiod so that flowering occurs during favorable conditions (Wilczek *et al.* 2010). Notably, rewiring of a conserved pathway can reverse photoperiod response. In *Arabidopsis thaliana*, a long-day plant that flowers earlier when day length exceeds a particular threshold, circadian and light signals interact so that *CONSTANS* (*CO*) only promotes expression of the mobile floral signal *FLOWERING LOCUS T* (*FT*) in long days (Corbesier *et al.* 2007; Valverde *et al.* 2004). In short-day sensitive rice, where flowering occurs earlier in day lengths below a threshold, *CO* represses *FT* in long days but promotes *FT* in short days (Hayama *et al.* 2003). Geographic clines in flowering time and photoperiod plasticity are often reported, but the adaptive nature and genetic causes of this variation are rarely determined (Bohlenius *et al.* 2006; Takahashi *et al.* 2009; Zhang *et al.* 2008). Consequently, the degree to which network structure predicts what genetic changes can yield such evolutionary shifts in plasticity is unknown (Erwin & Davidson 2009; Stern & Orgogozo 2009). Here, we circumscribe components of the flowering-time gene regulatory network likely contributing to adaptive evolution of flowering time plasticity in the common sunflower, *Helianthus annuus*, and discuss what these findings reveal about the interplay between natural selection and aspects of network structure that confer evolutionary flexibility or constraint.

Materials and Methods

Phenology Characterization

Six populations along a latitudinal transect (Fig. 1a)—Manitoba (MB), South Dakota (SD), Nebraska (NE), Kansas (KS), Oklahoma (OK), and Texas (TX)—were chosen on the basis of availability of seeds with family structure, absence of recent nearby sunflower cultivation, and previous use in genetic crosses (Table S1). Seeds from five heads per population were scarified, germinated in the dark on moistened filter paper for five days, and exposed to a single light-dark cycle to allow greening before sowing. Maternal half-sibs were evenly distributed among photoperiod treatments. TX seed with family structure was unavailable; seeds of unknown family structure were provided by the United States Department of Agriculture (PI494567). Seedlings were sown in 5" pots containing a 1:1 mixture by weight of sand and Metro Mix (Sun Gro Horticulture). Growth chambers were maintained at 25.5°C and 8 hours light : 16 hours dark, 12 hours light : 12 hours dark, or 16 hours light : 8 hours dark. Light intensity was adjusted so plants in different photoperiods received equivalent daily photosynthetically active radiation. Days to budding (R1 stage; Schneiter & Miller 1981) and anthesis were measured for one plant per maternal half-sib family in each chamber. Plant positions were randomized biweekly. Log-transformed data were analyzed by balanced ANOVA in Minitab specifying photoperiod treatment, latitude, and their interaction as fixed effects.

Qst Measurement

Seeds from ten maternal half-sib families per population were germinated. OK was excluded due to insufficient germination; too few seeds per family remained from the previous collection, and newly collected seeds were immature as the population had just reached peak flowering at collection (Fig. 1c). TX families came from new collections. Five seedlings per family were sown into 6" pots and grown in 16-hour days in the greenhouse under natural light supplemented by artificial lights. Plants were distributed one individual per family per block over five blocks. Every 7–10 days, plant positions within blocks and block locations in the greenhouse were randomized.

Days to budding and days to anthesis were measured with germination date as day 0. Leaf traits were measured at 28 days after sowing on mature leaves subtending the third node.

Floral traits were measured at anthesis. Leaf area was measured using a LI-COR LI-3000A Portable Area Meter. Leaf wet weights and dry weights were measured on a scientific balance, and specific leaf area was calculated as area / dry weight. Succulence was calculated as (wet weight - dry weight) / area. Ray, bract, and leaf shapes were calculated as the ratio of maximum length to maximum width of that structure. An outlier for specific leaf area was removed from the analysis as it was very aberrant and strongly affected the results. Bract shape, ray shape, ray number, and specific leaf area were log-transformed to improve normality of the residuals.

Qst values were calculated from variance components obtained using a restricted maximum likelihood mixed-linear model implemented with the lme function in R (The R Project for Statistical Computing) specifying block as a fixed effect and population and family nested within population as random effects (Pinheiro & Bates 2000). Variance components were extracted with the *varcomp* function. Qst was calculated as $V_P / (V_P + 2 * (4 * V_F))$ where V_P is the variance between populations, and V_F is the variance among maternal half-sib families within populations. Multiplying V_F by four yields the additive genetic variance (Lynch & Walsh 1998). Theory predicts dominance and epistasis only reduce Qst relative Fst (Whitlock 2008). Thus, if these factors impact the variation observed, they would only reduce our power to detect Qst >> Fst. Since we used a maternal half-sib design, our model cannot account for maternal effects. Confidence intervals were estimated by simulation (O'Hara & Merila 2005) using the maximum likelihood estimates for the fitted values and dispersion matrices obtained from the model. Qst values were calculated for 1000 simulations. Upper and lower percentile cutoffs for 95% confidence intervals were obtained using the BCa method (Efron & Tibshirani 1993).

Fst Measurement

DNA was extracted from one seedling from each of 16 families per population—including all families used for Qst measurement—with the Qiagen Plant DNeasy 96 Kit. Individuals were genotyped for 11 microsatellite markers (Table S3) as described previously (Heesacker *et al.* 2008). Fst was calculated with Microsatellite Analyzer (Dieringer & Schlötterer 2003), and confidence intervals were obtained by bootstrapping over loci 1000 times.

Transformation

HaCOL1 and *HaCOL2* cDNAs were amplified by PCR from cultivar HA274—*HaCOL1*: forward – CACCATGTTAAATGAAGATCTCTACTAG, reverse – TTTGATCCGGAGCATTGCTTAAA; *HaCOL2*: forward – CACCATGTTGGATCACACCGGTACCTTATG, reverse – CGTCTTTAAACGAGGGTACAATTCC. MB, KS, and TX *HaCOL2* cDNA sequences were also obtained with these primers.

Amplified fragments were introduced by Gateway™ cloning (Invitrogen) into vector pMDC32 (Curtis & Grossniklaus 2003). Columbia-0 and *co*-mutant (SAIL24H04) *A. thaliana* were transformed by the floral dip method (*Agrobacterium tumefaciens* strain GV1301). After two days stratification at 4°C, seeds were germinated at 23°C on 50µg/ml hygromycin MS plates. Resistant transformants were transplanted to Metro Mix five days later. Successful transformation was confirmed by RT-PCR.

Gene Expression

Leaves were collected every four hours over a single day ~5 weeks after sowing from plants grown alongside the phenotyped plants in the 8-hour and 16-hour treatments. For each time point within each treatment, two biological replicates consisting of leaves pooled from two to three plants from the same population were collected.

To determine the diurnal phase of expression for photoperiod pathway homologs, we germinated intrapopulation F1 seed generated from plants raised for Qst measurement. For OK, field collected seeds were used. Leaf samples were collected at 6, 4, 3, 2, and 1 hours before dawn, dawn, and 2, 4, 5, 6, 7, 8, 10, 12, and 15 hours after dawn ~5 weeks after sowing under 8-hour days. The periods of most frequent sampling correspond to the periods when *CO*, *GIGANTEA (GI)*, and *FLAVIN-BINDING KELCH-REPEAT F-BOX 1 (FKF1)* paralog up-regulation occurs. Three biological replicates were collected, and replicate leaves were harvested from different nodes to average out developmental differences. Leaves were never collected from the same plant within a two-hour period.

For shoot apex expression, MB, KS, and TX maternal half-sibs were grown under 8- and 16-hour days. Three biological replicates consisting of shoot apices pooled from two to three plants were collected ~5 weeks after sowing.

RNA extraction, DNase treatment and quantitative RT-PCR were conducted as previously described (Blackman *et al.* 2010). Primers are listed in Table S3. Two to three technical replicates were performed for each reaction and averaged. Relative expression was calculated as $2^{-\Delta\Delta C_t}$ —normalizing to *Ha60S rRNA*—and converted to a percentage of the highest expression value measured in either light treatment. For several genes, transcript abundance was assayed by RT-PCR and visualized by ethidium bromide staining on an agarose gel (28 cycles for *Ha60S rRNA*; 30 cycles for all other genes).

Regressions of total daily gene expression on latitude or mean flowering time on mean gene expression were completed in Minitab. Total daily gene expression was calculated as the sum of expression means for the biological replicates at the six time points assayed. For the diurnal phase grow-out, hourly transcript abundances were interpolated into intervals not sampled at that frequency, and total gene expression was calculated over the time course's duration.

Results

Diversity in Flowering Time and Photoperiod Response

We investigated the diversity of photoperiodic flowering in wild sunflower, *Helianthus annuus*, because the existence of long-day, short-day, and photoperiod insensitive cultivars suggested an opportunity to identify mechanisms through which such variation can evolve within species (Goyné & Schneiter 1987). Seeds from six populations sampled along a latitudinal transect from Manitoba (49.23°N) to Texas (28.13°N) were germinated and grown under 8-, 12-, or 16-hour days (Fig. 1a and Table S1). Day length, latitude, and their interaction explained nearly all the observed variation (balanced ANOVA; budding: $R^2 = 0.84$, $F_{\text{day length}}(5,89) = 45.62$, $F_{\text{latitude}}(2,89) = 30.21$, $F_{\text{interaction}}(10, 89) = 8.66$; anthesis: $R^2 = 0.88$, $F_{\text{day length}}(5,89) = 59.46$, $F_{\text{latitude}}(2,89) = 77.68$, $F_{\text{interaction}}(10, 89) = 11.75$; $p < 0.001$ for all terms for both traits). The number of days to budding and days to anthesis increased as latitude decreased in all treatments, and photoperiod response transitions occurred at both ends of the cline (Fig. 1b). All three major photoperiod response classes were observed. Manitoba (MB), the northernmost population, was day neutral, flowering earliest and at the same age under all conditions. Mid-latitude populations—South Dakota (SD), Nebraska (NE), Kansas (KS), and Oklahoma (OK)—were short-day sensitive, flowering ~0.5 to three months earlier under 8- or 12-hour days than under 16-hour days. In contrast, plants from the southernmost population, Texas (TX), were long-day sensitive, flowering about a month earlier under 12- and 16-hour days than under 8-hour conditions.

Variation was also observed in natural habitats. For instance, Oklahoman populations reached peak flowering during the shortening days of early fall when populations in central

Texas had already senesced, having flowered months earlier under long days (Fig. 1, c and d). This differentiation is likely adaptive. Short growing seasons in Manitoba may impose selection for initiating reproduction prior to experiencing photoperiods that induce short-day populations to flower. In Texas, warm, wet winters and severely dry summers shift the growth season; populations germinate in late February and flower prior to summer drought.

Comparison of Qst and Fst

We raised maternal half-sib families from all populations, except OK, under long days and measured phenological traits in addition to leaf, floral, and physiological characters uncorrelated with flowering to obtain estimates of Qst, a measure of quantitative trait differentiation. Populations were also genotyped for 11 microsatellite markers to estimate Fst, a measure of neutral genetic differentiation. As neutrally evolving traits and neutral genetic markers should accumulate similar levels of differentiation, if a trait's Qst is much greater than Fst, then a hypothesis of neutral evolution can be rejected (Spitze 1993; Whitlock 2008). Qst for flowering-time measures were 4–6 times as high as mean Fst and also higher than nearly all other traits (Table 1), consistent with adaptive differentiation in response to geographic variation in selection. Qst for succulence was also high, which we attribute to strong differentiation of TX from other populations and little variation among families within populations.

Gene Expression Correlates of Flowering Time Variation

We next examined gene expression variation in the photoperiod and gibberellin pathways because these cues have demonstrated effects on flowering in sunflower (Almeida & Pereira 1996; Goyné & Schneiter 1987). Gibberellins (GAs) promote flowering by antagonizing DELLA-domain proteins, transcriptional repressors that suppress expression of the floral inducers *SUPPRESSOR OF OVEREXPRESSION OF CONSTANS 1* (*SOC1*) and *LEAFY* (*LFY*) (Achard *et al.* 2004; Ueguchi-Tanaka *et al.* 2005). We drew on these known regulatory pathways by inferring that if gene expression and phenotype were associated, then the changes responsible for both likely act upstream in the network or in *cis*- to expression of that gene. If upstream genes lacked these associations, we inferred the changes responsible for the phenotype act downstream or in a parallel pathway to expression of these genes. Thus, we bracketed network sections likely contributing to the cline and evolutionary shifts between photoperiod response types.

Based on their evolutionary conservation among angiosperms, we predicted that the network's gene content and hierarchical structure are conserved in sunflower (Hayama *et al.* 2003; Maizel *et al.* 2005; Serrano *et al.* 2009; Tamaki *et al.* 2007; Yasumura *et al.* 2007). Sunflower homologs of many flowering genes are known, though the genome is not sequenced and additional paralogs may exist (Blackman *et al.* 2011). Even so, functional equivalence of three sunflower *FT* homologs was confirmed by heterologous complementation in *Arabidopsis* (Blackman *et al.* 2010). We also overexpressed two *CO* homologs in wild-type and *co*-mutant *A. thaliana*. Although neither paralog accelerated flowering in wild-type plants, *HaCOL2* overexpression complemented the *co* mutation while *HaCOL1* overexpression did not (Fig. S1). For functionally uncharacterized paralog sets, we measured expression of all known copies.

We first assayed transcript levels throughout the diurnal cycle for the downstream floral inducers *FT* and *SOC1* in leaves from sunflowers grown in 8- or 16-hour days. Two *FT* homologs exhibited clines in expression under short days. Higher total daily gene expression levels were significantly associated with higher latitude (Fig. 2, a and b; regression, *HaFT2*: $R^2=79.1\%$, $F_{1,5}=15.11$, $p=0.018$; *HaFT4*: $R^2=80.4\%$, $F_{1,5}=16.40$, $p=0.015$) and were also predictive for earlier flowering (*HaFT2*: $R^2=60.8\%$, $F_{1,5}=6.22$, $p=0.067$; *HaFT4*: $R^2=64.1\%$,

$F_{1,5}=7.13$, $p=0.056$). In both photoperiods, *HaSOC1A* expression was higher in more northern populations (Fig. 2c; short days: $R^2=80.1\%$, $F_{1,5}=16.07$, $p=0.016$; long days: $R^2=64.4\%$, $F_{1,5}=7.23$, $p=0.055$) and positively associated with flowering time (short days: $R^2=75.2\%$, $F_{1,5}=12.13$, $p=0.025$; long days: $R^2=67.7\%$, $F_{1,5}=8.37$, $p=0.044$). Similar trends were not observed for *HaSOC1B* (Fig. S2 and Table S2). These patterns suggest that molecular changes contributing to the overall cline in flowering time may involve both the photoperiod and gibberellin pathways and most likely act upstream or in *cis*- to *HaFT2*, *HaFT4*, and *HaSOC1A* expression (Fig. 2f).

Although neither *HaFT* paralog was expressed in long days in most populations, consistent with short-day response, *HaFT4* was expressed in long days in MB, consistent with day-neutrality (Fig. 2b). Therefore, changes upstream or in *cis*- to this *HaFT* paralog appear to cause a gain of expression likely involved in loss of photoperiod-responsive flowering in *H. annuus*' northern range.

We next examined expression of *CO* and DELLA protein homologs. In long days, one of two leaf-expressed DELLA-like genes was expressed at higher levels in later flowering populations, indicating the genetic changes contributing the overall cline in long days may act upstream or in *cis*- to its expression (Fig. 2, e and f, and Fig. S3). This finding was surprising and must be treated tentatively since functional consequences of variation in DELLA transcription for flowering have not been reported; however, it is noteworthy that DELLA gene expression in *A. thaliana* responds to a wide variety of cues (Gallego-Bartolomé *et al.* 2010) and transcriptional regulation of DELLA genes is an important mediator of light signaling on germination (Oh *et al.* 2007).

HaCOL2 daily expression abundance was neither associated with latitude (Fig. 2d; short days: $R^2<0.1\%$, $F_{1,5}<0.01$, $p=0.986$; long days: $R^2=31.6\%$, $F_{1,5}=1.84$, $p=0.246$) nor associated with flowering time (short days: $R^2=23.7\%$, $F_{1,5}=1.24$, $p=0.328$; long days: $R^2=52.1\%$, $F_{1,5}=4.36$, $p=0.105$), and MB expression levels did not differ notably. These findings suggested that the changes responsible for the overall cline in short days and the transition to day neutrality lie downstream or act in parallel to *HaCOL2* expression (Fig. 2f). We subsequently assayed *HaCOL2* expression at finer temporal resolution to better characterize its phase of expression relative to daylight, a parameter critical in determining whether *CO* promotes or represses flowering. Unlike northern populations where *HaCOL2* expression peaks before dawn, the peak occurs at dawn in OK and TX (Fig. S4), and this shifted coincidence of *HaCOL2* abundance and light may partly explain the reduced *HaFT* expression in these populations (Fig. 2f). This trend may also be related to the observation that peak *HaFT4* expression occurred later in the day as latitude decreased from MB to KS (Fig. S2a), which may parallel similar findings in poplar (Bohlenius *et al.* 2006); however, we note this trend was only observed in one of our two experiments. No nonsynonymous *HaCOL2* sequence changes are fixed between populations. Expression assays of additional genes homologous to factors acting upstream of *CO*—e.g. *G1* and *FKF1*—or upstream of *FT* and downstream or in parallel to *CO* revealed no clines in transcript abundance or diurnal phase (Fig. S3–S6 and Table S2).

We also examined expression of floral inducers further downstream in the regulatory network that act in the shoot apex. As in leaves, *HaSOC1A* expression in short days was clinal; however, its expression pattern differed dramatically under long days: expression in TX was as high as expression in MB, consistent with long-day response (Fig. 3 and Table S2). *HaSOC1B* exhibited similar expression variation, and likewise, some of this variation is not mirrored in the leaf (Fig. S2 and Table S2). Therefore, tissue-specific regulatory changes acting upstream or in *cis*- to shoot apex expression of *HaSOC1* genes likely contribute to the transition from short-day to long-day response (Fig. 2F). Expression of downstream genes

HaLFY and *HAM75*, an *APETALA1* (*API*) homolog, was consistent with this interpretation (Fig. S7). *HaFT1* and the *FRUITFULL* homolog *HaFUL* exhibited shoot apex expression patterns similar to the *SOC1* homologs, though less robust (Fig. S7). Since conservation of the regulatory hierarchy among these genes is ambiguous, positioning of important evolutionary changes in TX relative to these genes is uncertain. Shoot-apex expression levels of three DELLA paralogs were similar across populations, suggesting that relevant genetic changes act downstream or in parallel to their expression, although upstream factors affecting DELLA protein levels cannot be excluded.

Discussion

Together, these results implicate multiple upstream and downstream changes in both the photoperiod and gibberellin pathways as potential contributors to the observed diversity (Fig. 2f), suggesting that natural variation is maintained throughout the network over sunflower's geographic range. Two aspects of gene regulatory network structure, redundancy and modularity, appear particularly important for the evolution of plasticity. In multiple contexts—e.g. *HaFT4* for day neutrality—paralog-specific expression divergence appears responsible for flowering-time differentiation, indicating that the redundancy afforded by gene duplication makes certain nodes of a gene network more evolutionarily labile. Tissue-specific changes in *SOC1* homolog expression in TX were associated with the transition to long-day response, supporting the idea that modularity in gene networks provided by *cis*- or *trans*-acting factors may foster evolutionary change.

Our understanding of the molecular underpinnings of many plastic traits and the evolutionary conservation of these networks is ever increasing (e.g. germination - Penfield & King 2009; sex-determination - Shoemaker & Crews 2009, Valenzuela 2008; and plant-microbe mutualisms - Heath *et al.* 2010, Kouchi *et al.* 2010). In addition, recent technological innovations now allow gene expression to be measured cost-effectively at tens of loci over many samples from diverse genotypes and treatments (e.g. Wang *et al.* 2011). Therefore, we expect network-informed gene expression approaches to become increasingly more common and productive means of identifying the mechanistic correlates to adaptive natural variation in plasticity.

As noted above, there are several sources of uncertainty with this approach that impact our findings. For instance, while we have assayed expression of all known orthologous copies for many gene families, we cannot exclude the possibility that additional relevant but undiscovered orthologs were not characterized. However, since we find expression patterns correlated with phenotype for one or multiple paralogs in several gene families, and since expression patterns of downstream targets corroborate these findings, we would argue that we have identified multiple relevant readouts and contextualized them appropriately.

The degree to which the flowering time gene regulatory network is conserved between sunflower and other plants is another source of uncertainty. Although the gene content of the gibberellin pathway and the *CO*-dependent and *CO*-independent photoperiod pathways are widely conserved, there are some species-specific differences or unresolved aspects of the flowering time network—e.g. regulatory interactions of *FT*, *SOC1*, and *FUL* in the shoot apex, the role of DELLA transcription in flowering time regulation, etc.—that lead to ambiguity in interpreting our results. In addition, we cannot account for the evolution of regulatory pathways specific to the sunflower lineage, as have been observed for the regulation of flowering in rice and maize (Colasanti *et al.* 1998; Doi *et al.* 2004; Itoh *et al.* 2010). If genetic variation is maintained throughout the network, then differential and combinatorial contributions of various upstream factors across populations could also lead to clines in expression of downstream factors without exhibiting clinal expression patterns.

themselves. Finally, several of the observed clinal gene expression trends are relatively subtle and while worth noting, they must be treated tentatively. Thus, we stress that while our work provides preliminary evidence attributing important evolutionary changes to particular pathways (Fig. 2f) and sufficient data to interpret broad trends, development and application of suitable functional methods will be necessary for confirmation of causal relationships. Nevertheless, we would argue that our conclusion that redundancy and modularity are key properties of gene networks that foster the evolution of plasticity is particularly strongly grounded as it derives solely from findings involving homologs to two broadly conserved downstream floral integrators, *FT* and *SOC1*.

These network properties may also relate to our finding that convergent evolution of long-day response in cultivated and wild sunflower involved different genetic changes. Long-day response evolved independently in the lineage leading to the cultivar CMSHA89, as sunflower was domesticated from short-day populations in Eastern North America (Harter *et al.* 2004). In CMSHA89, changes upstream to *FT* homolog expression in the leaf—rather than the *SOC1* homolog expression changes observed in TX—were associated with the shift to long-day response (Blackman *et al.* 2010). CMSHA89 also flowers earlier and has a higher threshold day length than TX. These phenotypic differences and the distinct mechanisms underlying these short-day to long-day transitions may be consequences of different initial conditions or dissimilar selective pressures during wild *H. annuus*' range expansion southward into Texas versus modern crop improvement.

Our findings add to the growing variety of mechanisms implicated in flowering-time variation in diverse species (Bohlenius *et al.* 2006; Turner *et al.* 2005; Zhang *et al.* 2008), and they strongly contrast with the repeated involvement of the same genes in evolution of similar phenotypes often observed for more canalized developmental and metabolic traits (Stern & Orgogozo 2009). This evolutionary flexibility may arise from the broader multiplicity of regulatory inputs into the network, which could both increase mutational target size and reduce deterministic developmental constraints. Our findings also suggest that for phenotypically plastic traits, stochastic differences in initial genetic and demographic conditions or differences in the balance of environmental factors exerting selection pressures play more pivotal roles than developmental constraint in evolution (Abouheif & Wray 2002).

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by NIH (GM059065) and NSF (DBI0421630) grants to L.H.R.; NSF (IOB0447583) and NIH (GM075060) to S.D.M.; and NSF (DEB0608118, DBI0905958) to B.K.B. B.K.B., S.D.M., and L.H.R. designed the research; B.K.B. performed the research, analyzed the data, and wrote the paper. We thank E. Baack, N. Kane, S.-C. Kim, and the USDA for seed provision; S. Karrenberg, S. Dickinson, E. Houseworth, H. Luton, Z. Lai, D. Rasmussen, K. Turner, A. Posto, L. Moretz, R. Koucouthakis, J. Hvala, A. Saberi, L. Washington, and the Indiana University Greenhouse Staff for statistical and technical assistance; L. Delph and the Indiana University Center for Genomics and Bioinformatics for equipment; and L. Moyle, A. Moczek, M. Lynch, E. Snell-Rood, the Rieseberg, Michaels, and Moyle labs, and three anonymous reviewers for improvements to the manuscript.

References and Notes

- Abouheif E, Wray GA. Evolution of the gene network underlying wing polyphenism in ants. *Science*. 2002; 297:249–252. [PubMed: 12114626]
- Achard P, Herr A, Baulcombe DC, Harberd NP. Modulation of floral development by a gibberellin-regulated microRNA. *Development*. 2004; 131:3357–3365. [PubMed: 15226253]

- Almeida JAS, Pereira MFDA. The control of flower initiation by gibberellin in *Helianthus annuus* L. (sunflower), a non-photoperiodic plant. *Plant Growth Regulation*. 1996; 19:109–115.
- Blackman BK, Rasmussen DA, Strasburg JL, et al. Contributions of flowering time genes to sunflower domestication and improvement. *Genetics*. 2011; 187:271–287. [PubMed: 20944017]
- Blackman BK, Strasburg JL, Raduski AR, Michaels SD, Rieseberg LH. The role of recently derived FT paralogs in sunflower domestication. *Current Biology*. 2010; 20:629–635. [PubMed: 20303265]
- Bohlenius H, Huang T, Charbonnel-Campaa L, et al. CO/FT regulatory module controls timing of flowering and seasonal growth cessation in trees. *Science*. 2006; 312:1040–1043. [PubMed: 16675663]
- Bradshaw WE, Holzapfel CM. Genetic response to rapid climate change: it's seasonal timing that matters. *Molecular Ecology*. 2008; 17:157–166. [PubMed: 17850269]
- Colasanti J, Yuan Z, Sundaresan V. The indeterminate gene encodes a zinc finger protein and regulates a leaf-generated signal required for the transition to flowering in maize. *Cell*. 1998; 93:593–603. [PubMed: 9604934]
- Corbesier L, Vincent C, Jang S, et al. FT protein movement contributes to long-distance signaling in floral induction of *Arabidopsis*. *Science*. 2007; 316:1030–1033. [PubMed: 17446353]
- Curtis MD, Grossniklaus U. A Gateway cloning vector set for high-throughput functional analysis of genes in planta. *Plant Physiology*. 2003; 133:462–469. [PubMed: 14555774]
- Dieringer D, Schlötterer C. Microsatellite analyser (MSA): a platform independent analysis tool for large microsatellite data sets. *Molecular Ecology Notes*. 2003; 3:167–169.
- Doi K, Izawa T, Fuse T, et al. Ehd1, a B-type response regulator in rice, confers short-day promotion of flowering and controls FT-like gene expression independently of Hd1. *Genes & Development*. 2004; 18:926–936. [PubMed: 15078816]
- Efron, B.; Tibshirani, RJ. *An Introduction to the Bootstrap*. New York: Chapman & Hall; 1993.
- Erwin DH, Davidson EH. The evolution of hierarchical gene regulatory networks. *Nature Reviews Genetics*. 2009; 10:141–148.
- Gallego-Bartolomé J, Minguet EG, Marín JA, et al. Transcriptional diversification and functional conservation between DELLA proteins in *Arabidopsis*. *Molecular Biology and Evolution*. 2010; 27:1247–1256. [PubMed: 20093430]
- Goyné PJ, Schneiter AA. Photoperiod influence on development in sunflower genotypes. *Agronomy Journal*. 1987; 79:704–709.
- Harter AV, Gardner KA, Falush D, et al. Origin of extant domesticated sunflowers in eastern North America. *Nature*. 2004; 430:201–205. [PubMed: 15241413]
- Hayama R, Yokoi S, Tamaki S, Yano M, Shimamoto K. Adaptation of photoperiodic control pathways produces short-day flowering in rice. *Nature*. 2003; 422:719–722. [PubMed: 12700762]
- Heath KD, Stock AJ, Stinchcombe JR. Mutualism variation in the nodulation response to nitrate. *Journal of Evolutionary Biology*. 2010; 23:2494–2500. [PubMed: 20825525]
- Heesacker A, Kishore VK, Gao W, et al. SSRs and INDELs mined from the sunflower EST database: abundance, polymorphisms, and cross-taxa utility. *Theoretical and Applied Genetics*. 2008; 117:1021–1029. [PubMed: 18633591]
- Itoh H, Nonoue Y, Yano M, Izawa T. A pair of floral regulators sets critical day length for Hd3a florigen expression in rice. *Nature Genetics*. 2010; 42:635–638. [PubMed: 20543848]
- Kouchi H, Imaizumi-Anraku H, Hayashi M, et al. How many peas in a pod? Legume genes responsible for mutualistic symbioses underground. *Plant and Cell Physiology*. 2010; 51:1381–1397. [PubMed: 20660226]
- Lynch, M.; Walsh, B. *Genetics and Analysis of Quantitative Traits*. Sunderland: Sinauer Associates, Inc.; 1998.
- Maizel A, Busch MA, Tanahashi T, et al. The floral regulator LEAFY evolves by substitutions in the DNA binding domain. *Science*. 2005; 308:260–263. [PubMed: 15821093]
- Moczek AP, Nijhout HF. Rapid evolution of a polyphenic threshold. *Evolution & Development*. 2003; 5:259–268. [PubMed: 12752765]
- O'Hara RB, Merila J. Bias and precision in QST estimates: problems and some solutions. *Genetics*. 2005; 171:1331–1339. [PubMed: 16085700]

- Oh E, Yamaguchi S, Hu J, et al. PIL5, a phytochrome-interacting bHLH Protein, regulates gibberellin responsiveness by binding directly to the GAI and RGA promoters in Arabidopsis seeds. *Plant Cell*. 2007; 19:1192–1208. [PubMed: 17449805]
- Penfield S, King J. Towards a systems biology approach to understanding seed dormancy and germination. *Proceedings of the Royal Society B: Biological Sciences*. 2009; 276:3561–3569.
- Pinheiro, JC.; Bates, DM. *Mixed-Effects Models in S and S-PLUS*. New York: Springer-Verlag; 2000.
- Schlichting, CD.; Pigliucci, M. *Phenotypic Evolution: A Reaction Norm Perspective*. Sunderland, MA: Sinauer; 1998.
- Schneiter AA, Miller JF. Description of sunflower growth stages. *Crop Science*. 1981; 21:901–903.
- Serrano G, Herrera-Palau R, Romero JM, et al. Chlamydomonas CONSTANS and the evolution of plant photoperiodic signaling. *Current Biology*. 2009; 19:359–368. [PubMed: 19230666]
- Shoemaker CM, Crews D. Analyzing the coordinated gene network underlying temperature-dependent sex determination in reptiles. *Seminars in Cell & Developmental Biology*. 2009; 20:293–303. [PubMed: 19022389]
- Spitze K. Population structure in *Daphnia obtusa*: quantitative genetic and allozymic variation. *Genetics*. 1993; 135:367–374. [PubMed: 8244001]
- Stern DL, Orgogozo V. Is genetic evolution predictable? *Science*. 2009; 323:746–751. [PubMed: 19197055]
- Takahashi Y, Teshima KM, Yokoi S, Innan H, Shimamoto K. Variations in Hd1 proteins, Hd3a promoters, and Ehd1 expression levels contribute to diversity of flowering time in cultivated rice. *Proceedings of the National Academy of Sciences, USA*. 2009; 106:4555–4560.
- Tamaki S, Matsuo S, Wong HL, Yokoi S, Shimamoto K. Hd3a protein is a mobile flowering signal in rice. *Science*. 2007; 316:1033–1036. [PubMed: 17446351]
- Turner A, Beales J, Faure S, Dunford RP, Laurie DA. The Pseudo-Response Regulator Ppd-H1 provides adaptation to photoperiod in barley. *Science*. 2005; 310:1031–1034. [PubMed: 16284181]
- Ueguchi-Tanaka M, Ashikari M, Nakajima M, et al. GIBBERELLIN INSENSITIVE DWARF1 encodes a soluble receptor for gibberellin. *Nature*. 2005; 437:693–698. [PubMed: 16193045]
- Valenzuela N. Evolution of the gene network underlying gonadogenesis in turtles with temperature-dependent and genotypic sex determination. *Integrative and Comparative Biology*. 2008; 48:476–485. [PubMed: 21669808]
- Valverde F, Mouradov A, Soppe W, et al. Photoreceptor regulation of CONSTANS protein in photoperiodic flowering. *Science*. 2004; 303:1003–1006. [PubMed: 14963328]
- Wang W, Barnaby JY, Tada Y, et al. Timing of plant immune responses by a central circadian regulator. *Nature*. 2011; 470:110–114. [PubMed: 21293378]
- Whitlock MC. Evolutionary inference from QST. *Molecular Ecology*. 2008; 17:1885–1896. [PubMed: 18363667]
- Wilczek AM, Burghardt LT, Cobb AR, et al. Genetic and physiological bases for phenological responses to current and predicted climates. *Philosophical Transactions of the Royal Society B: Biological Sciences*. 2010; 365:3129–3147.
- Yasumura Y, Crumpton-Taylor M, Fuentes S, Harberd NP. Step-by-step acquisition of the gibberellin-DELLA growth-regulatory mechanism during land-plant evolution. *Current Biology*. 2007; 17:1225–1230. [PubMed: 17627823]
- Zhang Q, Li H, Li R, et al. Association of the circadian rhythmic expression of GmCRY1a with a latitudinal cline in photoperiodic flowering of soybean. *Proceedings of the National Academy of Sciences, USA*. 2008; 105:21028–21033.

Fig. 1. Clinal differentiation of flowering and changes in photoperiod response in wild sunflower. (a) Locations of the populations sampled. (b) Flowering responses to three photoperiod conditions (mean $\pm$ SE). (c) Sunflowers near Norman, OK at peak flowering on September 19, 2006. (d) Senescent sunflowers near Austin, TX on September 21, 2006.

Fig. 2.

Natural variation in flowering is attributable to specific portions of the underlying gene regulatory network. (a, b, c, and d) Relative expression (mean $\pm$ SE) of *HaFT2* (a), *HaFT4* (b), *HaSOC1A* (c), and *HaCOL2* (d) assayed by qRT-PCR in leaves of six wild populations monitored over a diurnal cycle in plants raised in short or long days. Bars along the x-axis illustrate periods of light and dark in each condition relative to time (hours after dawn). Expression was normalized to *Ha60S rRNA* expression (see Materials and Methods). (e) Expression of *HaDELLA2* and *Ha60S rRNA* in leaves under long days. (f) Inferred regulatory network locations of genetic changes contributing to flowering-time variation in sunflower, highlighted in blue. Question marks note regulatory relationships differing

between *A. thaliana* and rice. The wavy arrow represents transport of *FT* from the leaf to the shoot apical meristem (SAM).

Fig. 3. *HaSOC1A* expression (mean $\pm$ SE), assayed by qRT-PCR, in the shoot apex in long days is associated with TX transition to long-day response.

Table 1

Quantitative trait differentiation relative to neutral genetic differentiation. Q_{st} is the proportion of the additive genetic variation for a trait partitioned between populations while F_{st} is proportion of the total genetic variation at 11 microsatellite loci partitioned between populations. Confidence intervals were calculated from 1000 simulations of the data. No confidence interval is provided for ray number due to non-normality of the data (Kolmogorov-Smirnov test, $p < 0.001$).

Trait	Q_{st}	95% CI
Succulence	0.756	0.280–1.000
Days to Anthesis	0.513	0.136–0.736
Days to Budding	0.463	0.106–0.741
Disk Diameter	0.363	0.022–0.669
Bract Number	0.323	0.036–0.594
Ray Number	0.258	
Specific Leaf Area	0.253	0.001–0.645
Ray Shape	0.163	0.000–0.470
Leaf Shape	0.120	0.000–0.386
Bract Shape	0.077	0.000–0.351
Petiole Length	0.025	0.000–0.141
<i>Neutral Markers</i>	<i>0.087 (F_{st})</i>	<i>0.062–0.118</i>