Published in final edited form as:

Curr Stem Cell Res Ther. 2010 September ; 5(3): 268–272.

Clinical Applications of Mesenchymal Stem Cells in Laryngotracheal Reconstruction

Summer Hanson, MD^a, Susan L. Thibeault, PhD^{b,d}, and Peiman Hematti, MD^{c,d}

^aDivision of Plastic and Reconstructive Surgery, University of Wisconsin-Madison, School of Medicine and Public Health, Madison, Wisconsin

^bDivision of Otolaryngology – Head and Neck Surgery, University of Wisconsin-Madison, School of Medicine and Public Health, Madison, Wisconsin

^cDepartment of Medicine, University of Wisconsin-Madison, School of Medicine and Public Health, Madison, Wisconsin

^dUniversity of Wisconsin Carbone Cancer Center, Madison, Wisconsin

Abstract / Summary

During the past several years, mesenchymal stem cells (MSCs) derived from adult tissue have rapidly moved from *in vitro* and animal studies into clinical trials as a therapeutic modality for a diverse group of clinical applications, including head and neck reconstruction. For many diseases, cell therapy could affect the underlying pathophysiologic processes through multiple pathways providing an advantage over current treatment modalities. There is an emerging body of evidence that MSCs have unique immunomodulatory properties in addition to the ability to differentiate into multiple tissue lineages which make them even more attractive for regenerative medicine. A variety of pre-clinical and clinical studies have shown that MSCs may have a useful role in tissue repair as well as engineering strategies in head and neck reconstructive surgery. Clinically, this has ranged from injection laryngoplasty to the implantation of a tracheal construct seeded with MSC-derived chondrocytes. Recent advances in stem cell immunobiology can offer insight to the multiple mechanisms through which MSCs could affect underlying pathophysiologic processes ranging from vocal fold scarring to composite tissue defects. Thorough evaluation of the current literature is necessary in understanding how MSCs could potentially revolutionize our approach to head and neck defects. The purpose of this review is to highlight the advances in MSC-based therapies in head and neck surgery, specifically laryngotracheal reconstruction. The clinical role of tissue-derived MSCs, though not well understood, holds promise for many therapeutic applications in regenerative medicine and reconstruction.

Keywords

mesenchymal stem cells; human vocal fold stem cells; laryngotracheal reconstruction; immune modulation

Introduction

Laryngotracheal disease, from traumatic injury, scarring, tumor resection, neuromuscular disorders or radiation among other causes, can be devastating. The functions maintained by

an intact larynx include swallowing, coughing, secretion clearance, smelling, sniffing, tasting and speech. While there are special considerations associated with pathology along the laryngotracheal tree, the social implications associated with a clinically deficient larynx, from glottic insufficiency to large defects requiring tracheostomy or reconstruction, are immeasurable and debilitating. Much attention has been directed at improving treatment options and outcomes for the many disorders that fall along the spectrum of laryngotracheal dysfunction. Now is the time when we will witness an increase in translation of the basic science advances seen in a variety of animal models to the clinical arena. Showing great clinical promise in the area of tissue reconstruction are initial reports of cell based therapies ranging from injection laryngoplasty to the implantation of a tissue engineered airway. This review serves to highlight advances in cell based therapies for laryngotracheal reconstruction and propose additional applications of mesenchymal stem cells (MSCs) in particular.

Cellular Therapies

Cell therapy involves the transplantation of autologous or allogeneic cells to patients through local delivery or systemic infusion and is now being widely adopted as a novel therapeutic modality for a variety of diseases. For many of these diverse clinical diseases, cell therapy could affect the underlying pathophysiology through multiple pathways providing an advantage over current treatments. This is especially true for diseases in which replacement of the cellular component of the tissue could restore normal physiology or functionality, and is the primary objective of regenerative medicine (1). Although a wide range of cells, from progenitor/stem cells to terminally differentiated cells, could be used for tissue regeneration, stem cells have the advantages of being capable of not only differentiating along multiple tissue lineages but also self-renewing over time, thereby providing new cells to replace those in tissues with high cellular turnover.

During the past several years, MSCs derived from adult tissues have rapidly moved from *in vitro* and animal studies into clinical trials as a therapeutic modality potentially applicable to a wide range of disorders (2). Recent studies show that cells with similar characteristics can be isolated from most adult tissues, including fat (3,4), though these cells were originally isolated from the stromal vascular fraction of adipose tissue more than three decades ago (5). Mesenchymal stem cells have also been derived from embryonic stem cells (6,7). MSCs isolated from different tissue sources all have the capability to differentiate into multiple tissue lineages (8,9) and specific immunomodulatory properties (10,11) though there is evidence to suggest these cells are phenotypically different based on their tissue source (12). While the differentiation potential of MSCs has best been shown in culture-controlled derivation of mesodermal lineages, the degree of *in vivo* contribution of their trans-differentiation is still being debated.

Our group has recently characterized previously isolated human vocal fold fibroblasts (hVFF) derived from adult vocal fold lamina propria, based on the consensus criteria developed to define MSCs (13), and compared them to MSCs derived from adipose tissue (AT) and bone marrow (BM). We have shown that hVFF share similar multipotency and immunophenotype as MSCs (14). From a clinical standpoint, the similarity in BM- and AT-derived MSCs and hVFF, now characterized as VF-MSCs in our recent study, indicates that MSCs derived from different tissue sources could be suitable for novel therapies in vocal fold repair and perhaps laryngotracheal regeneration. Nevertheless, it is quite possible that VF-MSCs will have different gene expression signatures compared to BM and AT; these investigations are currently underway in our lab.

A variety of *in vitro* studies with bone marrow derived MSCs have demonstrated that they possess characteristics that make them a potential commodity for tissue repair including their ability to migrate to injured tissue or inflammation, stimulate resident progenitor cells to proliferate or differentiate, and promote recovery of injured cells through growth factor secretion and matrix remodeling (10,11,15-17). Such properties make these cells very attractive for use in regenerative medicine. Currently, MSCs derived from bone marrow and fat are being actively pursued as an alternative for soft tissue reconstruction or as an augmentation to standard treatment modalities. For example, autologous bone marrow MSCs have been reported for use in topical treatment of non-healing wounds (14,18,19). While this data is limited to case reports or small series of patients they are both novel in their approach and promising in their results. Of particular note is a severe radiation burn treated with a series of injections of culture expanded bone marrow MSCs and split thickness skin grafts (20). Culture expanded autologous adipose tissue derived MSCs were used in combination with a fibrin glue sealant in complex perianal fistula tracts (21,22). The phase I clinical trial was the first to show safety of culture expanded MSCs derived from lipoaspirate (21) and the follow up study showed a healing rate of 71% (22). Additionally, cells isolated from lipoaspirate have been used to augment fat grafting in facial recontouring (23), though this is a heterogeneous mixture containing other cells, cytokines and growth factors.

Moreover, both BM- and AT- derived MSCs have shown the ability to modulate the activation and proliferation of immune cells such as T lymphocytes, B lymphocytes, natural killer (NK) cells and dendritic cells (24-28). Such properties have been the basis of using allogeneic MSCs without HLA matching in a variety of clinical applications such as graft versus host disease (7). Furthermore, allogeneic MSCs are currently being investigated for their potential efficacy for treatment of disorders such as myocardial infarction, newly diagnosed diabetes mellitus, Crohn's disease, chronic obstructive pulmonary disease, and amyotrophic lateral sclerosis. While it is not clear for which of these disorders MSCs will provide a therapeutic advantage, it is expected that use of tissue-derived MSCs will only expand in regenerative medicine and tissue engineering.

Airway Defects of the Larynx and Trachea

Millions of reconstructive surgical procedures are performed each year to address congenital anomalies, traumatic defects, tumor resection and cosmetic augmentation (29). Head and neck surgeons are faced with the additional challenge of complex tissue composition and unique functional requirements associated with the larynx, trachea, and vocal folds. The current methods of reconstruction available in head and neck surgery are not without limitations. Vocal folds are composed of a combination of extracellular matrix (ECM) components that account for the unique biomechanical characteristics not adequately matched by current replacement modalities (30-34). Furthermore, loss of any portion of the laryngotracheal tree is no small deficit. While the mainstays of laryngotracheal management and reconstruction, namely tracheostomy, autogenous tissue transfer and stenting, have provided patients with reliable outcomes, these methods do not offer functional restoration or improved quality of life. To this end, the first attempt at replacing the larynx with a partial transplantation in a patient with cancer was performed in 1969 and associated with both recurrence and poor recovery (35). Nearly 20 years later, a report of laryngeal transplantation for a traumatic injury showed near-total functional recovery of not only speech and swallowing but taste and smell as well (36). An additional 13 cases of laryngeal transplantation from Columbia have been described (37) though long-term results remain to be reported. Although the operations are complex, the recovery is challenging, and the long-term immunosuppression is not without risk, subsequently, composite laryngeal

transplantation may be the best option for functional recovery with enhanced quality of life (38).

As an alternative method of airway replacement, tissue engineering strategies have been explored more recently in select clinical scenarios (39,40). While previous reports have illustrated successful, clinically useful tissue-engineered human organs elsewhere, including bladder (41) and skin (42), the airway defect represents a complex construct with mechanical and functional demands not addressed until this time. First reported was a tissue engineered patch made from autologous cells and a collagen matrix (39). The patient had recurrent non-small cell lung cancer who was initially treated with lung resection and radiation, and later completion carinal pneumonectomy. However, he returned with breakdown of his tracheal anastomosis, aspiration pneumonia and empyema. During subsequent procedures to treat his pulmonary infection, muscle and skin biopsies were taken and muscle cells and fibroblasts were isolated and expanded to three passages. These cells were then seeded on a collagen matrix derived from decellularized porcine jejunum and cultured in a bioreactor for an additional three weeks while functional tissue was formed. The engineered segment was implanted over the dehissed tracheal segment and the pleural space was filled with omentum and subscapular flaps. According to the authors, postoperative endoscopy shows an airtight graft and respiratory endothelium on the surface. Moreover, the patient's vocal quality returned and he was able to perform physical activity at 24 weeks follow up.

More recently, Macchiarini and colleagues published their results with the transplantation of a tissue-engineered tracheal segment, the first of its kind (40) in a young woman with bronchomalacia and expiratory collapse refractory to standard treatment and reconstruction. Autologous bronchial epithelial cells were isolated from bronchoscopic mucosal biopsies and expanded in supplemented media, and autologous chondrocytes were derived from bone marrow MSCs treated with chondrogenic media. The scaffold was a decellularized allograft from a cadaveric trachea. It was seeded with the epithelial cells on the inner surface and the chondrocytes on the outer surface, and maintained in a bioreactor to introduce biomechanical cues that would need to be tolerated by the engineered construct. This *ex vivo* generated graft was then implanted end-to-end to normal tracheal tissue to replace the stenotic segment. According to the authors, the lung immediately ventilated well and subsequent evaluations showed no indication of inflammation related to the donor trachea, normal pulmonary function tests and an immeasurable improvement in quality of life. Both reports provide novel solutions for serious clinical disorders and usher in a new era in regenerative medicine.

Vocal Fold Scarring and Glottic Insufficiency

Voice disorders affect 3-9% Americans yearly and the most serious is attributed to scar of the vocal fold (43). Vocal fold scarring causes a deformity of the vocal fold edge, a disruption of the viscoelastic layered structure of the lamina propria and increase in stiffness of the vibratory structure and glottic incompetence (34). Many technological advances have renewed interest in injection laryngoplasty and vocal fold regeneration. Originally described in the early 20th century, early techniques of injection laryngoplasty using synthetic materials, such as paraffin and polytetrafluoroethylene, or biologic materials, such as bone dust, were limited by inflammation, migration or extrusion of the material, or worsening scar (44). Dermal fillers are now being used for unilateral vocal fold paralysis, including collagen (45), calcium hydroxylapatite (46), hyaluronic acid (Restylane) (47), decellularized dermal matrix (48,49) and autologous tissues (50-52). Autologous fat injection is of interest in laryngoplasty as it potentially takes advantage of the undifferentiated MSC population we now know is present in adipose tissue (53,54). First reported in 1991, intracordal fat

injection (55) has been clinically useful though the amount of adipose resorption is unpredictable. While there are advantages to using autologous fat for head and neck reconstruction, as with other autologous tissue grafts, the technique is limited by mechanical damage during harvest, cyst formation, and localized necrosis (56).

For these reasons, long-term outcomes of fat injection laryngoplasty have varied in the literature. Dursun and colleagues compared patients treated with medialization surgery using a synthetic implant with those treated with injection augmentation of either autologous fat or synthetic calcium hydroxylapatite (57). While the data does not distinguish among the two injectable treatments compared to the silastic implant, the authors do comment that voice improvement has not remained stable in follow up evaluation. Others however, have reported improvement of vocal function from 50-100% in long term follow up (51,54,56,58). Furthermore, Lo Cicero et al. (54) expanded MSCs from adipose samples used in laryngoplasty, illustrating the tissue harvest technique does not impair the MSC population. One can anticipate that a next step in the progression of injection laryngoplasty be either direct injection of MSCs or in combination with ECM proteins already being used for injection laryngoplasty (collagen, hyaluronic acid, hydroxylapatite) as an engineered construct to serve as a filler and facilitate vocal fold lamina propria regeneration, though this has only been reported in animals (50,59-63). Current animal models include mice, rabbits, pigs and canines (34). The development of such a treatment strategy that could be used to minimize scarring and improve tissue repair would have far reaching implications and applicability.

Discussion

Emerging clinical data on the safety of bone marrow-derived MSCs, as well as adipose-derived MSCs, in a variety of inflammatory or immune system disorders as well as their use in wound healing warrants considering the use of these cells in a wider range of applications, including laryngotracheal reconstruction. There are several prospective roles MSCs could assume in treatment of the disorders approached above. These multipotent progenitor cells will likely be the cell of choice, whether from BM, AT or other head and neck specific tissues, for tissue engineering strategies to replace a segment of trachea or augment a scarred vocal fold.

Many recent studies confirm that MSCs from different tissues share similar differentiation potential and immunological properties (27,28) though their gene expression may be different. Thus, there is increasing interest in using adipose-derived MSCs, instead of bone marrow-derived MSCs, in clinical studies for treatment of complications, such as graft versus host disease after bone marrow transplantation or complex cutaneous wound healing. It is possible, however, that the tissue source used to generate MSCs could have an impact on the outcome based on the applications used (12). The MSCs our group has derived from vocal folds may provide insight to the potential advantages of MSCs derived from one tissue source over another for repair of vocal fold maladies and other head and neck reconstruction (14). For example, just as ease of expansion, differentiation potential and immunological characteristics similar to bone marrow MSCs makes adipose-derived MSCs appealing in many clinical applications, perhaps these shared functional and phenotypic qualities of vocal fold MSCs will make them useful in tissue specific repair or reconstruction. Given that these cells reside within the unique environmental and functional niche of the laryngotracheal tree and demonstrate analogous immunobiology of the more well-characterized MSCs, further study into the possible clinical applications of laryngotracheal derived MSCs is certainly warranted.

There are several important questions that remain unanswered in using MSCs in such a variety of clinical applications. Not all reports are optimistic. When considering MSCs for novel cell-based therapies it is necessary to address migration, engraftment, and tumorigenicity. The exact fate of locally or systemically delivered MSCs has not been clearly established. Many pre-clinical and clinical studies show variations in engraftment of MSCs in a variety of tissues (64) potentially limiting the anti-inflammatory or regenerative effect. It is thought that sites of injury or pathology elicit enhanced MSC migration through a variety of inflammatory markers (65). While this is advantageous in a wound healing model, similar inflammatory markers could promote MSCs to a tumor bed. Moreover, it has been speculated that bone marrow derived MSCs are a source of tumor-associated stromal cells and, as such, can enhance tumor growth (65,66). Further understanding of not only how MSCs can effect the underlying pathophysiology of head and neck specific disorders but how they in turn are affected by the wounded or diseased tissues of interest are needed to clarify the clinical utility of MSCs in laryngotracheal repair and reconstruction.

In cases where both form and function are more complex, as in laryngeal reconstruction, perhaps tissue transplantation is the best option available; there is evidence to suggest that MSCs may be used in conjunction with such methods to augment healing and allograft survival and mediate the inflammatory response (7). A thorough understanding of the immunobiology of mesenchymal stem cells is necessary to realize the complement of head and neck pathology that could be affected by MSC-based therapy. The novel methods reviewed are highly promising, with the collective goal of identifying new therapeutic approaches to head and neck healing that are broadly applicable to many chronic diseases, and can safely accelerate the transition of basic research findings into clinical advances in many areas of regenerative medicine and reconstructive surgery.

Acknowledgments

Source of Funding: NIDCD - NIH grant R01 DC4336; NHLBI – NIH grant K08 HL081076; NCI – NIH grant T32 CA009614

References

- (1). Mason C, Dunnill P. A brief definition of regenerative medicine. *Regen Med.* Jan; 2008 3(1):1–5. [PubMed: 18154457]
- (2). Giordano A, Galderisi U, Marino IR. From the laboratory bench to the patient's bedside: an update on clinical trials with mesenchymal stem cells. *J Cell Physiol.* Apr; 2007 211(1):27–35. [PubMed: 17226788]
- (3). Zuk PA, Zhu M, Mizuno H, Huang J, Futrell JW, Katz AJ, et al. Multilineage cells from human adipose tissue: implications for cell-based therapies. *Tissue Eng.* Apr; 2001 7(2):211–228. [PubMed: 11304456]
- (4). Beahm EK, Walton RL, Patrick CW Jr. Progress in adipose tissue construct development. *Clin Plast Surg.* Oct; 2003 30(4):547–58. viii. [PubMed: 14621302]
- (5). Van RL, Bayliss CE, Roncari DA. Cytological and enzymological characterization of adult human adipocyte precursors in culture. *J Clin Invest.* Sep; 1976 58(3):699–704. [PubMed: 956396]
- (6). Trivedi P, Hematti P. Derivation and immunological characterization of mesenchymal stromal cells from human embryonic stem cells. *Exp Hematol.* Mar; 2008 36(3):350–359. [PubMed: 18179856]
- (7). Hematti P. Role of mesenchymal stromal cells in solid organ transplantation. *Transplant Rev (Orlando).* Oct; 2008 22(4):262–273. [PubMed: 18656340]
- (8). Caplan AI. Mesenchymal stem cells. *J Orthop Res.* Sep; 1991 9(5):641–650. [PubMed: 1870029]
- (9). da Silva Meirelles L, Chagastelles PC, Nardi NB. Mesenchymal stem cells reside in virtually all post-natal organs and tissues. *J Cell Sci.* Jun 1; 2006 119(Pt 11):2204–2213. [PubMed: 16684817]

- (10). Caplan AI. Adult mesenchymal stem cells for tissue engineering versus regenerative medicine. *J Cell Physiol.* Nov; 2007 213(2):341–347. [PubMed: 17620285]
- (11). Uccelli A, Pistoia V, Moretta L. Mesenchymal stem cells: a new strategy for immunosuppression? *Trends Immunol.* May; 2007 28(5):219–226. [PubMed: 17400510]
- (12). Noel D, Caton D, Roche S, Bony C, Lehmann S, Casteilla L, et al. Cell specific differences between human adipose-derived and mesenchymal-stromal cells despite similar differentiation potentials. *Exp Cell Res.* Apr 15; 2008 314(7):1575–1584. [PubMed: 18325494]
- (13). Dominici M, Le Blanc K, Mueller I, Slaper-Cortenbach I, Marini F, Krause D, et al. Minimal criteria for defining multipotent mesenchymal stromal cells. The International Society for Cellular Therapy position statement. *Cytotherapy.* 2006; 8(4):315–317. [PubMed: 16923606]
- (14). Hanson SE, Kim J, Johnson BH, Bradley B, Breunig MJ, Hematti P, et al. Characterization of mesenchymal stem cells from human vocal fold fibroblasts. *Laryngoscope.* Mar; 2010 120(3): 546–551. [PubMed: 20131365]
- (15). Phinney DG, Prockop DJ. Concise review: mesenchymal stem/multipotent stromal cells: the state of transdifferentiation and modes of tissue repair--current views. *Stem Cells.* Nov; 2007 25(11): 2896–2902. [PubMed: 17901396]
- (16). Chamberlain G, Fox J, Ashton B, Middleton J. Concise review: mesenchymal stem cells: their phenotype, differentiation capacity, immunological features, and potential for homing. *Stem Cells.* Nov; 2007 25(11):2739–2749. [PubMed: 17656645]
- (17). Dazzi F, Horwood NJ. Potential of mesenchymal stem cell therapy. *Curr Opin Oncol.* Nov; 2007 19(6):650–655. [PubMed: 17906466]
- (18). Falanga V, Iwamoto S, Chartier M, Yufit T, Butmarc J, Kouttab N, et al. Autologous bone marrow-derived cultured mesenchymal stem cells delivered in a fibrin spray accelerate healing in murine and human cutaneous wounds. *Tissue Eng.* Jun; 2007 13(6):1299–1312. [PubMed: 17518741]
- (19). Yoshikawa T, Mitsuno H, Nonaka I, Sen Y, Kawanishi K, Inada Y, et al. Wound therapy by marrow mesenchymal cell transplantation. *Plast Reconstr Surg.* Mar; 2008 121(3):860–877. [PubMed: 18317135]
- (20). Lataillade JJ, Doucet C, Bey E, Carsin H, Huet C, Clairand I, et al. New approach to radiation burn treatment by dosimetry-guided surgery combined with autologous mesenchymal stem cell therapy. *Regen Med.* Sep; 2007 2(5):785–794. [PubMed: 17907931]
- (21). Garcia-Olmo D, Garcia-Arranz M, Herreros D, Pascual I, Peiro C, Rodriguez-Montes JA. A phase I clinical trial of the treatment of Crohn's fistula by adipose mesenchymal stem cell transplantation. *Dis Colon Rectum.* Jul; 2005 48(7):1416–1423. [PubMed: 15933795]
- (22). Garcia-Olmo D, Herreros D, Pascual I, Pascual JA, Del-Valle E, Zorrilla J, et al. Expanded adipose-derived stem cells for the treatment of complex perianal fistula: a phase II clinical trial. *Dis Colon Rectum.* Jan; 2009 52(1):79–86. [PubMed: 19273960]
- (23). Yoshimura K, Sato K, Aoi N, Kurita M, Inoue K, Suga H, et al. Cell-Assisted Lipotransfer for Facial Lipoatrophy: Efficacy of Clinical Use of Adipose-Derived Stem Cells. *Dermatol Surg.* May 28, 2008
- (24). Le Blanc K, Ringden O. Immunomodulation by mesenchymal stem cells and clinical experience. *J Intern Med.* Nov; 2007 262(5):509–525. [PubMed: 17949362]
- (25). Le Blanc K, Samuelsson H, Gustafsson B, Remberger M, Sundberg B, Arvidson J, et al. Transplantation of mesenchymal stem cells to enhance engraftment of hematopoietic stem cells. *Leukemia.* Aug; 2007 21(8):1733–1738. [PubMed: 17541394]
- (26). Nauta AJ, Fibbe WE. Immunomodulatory properties of mesenchymal stromal cells. *Blood.* Nov 15; 2007 110(10):3499–3506. [PubMed: 17664353]
- (27). Puissant B, Barreau C, Bourin P, Clavel C, Corre J, Bousquet C, et al. Immunomodulatory effect of human adipose tissue-derived adult stem cells: comparison with bone marrow mesenchymal stem cells. *Br J Haematol.* Apr; 2005 129(1):118–129. [PubMed: 15801964]
- (28). Tholpady SS, Ogle RC, Katz AJ. Adipose stem cells and solid organ transplantation. *Curr Opin Organ Transplant.* Feb; 2009 14(1):51–55. [PubMed: 19337147]
- (29). American Society of Plastic Surgeons. National Plastic Surgery Statistics. 2006. 2006; Available at: www.plasticsurgery.org/media/statistics.

- (30). Titze IR. The physics of small-amplitude oscillation of the vocal folds. *J Acoust Soc Am.* Apr; 1988 83(4):1536–1552. [PubMed: 3372869]
- (31). Gray SD, Duflo S, Thibeault SL, Li W, Shu XZ, Prestwich G. Cellular physiology of the vocal folds. Effect of a synthetic extracellular matrix on vocal fold lamina propria gene expression in early wound healing. *Otolaryngol Clin North Am. Tissue Eng.* Aug; Nov; 2000 2006 3312(4) (11):679–698. [PubMed: 10918654]
- (32). Duflo S, Thibeault SL, Li W, Shu XZ, Prestwich GD. Vocal fold tissue repair in vivo using a synthetic extracellular matrix. *Tissue Eng.* Aug; 2006 12(8):2171–2180. [PubMed: 16968158]
- (33). Hansen JK, Thibeault SL. Current understanding and review of the literature: vocal fold scarring. *J Voice.* Mar; 2006 20(1):110–120. [PubMed: 15964741]
- (34). Kluyskens P, Ringoir S. Follow-up of a human larynx transplantation. *Laryngoscope.* Aug; 1970 80(8):1244–1250. [PubMed: 5469227]
- (35). Strome M, Stein J, Esclamado R, Hicks D, Lorenz RR, Braun W, et al. Laryngeal transplantation and 40-month follow-up. *N Engl J Med.* May 31; 2001 344(22):1676–1679. [PubMed: 11386266]
- (36). Duque E, Duque J, Nieves M, Mejia G, Lopez B, Tintinago L. Management of larynx and trachea donors. *Transplant Proc.* Sep; 2007 39(7):2076–2078. [PubMed: 17889100]
- (37). Birchall M. Human laryngeal allograft: shift of emphasis in transplantation. *Lancet.* Feb 21; 1998 351(9102):539–540. [PubMed: 9492768]
- (38). Macchiarini P, Walles T, Biancosino C, Mertsching H. First human transplantation of a bioengineered airway tissue. *J Thorac Cardiovasc Surg.* Oct; 2004 128(4):638–641. [PubMed: 15457176]
- (39). Macchiarini P, Jungebluth P, Go T, Asnaghi MA, Rees LE, Cogan TA, et al. Clinical transplantation of a tissue-engineered airway. *Lancet.* Dec 13; 2008 372(9655):2023–2030. [PubMed: 19022496]
- (40). Atala A, Bauer SB, Soker S, Yoo JJ, Retik AB. Tissue-engineered autologous bladders for patients needing cystoplasty. *Lancet.* Apr 15; 2006 367(9518):1241–1246. [PubMed: 16631879]
- (41). Pham C, Greenwood J, Cleland H, Woodruff P, Maddern G. Bioengineered skin substitutes for the management of burns: a systematic review. *Burns.* Dec; 2007 33(8):946–957. [PubMed: 17825993]
- (42). Woo P, Casper J, Colton R, Brewer D. Diagnosis and treatment of persistent dysphonia after laryngeal surgery: a retrospective analysis of 62 patients. *Laryngoscope.* Sep; 1994 104(9):1084–1091. [PubMed: 8072354]
- (43). O'Leary MA, Grillone GA. Injection laryngoplasty. *Otolaryngol Clin North Am.* Feb; 2006 39(1):43–54. [PubMed: 16469654]
- (44). Kwon TK, Buckmire R. Injection laryngoplasty for management of unilateral vocal fold paralysis. *Curr Opin Otolaryngol Head Neck Surg.* Dec; 2004 12(6):538–542. [PubMed: 15548914]
- (45). Rosen CA, Thekdi AA. Vocal fold augmentation with injectable calcium hydroxylapatite: short-term results. *J Voice.* Sep; 2004 18(3):387–391. [PubMed: 15331112]
- (46). Hirano S, Bless DM, Heisey D, Ford CN. Effect of growth factors on hyaluronan production by canine vocal fold fibroblasts. *Ann Otol Rhinol Laryngol.* Jul; 2003 112(7):617–624. [PubMed: 12903682]
- (47). Karpenko AN, Dworkin JP, Meleca RJ, Stachler RJ. Cymetra injection for unilateral vocal fold paralysis. *Ann Otol Rhinol Laryngol.* Nov; 2003 112(11):927–934. [PubMed: 14653360]
- (48). Lundy DS, Casiano RR, McClinton ME, Xue JW. Early results of transcutaneous injection laryngoplasty with micronized acellular dermis versus type-I thyroplasty for glottic incompetence dysphonia due to unilateral vocal fold paralysis. *J Voice.* Dec; 2003 17(4):589–595. [PubMed: 14740939]
- (49). Glatz FR, Kalkanis J, Neumeister M, Suchy H, Lyons S, Mowlavi A. Volume analysis of preadipocyte injection for vocal cord medialization. *Laryngoscope.* Jul; 2003 113(7):1113–1117. [PubMed: 12838006]

- (51). Laccourreye O, Papon JF, Kania R, Crevier-Buchman L, Brasnu D, Hans S. Intracordal injection of autologous fat in patients with unilateral laryngeal nerve paralysis: long-term results from the patient's perspective. *Laryngoscope*. Mar; 2003 113(3):541–545. [PubMed: 12616211]
- (52). Rihkanen H. Vocal fold augmentation by injection of autologous fascia. *Laryngoscope*. Jan; 1998 108(1 Pt 1):51–54. [PubMed: 9432067]
- (53). Moseley TA, Zhu M, Hedrick MH. Adipose-derived stem and progenitor cells as fillers in plastic and reconstructive surgery. *Plast Reconstr Surg*. Sep; 2006 118(3 Suppl):121S–128S. [PubMed: 16936551]
- (54). Lo Cicero V, Montelatici E, Cantarella G, Mazzola R, Sambataro G, Rebulli P, et al. Do mesenchymal stem cells play a role in vocal fold fat graft survival? *Cell Prolif*. Jun; 2008 41(3): 460–473. [PubMed: 18435791]
- (55). Mikaelian DO, Lowry LD, Sataloff RT. Lipoinjection for unilateral vocal cord paralysis. *Laryngoscope*. May; 1991 101(5):465–468. [PubMed: 1827654]
- (56). Shindo ML, Zaretsky LS, Rice DH. Autologous fat injection for unilateral vocal fold paralysis. *Ann Otol Rhinol Laryngol*. Aug; 1996 105(8):602–606. [PubMed: 8712629]
- (57). Dursun G, Boynukalin S, Ozgursoy OB, Coruh I. Long-term results of different treatment modalities for glottic insufficiency. *Am J Otolaryngol*. Jan-Feb; 2008 29(1):7–12. [PubMed: 18061825]
- (58). Hsiung MW, Woo P, Minasian A, Mojica J Schaefer. Fat augmentation for glottic insufficiency. *Laryngoscope*. Jun; 2000 110(6):1026–1033. [PubMed: 10852525]
- (59). Kanemaru S, Nakamura T, Omori K, Kojima H, Magrufov A, Hiratsuka Y, et al. Regeneration of the vocal fold using autologous mesenchymal stem cells. *Ann Otol Rhinol Laryngol*. Nov; 2003 112(11):915–920. [PubMed: 14653358]
- (60). Kanemaru S, Nakamura T, Yamashita M, Magrufov A, Kita T, Tamaki H, et al. Destiny of autologous bone marrow-derived stromal cells implanted in the vocal fold. *Ann Otol Rhinol Laryngol*. Dec; 2005 114(12):907–912. [PubMed: 16425555]
- (61). Hertegard S, Cedervall J, Svensson B, Forsberg K, Maurer FH, Vidovska D, et al. Viscoelastic and histologic properties in scarred rabbit vocal folds after mesenchymal stem cell injection. *Laryngoscope*. Jul; 2006 116(7):1248–1254. [PubMed: 16826069]
- (62). Thibeault SL, Klemuk SA, Smith ME, Leugers C, Prestwich G. In Vivo Comparison of Biomimetic Approaches for Tissue Regeneration of the Scarred Vocal Fold. *Tissue Eng Part A*. Dec 10.2008
- (63). Quinchia-Johnson BH, Fox R, Chen X, Thibeault SL. Tissue regeneration of the vocal fold using bone marrow mesenchymal stem cells and synthetic extracellular matrix injections in rats. *Laryngoscope* In Review.
- (64). Laurila JP, Laatikainen L, Castellone MD, Trivedi P, Heikkila J, Hinkkanen A, et al. Human embryonic stem cell-derived mesenchymal stromal cell transplantation in a rat hind limb injury model. *Cytotherapy*. 2009; 11(6):726–737. [PubMed: 19878059]
- (65). Kidd S, Spaeth E, Klopp A, Andreeff M, Hall B, Marini FC. The (in) auspicious role of mesenchymal stromal cells in cancer: be it friend or foe. *Cytotherapy*. 2008; 10(7):657–667. [PubMed: 18985472]
- (66). Mishra PJ, Mishra PJ, Glod JW, Banerjee D. Mesenchymal stem cells: flip side of the coin. *Cancer Res*. Feb 15; 2009 69(4):1255–1258. [PubMed: 19208837]