Reminder of important clinical lesson

A renal transplant patient with abdominal discomfort, vomiting and diarrhoea for 1 week

Nancy Lutwak,¹ Curt Dill²¹Department of Emergency Services New York, VA New York Harbor Healthcare Center, New York, United States;²Department of Emergency Medicine New York, VA New York Harbor Healthcare Center, NYU School of Medicine, New York, United States**Correspondence to** Dr Nancy Lutwak, nancy.lutwak@gmail.com**Summary**

The patient is a 61-year-old diabetic male with history of renal transplant who presented to the emergency department with complaints of intermittent abdominal discomfort accompanied by multiple episodes of vomiting and diarrhoea. He had delayed seeking medical attention until his friends insisted that he come to the emergency department, since the abdominal discomfort was worsening. The patient's ECG revealed an ST-segment elevation myocardial infarction.

BACKGROUND

Patients with diabetes and renal disease are at great risk for cardiovascular events. It is now recognised that even moderate renal disease is an independent risk factor. Patients with diabetes and end stage renal disease are in the highest risk group for death from cardiac causes. Nevertheless, there have been multiple recent publications demonstrating that patients with renal disease, who frequently have multiple co-morbidities including diabetes, are less likely to receive evidence-based procedures and therapies in the setting of an acute myocardial infarction.

CASE PRESENTATION

The patient is a 61-year-old male with history of insulin dependent diabetes and renal transplant 5 years earlier who presented to the emergency department with complaints of abdominal discomfort and several episodes of

vomiting and diarrhoea. He noted that this started 1 week prior to arrival and had begun with two short episodes of a choking sensation in his throat and shortness of breath. He denied chest pain, chest tightness or back pain. The patient was pain-free in the emergency department.

Medical history is significant for insulin dependent diabetes, renal transplant, hypertension and cerebral vascular accident. He denied angina or myocardial infarction in the past.

Medications included furosemide 40 mg daily, tacrolimus 5 mg daily, prednisone 5 mg daily, atenolol 100 mg daily, insulin glargine 25 units daily, mycophenolate mofetil 200 mg daily, lisinopril 5 mg daily and aspirin 81 mg daily. The patient denied cigarettes, use of illicit drugs and drinking alcohol. He is retired and lives alone.

On arrival to the emergency department, the patient was calm, comfortable and in no distress. Blood pressure

Figure 1 ECG of renal transplant patient demonstrating 1 mm ST-segment elevations in leads II, III, AVF and V6.

Figure 2 Previous ECG of same patient.

was 115/73, pulse 72, respiratory rate 18 and temperature was 96.3 degrees F. Oxygen saturation was 100% on room air. Physical examination was unremarkable except for obesity.

INVESTIGATIONS

The ECG revealed normal sinus rhythm at rate of 64, with 1 mm ST elevations in leads II, III, AVF and V6 (figure 1). The patient's previous ECG (figure 2) was normal. Chest radiograph was negative. Laboratory tests revealed troponin-T point-of-care (Biosite) of 7.61 ng/ml (negative less than 0.05), troponin ultra of 11.14 ng/ml (negative <0.7 ng/ml, positive >78 ng/ml), blood urea nitrogen of 56 mg/dl (nl 6–22), creatinine of 2.9 mg/dl (nl 4–1.2), glucose of 160 mg/dl (nl 65–115), haemoglobin of 10.1 g/dl (nl 13–18). Other values were normal.

TREATMENT

The patient was treated with clopidogrel, aspirin, nitroglycerin, metoprolol and heparin in the emergency department. It was decided to treat the patient conservatively since he was asymptomatic in the emergency department, presented late and had chronic renal insufficiency post-transplant. The patient was admitted to the cardiac care unit. Echocardiogram revealed minimal inferolateral hypokinesis with a preserved normal ejection fraction.

OUTCOME AND FOLLOW-UP

The patient has been doing well and is followed in cardiology clinic.

DISCUSSION

Mild chronic kidney disease is a coronary risk factor. Even moderate renal disease is a predictor of myocardial

infarction.¹ There seems to be 'renal-specific' factors that cause increasing levels of cardiovascular disease.² The incidence and severity of cardiovascular events in patients with renal disease is not attributable solely to the traditional risk factors of hypertension, hyperlipidemia, diabetes and smoking. Vascular calcification is a pathogenesis being investigated. Medial artery calcification is very common in patients with renal disease and is associated with high mortality. It leads to development of left ventricular hypertrophy, myocardial ischemia, worsening of coronary artery perfusion and arterial inelasticity.³ Promoters of this calcification in dialysis patients include uremic toxins, inflammatory cytokines, tumour necrosis factor α and altered calcium/phosphate metabolism.⁴

A recent review of patients with ST-segment elevation myocardial infarction and non-ST-segment elevation myocardial infarction revealed that patients with the most severe chronic renal disease had highest rates of death. Regardless of the higher mortality of this group of patients, they are less likely to be given thrombolytics, aspirin, intravenous heparin and β blockers immediately for acute myocardial infarction.^{5 6} Patients with renal disease who had myocardial infarctions were also less likely to have discharge counselling regarding smoking cessation, diet and exercise.⁵ Patients with renal disease definitely need secondary prevention strategies.⁷

Furthermore, patients with chronic kidney disease are infrequently prescribed medications to reduce the risk of cardiovascular events. Pravastatin reduces the risk of cardiovascular events in patients with chronic mild to moderate kidney disease.⁸ ACE inhibitors, as well as statins, are recommended for patients with renal disease to reduce the risk of cardiovascular events.⁹ Physicians' awareness of these treatment patterns may prove of benefit.

Learning points

- ▶ Diabetic patients often present atypically when having a myocardial infarction.
- ▶ Renal insufficiency is now recognised as an independent risk factor for myocardial infarction.
- ▶ Patients with renal disease are less likely to receive adjunctive treatment for acute myocardial infarction despite their high mortality.
- ▶ Patients with renal disease are less likely to be discharged on clopidogrel, β -blockers and statins after a myocardial infarction.
- ▶ Patients with renal disease are less likely to receive discharge counselling regarding diet, exercise and smoking cessation after a myocardial infarction.

Competing interests None.

Patient consent Obtained.

REFERENCES

1. **Williams ME.** Coronary revascularization in diabetic chronic kidney disease/end-stage renal disease: a nephrologist's perspective. *Clin J Am Soc Nephrol* 2006;**1**:209–20.
2. **Bhandari S.** Risk factors and metabolic mechanisms in the pathogenesis of uraemic cardiac disease. *Front Biosci* 2011;**16**:1364–87.
3. **Mizobuchi M,** Towler D, Slatopolsky E. Vascular calcification: the killer of patients with chronic kidney disease. *J Am Soc Nephrol* 2009;**20**:1453–64.
4. **Checherita IA,** David C, Stoica L, *et al.* New mediators of vascular damage in dialysed patients. *Rom J Morphol Embryol* 2011;**52**:533–6.
5. **Fox CS,** Muntner P, Chen AY, *et al.* Use of evidence-based therapies in short-term outcomes of ST-segment elevation myocardial infarction and non-ST-segment elevation myocardial infarction in patients with chronic kidney disease: a report from the National Cardiovascular Data Acute Coronary Treatment and Intervention Outcomes Network registry. *Circulation* 2010;**121**:357–65.
6. **Wright RS,** Reeder GS, Herzog CA, *et al.* Acute myocardial infarction and renal dysfunction: a high-risk combination. *Ann Intern Med* 2002;**137**:563–70.
7. **Fort J.** Chronic renal failure: a cardiovascular risk factor. *Kidney Int Suppl* 2005;**99**:S25–9.
8. **Tonelli M,** Isles C, Curhan GC, *et al.* Effect of pravastatin on cardiovascular events in people with chronic kidney disease. *Circulation* 2004;**110**:1557–63.
9. **Lourengo C,** Teixeira R, António N, *et al.* Impact of renal function on mortality and incidence of major adverse cardiovascular events following acute coronary syndromes. *Rev Port Cardiol* 2010;**29**:1331–52.

This pdf has been created automatically from the final edited text and images.

Copyright 2011 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Please cite this article as follows (you will need to access the article online to obtain the date of publication).

Lutwak N, Dill C. A renal transplant patient with abdominal discomfort, vomiting and diarrhoea for 1 week. *BMJ Case Reports* 2011; 10.1136/bcr.07.2011.4529, date of publication

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow