Images in...

Sister Mary Joseph's nodule

Javier Munoz,¹ Wadih Chacra,² Amr Hanbali,¹ Haythem Ali¹

¹Hematology-Oncology Department, Henry Ford Hospital, Detroit, Michigan, USA;

²Department of Internal Medicine, Henry Ford Hospital, Detroit, Michigan, USA

Correspondence to Dr Javier Munoz, jlmunoz@mdanderson.org

DESCRIPTION

A 44-year-old female presented with 1-year history of progressive abdominal distension (figure 1A) which she thought might be related to pregnancy and a tumour protruding through her umbilical area that was oozing serosanguinous fluid. The patient did not seek medical attention initially due to lack of insurance. CT of the abdomen (figure 1B,C) revealed a large mass with loculated ascites. Multiple lung nodules were identified as well. The patient underwent tumour debulking that revealed a poorly

differentiated malignant neoplasm (figure 1D) involving ovaries, the umbilical area and widespread omental metastases. Postoperatively the patient continued to produce massive amounts of ascites and ultimately succumbed due to sepsis.

Sister Mary Joseph's nodule, or a metastatic lesion of the umbilicus, was coined in 1949 by the English surgeon Sir Hamilton Bailey¹ after Sister Mary Joseph (1856–1939), a surgical assistant for Dr William Mayo², who was the first to associate intra-abdominal malignancy and

Figure 1 (A) Tumour protruding through her umbilical area that was oozing serosanguinous fluid. (B,C) CT of the abdomen revealing a large mass with loculated ascites. Multiple lung nodules were identified as well. (D) A poorly differentiated malignant neoplasm involving ovaries, the umbilical area and widespread omental metastases following tumour debulking.

the umbilical lesions while cleaning the skin of patients before surgery.

Competing interests None.

Patient consent Obtained.

REFERENCES

1. **Abu-Hilal M**, Newman JS. Sister Mary Joseph and her nodule: historical and clinical perspective. *Am J Med Sci* 2009;**337**:271–3.
2. **Albano EA**, Kanter J. Images in clinical medicine. Sister Mary Joseph's nodule. *N Engl J Med* 2005;**352**:1913.

This pdf has been created automatically from the final edited text and images.

Copyright 2011 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.

BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Please cite this article as follows (you will need to access the article online to obtain the date of publication).

Munoz J, Chacra W, Hanbali A, Ali H. Sister Mary Joseph's nodule. *BMJ Case Reports* 2011;10.1136/bcr.06.2011.4349, date of publication

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow