Published in final edited form as:

Tetrahedron Lett. 2011 September 21; 52(38): 4941–4943. doi:10.1016/j.tetlet.2011.07.068.

An unexpected synthesis of 3,5-diaryl-1,2,4-thiadiazoles from thiobenzamides and methyl bromocyanoacetate

Abdelrahman S. Mayhoub[§], Evgeny Kiselev, and Mark Cushman^{*}

Department of Medicinal Chemistry and Molecular Pharmacology, College of Pharmacy, and the Purdue University Center for Cancer Research, Purdue University, West Lafayette, Indiana 47907

Abstract

Aryl thioamides undergo a very rapid condensation in the presence of methyl bromocyanoacetate to provide quantitative yields of 3,5-diaryl-1,2,4-thiadiazoles with easy work-up and a high degree of product purity. The method can be scaled up with no loss in efficiency.

Keywords

1,2,4-Thiadiazoles; Methyl bromocyanoacetate; Thioamides; Condensation

1. Introduction

A number of methyl 2-aryl-4-methylthiazole-5-carboxylates **3** and their derivatives were previously prepared by the condensation of methyl α -chloroacetoacetate (**1**) with thiobenzamides **2** (Scheme 1).¹ In order to further study the potential of the arylthiazole class of compounds as antiviral agents, the preparation of methyl 4-amino-2-(4-chlorophenyl)thiazole-5-carboxylate (**6**) was attempted (Scheme 2).

Compound **6** was originally intended to be obtained by the condensation of methyl bromocyanoacetate (**4**) with thiobenzamide **2a** (Scheme 2). Compound **4**, prepared by bromination of the commercially available methyl cyanoacetate with *N*-bromosuccinimide (NBS), was allowed to react with thiobenzamide **2a**, but instead of the expected 4-amino-2-arylthiazole **6**, the diarylthiadiazole **5a** formed *instantaneously* and was isolated simply by filtration in *near quantitative* yield. A mechanism for the expected conversion of **2a** and **4** to **6** is outlined in Scheme 3. The structure of **5a** was confirmed by the spectral data (see supporting information). To the best of our knowledge, this reaction has never been reported before. However, the bromocyanoacetate **4**, together with other α -halocarbonyl compounds, have been reported as mild oxidizing agents in the conversion of thiols into disulfides.²

© 2011 Elsevier Ltd. All rights reserved.

^{*}Corresponding author. Tel.: +1-765-494-1465; fax: +1-765-494-6790; cushman@purdue.edu.

[§]On leave from Al-Azhar University, Collage of Pharmacy, Cairo, Egypt.

Supplementary Material

Detailed experimental data and NMR spectra for all new and known compounds. Supplementary data associated with this article can be found in the online version.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

The goal of the present study was to determine the scope of the reaction of aryl thioamides **2** with methyl bromocyanoacetate (**4**) to determine if it could constitute a general method for synthesis of 3,5-diaryl-1,2,4-thiadiazoles.

2. Results and Discussion

First, to determine the solvent effects on the yield of compound **5a**, methyl bromocyanoacetate (**4**) was allowed to react with the thiobenzamide **2a** using different solvents (methanol, ethanol, dioxane, and dichloromethane). The thiadiazole **5a** was obtained without any significant differences in the reaction time or yield. Methanol was generally used as the solvent throughout this work as mentioned in the supporting information.

Next, a variety of commercially available aryl thioamides **2** were allowed to react with methyl bromocyanoacetate (**4**) at room temperature (Scheme 4). Among the 1,2,4-thiadiazoles reported in the chemical literature, the ortho-substituted derivatives have been more difficult to prepare and there are fewer examples of them.³ Therefore, in this study more attention was given to synthesis of products containing substituents in the ortho positions of the phenyl rings and six ortho substituted derivatives **5c**, **5d**, **5h**, **5i**, **5j** and **5k** were prepared. The reaction of *o*-substituted thiobenzamides **2c**, **2h**, **2i**, **2j** and **2k** proceeded in quantitative yields; however, the *o*-trifluoromethyl derivative **5d** was obtained in low yield (10%). Most of the other meta, para, and disubstituted thiobenzamide derivatives afforded the corresponding 1,2,4-thiadiazoles in essentially quantitative isolated yields of pure products regardless of differences in electronic and/or steric properties of the substituent(s). The 3- and 4-pyridyl derivatives **5t**, **5u** were obtained in moderate yields (65–69%). However, the corresponding 2-pyridyl derivative could not be obtained in any yield and the starting materials decomposed under the reaction conditions. The condensation reaction to produce 3,5-diaryl-1,2,4-thiadiazoles did not work when 2,6-disubstituted thiobenzamides were used. To date, there are no efficient methods reported for the synthesis of 3,5-bis(2,6-dichlorophenyl)-1,2,4-thiadiazole, although it has been obtained as a by-product from the reaction of thioamides with nitrous acid.⁴ Moreover, oxidation of 2,6-dichlorothiobenzamide was reported to yield the corresponding isothiocyanate.⁵ We are unaware of any efficient methods reported for preparation of 3,5-di(2-pyridyl)-1,2,4-thiadiazole or any 3,5-[(2,6-disubstituted)aryl]-1,2,4-thiadiazole, including the presently reported method. Thioacetamide was tried as an example of an aliphatic thioamide and it was unsuccessful.

To investigate the applicability of this method for large-scale synthesis, four grams of thiobenzamides **2a** and **2g** were allowed to react with a slight excess (1.3 equivalent) of methyl bromocyanoacetate (**4**). The reaction mixtures were stirred vigorously for 2–3 min to ensure consumption of all of the starting materials, and the corresponding thiadiazoles **5a** and **5g** were isolated in a high degree of purity in 95 and 97% yields, respectively.

Several literature precedents suggest a mechanism for the condensation of two molecules of the aryl thioamides **2** to form 3,5-diaryl-1,2,4-thiadiazoles **5** in the presence of **4** (Scheme 5). The question of S-Br vs. S-C bond formation resulting from the reaction of thiols with α -bromo esters has been studied before, with S-Br bond formation being documented in cases in which the anion is more highly resonance stabilized (e.g. diethyl bromomalonate).² This suggests the initial bromination of the sulfur of the thioamide with methyl bromocyanoacetate to form intermediate **9**. Treatment of thioamides with alkylating agents that form stable carbocations was reported to afford the more thermodynamically stable *N*-alkylated product instead of the less stable *S*-analogue.⁶ Therefore, the formation of intermediate **10** from the stable carbocation **9** is suggested. Elimination of HSB_r will afford

aminoarylmethylenethioamide **12**. Oxidation of the sulfur of thioamides is reported to facilitate N-S bond formation,⁷ which, in this case, could be achieved by a second bromocyanoacetate molecule. Therefore, *S*-bromination of **12** followed by tautomerization and elimination of HBr would provide the diarylthiadiazole **5**. The reaction stoichiometry supports this mechanism. When thiobenzamide **2a** was allowed to react with 0.5 molar equivalent of **4**, the thiadiazole product **5a** was obtained in about 50% yield and approximately 50% of the thiobenzamide starting material was left unchanged in the reaction mixture as estimated by NMR. Using of 1.2 equivalents of **4** results in complete conversion of the thiobenzamides **2** to the final products.

The reported methods for the preparation of 3,5-diaryl-1,2,4-thiadiazoles from the corresponding thioamides include using different oxidants such as hypervalent iodine,³ dimethyl sulfoxide (DMSO)/2-chloro-1,3-dimethylimidazolinium chloride (DMC),⁸ DMSO/haloiminium salt,⁹ DMSO/HCl,⁷ methanesulfonic acid derivative with benzenetellurinic acid,¹⁰ and telluroxide or selenoxide.¹¹ Other more complicated methods are also reported: from the corresponding oxathiazolone,¹² benzenecarboximidamide and diethyl azodicarboxylate,¹³ or from 1,2,4-oxathiazoles and Lewis acid.¹⁴ Thermal cycloreversion of 6*H*-1,3,5-oxathiazine *S*-oxides followed by ring closure of the resulting intermediates is also reported to yield 1,2,4-thiadiazoles.¹⁵ More recently, Khosropour and Noei described the preparation of 1,2,4-thiadiazoles using 2,4,6-trichloro-1,3,5-triazine, DMSO and polyethylene glycol 400.¹⁶ Many of the reported methods that provide high yields (> 80%) require specific alkaline or acidic reagents that might be incompatible with some sensitive groups, tedious work up and longer reaction times than the presently reported method, or cannot be applied for larger-scale reactions.

3. Conclusion

In conclusion, a new and efficient method for the synthesis of the 3,5-diaryl-1,2,4-thiadiazole system was investigated using methyl bromocyanoacetate (**4**), which can be prepared easily in one step. This method was found to be applicable with sensitive groups such as the hydroxyl group, ortho derivatives, and on gram-scale reactions with easy work-up, high yield, very short reaction times, and a very high degree of product purity. These features compare favorably with many of the other methods of 3,5-diaryl-1,2,4-thiadiazoles synthesis.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This research was supported by a fellowship to A.S.M. from the Egyptian government and by National Institutes of Health (NIH) Research Grant U01 CA89566.

References and Notes

1. Ze L, Khaliq M, Zhou Z, Post CB, Kuhn RJ, Cushman M. *J Med Chem*. 2008; 51:4660–4671. [PubMed: 18610998]
2. Kato E, Oya M, Iso T, Iwao J. *Chem Pharm Bull*. 1986; 34:486–485. and references cited within
3. A) Shah AA, Khan ZA, Choudhary N, Loholter C, Schafer S, Marie GPL, Farooq U, Witulski B, Wirth T. *Org Lett*. 2009; 11:3578–3581. [PubMed: 19610610] B) Cheng D, Chen Z. *Synthetic Commun*. 2002; 32:2155–2159. C) Yan M, Chen Z, Zheng Q. *J Chem Res (S)*. 2003:618–619. D) Patil PC, Bhalerao DS, Dangate PS, Akamanchi KG. *Tetrahedron Lett*. 2009; 50:5820–5822.
4. Yusuf I, Harun P. *Chemiker-Zeitung*. 1980; 104:365–367.

5. Bahadir M, Nitz S, Parlar H, Korte F. *Z Naturforsch, B: Anorg Chem, Org Chem.* 1979; 5:768–769.
6. Katritzky AR, Drevniak M. *Tetrahedron Lett.* 1988; 29:1755–1758.
7. Forlani L, Lugli A, Boga C, Corradi AB, Sgarabotto P. *J Heterocyclic Chem.* 2000; 37:63–69.
8. Isobe T, Ishikawa T. *J Org Chem.* 1999; 64:6989–6992.
9. Isobe, T. *Jpn Kokai Tokkyo Koho. Heisei:* 1994. p. 06128243
10. Fukumoto T, Matsuki T, Hu NX, Aso Y, Otsubo T, Ogura F. *Chem Lett.* 1990:2269–72.
11. A) Hu NX, Aso Y, Otsubo T, Ogura F. *Bull Chem Soc Japan.* 1986; 59:879–84. b) Hu NX, Aso Y, Otsubo T, Ogura F. *Chem Lett.* 1985:603–606.
12. A) Howe RK, Shelton BR. *J Org Chem.* 1981; 46:771–5. B) Hojo M, Masuda R. *J Org Chem.* 1975; 40:963–5.
13. Kihara Y, Kabashima S, Uno K, Okawara T, Yamasaki T, Furukawa M. *Synthesis.* 1990:1020–1023.
14. Rafiqul-Islam M, Shimada K, Aoyagi S, Takikawa Y, Kabuto C. *Heteroatom Chem.* 2004; 15:175–186.
15. Shimada K, Rafiqul-Islam M, Sato M, Aoyagi S, Takikawa Y. *Tetrahedron Lett.* 2003; 44:2517–2519.
16. Khosropour AR, Noei J. *Montsh Chem.* 2010; 141:649–651.

Scheme 1.
Preparation of thiazoles.

Scheme 2.
Condensation of thiobenzamide **2a** with methyl bromocyanoacetate.

Scheme 3.
Mechanism for expected formation of **6**.

b, R = C₆H₅

c, R = 2-IC₆H₄

d, R = 2-CF₃C₆H₄

e, R = 3-BrC₆H₄

f, R = 3-IC₆H₄

g, R = 3-FC₆H₄

h, R = 2,3-Cl₂C₆H₃

i, R = 2,5-Cl₂C₆H₃

j, R = 2-Br-*p*-tolyl

k, R = 5-Cl-*o*-tolyl

l, R = 3,4-Cl₂C₆H₃

m, R = 4-CNC₆H₄

n, R = 4-CH₃C₆H₄

o, R = 4-*t*BuC₆H₄

p, R = 4-*n*BuC₆H₄

q, R = 4-NO₂C₆H₄

r, R = 4-HOC₆H₄

s, R = 4-CH₃OC₆H₄

t, R = 3-pyridyl

u, R = 4-pyridyl

Scheme 4.

Condensation of aryl thioamides with methyl bromocynoacetate.

Scheme 5.
Mechanism for formation of **5**.