Published in final edited form as:

J Org Chem. 2011 September 16; 76(18): 7341–7351. doi:10.1021/jo2012434.

Design and Synthesis of C₂-Symmetric *N*-Heterocyclic Carbene Precursors and Metal Carbenoids

Abigail Albright^a, Daniel Eddings^a, Regina Black^b, Christopher J. Welch^b, Nikolay N. Gerasimchuk^c, and Robert E. Gawley^{a,*}

^aUniversity of Arkansas, Department of Chemistry and Biochemistry; Fayetteville, Arkansas 72701

^bMerck & Co., Inc., One Merck Dr, P O Box 100, Whitehouse Station, New Jersey 08889

^cMissouri State University, Chemistry Department, Springfield, MO 65897

Abstract

Chiral, C₂-symmetric imidazolium and imidazolinium ions, as well as the corresponding copper or silver bound carbenoids, have been prepared. Structural study of these compounds by X-ray crystallography reveals a chiral pocket that surrounds the putative carbene site or the metal-carbene bond, at carbon 2, in three of the four ligands prepared. Preliminary investigation into the application of these complexes has shown one of them to be highly enantioselective in the hydrosilylation of acetophenone.

Introduction

N-Heterocyclic carbenes (NHCs) and their organometallic complexes have risen to prominence in recent years, in part due to their broad utility in the catalysis of numerous bond forming reactions.¹ The field of NHC catalysis has evolved rapidly and a variety of practical uses in synthetic chemistry have been developed.²

The immediate precursors to many of the useful NHC catalysts contain *N,N'*-diarylimidazolium and -imidazolinium rings. Removal of the C-2 proton of an imidazolium results in the formation of a carbene and retention of aromaticity, deprotonation of an imidazolinium affords a saturated carbene, and complexation of either with a metal affords a metal carbenoid (Figure 1). Consideration of the electronic attributes of each structure can prove useful in the rational design of an NHC catalyst for a given application.

*Corresponding author: bgawley@uark.edu.

SUPPORTING INFORMATION

Spectroscopic data for all compounds synthesized in the course of this investigation but not included in the text of the paper, as well as all XRD data and details of the DFT calculation. This information is available free of charge via the Internet at <http://pubs.acs.org>.

One of the most versatile (achiral) imidazolium NHCs is the *N,N'*-bis(2,6-diisopropylphenyl)imidazol-2-ylidene (IPr) ligand, in which 2,6-diisopropylphenyl groups comprise the aryl substituent on the heterocycle.³ The X-ray crystal structure of the IPrCuCl complex is displayed in Figure 2, in both wire and space-filling representations. It can be seen that the four methyls of the isopropyl groups are in close proximity to the copper atom, with an average distance of 3.840 Å. One can reasonably speculate that when the IPr NHC is used as a free carbene (organocatalyst), the catalyst-coordinated reactant would be in approximately the same place as the copper, while in metal-mediated catalysis, the position of the reactant would be approximately in the position of the chlorine atom, which is ~4.8 Å distant from the four methyl groups. Analysis of the crystal structure of CuIPr led us to speculate that an NHC stereocenter at the location of the isopropyl methine could produce a compound with a well-formed chiral pocket around the catalytic reactive site. A similar chiral pocket could ensue in a C₂-symmetric imidazolinium having *trans*-2,3-diphenyl substituents on the heterocyclic ring and a single *ortho*-substituted *N*-phenyl substituent.⁴ If such compounds retain synthetic versatility, they could constitute a useful class of enantioselective NHC catalysis.

A design in which the stereogenic center is γ to the annular nitrogen could produce an NHC that retains the synthetic versatility of the *N*-aryl NHC ligands and simultaneously place the reactive site in a chiral pocket if a C₂-symmetric conformation is retained. To our knowledge, NHC imidazole-derived ligands having stereocenter(s) γ to the nitrogen(s) have not yet been reported. A C₂-symmetric conformation would place two or more stereogenic centers in close proximity to the putative carbene site, whereby large groups would occupy diagonally opposite quadrants around the carbene. We therefore decided to investigate the preparation of such species, characterize them structurally by X-ray crystallography, and test them in the hydrosilylation of acetophenone.

Our decision to pursue asymmetric hydrosilylation as a test of the synthetic design was guided by the high industrial and academic utility of this transformation. The hydrosilylation of ketones provides an attractive alternative to hydrogenation, avoiding the use of expensive precious metal catalysts and harsh reaction conditions which can lead to over-reduction of the product. The resultant silyl ethers are readily cleaved to the corresponding alcohol, or alternatively the silyl group can be retained as a protecting group. Only a small number of NHC hydrosilylation catalysts have been reported to date.⁵

Results and Discussion

The design we envision for chiral, C₂-symmetric imidazoliums and imidazoliniums is shown in Scheme 1, along with the retrosynthetic plans for their preparation. Cyclization of diamine **2** with an orthoformate affords imidazolinium **1**, whereas cyclization of a diimine, **6**, with a formaldehyde equivalent affords the imidazoliums **7** and **8**. The plan for the preparation of geared imidazoliniums involved Buchwald-Hartwig coupling of an orthosubstituted bromobenzene such as **3** with 1,2-diphenylethylenediamine. We planned to make the bromobenzene by Sandmeyer substitution of a 2-substituted aniline, **5**, which is available by dynamic thermodynamic resolution (DTR). The diimine precursor, **6**, of the imidazoliums **7** and **8** could be prepared by condensation of anilines **4** or **5** with glyoxal. The 2,6-disubstituted aniline **4** could be prepared with Friedel-Crafts type chemistry.

According to this synthetic plan, we have succeeded in the preparation of four novel NHCs and NHC-precursors, depicted in Figure 5 as compounds **9–12**. Each of these compounds were characterized by X-ray diffraction and, with the exception of **10**, were found to beautifully illustrate the goal of our synthetic design. The bulk of the *o*-aryl substituents or, in the case of imidazolinium **9**, the gearing of the backbone phenyl moieties, induced the

formation of chiral pocket around C-2, with two stereocenters residing less than 5 Å from the reactive site of the catalyst.

Individual Syntheses: Imidazolinium **9**.

Imidazolinium **9** was derived from the enantioenriched aniline, **16** (Scheme 2). This compound is in turn obtained by the acylation of commercially available 2-benzylaniline with pivaloyl chloride, followed by enantioselective methylation via dynamic thermodynamic resolution, and lastly hydrolysis. Compounds **15** and **16** have previously been synthesized by Wilkinson et al.,⁶ however in much lower yield and selectivity. By conducting the DTR in methyl *tert*-butyl ether (MTBE) rather than diethyl ether and employing methyl iodide rather than methyl tosylate as the electrophile, we were able to obtain **15** in 89% yield and 95:5 er, improved from 41% yield and 78:22 er.⁶ The improved enantioenrichment obtained by solvation with MTBE was first noted by Beak with the same substrate, **14**, although a methylating electrophile was not included in the scope.⁷ Furthermore, the hydrolysis conditions employed below differ from those published by the Hussain group, which removed the pivaloyl group in THF using LiAlH₄. Refluxing in absolute ethanol and concentrated hydrochloric acid afforded aniline **16** in 83% yield (63% from **13**).⁸ Lastly, it should be noted that our assignment of the absolute configuration as *R* was confirmed by the subsequent XRD analysis of both imidazolinium **9** and silver carbenoid **11**, although this assignment differs from that reported by the Hussain group (See Experimental for further details). Our assignment is consistent with the report by Beak on the (–)- sparteine-mediated DTR of compound **14**, in which electrophilic addition to the *Re* face of the benzylic methylene was observed.⁷

Compound **16** was then converted to the corresponding aryl bromide via traditional Sandmeyer reaction conditions⁹ in nearly quantitative yield and subjected to a palladium catalyzed coupling^{4a} with *S,S*-1,2-diphenylethane-1,2-diamine to yield the chiral diamine **18** in 85% yield as a single diastereomer.

Diamine **18** was subjected to Grubbs' cyclization protocol^{4b} and gave a 37% yield of the hexafluorophosphate salt **9**, which was purified by recrystallization from ethanol. None of the numerous metallation protocols investigated allowed for the generation of an isolatable metal carbenoid from **9**.

Crystals of **9** suitable for XRD were grown by slow diffusion of dichloromethane into pentane. The monoclinic crystal ($\beta = 94.5024^\circ$) was determined to have the P2₁ space group and a final R factor of 0.0433 was obtained. The crystal structure is shown in Figure 6. The Flack Parameter was refined to a value of zero, providing confirmation of the absolute configuration as the (*S,S,R,R*) enantiomer. The phenyl groups of the *ortho*-aryl substituent are geared to the front of the imidazolinium ring to avoid steric interaction with the phenyl groups of the imidazolinium backbone, thereby blocking two faces of the carbene center. Encouraged by these results, we sought to achieve the synthesis of an unsaturated NHC containing a similar chiral pocket, formed via the steric interaction of bulky *ortho*-aryl substituents across the face of the heterocycle rather than by the gearing effect.

Individual Syntheses: Imidazolium **10** and Silver Carbenoid **11**

Imidazolium **10** was derived from the enantioenriched aniline **22** (Scheme 5). This compound was in turn synthesized by the acylation of commercially available 2-ethylaniline with pivaloyl chloride followed by dynamic thermodynamic resolution according to the method of Beak,¹⁰ and lastly hydrolysis. While this nearly quantitative hydrolysis of the enriched pivanolide is a new protocol, the DTR yielding **21** was performed with comparable yield and selectivity by Beak et al.

The acid catalyzed condensation of chiral anilines **16** and **22** with glyoxal was achieved in excellent yield (Scheme 6). Subsequent cyclization of diimines **23** and **24** with chloromethylethylether was achieved in 47% and 45% yield, respectively. Imidazolium chlorides **10** and **25** were purified by column chromatography and obtained as single diastereomers.

Characterization

Crystals suitable for x-ray crystallography were obtained for **10** through slow evaporation of a THF solution. The crystal was found to be orthorhombic, and determined to have the P212121 space group. A final R factor of 0.0354 was obtained. The crystal structure is illustrated in Figure 7. The Flack parameter was refined to a value of zero, providing confirmation of the absolute configuration as the (*R,R*) enantiomer. We were chagrined to observe that the aryls groups reside in a conformation lacking C₂-symmetry. Both trimethylsilyl moieties reside on the same face of the heterocycle, indicating that the steric bulk of the *ortho*-aryl trimethylsilyl substituents is insufficient to enforce C₂-symmetry and create the desired chiral pocket. Our work was consequently focused on the metallation and characterization of **25**.

The metallation of imidazolium chloride **25** was complicated by its decomposition to a formamidine in the presence of base. Even mild organic bases induced decomposition, but a protocol by Wang and Lin¹¹ utilizing silver (II) oxide as both base and metal source provided for the isolation of **11** in low yield (Scheme 7).

Characterization

Crystals of **11** suitable for x-ray diffraction were grown by slow evaporation of a dichloromethane solution. The monoclinic crystal ($\beta = 116.653^\circ$) was determined to have the C2 space group a final R factor of 0.0409 was obtained. The crystal structure is presented in Figure 8. The Flack Parameter was refined to a value of zero, providing confirmation of the absolute configuration as the (*R,R*) enantiomer. The crystal structure of the silver carbenoid reveals a nicely formed chiral pocket and served to confirm the rationale of our synthetic design. To our knowledge, compound **11** represents the first example of a structural gearing effect induced not by placement of bulky substituents on the backbone of a saturated imidazolinium but via trans-annular steric interaction of bulky substituents γ to nitrogen in an unsaturated imidazolium. Nevertheless, the low yielding metallation step provides a major limitation in the future utility of compound **11** in asymmetric catalysis. Our efforts were thus concentrated on the synthesis of an enantioenriched 2,6-disubstituted aniline.

Individual Syntheses: Copper Carbenoid **12**

We surmised that increasing structural rigidity via the use of a di-*ortho*-substituted aniline precursor could yield a more promising final structure. Initial investigation led us to attempt a lithiation/DTR of 2,6-diethylpivaniide (Scheme 8). While this transformation proceeds in comparable yield to the singly substituted pivaniides, the enantioselectivity is poor (60:40er).

Numerous methodologies were investigated in attempts to synthesize an enantioenriched 2,6-disubstituted aniline. Ultimately, Friedel-Crafts type chemistry was employed for the di-*ortho* substitution of toluidine with either phenylacetylene followed by reduction, or styrene, protocols published by Tarantola¹² and Coates,¹³ respectively (Scheme 9). Although lengthier by one step, the Tarantola route was found to be a higher yielding and cleaner way to access compound **28** as a mixture 15 of a chiral racemate and a *meso* diastereomer. This

mixture was separated via stacked injection into a semi-preparative chiral stationary phase SFC (Figure 9).

Each of the stereoisomers were independently analyzed: It was found that the first to elute is (+)-**28**, the second (–)-**28** and the third is the optically inactive *meso* **28**. Subsequent XRD analysis of the copper carbenoid **12**, prepared from (+)-**28**, revealed it to be the (*S,S*) enantiomer (see below).

The following synthesis was optimized with *meso*-**28**, as it has no potential in asymmetric catalysis. For clarity's sake, we will illustrate the synthesis with (+)-(*S,S*)-**28**. Enantioenriched aniline (+)-(*S,S*)-**28** was condensed with glyoxal in quantitative yield to give diimine (*S,S*)-**29** (Scheme 10). The cyclization and purification of this product, however, required extensive optimization. Fortunately, minimal revisions to a cyclization protocol by Markó,¹⁴ which utilizes zinc chloride to coordinate the diimine in the reactive *s-cis* conformation, provided for isolation of pure (*S,S,S,S*)-**30** in 87% yield after recrystallization.

When *meso*-**28** was condensed and cyclized, some interesting stereochemical features were discovered in the characterization of *meso*-**30**: The NMR spectra exhibited quadrupling of the aliphatic signals. Further consideration of this phenomenon led us to conclude that *meso*-**30** exists as a mixture of three rotamers (Figure 10). We hypothesize that conformer A is the most highly populated species: one can imagine that the cyclization of the diimine to the imidazolium is facilitated by a conformation which places the forming imidazole carbon in the least sterically hindered environment. Rotation around both *N*-aryl bonds would give conformer B,¹⁵ likely the least populated species because carbon 2 is in a highly crowded environment in the midst of four bulky phenyl groups. Rotation around only one of the rightmost *N*-aryl bonds of A or B gives compound C. The two depictions of compound C are identical, however the methyl groups γ to the heterocyclic nitrogen are diastereotopic, resulting in their non-equivalence in the proton NMR spectrum.

Imidazolium chlorides (*R,R,R,R*)-**30** and (*S,S,S,S*)-**30** were both subjected to metallation in the presence of copper (I) chloride and NaOt-Bu (Scheme 11). Copper carbenoids (*R,R,R,R*)-**12** and (*S,S,S,S*)-**12** were obtained in good yield via this protocol.

Crystals of (*S,S,S,S*)-**12** suitable for x-ray diffraction were grown by slow diffusion of hexanes into ether. The monoclinic crystal ($\beta = 115.097^\circ$) was determined to have the C2 space group; a final R factor of 0.0438 was obtained. The crystal structure is presented in Figure 11. The Flack Parameter was refined to a value of zero, establishing the absolute configuration as the (*S,S,S,S*). We were pleased to observe the formation of a chiral pocket at the site of the putative carbene. Compound **12** successfully places four stereocenters in close proximity to the C-2 reactive site.

Application

Of the four novel NHC precursors and metal carbenoids we were able to access synthetically, three served to confirm the feasibility of the design motif. With these compounds in hand, we set out to test this new class of NHCs in an enantioselective reaction. The hydrosilylation of prochiral ketones provides access to enantioenriched alcohols without the use of high pressure reactors and expensive metal catalysts. Simmons and Hartwig recently showed that racemic (hydrido)silyl ethers of 1-phenyl alcohols can be used to activate the ortho C–H bond to make benzoxasiloles, which can then be further functionalized in a number of useful ways.¹⁶ Furthermore, the catalytic cycle using the CuIPr NHC has been studied in detail.¹⁷ As our novel class of metal bound NHCs and NHC precursors can be considered chiral analogs of the CuIPr NHC, with chiral centers

incorporated at the location of the isopropyl methine, we chose this transformation for the investigation of compounds **9**, **11**, and **12** as asymmetric catalysts (Scheme 12 and Table 1).

Imidazolinium **9** was found to be the least effective in the hydrosilylation of acetophenone, achieving only minimal conversion to the silylether after 24h. The enantioselectivity of the transformation was very poor. Unsaturated imidazoliums **25** and **12** were found to give nearly quantitative conversion to the silylether at room temperature in 4 and <1h, respectively. And while **25** yields the product with only very moderate enantioenrichment, **12** provides for isolation of nearly enantiopure silylether. The absolute configuration of the silylether was determined by coinjection of an enantiopure authentic sample and found to be *S* when using the *R,R,R,R* stereoisomer of **12**.

The usually accepted¹⁸ catalytic cycle (Scheme 13) proceeds via the formation of a copper hydride followed by coordination of the ketone and sigma bond metathesis. There is precedent for the in-situ formation of the copper carbenoid,¹⁹ so imidazolinium **9** and imidazolium **25** – for which we lack efficient metallation procedures – were investigated in this manner.

In order to obtain better insight into the steric course for the hydrosilylation of acetophenone catalyzed by **12**, a DFT minimization (B3LYP 3-21g-d) was carried out on two conformations of intermediate **B** in the catalytic cycle using the PQS ab-initio molecular modeling software.²⁰ The minimized structures are shown in Figure 12, with two of the *ortho* *N*-aryl substituents drawn as R groups for clarity. In the lowest energy conformation of **B**, the copper, the hydride, and the carbonyl of acetophenone lie in a common plane, and more critically, this plane is roughly orthogonal to the plane of the heterocycle. Recalling that **12** has C₂-symmetry, examination of the minimized structures reveals that the phenyl group of acetophenone preferentially resides in the less crowded upper right quadrant and thereby exposes the *Re* face of the carbonyl to the hydride residing on copper. Delivery of the hydride in this manner would lead to formation of the *S*-enantiomer of the silyl ether, as is observed experimentally. The relative energy between these two intermediates was found to be a 3.6 kcal/mol.

Summary and Conclusion

New, chiral C₂-symmetric NHC ligands have been prepared. These NHC's and metal carbenoids exhibit the structural motif of having two or four stereocenters γ to the heterocyclic nitrogen. The imidazolinium **9** induces these chiral "arms" to reside in close proximity to the putative carbene via the gearing effect, while the conformations of metal carbenoid **11** is geared by transannular steric interaction of the bulky *ortho*-aryl substituents. Copper carbenoid **12**, CuPhEt, has little conformational mobility and so the chiral pocket created by the α -phenethyl arms is essentially fixed. Initial investigation into the application of this novel class of NHCs has revealed **12** to be a powerful catalyst for the hydrosilylation of acetophenone, achieving this transformation at room temperature in 45 minutes in quantitative conversion and high enantiopurity. Further investigation into the mechanism, stereoselectivity and application of these catalysts is ongoing and will be reported in due course.

EXPERIMENTAL

General

All reagents and solvents were used as received, unless indicated below. Diethyl ether (Et₂O), *t*-butyl methyl ether (MTBE), and tetrahydrofuran (THF) were distilled over Na/benzophenone ketyl under N₂ immediately prior to use. Toluene was dried over CaH₂

followed by distillation and storage over 3 Å molecular sieves. Methylene chloride was distilled over CaH₂ under N₂ immediately prior to use. *N,N,N',N'*-tetramethylethylenediamine (TMEDA) was vacuum distilled over CaH₂ and stored over 4 Å molecular sieves. (–)-Sparteine was vacuum distilled over CaH₂ and stored over 4 Å molecular sieves. Solutions of *s*-BuLi in cyclohexane were titrated against *N*-pivaloyl-*o*-benzaniline. ¹H NMR spectra were collected at 300 or 400 MHz in CDCl₃ unless otherwise noted. Coupling constants (J), are reported in Hz. Supercritical fluid chromatography was used to determine enantiomer ratios on a Daicel Chiracel OD-H column or a Whelk-O column using absolute ethanol as modifier. Details of these separations can be found in the Supporting Information.

Compounds **13** and **19** are commercially available and were used as received without further purification. The following compounds have been previously reported and their spectra were consistent with that of the published data: Pivanilides **14**, **15**, **20**, **21**,^{6,10} anilines **16**,⁶ **28**,¹³ aryl bromide **17**,²¹ and diphenylsilyl ether **31**.²²

The characterization of the following compounds is detailed below: Imidazolium **9**, imidazoliums **10**, **25**, **30**, silver carbenoid **11**, copper carbenoid **12**, diamine **18**, aniline **22**, diimines **23**, **24**, **29**, pivanilide **27**, **29**, **30**. If synthesized according to a known procedure, the reference is given. Any alterations to the published procedure are noted. Otherwise details of the synthetic protocol can be found below.

General Procedure for Dynamic Thermodynamic Resolution

To an oven dried, Argon purged, 250 mL round bottom flask was added the pivanilide **14**, **20** or **26** (14.6 mmol) in 120 mL freshly distilled MTBE or Et₂O and cooled to –25 °C. Some crystallization occurs. To this was added dropwise 35 mL of 1.4 M *s*-BuLi in cyclohexane (49.0 mmol). After 2 h, (–)-Sparteine (50.0 mmol) was added, stirred 45 min, then rapidly cooled to –78 °C. After 30 min, the electrophile (TMSCl or MeI, 29.5 mmol) was added and stirred 16 h. The reaction was quenched with 15 mL MeOH followed by 15 mL water and warmed to room temperature. The contents were transferred to a separatory funnel, diluted with 50 mL water, and washed 3 × 25 mL EtOAc. The combined organic layers were washed with 3 × 20 mL 5% H₃PO₄, 3 × 20 mL brine, dried over MgSO₄, filtered and concentrated in vacuo to give a pale tan oil which was used without further purification for subsequent reactions (89–99% yield). An analytical sample was prepared by radial chromatography on silica, eluting with 95% hexanes/4.5% EtOAc/0.5% NH₄OH followed by recrystallization via slow evaporation of hexanes. NMR data of **15** and **21** match the literature data.^{6,10}

***N*-(2-ethyl-6-(1-(trimethylsilyl)ethyl)phenyl)pivalamide (*R*)-27**—¹H NMR (CDCl₃): –0.0 (9H, s), 1.2 (3H, t, J=7.5), 1.3 (3H, d, J=7.5), 1.4 (9H, s), 2.3 (1H, q, J=7.5), 2.5 (2H, q, J=7.5), 6.7 (1H, bs), 7.0 (2H, m), 7.2 (1H, t, J=7.6). ¹³C NMR (CDCl₃): –2.8 (CH₃), 14.7 (CH₃), 15.7 (CH₃), 23.6 (CH), 25.5 (CH₂), 28.0 (CH₃), 39.4 (C), 124.9 (2-CH), 127.7 (CH), 131.9 (C), 141.5 (C), 144.0 (C), 176.6 (C). MS (CI/MH⁺): 306. Anal calcd for C₁₈H₃₁NOSi: C, 70.76; H, 10.23. Found: C, 70.62; H, 10.26. er=60:40.

General Procedure for Pivanilide Hydrolysis

To a 100mL round bottom flask equipped with a stir bar was added the pivanilide (2.88mmol), absolute ethanol (30mL) and concentrated hydrochloric acid (30mL). The reaction solution was fitted with a condenser and refluxed at 100 °C for 2d after which time the reflux condenser was replaced with a distillation short path and the reaction volume was reduced to approximately 5mL. The crude reaction solution was then cooled to room temperature, diluted with diethyl ether and neutralized with potassium hydroxide. The

organic layer was extracted thrice with diethyl ether (10mL) and washed with distilled water before drying over MgSO₄. The reaction solution was filtered and concentrated in vacuo to give the corresponding anilide.

2-(1-phenylethyl)benzenamine (R)-16—¹H NMR (CDCl₃): 1.7 (3H, d, J=7.2), 3.5 (2H, s), 4.1 (1H, q, J=7.2), 6.7 (1H, dd, J=1.2, 7.8), 6.9 (1H, dt, J=1.2, 7.5), 7.1 (1H, dt, J=1.5, 7.5), 7.2–7.3 (3H, m), 7.3–7.4 (3H, m). ¹³C (CDCl₃): 22.0 (CH₃), 40.4 (CH), 116.4 (CH), 118.9 (CH), 126.6 (CH), 127.4 (CH), 127.6 (CH), 128.9 (CH), 129.9 (C), 144.5 (C), 145.8 (C). MS (CI/MH⁺): 198. Anal calcd for C₁₄H₁₅N: C, 85.24; H, 7.66. Found: C, 85.27; H, 7.75. ⁶er=95:5.

2-(1-(trimethylsilyl)ethyl)benzenamine (R)-22—¹H NMR (CDCl₃): 0.0 (9H, s), 1.4 (3H, d, J=7.5), 2.1 (1H, q, J=7.5), 3.5 (2H, bs), 6.7 (1H, dd, J=1.2, 7.8), 6.8 (1H, dt, J=1.2, 7.5), 6.9–7.0 (2H, m). ¹³C NMR (CDCl₃): -2.7 (CH₃), 15.4 (CH₃), 22.3 (CH), 116.0 (CH), 119.3 (CH), 125.1 (CH), 127.0 (CH), 131.2 (C), 143.1 (C). MS (CI/MH⁺): 194. Anal calcd for C₁₁H₁₉NSi: C, 68.33; H, 9.90. Found: C, 68.23; H, 9.93. er=94:6.

General Procedure for Sandmeyer Reaction

The following is revised from the published protocol for a different aniline substrate.⁹ To a 50mL round bottom flask was added NaNO₂ (14.5mmol) in 10mL DMSO. To this solution was added CuBr (7.0mmol) and the aniline substrate **16**, (3.9mmol). The resulting suspension was cooled to 0 °C and treated dropwise with a solution of HBr (5mL, 48%) in DMSO (10mL). The reaction solution was stirred for an additional 30 min before warming to room temperature and stirring for 1h. The reaction solution was then transferred to a separatory funnel, diluted with water (50mL) and washed with diethyl ether (3×20mL). The combined organic layers were washed with water (3×15mL) and then dried over MgSO₄, filtered and concentrated in vacuo.

1-(1-(2-bromophenyl)ethyl)benzene (R)-17—¹H NMR (CDCl₃): 1.7 (3H, d, J=7.2), 4.7 (1H, q, J=7.2), 7.1 (1H, m), 7.2–7.4 (7H, m), 7.6 (1H, dd, J=1.2, 7.8). ¹³C NMR (CDCl₃): 21.5 (CH₃), 43.7 (CH), 125.0 (C), 126.4 (CH), 127.8 (CH), 127.9 (CH), 128.0 (CH), 128.5 (CH), 129.0 (CH), 133.1 (CH), 145.1 (C), 145.6 (C). MS (EI): 260.262. er=95:5. As compound **17** was reported in 1973,²¹ we have included our spectra in the supplemental information to provide more detailed characterization.

1,2-Diphenyl-*N,N'*-bis-[(R)-2-(1-phenyl-ethyl)-phenyl]-ethane-(S,S)-1,2-diamine, 18—(New compound synthesized with a previously published procedure^{4a}) ¹H NMR (CDCl₃): 1.6 (6H, d, J=7.2), 3.9 (2H, q, J=7.2), 4.2 (2H, d, J=6.3), 4.5 (2H, d, J=6.9), 6.4 (6H, d, J=7.5), 6.7 (2H, dt, J=0.6, 7.5), 6.9 (4H, m), 7.0 (4H, m), 7.1–7.2 (8H, m), 7.3 (4H, m). ¹³C NMR (CDCl₃): 22.3 (CH₃), 40.1 (CH), 61.9 (CH), 112.1 (CH), 117.4 (CH), 126.3 (CH), 126.8 (CH), 127.4 (CH), 127.5 (CH), 127.8 (CH), 128.0 (CH), 128.2 (CH), 128.8 (CH), 130.0 (C), 139.2 (C), 144.0 (C), 145.7 (C). MS (ESI/MH⁺): 573. Anal calcd for C₄₂H₄₀N₂: C, 88.07; H, 7.04. Found: C, 88.25; H, 7.20. Single diastereomer.

(S,S)-4,5-Diphenyl-1,3-bis-[(R)-2-(1-phenyl-ethyl)-phenyl]-4,5-dihydro-3*H*-imidazol-1-ium; hexafluorophosphate – 9—(New compound synthesized with a previously published procedure^{4b}) ¹H NMR (CDCl₃): 1.2 (6H, d, J=6.9), 3.9 (2H, q, J=6.9), 5.8 (2H, s), 6.9 (4H, d, J=7.5), 7.1–7.4 (22H, m), 7.5 (1H, s). ¹³C NMR (CDCl₃): 22.8 (CH₃), 39.5 (CH), 76.5 (CH), 127.2 (CH), 127.5 (CH), 128.6 (CH), 128.7 (CH), 128.8 (CH), 129.2 (CH), 129.3 (CH), 130.0 (CH), 130.8 (CH), 130.8 (CH), 132.1 (C), 132.4 (C), 140.4 (C), 144.8 (C), 157.3 (CH). MS (ESI/-PF₆): 583. Anal calcd for C₄₂H₄₀N₂·1/2 H₂O: C, 70.00; H, 5.46. Found: C, 69.83; H, 5.33. XRD analysis of **9** resulted in refinement of the

Flack Parameter to a value of zero, providing confirmation of the absolute configuration as (*S,S,R,R*).

General Procedure for Glyoxal Condensation and Cyclization of **16** and **22**: To a 25 mL round bottom flask was added chiral aniline **16** or **22** (5.80 mmol) in 3 mL EtOH. To this was added 40% glyoxal (2.9 mmol). The reaction mixture was refluxed and monitored by TLC until all starting material had been consumed, taking approximately 2 h. The contents were then concentrated in vacuo to a dark purple foam. After drying, the product (**23** or **24**) was recovered (93–96% yield) and 2.7mmol were added to a 50 mL round bottom flask under inert atmosphere. To this was added chloromethylethylether (3.5 mmol) in 5 mL THF and a couple of drops of water. This was stirred at 50 °C for 16 h. The contents were concentrated in vacuo and purified by silica column chromatography 2% MeOH/DCM. The resulting residue was sonicated in ether and filtered (45–47% yield). These diimines were used without further purification.

***N,N*-Bis-[2-[1-(trimethyl-silanyl)-ethyl]-phenyl]-ethane-1,2-diylidenediimine (*R,R*)-23**—¹H NMR (CDCl₃): −0.098 (18H, s), 1.4 (6H, d, *J*=7.8), 3.1 (2H, q, *J*=7.8), 6.9–7.1 (2H, m), 7.1–7.3 (6H, m), 8.3 (2H, s).

***N,N'*-Bis-[2-(1-phenyl-ethyl)-phenyl]-ethane-1,2-diylidenediimine (*R,R*)-24**—¹H NMR (CDCl₃): 1.7 (6H, d, *J*=7.2), 4.8 (2H, *J*=7.2), 6.9–7.4 (18H, m), 8.2 (2H, s).

1,3-Bis-[2-[1-(trimethyl-silanyl)-ethyl]-phenyl]-1*H*-imidazolium chloride, (*R,R*)-10—¹H NMR (CDCl₃): −0.066 (18H, s), 1.4 (6H, d, *J*=7.2), 1.9 (2H, q, *J*=7.2), 7.2–7.3 (4H, m), 7.4–7.5 (2H, m), 7.6–7.7 (2H, m), 8.2 (2H, d, *J*=7.8). 9.4 (1H, s). ¹³C NMR (CDCl₃): −2.7 (CH₃), 16.6 (CH₃), 23.8 (CH), 124.8 (CH), 126.7 (CH), 128.2 (CH), 128.8 (CH), 131.4 (CH), 132.4 (CH), 137.9 (C), 141.7 (C). ESI (MS/−Cl): 421. Anal calcd for C₂₅H₃₇ClN₂Si₂: C, 65.68; H, 8.16. Found: C, 65.41; H, 8.23. Single diastereomer. XRD analysis of **10** resulted in refinement of the Flack Parameter to a value of zero, providing confirmation of the absolute configuration as (*R,R*).

1,3-Bis-[2-(1-phenyl-ethyl)-phenyl]-1*H*-imidazolium chloride (*R,R*)-25—¹H NMR (CDCl₃): 1.6 (6H, d, *J*=6.9), 4.3 (2H, q, *J*=6.9), 6.9 (4H, m), 7.1–7.3 (8H, m), 7.4–7.5 (4H, m), 7.5–7.7 (4H, m), 8.9 (1H, s). ¹³C NMR (CDCl₃): 22.8 (CH₃), 39.9 (CH), 124.7 (CH), 127.0 (CH), 127.1 (CH), 127.9 (CH), 128.5 (CH), 129.0 (CH), 129.2 (CH), 131.9 (CH), 133.7 (C), 140.6 (C), 145.6 (C). ESI-FTMS Calc. [M]⁺ C₃₁H₂₉N₂ [M]⁺ = 429.2331 Found [M]⁺ = 429.2328. Single Diastereomer.

4-methyl-2,6-bis(1-phenylethyl)benzenamine (mixture of stereoisomers)-28—A 25mL round bottom flask equipped with a stir bar was charged with toluidine (10.0mmol), KSF montmorillonite (1.0g), and phenyl acetylene (40.0mmol). The round bottom was fitted with a reflux condenser and the heterogeneous slurry was refluxed with vigorous stirring at 140°C for 8h. The reaction vessel was allowed to cool to room temperature before dilution with ethyl acetate and filtration. The solvent was removed from the mother liquor under reduced pressure and the resultant red oil was purified via column chromatography with 50:50 dichloromethane and hexanes. The product (4-methyl-2,6-bis(1-phenylvinyl)benzenamine) was obtained as an off-white solid in 88% yield. ¹H NMR (CDCl₃): 2.2 (3H, s), 3.3 (2H, s), 5.3 and 5.7 (4H, dd), 6.9 (2H, s), 7.1–7.4 (10H, m). ¹³C NMR (CDCl₃): 20.5, 116.2, 126.5, 127.8, 128.3, 128.7, 131.0, 139.3, 139.9, 147.5. A Parr shaker was charged with 4-methyl-2,6-bis(1-phenylvinyl)benzenamine (8.8mmol), 10w/w% Pd/C (0.88mmol), and absolute ethanol (25mL). The shaker was thrice purged with hydrogen before it was pressurized to 90psi and shaken overnight. The black heterogeneous solution was filtered and the solvent removed from the mother liquor under reduced pressure

to give compound **28** in nearly quantitative yield (99%) with a diastereomer ratio of 60:20:20. The details of the preparative separation of **28** can be found in the Supporting Information.

N,N'-Bis-[4-methyl-2,6-bis(-1-phenylethyl)phenyl]-ethane-1,2-diylidenediimine-29—To a 25 mL round bottom flask was added chiral or meso aniline **28** (5.80 mmol) in 3 mL EtOH. To this was added 40% glyoxal (2.9 mmol) and one drop of formic acid. The reaction mixture was sonicated overnight at room temperature. The contents were then filtered to give a bright yellow solid. The mother liquor was concentrated and recrystallized from ethanol. These combined batches yielded the product diimine **29** quantitatively (2.88mmol) as a single diastereomer. ¹HNMR (CDCl₃): 1.6 (12H, d, J=6.3), 2.3 (6H, s), 4.0 (4H, q, J=6.5), 7.0 (s, 4H), 7.2 (m, 20H), 7.4 (s, 2H). ¹³CNMR (CDCl₃): 21.4, 22.1, 39.1, 125.9, 126.2, 127.8, 128.6, 134.5, 146.6, 164.0. [α]_D = 66.3, (c = 1.76, dichloromethane). ESI-FTMS Calc. [M+H]⁺ C₄₈H₄₉N₂ = 653.38957. Found [M+H]⁺ = 653.38923.

1,3-Bis-[2,6-(S,S)-(1-phenyl-ethyl)-phenyl]-1H-imidazolium chloride-30—The following procedure is a revision of the protocol published by Markó.¹⁴ A solution of zinc (II) chloride and paraformaldehyde in concentrated hydrochloric acid was prepared (1:1:2, ~4.2M in paraformaldehyde). A test tube with a screw cap was charged with the diimine (**31**, 0.31mmol) in freshly distilled THF (8mL) under inert atmosphere and was treated with the acid solution (0.372mmol paraformaldehyde, 1.2eq). The reaction solution was then heated to 70 °C in the sealed tube for 1h before cooling to room temperature and removal of the solvent in vacuo. The resultant residue was then dissolved in dichloromethane (2mL) and washed with water (3×1mL) and saturated NaHCO₃ (1×1mL) before drying over MgSO₄, filtration and removal of the solvent in vacuo. The off white solid was then recrystallized from diethyl ether and dichloromethane. ¹HNMR (CDCl₃): 1.4 (6H, d, J=6.7), 1.6 (6H, d, J=6.7), 2.3 (6H, s), 3.75 (2H, q, J=6.6), 3.85 (2H, q, J=6.6), 6.5 (2H, s), 6.7 (4H, d, J=7.5), 6.9 (2H, s), 7.0–7.5 (18H, m), 11.6 (1H, s). ¹³CNMR (CDCl₃): 21.8, 21.85, 22.7, 38.5, 40.0, 124.3, 126.5, 126.8, 127.1, 127.6, 127.8, 128.3, 128.5, 129.1, 141.3, 141.9, 142.8, 143.5, 145.6. ESI-FTMS Calc. [M⁺] C₄₉H₄₉N₂ = 665.38957. Found [M⁺] = 665.38957. Single Diastereomer.

Copper carbenoid (S,S,S,S)-12—¹HNMR (CDCl₃): 1.25 (6H, d, J=8.0), 1.4 (6H, d, J=8.0), 2.15 (6H, s); 3.65 (m, overlapping quartets, J=5.3); 6.3 (2H, s), 6.6–7.3 (m, 24H). ¹³CNMR (CDCl₃): 21.7, 21.9, 22.7, 38.0, 39.6, 123.0, 126.0, 126.5, 127.0, 127.5, 128.0, 129.1, 133.2, 140.5, 142.3, 144.0, 144.3, 145.9, 181.2. ESI-FTMS Calc [M⁺] C₄₉H₄₈CuN₂ = 727.31135 Found [M⁺] = 727.31176. XRD analysis of **12** resulted in refinement of the Flack Parameter to a value of zero, providing confirmation of the absolute configuration as (S,S,S,S).

General Procedure for Hydrosilylation

(Following the procedure of Nolan¹⁸) To an oven dried vial fitted with a septum screw cap was added *either* copper carbenoid **12** (6.0μmol) and sodium *tert*butoxide (0.024mmol) *or* imidazolium **25** or imidazolinium **9** (6.0μmol), Cu(I)Cl (6.0μmol) and sodium *tert*-butoxide (0.024mmol) in freshly distilled toluene (2mL). The cloudy white reaction solution was stirred for 10 min at room temperature before addition of the silane (1.0 mmol). An immediate color change to bright yellow was observed and the reaction was stirred for an additional 20 min before addition of acetophenone (0.2mmol). The reaction progress was monitored by GC. Upon completion (or observation of the cessation of product formation) the reaction solution was treated with charcoal and filtered first through a plug of celite and then a plug of silica gel. The solvent was removed under reduced pressure and the residue

was analyzed by supercritical fluid chromatography. Diphenylsilyl ether **31** was separated on a Daicel Chiralcel OD-H (Chiral Stationary Phase: 4-Cellulose tris-(3,5-dimethylphenylcarbamate) coated on 5 μ m silica-gel) at a flow rate of 2.0 mL/min, with polarity modifier = 2.0% EtOH and outlet pressure and oven temperature of 150psi and 35°C, respectively. Alternatively, the enantiomers of **31** were resolved via CSP-GC with a β -Dex column (temperature program 100 °C isothermal).

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank the Arkansas Biosciences Institute and the National Science Foundation (CHE 1011788) for direct support of this work. Core facilities were funded by the Arkansas Biosciences Institute and the National Institutes of Health (1P20RR 15569 and 1P30RR 031154). The authors thank Kirsten Zeitler and Oliver Reiser for their help in obtaining the crystal structure of **9** and Bruce Noll for the crystal structure of **10**. We thank Peter Pulay for his help with the DFT study of intermediate **B**, and Chris Fitchett, Peter Steel, and Robert Grubbs for helpful discussions.

References and Notes

1. (a) Glorius, F., editor. N-Heterocyclic carbenes in transition metal catalysis. Vol. 21. Springer-Verlag; Berlin: 2004. (b) Nolan, SP., editor. N-Heterocyclic carbenes in synthesis. Wiley-VCH; Weinheim: 2006.
2. (a) Herrmann WA. Angew Chemie Int Ed. 2002; 41:1290–1239.(b) Enders D, Balensiefer T. Acc Chem Res. 2004; 37:534–541. [PubMed: 15311952] (c) Crudden CM, Allen DP. Coord Chem Rev. 2004; 248:2247–2273.(d) César V, Bellemin-Laponnaz S, Gade LH. Chem Soc Rev. 2004; 33:619–636. [PubMed: 15592627] (e) Mauduit, M.; Clavier, H. N-Heterocyclic Carbenes in Synthesis. Nolan, SP., editor. Wiley-VCH; Weinheim: 2006. p. 183-222.(f) Gade, LH.; Bellemin-Laponnaz, S. N-Heterocyclic Carbenes in Transition Metal Catalysis. Glorius, F., editor. Vol. 21. Springer-Verlag; Berlin: 2007. p. 117-157.(g) Jones WD. J Am Chem Soc. 2009; 131:15075–15077. [PubMed: 19919151] (h) Bantreil X, Broggi J, Nolan SP. Annu Rep Prog Chem, Sect B: Org Chem. 2009; 105:232–263.(i) Díez-González; Marion N, Nolan SP. Chem Rev. 2009; 109:3612–3676. [PubMed: 19588961]
3. Kaur H, Zinn RK, Nolan SP. Organometallics. 2004; 23:1157–1160.
4. (a) Seiders TJ, Ward DW, Grubbs RH. Org Lett. 2001; 3:3225–3228. [PubMed: 11574037] (b) Funk TW, Berlin JM, Grubbs RH. J Am Chem Soc. 2006; 128:1840–1846. [PubMed: 16464082]
5. (a) Duan WL, Shi M, Rong GB. Chem Commun. 2003:2916–2917.(b) Gade LH, César V, Bellemin-Laponnaz S. Angew Chem Int Ed. 2004; 43:1014–1017.(c) Song C, Ma C, Ma Y, Feng W, Ma S, Chai Q, Andrus MB. Tetrahedron Lett. 2005; 46:3241–3244.(d) César V, Bellemin-Laponnaz S, Wadepohl H, Gade LH. Chem Eur J. 2005; 11:2862–2873.(e) Liu, L-j; Wang, F.; Shi, M. Organometallics. 2009; 28:4416–4420.
6. Wilkinson JA, Rossington SB, Ducki S, Leonard J, Hussain N. Tetrahedron. 2006; 62:1833–1844.
7. Park YS, Yum EK, Basu A, Beak P. Org Lett. 2006; 8:2667–2670. [PubMed: 16774227]
8. We assume that this acid hydrolysis does not alter the enantiomer ratio measured for **15**.
9. Hartwell JL. Org Synth. 1955Coll. 3:185–187.
10. (a) Basu A, Beak P. J Am Chem Soc. 1996; 118:1575–1576.(b) Thayumanavan S, Basu A, Beak P. J Am Chem Soc. 1997; 119:8209–8216.
11. Wang HMJ, Lin IJB. Organometallics. 1998; 17:972–975.
12. Arienti A, Bigi F, Maggi R, Marzi E, Moggi P, Rastelli M, Sartori G, Tarantola F. Tetrahedron. 1997; 53:3795–3804.
13. Cherian AE, Domski GJ, Rose JM, Lobkovsky EB, Coates GW. Org Lett. 2005; 7:5135–5137. [PubMed: 16268521]

14. Berthon-Gelloz G, Siegler MA, Spek AL, Tinant B, Reek JNH, Markó IE. Dalton Trans. 2010; 39:1444–1446. [PubMed: 20104298]
15. Alternatively, the same interconversion could be achieved by rotation around all four aryl-benzylic carbon bonds. Due to A^{1,3}-strain, these bonds are likely to have a very high rotational barrier.
16. Simmons EM, Hartwig JF. J Am Chem Soc. 2010; 132:17092–17095.
17. (a) Lorenz C, Schubert U. Chem Berichte. 1995; 128:1267–1269. (b) Mankad NP, Laitar DS, Sadighi JP. Organometallics. 2004; 23:3369–3371.
18. Díez-González S, Kaur H, Zinn FK, Stevens ED, Nolan SP. J Org Chem. 2005; 70:4784–4796. [PubMed: 15932319]
19. Díez-González S, Scott NM, Nolan SP. Organometallics. 2006; 25:2355–2358.
20. Baker, J.; Wolinski, K.; Janowski, T.; Saebo, S.; Füsti-Molnár, L.; Magyarfalvi, G.; Pulay, P. Parallel Quantum Solutions LLC. Fayetteville, AR: <http://www.pqs-chem.com>
21. Jongsma C, Bickelha F. Recueil. 1973; 92:1143–1151.
22. Tondreau AM, Lobkovsky E, Chirik PJ. Org Lett. 2008; 10:2789–2792. [PubMed: 18537247]

Figure 1.
Common NHC Structural Motifs

Figure 2.
The crystal structure of IPrCuCl.³

Figure 5.
NHC-Precursors and Metal Carbenoids

Figure 6.
Crystal structure of **9**

Figure 7.
Crystal structure of **10**

Figure 8.
Crystal structure of **11**

Figure 9.
Analytical results of the preparative separation of the stereoisomers of **28**

Figure 10.
Rotamers of **30-meso**

Figure 11.
Crystal structure of (S,S,S,S)-12

Figure 12.
DFT Minimization of Intermediate B

Scheme 1.
Retrosynthesis plan

Scheme 2.
Synthesis of Enantioenriched Aniline **16**

Scheme 3.
Synthesis of Chiral Diamine **18**

Scheme 4.
Synthesis of Imidazolinium **9**

Scheme 5.
Synthesis of Enantioenriched Aniline **22**

Scheme 6.
Synthesis of Imidazoliums **10** and **25**

Scheme 7.
Synthesis of Silver Carbenoid **11**

Scheme 8.
DTR of 2,6-diethylpivaniide

Scheme 9.
Syntheses of **28** as a Mixture of Stereoisomers

Scheme 10.

Scheme 11.
Synthesis of Copper Carbenoid **12**

Scheme 12.
Hydrosilylation of acetophenone

Scheme 13.
Hydrosilylation catalytic cycle¹⁸

Table 1

Hydrosilylation of acetophenone

Catalyst	Time (h)	Silane	Product	%conversion [*]	er (S:R)
9	24	Ph ₂ SiH ₂		29	60:40
25	4	Ph ₂ SiH ₂		95	73:27
12	0.75	Ph ₂ SiH ₂		100	98:2

^{*} Determined by GC