A common SacI polymorphism in the gene for the M1 subunit of ribonucleotide reductase (RRM1)

L. Battat and U. Francke

Department of Human Genetics, Yale University School of Medicine, PO Box 3333, New Haven, CT 06520, USA

SOURCE AND DESCRIPTION OF CLONE: A 1.2 kb cDNA from the 3' end of the mouse RRM1 gene (1).**POLYMORPHISM:** The SacI restriction site polymorphism has two alleles: A1 (15 kb), A2 (10 kb / 5 kb). In addition, two larger invariant fragments (25 and 18 kb) are present (Figure 1).**FREQUENCIES:** In 15 unrelated Caucasians the allele frequencies were A1: 0.53 and A2: 0.47.**NOT POLYMORPHIC FOR:** BamHI, DraI, EcoRV, HincII, HindIII, KpnI, NsiI, PstI, PvuII, RsaI and TaqI in 8-10 unrelated Caucasians.**CHROMOSOMAL LOCALIZATION:** In situ hybridization to 11p15 (2) has confirmed the earlier assignment to 11p by expression studies in somatic cell hybrids (3).**MENDELIAN INHERITANCE:** Co-dominant inheritance was shown in one three generation family with 13 members.**PROBE AVAILABILITY:** Contact Ingrid Caras, Genentech Inc., South San Francisco or Lars Thelander, Univ. of Umea, Sweden.

- REFERENCES:** (1) Caras et al. (1985) J. Biol. Chem 260, 7015.
(2) Brissenden et al. (1988) Exptl. Cell Res. 174, 302-308.
(3) Engström Y. and Francke U. (1985) Exptl. Cell Res. 158, 477-483.

ACKNOWLEDGEMENTS: We thank Ingrid Caras for the mouse M1 cDNA probe. Supported by NIH research grant GM26105.

Figure 1
Lane 3: A1/A1
Lane 7: A2/A2
all other lanes: A1/A2