Published in final edited form as:

J Adolesc Health. 2011 October ; 49(4): 337–346. doi:10.1016/j.jadohealth.2011.01.005.

Who decides? Decision making and Fertility Preservation in Teens with Cancer: A Review of the literature

Gwendolyn P. Quinn, Ph.D.^{*,1,2}, Devin Murphy, MSW², Caprice Knapp, Ph.D.³, Daniel K. Stearsman, Pharm.D.², Kathy L. Bradley-Klug, Ph.D.⁴, Kelly Sawczyn, MD⁵, and Marla L. Clayman, Ph.D., M.P.H.⁶

¹College of Medicine, University of South Florida, Tampa, FL, United States

²Health Outcomes and Behavior, H. Lee Moffitt Cancer Center and Research Institute, Tampa, FL, United States

³College of Medicine, University of Florida, Gainesville, FL, United States

⁴College of Education, University of South Florida, Tampa, FL, United States

⁵All Children's Hospital, St. Petersburg, FL, United States

⁶Center for Communication in Healthcare, Northwestern University Feinberg School of Medicine, Robert H. Lurie Comprehensive Cancer Center, Chicago, Illinois, United States

Abstract

Purpose—The knowledge that cancer treatment may impair fertility in pediatric populations is an emerging aspect of quality of life in this population. However, decision making and use of fertility preservation among adolescent cancer patients and their families has not been well studied. This review summarizes the available literature on aspects of decision making and fertility preservation in adolescent cancer patients.

Methods—An electronic search was performed to identify peer reviewed studies published between 1999-2009 using key MESH terms and inclusion criteria. Inclusion criteria limited eligible studies to those that focused on adolescent decision-making in cancer treatment or fertility preservation, fertility concerns in pediatric oncology, capacity for decision-making, and health decision making in pediatrics. Studies were excluded that did not meet at least one of these criterion.

Results—A total of 29 articles were reviewed and summarized. Three categories of results were seen: a focus on adolescent decision making in oncology, decision making in chronic illness, and decision making in cancer related infertility and preservation.

Conclusion—The majority of studies showed adolescents have a strong desire to participate in decisions about their cancer treatment and many have concerns regarding their future fertility although barriers often prevented these discussions. More research is needed to explore the role of teens and parents in decisions about fertility in relation to cancer treatment.

© 2011 Society for Adolescent Medicine. Published by Elsevier Inc. All rights reserved.

*Corresponding Author: Gwendolyn Quinn, 12902 Magnolia Drive, MRC CANCONT, Tampa FL 33612, 813-745-1359, gwen.quinn@moffitt.org.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Keywords

Fertility; Decision Making; Teens; Cognition; Cancer

Approximately 16 cases per 100,000 people aged 0-19 are newly diagnosed with cancer in the US.[1] Fortunately, medical advances have resulted in improved survival and the average 5-year survival rate for pediatric cancer is currently 78%.[2] Cancer during adolescence presents challenges regarding treatment and late effects, and as the population grows, the focus has shifted toward assessments and interventions that improve the quality of survivorship. Research on quality of life during survivorship generates options to improve daily living while regaining a sense of normalcy. However, these options also give rise to the dilemma of the adolescent's role in making decisions regarding long-term survivorship issues, like infertility.

Cancer related infertility is an important issue because decisions about preserving future fertility will impact the adolescent as an adult survivor, though these decisions may not seem relevant during the time of treatment. While cancer treatments such as chemotherapy, radiation, or surgery are designed to save lives, there are risks of side effects such as infertility. The likelihood of infertility is dependent on age, type of therapy, as well as site, stage, and grade of the cancer.[3,4]

Infertility affects the future quality of life of cancer survivors of reproductive age and is a predictor of stress in current and future relationships. Janson et al. (2009) showed that even a perceived loss of fertility resulted in the reduced likelihood of being married and a higher likelihood of divorce.[5] Some adolescent survivors have revealed that they do not disclose their actual or suspected infertility for fear of rejection.[6] These concerns were recognized in 2006 when the American Society of Clinical Oncology (ASCO) published guidelines suggesting all newly diagnosed cancer patients of childbearing age be informed about potential loss of fertility and receive referrals to infertility specialists.[7]

Fertility preservation (FP) technologies are available for adolescent cancer patients and in most cases should be pursued prior to treatment. The established (non-experimental) option for males is sperm cryopreservation, and for females the established options are ovarian transposition and embryo cryopreservation. There are experimental options available for both males and females including oocyte, testicular, and ovarian tissue cryopreservation. [8,9] In some cases, the physician may decide an adolescent patient is unable to delay cancer treatment in order to pursue an FP option. [9] However, the need to consider FP depends on physical and developmental capacity, prognosis, financial circumstances, and religious, cultural, and ethical values.

The values and duties of the relevant parties and the potential conflicts that may arise must be examined. The Ethics Committee of the American Society for Reproductive Medicine (ASRM) extended physicians' fiduciary duty in a 2005 statement noting that, "Physicians should inform cancer patients about options for FP and future reproduction prior to treatment." [3] In this view, they opine that preserving future reproductive choices and making accommodations so that later autonomous decisions may be made, is in the best interest of the preautonomous minor with cancer. The principle of nonmaleficence limits potential harm to future fertility and reduces the threat of eliminating these later autonomous choices.

Conflicting evidence exists as to whether adolescents possess the capacity to make health-related decisions that they will be satisfied with in the future. Fueling this conflict is the lack of standardized definitions of children, minors, adolescents, and young adults. The National

Cancer Institute (NCI) defines children as persons between the ages of 0-14, and adolescents between 15 and 19.[10] The World Health Organization (WHO) defines adolescents between the ages of 10 and 19 [11] and the Center for Disease Control and Prevention (CDC) defines adolescents as 12 to 19.[12] This has implications for cancer registries and national data collection efforts. More recently, the term AYA (Adolescent/Young Adult) has been suggested for persons 15-39 years of age.[13-15]

In the U.S., individuals are considered minors until 18, however recent studies have challenged this.[16-19] There is disagreement regarding an adolescent's competency to make healthcare decisions with long-term consequences, though research has shown that adolescents have the ability and desire to participate in their own medical decisions. [20-22]

Adolescent brain development contributes to the law for legal age of consent.[23] Normal brain development posits the prefrontal cortex as a necessary region for decision making. [24-27] The amygdala, part of the limbic system and responsible for regulating emotions, grows at a much faster rate than the prefrontal cortex. Some studies of adolescent brain development show that judgment and responsibility do not increase until age 19, suggesting younger adolescents may be less competent decision-makers and are more likely to make decisions based on emotions versus facts.[28]

Theories on the biological capabilities of adolescent decision making has an impact in healthcare.[29] There is research that addresses the components of adolescent decision making with cancer treatment, fertility, and reproduction; however, studies combining these topics are limited. This is an important issue considering retrospective studies on survivors of childhood cancer suggest a reduced quality of life and increased regret for those who experienced a loss of fertility.[30] Coping skills and interactions within adult relationships can be directly affected by the loss of fertility.[5] The objective of this review is to summarize the available literature on decision making aspects of FP among adolescent oncology patients and recommend strategies for participation.

Methods: Search Strategy

An electronic search was performed to identify studies addressing adolescent oncology patients and fertility decision-making. Databases searched included Science Direct, Wiley Interscience, PsycINFO and Pubmed. Search criteria were: peer reviewed prospective studies published between 1999-2009, in English, and limited to the age categories of 6-12 years and 13-18 years. Terms searched were:

[adolescent AND decision making AND ability OR cognitive development OR culture AND neoplasms OR consent AND cancer AND treatment AND participation OR cancer treatment AND Fertility OR decision making AND reproduction AND chronic disease OR illness OR health AND pediatric]

Given the inconsistencies in defining adolescents, we chose to focus this paper on the WHO definition (ages 10-19). It should be noted that some studies included subjects younger than 10. For example, one article included participants ages 8 to 17. In those cases, we still included these articles in our review. We acknowledge that this may bias our conclusions, but it also reiterates the need for standardization.

Selection Strategy

Limiting the search to the last ten years ensured this review summarized the most current literature, especially with the onset of new FP technologies that may influence an adolescent's outlook on decision making. Initially, 226 titles were identified. The retrieved

studies were subjected to 4 inclusion criterion agreed upon by GQ and DM to encompass the interplay between factors related to FP and decision making. Each study was focused on **at least one** of the following: adolescent decision making in cancer treatment or FP; fertility concerns in pediatric oncology; cognitive development and capacity for decision making; and health decision making in pediatrics. Review articles were excluded (n=56), while 141 additional articles were excluded that did not meet the inclusion criteria. Overall, 29 studies were reviewed. Information on the purpose, method, and results were extracted (See Table 1).

Results

Twenty-nine studies were reviewed by the research team. Factors regarding adolescent decision-making and fertility preservation in oncology emerged as: 1) Factors unique to oncology, 2) Similarities between adolescents with cancer and other chronic illnesses, and 3) Factors specific to fertility preservation. The available literature was organized into these three categories.

Adolescent Decision Making in Oncology

Ten articles focused on participation in oncology decisions. A 2007 qualitative study showed six oncology patients aged 14-21 years unanimously wished to participate in their healthcare decisions; however, did not feel included in this process. The majority felt their health care providers did not provide any or adequate information about their treatment or side effects.[31]

In a study of 25 cancer patients and survivors ages 8-17 participating in online focus groups, the majority expressed a strong desire to have information on their disease and statistics on survival; however, results varied on desire for their parent's involvement in making decisions.[32] Thus adolescent patients wish to be well-informed regardless of whether or not they are the primary decision-maker.

Further, factoring in religious beliefs may have significant ethical impacts on the adolescent such as two cases of native Chinese cancer patients ages 15 and 17. Despite the adolescents' desire to engage in standard care, their parents refused treatment due to their cultural beliefs. Both situations resulted in negative health consequences.[33] The turbulence of a cancer diagnosis can overshadow an adolescent's independent desires. In some situations a court may override decisions made in agreement by parents and the minor.[34] Such was the recent case of Daniel Hauser in May, 2009. Daniel, a 13-year old Leukemia patient, and his parents desired alternative treatments that were in line with their religious beliefs in lieu of chemotherapy. Minnesota courts overruled the family's authority and Daniel was treated with modern medicine.[34]

Conflict may arise over an adolescent's competency to make decisions during a time of crisis such as a cancer diagnosis. Some studies indicate that adolescent cancer patients have struggled to plan appropriately for the future. Hokkanen et al. (2004) reported that of 20 adolescents aged 13 to 18 years attending a cancer adjustment camp, the majority made little if any attempt to plan for the future.[35] If an adolescent treated for cancer does exhibit difficulties for long-term planning parents may be less likely to allow autonomous decision-making. Desires for autonomous decision making was assessed in a 2009 study of cancer survivors aged 11-18 and their views on whether a physician should adhere to an adolescent's desire for legal euthanasia. It was concluded that the patient's own experience with illness impacted decision-making and planning when considering end-of-life care, therefore not all adolescent oncology patients are ill-equipped for long term planning. Over half of cancer-surviving adolescents felt that their decision for legal euthanasia should be

respected by the physician, though they identified safeguards that should be in place. Such safeguards include that the adolescent should be well-informed about their prognosis and process of euthanasia, parents must be consulted, and the adolescent cannot make this decision if exhibiting symptoms of depression. Fifty percent of adolescent survivors felt that their decision for euthanasia should still be respected even if the parent does not agree.[36]

An adolescent's desire for joint decision-making cannot always be predicted, though some studies have assessed this. Tyc et al. (2001) found that among 46 adolescent cancer survivors ages 10-18, decisions for health-protective, or less risky health behaviors were more predictable in younger subjects.[37] Comparatively, Geller et al. (2003) report that among adolescents with a family history of cancer ages 10-17, the largest predictors of joint decision-making to enroll in clinical trials were age and the adolescent- parent relationship. The majority of adolescents emphasized that the final decision should be their own.[38] This shows that an adolescent's desire to participate in healthcare decisions may not always be predictable and preferences should be inquired directly.

Not all adolescent oncology patients desire autonomous or participatory decision-making. Stegenga & Ward-Smith (2008) found from three individual interviews with adolescents between 13-15 years old diagnosed with cancer, each felt that decisions were best left to the healthcare team. Despite this, all three agreed on the need for information and to be kept informed during the initial six months of treatment.[39] This is supported by a 2004 study wherein the majority of 72 adolescents with cancer rated their desire for information as high, stressing the need to include adolescents in initial and ongoing discussions related to treatment.[40]

Adolescent Decision Making with Chronic Illnesses

Oncology is not the only area where adolescent decision-making is controversial. Adolescents with chronic illnesses may also be faced with healthcare decisions. There are inconsistencies in the literature regarding their decision-making capabilities, desires, and satisfaction with participation during a time of illness, and how this is generalizable between adolescents with chronic illnesses and adolescents with cancer. Nine articles were reviewed on this topic.

Like many adolescents with cancer, chronically ill adolescents have also shown a desire to participate in decision-making. Ninety-six percent of 25 chronically ill adolescents aged 13-21 years desired to participate in treatment decision-making, and 80% believed their doctors would adhere to their decisions.[41] In two studies that interviewed a combination of pediatricians, parents, and adolescents with chronic conditions, the child was most often found not to have an equal part in decision-making, despite a desire to do so.[42,43] In Garth et al. (2003), the majority of patients with cerebral palsy aged 8-12 (n=10) desired to participate in treatment decision-making; however they were not included as an equal partner with parents and physicians.[43] Broome and Richards (2003) assessed decision-making among 34 individuals ranging from 8-22 years old with chronic illnesses. Their results indicate a desire to be involved in their own medical decisions and the ability to communicate their decisions articulately. The majority expressed that they had 'faith' in their parents to adhere to their decisions.[21]

Parents may allow participatory decision-making at their discretion. Runeson et al. (2002) conducted an observational study of 24 in-patients, ages 5 months to 18 years, for 135 total hours. There was a distinction between participatory decision-making among older patients aged 11-18 (n=5) versus younger patients aged 5 months to 10 (n=19). The parent refused their child's decisions or did not consult with them only one time in the older age group versus nine times in the younger age group. Of the 21 situations classified as poor

communication between parent and patient, the lowest level of participation for both groups was in the treatment decision-making process.[44] Parents also have discretionary authority to involve their child in the decision to enroll in clinical trials. Of 81 chronically ill minors ages 7-14, 90% of believed they should be involved in making research enrollment decisions, contrasted with 62% of parent respondents who believed children should be involved.[45]

Despite much of the available literature indicating an adolescent's preference for shared decision-making, Knopf et al. (2008) found that only 36% of 82 adolescents with chronic illnesses preferred shared decision-making, while 17% preferred sole decision-making authority. The highest rating was actually passive decision making at 46%; further evidence that parents and physicians need to inquire directly about the adolescent's preference.[46]

Quality decision-making stems from open communication, yet many barriers exist. In a 2003 study of 63 adolescents aged 11-16 years with a chronic condition, adolescents identified communication barriers with physicians as: frequency of contact, gender, physician's perceptions of adolescents, and the presence of parents and medical students. [47] The unease of parent's presence in the examining room was corroborated by another study with 13 families, including patients aged 8-17. Many adolescents were happy with their parents' involvement, but some identified their parent as restrictive.[48] Parent or physician presence may cause the adolescent to feel they have limited decision-making authority. Even if the adolescent does communicate their concerns, they are likely to feel discouraged if these concerns are overlooked. Coyne found that the majority of children and adolescents treated for chronic conditions (n=11) explained that despite wanting physicians to consult them about treatment decisions, many expressed feeling "ignored." [42]

Adolescent Decision making and Fertility Concerns

The public is becoming increasingly aware of the effects of cancer treatments on fertility. Despite this, little is known about what information is and should be provided to adolescent patients. Can adolescent cancer patients make decisions about consenting or refusing FP procedures that they will not regret in adulthood? Long-term studies of pediatric cancer survivors and their satisfaction with FP decisions made in childhood are rare. Nine articles on the concerns of fertility-related decision-making within pediatric oncology were reviewed.

Impact on fertility and FP options have often been neglecting during discussions of an adolescent's cancer treatment. Oosterhuis et al. (2008) report that less than 30% of parents of pediatric oncology patients over the age of 14 were satisfied with the amount of fertility information disclosed by their child's healthcare professional (n=134). Less than 11% recalled being given information regarding FP options, over 50% wanted to discuss FP with their oncologist or a specialist, and over 60% of adolescents wanted educational material. Approximately 50% of parents incorrectly affirmed that all cancer therapies cause damage to reproduction [49] indicating the need for education.

Absent or ambiguous FP discussions with adolescents may preclude them from being involved in decision-making. Edge et al. (2006) examined survey responses from 45 male cancer patients and survivors aged 12-21 that had fertility discussions documented in their hospital notes. Seventy-four percent of males did not have any documentation. Among the 26% with documentation, respondents indicated they had a higher concern for future fertility during the time of the interview, despite the fact that most had been off therapy about 2 years. Barriers that had an impact on the decision-making process to sperm bank included anxiety related to the cancer diagnosis, embarrassment, and feeling 'emotionally drained.' When asked what changes could be made to benefit future patients, subjects recommended

having more detailed FP discussions as many felt they had an inadequate understanding of how FP procedures were carried out.[50]

Not all studies have indicated low rates of FP discussions. Burns et al. (2006) found that of 50 female adolescent cancer patients and their parents, fertility was discussed in 49% and 69% of cases, respectively. Adolescent age groups were separated 10-14 and 15-21 years, and only 25% of the younger group indicated they had fertility discussions but were found just as likely as the older group to have fertility concerns. The majority of patients in both groups report they frequently think about having children in the future. Moreover, 94% said they were aware of the concept of infertility, but despite this less than 20% of parents and 30% of adolescents indicated they would delay treatment to undergo FP methods.[51] The authors argue in support of adolescents' competency to make FP decisions and the discussion of FP options.

Survivorship studies document the concern and struggles with having children post-treatment; however, adolescents' concern over future fertility at the time of diagnosis is not well-documented. Only one qualitative study examined survivors aged 16-30 and their concern with possible infertility (n=38). Sixteen participants were between the ages of 13 and 21 however age did not prove significant within the results. The majority reported at the time they were diagnosed having a strong desire to know as soon as possible how their fertility would be impacted. No participant had utilized FP options. The majority who were aware of their impaired fertility status were concerned about "finding a partner willing to enter a potentially childless relationship." The majority were pleased with their parents' involvement in the decision-making process, however a small number of males preferred to make the decision to sperm bank on their own.[30] This conflicts with another study of male adolescent oncology patients (aged 15-19), in which parents were reported to have significant influence on sperm banking, with 80% making joint-decisions with their sons. Similar to the previous study, almost half of patients said they preferred to have initial discussions without their parent present.[52]

At times, the pediatric oncologist may be a barrier to FP discussions. In a qualitative study of 21 males aged 16-26 who had either completed or were still in active treatment for cancer, the majority indicated a high concern about future fertility; however, felt the presence of many barriers to initiating these conversations. Individual interviews yielded concerns such as presenting sperm banking as a choice and having a desire for counseling but not being offered. Some males described their own unease of fertility discussions as the main barrier.[53] Schover et al. (2002) assessed the likelihood of oncologists recommending sperm banking with male patients and found that 37% felt parental consent should be required for a teen to sperm bank, and only 9% said they would initiate the conversation with the teen alone.[54] Gender may also influence the prevalence of fertility related discussions. A recent study of pediatric oncologists showed physicians perceived girls to be more open to discussing fertility than boys; however girls showed more characteristics of distress than boys.[55] Anderson et al. (2008) found that in a sample of 1,030 boys and girls with cancer in the U.K. aged 1-22, only 1% of post-pubertal girls were referred to a FP clinic. Comparatively, 83% of post-pubertal boys were referred for sperm cryopreservation, however this fell to 39% for pre-pubertal boys.[56]

Discussion

This review summarizes many issues related to adolescent decision-making regarding FP. Both the ASCO and AAP guidelines recommend oncologists discuss cancer treatments' impact on fertility with patients, regardless of age. These guidelines are strong statements;

however, they are not laws. Barriers surrounding these discussions may overshadow the guidelines.[55,57,58]

Research on an adolescent's desires for their parent to solely manage communication with healthcare providers is inconclusive.[40,59] Ellis & Leventhal showed that 63% of adolescent oncology patients wanted to make their own decisions about palliative therapy. Thirty-nine percent of patients and 16% of their parents however, viewed that decisions about curative therapies was not the role of the pediatric patient.[60] Fifty-eight parents of children with cancer under the age of 13 were surveyed for their preferences in participation in their children's treatment. Most preferred a collaborative role, while the second-most common preference was passivity.[61] In 2004, The Children's Voices Project from the Commission for Health Improvement in the UK compiled from 59 reports of children and adolescents admitted to hospitals. Most of the children and adolescents were explained treatment options using language they did not understand, not provided any information, and not included in the decision-making process.[62]

Teens may be selective about whom they wish to discuss fertility concerns with. In a 1999 study female teens more frequently discussed the physical aspects of puberty with their mothers, discussed refraining from sexual activity with their fathers, and discussed sexual intercourse with their friends.[63] Apprehension in discussing FP may be relevant if the child has not previously discussed sexuality with their parents. Further, because FP for males is less medically invasive and expensive than for females, gender can also influence the frequency of these discussions.[7] These barriers are likely to impact the adolescents' engagement in decision-making.

Scientific advances have made FP options available in cases previously not possible. Empirical data highlights the need for advocacy regarding the complexities of infertility related discussions and referrals between providers and patients. Adolescents provided developmentally appropriate language may be more equipped for long-term decision-making. Children as young as 11 have been found to understand their own illness in parallel to their parents' level of understanding.[64,65] Decision-making capacity of adolescents has been widely researched in neuroscience and behavioral contexts, but there is no agreement about whether adolescents can make sound, long-term decisions. Behavioral research has argued that though the brain has not fully developed, when provided with comprehensive information, adolescents can understand the ramifications of decisions.[39,66,67] Much of the research in this review indicates that adolescents prefer participatory decision-making concerning cancer treatment however these preferences are often not discussed. There is a clear lack of research on adolescents' decision-making preferences regarding FP options.

It is important to understand factors that may contribute to a parent's healthcare decisions on behalf of their child. Parents of terminally ill adolescents base many decisions on their past experiences with death and interpretation of their child's suffering.[68] Parents may be reluctant to include their child in decision-making, fearing he/she lacks the emotional strength to consider a variety of outcomes. Parents may also assume their child does not want to participate due to the overwhelming nature of a cancer diagnosis. If only parents are approached to consider treatment options, parents may assume the role as primary decision-maker instead of including the adolescent.[69-71]

Most teens rely on their parents to interpret and process information regarding serious health issues. Likewise, a parent's attitude towards FP options can impact the adolescent's view and perhaps the outcome. A 2002 study examined how the order of delivering information regarding risks and benefits can influence decisions. Participants hearing the risks after the benefits were twice as likely to refuse consent than participants who heard the benefits after

the risks.[72] An adolescent's decision may actually be a reflection of what the parent understands. Adolescents' decisions tend to reflect the values and morals of their parents and the healthcare system in which they are receiving care.[73] Thus, with regard to FP, if issues of infertility and cancer treatment have not been previously discussed, the teen may have little context to base a decision, or may understand their options as their parent understands them.

Evidence on adolescent decision-making in oncology suggests that FP is a suitable conversation for teens. The literature shows that adolescents with cancer have three main desires about fertility: for information on the potential impact to fertility by cancer treatment, for information on FP options, and to participate in making decisions. Clearly such desires mirror those of adults diagnosed with cancer. Despite that adolescents desire for information on fertility, young adult survivors of pediatric cancer have reported they were unsure about their fertility status and struggled to recall discussions on the impact of fertility as a result of treatment.[6]

Ethical questions may arise during the decision-making process if there are conflicting views between the child and the parent(s), child and provider, parents and provider, or between two parents. When autonomy is lacking, and parents do not know what a child would want for themselves, who will serve the best interest of the child and their potential offspring? Case law states that, "The child is not the mere creature of the state; those who nurture him and direct his destiny have the right, coupled with the high duty, to recognize and prepare him for additional obligations." [74] Tension may exist between a parent, who has a constitutional liberty to raise children as they see fit,[75] and case law that affirms a child's right to bear children.[76] At the state level the doctrine of *parens patriae* is invoked in cases when the state must intercede guardianship over minors if the parent is found negligent or incompetent.[77] Conversely, some states have adopted the *mature minors doctrine*, whereby an adolescent may independently provide consent for medical treatment based on age and maturity, subjectively determined by the court system.[78]

Concerns may continue as what to do with harvested eggs, tissue, or sperm if they are not used.[79] These considerations should be attended to at the time of selection of FP, weighed with the expressed desires of the patient, using a participatory decision-making process.[80]

This review synthesized the most current literature in order to better understand factors that affect the decision-making process regarding FP among adolescent oncology patients. We acknowledge several limitations. The 10 year age range limited the search results to a few eligible studies and some studies used small sample sizes [31,33,34,39]. Additionally, many studies varied widely on the ages that defined adolescence; therefore, it was left to subjective judgment on inclusion and this may bias our conclusions.

Conclusions

A variety of literature exists discussing decision-making, quality of life, and fertility among adolescents both with and without cancer. However, the quantity of empirical research is limited. This topic spans issues on health, sexuality, planning for the future, and decision-making in times of crisis. Existing literature evidences that adolescents prefer to discuss their cancer treatment's possible impact on fertility, yet further research is needed to explore how to determine personalized approaches for these patients. Additional FP areas of research include the impact of socioeconomic status and cost, decision-making strategies, satisfaction as adults with the decisions made as an adolescent, and reasons for not wanting information on FP. Appropriate timing for fertility related discussions is also an area of research that has been neglected.

Research is recommended to explore participatory decision-making regarding fertility by means of an assessment tool. Developing an assessment tool for use by health professionals would be beneficial by evaluating the adolescents': a) current health literacy, b) understanding of sexual reproduction, c) concern for future fertility, d) decision making competencies, and e) desire for physician/parental/legal guardian involvement. The tool will be best-served by pairing assessment items with AAP/ASCO guidelines. Determining how much an adolescent cancer patient knows, and is concerned with fertility, will help tailor discussions and language to fit the patient's needs.

Acknowledgments

This research was supported by the Oncofertility Consortium®, funded by the National Institutes of Health through the NIH Roadmap for Medical Research, Grant UL1DE19587 and UL1DE019587

References

1. Surveillance epidemiology and end results incidence statistics. [Accessed July 9, 2009]. 2009 Available from: <http://seer.cancer.gov/statistics/>
2. Osteosarcoma and Aligned Fibrous Histiocytoma of Bone. [Accessed July 9, 2009]. 2006 cited Available from: <http://www.cancer.gov/cancertopics/types/bone>
3. Lee SJ, Schover LR, Partridge AH, et al. American Society of Clinical Oncology recommendations on fertility preservation in cancer patients. *J Clin Oncol*. 2006; 24:2917–2931. [PubMed: 16651642]
4. Byrne J, Mulvihill JJ, Myers MH, et al. Effects of treatment on fertility in long-term survivors of childhood or adolescent cancer. *N Engl J Med*. 1987; 317:1315–1321. [PubMed: 3683460]
5. Janson C, Leisenring W, Cox C, et al. Predictors of marriage and divorce in adult survivors of childhood cancers: A report from the Childhood Cancer Survivor study. *Cancer Epidemiol Biomarkers Prev*. 2009; 18:2626–2635. [PubMed: 19815636]
6. Zebrack BJ, Casillas J, Nohr L, et al. Fertility issues for young adult survivors of childhood cancer. *Psychooncology*. 2004; 13:689–699. [PubMed: 15386645]
7. Fallat ME, Hutter J. Preservation of fertility in pediatric and adolescent patients with cancer. *Pediatrics*. 2008; 121:1461–1469.
8. Brinster RL, Zimmermann JW. Spermatogenesis following male germ-cell transplantation. *Proc Natl Acad Sci U S A*. 1994; 91:11298–11302. [PubMed: 7972053]
9. Lobo RA. Potential options for preservation of fertility in women. *N Engl J Med*. 2005; 353:64–73. [PubMed: 16000356]
10. Smith, MA.; Gurney, J.; Gloeckler-Ries, L. *Seer Pediatric Monograph*. the National Cancer Institute; Bethesda: 1999. Cancer among adolescents 15-19 years old.
11. World Health Organization. Adolescent Health and Development. 2009. cited August 30, 2010 Available from: http://www.searo.who.int/en/Section13/Section1245_4980.htm
12. Center for Disease Control and Prevention. Parent Portal. 2009. cited August 30, 2010 Available from: <http://www.cdc.gov/parents/>
13. Thomas DM, Seymour JF, O'Brien T, et al. Adolescent and young adult cancer: A revolution in evolution? *Intern Med J*. 2006; 36:302–307. [PubMed: 16650195]
14. Soliman H, Agresta SV. Current issues in adolescent and young adult cancer survivorship. *Cancer Control*. 2008; 15:55–62. [PubMed: 18094661]
15. Pollock BH, Birch JM. Registration and classification of adolescent and young adult cancer cases. *Pediatr Blood Cancer*. 2008; 50:1090–1093. [PubMed: 18360832]
16. Hockenberry-Eaton M, Kline NE. A minor's consent to treatment: An ethical dilemma. *J Pediatr Health Care*. 1995; 9:282–284. [PubMed: 8699315]
17. Hickey K. Minors' rights in medical decision making. *JONAS Healthc Law Ethics Regul*. 2007; 9:100–104. [PubMed: 17728582]
18. Hartman RG. Coming of age: Devising legislation for adolescent decision-making. *Am J Law Med*. 2002; 28:409–453. [PubMed: 12516175]

19. Grisso T, Vierling L. Minors' consent to treatment: A developmental perspective. *Prof Psychol.* 1978; 9:412–427. [PubMed: 11662545]
20. Weithron LC, Susan. The competency of children and adolescents to make informed treatment decisions. *Child Dev.* 1982; 53:1589–1598. [PubMed: 7172783]
21. Broome ME, Richards DJ. The influence of relationships on children's and adolescents' participation in research. *Nurs Res.* 2003; 52:191–197. [PubMed: 12792260]
22. Kaser-Boyd N, Adelman HS, Taylor L. Minors' ability to identify risks and benefits of therapy. *Prof Psychol Res Pr.* 1985; 16:411–417. [PubMed: 11650746]
23. What are the implications of adolescent brain development for juvenile justice?. Coalition for Juvenile Justice Emerging Concepts Brief. [Accessed November 3, 2009]. 2006 Available from: http://www.juvjustice.org/media/resources/public/resource_134.pdf
24. Dahl R. Adolescent brain development, a period of vulnerabilities and opportunities. Keynote address. *Ann N Y Acad Sci.* 2004; 1021:1–22. [PubMed: 15251869]
25. Giedd JN, Blumenthal J, Jeffries NO, et al. Brain development during childhood and adolescence: A longitudinal MRI study. *Nat Neurosci.* 1999; 2:861–863. [PubMed: 10491603]
26. Sowell ER, Thompson PM, Holmes CJ, et al. In vivo evidence for post-adolescent brain maturation in frontal and striatal regions. *Nat Neurosci.* 1999; 2:859–861. [PubMed: 10491602]
27. Thompson PM, Giedd JN, Woods RP, et al. Growth patterns in the developing brain detected by using continuum mechanical tensor maps. *Nature.* 2000; 404:190–193. [PubMed: 10724172]
28. Cauffman E, Steinberg L. (Im)maturity of judgment in adolescence: Why adolescents may be less culpable than adults. *Behav Sci Law.* 2000; 18:741–760. [PubMed: 11180420]
29. Cox, E.; Smith, M. Look who's talking: Shared decision-making with children; Abstr Academy Health Meet; 2004; p. 21
30. Crawshaw, M.; Sloper, P. A qualitative study of the experiences of teenagers and young adults when faced with possible or actual fertility impairment following cancer treatment. University of York; York: 2006.
31. Palmer S, Mitchell A, Thompson K, et al. Unmet needs among adolescent cancer patients: A pilot study. *Palliat Support Care.* 2007; 5:127–134. [PubMed: 17578063]
32. Zwaanswijk M, Tates K, van Dulmen S, et al. Young patients', parents', and survivors' communication preferences in paediatric oncology: Results of online focus groups. *BMC Pediatrics.* 2007; 7:35. [PubMed: 17996108]
33. Hui E. Parental refusal of life-saving treatments for adolescents: Chinese familism in medical decision-making re-visited. *Bioethics.* 2008; 22:286–295. [PubMed: 18447864]
34. Miller DM. Legal matters: Protecting the right to choose CAM over conventional medicine: The case of Daniel, Colleen, and Anthony Hauser. *Alternative and Complementary Therapies.* 2009; 15:261–266.
35. Hokkanen H, Eriksson E, Ahonen O, et al. Adolescents with cancer: Experience of life and how it could be made easier. *Cancer Nurs.* 2004; 27:325–335. [PubMed: 15292729]
36. Pousset G, Bilsen J, De Wilde J, et al. Attitudes of adolescent cancer survivors toward end-of-life decisions for minors. *Pediatrics.* 2009; 124:1142–1148.
37. Tyc VL, Hadley W, Crockett G. Prediction of health behaviors in pediatric cancer survivors. *Med Pediatr Oncol.* 2001; 37:42–46. [PubMed: 11466722]
38. Geller G, Tambor ES, Bernhardt BA, et al. Informed consent for enrolling minors in genetic susceptibility research: A qualitative study of at-risk children's and parents' views about children's role in decision-making. *J Adolesc Health.* 2003; 32:260–271. [PubMed: 12667730]
39. Stegenga K, Ward-Smith P. The adolescent perspective on participation in treatment decision making: A pilot study. *J Pediatr Oncol Nurs.* 2008; 25:112–117. [PubMed: 18270308]
40. Decker C, Phillips CR, Haase JE. Information needs of adolescents with cancer. *J Pediatr Oncol Nurs.* 2004; 21:327–334. [PubMed: 15475469]
41. Lyon ME, McCabe MA, Patel KM, et al. What do adolescents want? An exploratory study regarding end-of-life decision-making. *J Adolesc Health.* 2004; 35:521–529.
42. Coyne I. Research with children and young people: The issue of parental (proxy) consent. *Children & Society.* 2009; 24:227–237.

43. Garth B, Murphy GC, Reddihough DS. Perceptions of participation: Child patients with a disability in the doctor-parent-child partnership. *Patient Educ Couns.* 2009; 74:45–52. [PubMed: 18760558]
44. Runeson I, Hallström I, Elander G, et al. Children's participation in the decision-making process during hospitalization: An observational study. *Nurs Ethics.* 2002; 9:583–598. [PubMed: 12449996]
45. Varma S, Jenkins T, Wendler D. How do children and parents make decisions about pediatric clinical research? *J Pediatr Hematol Oncol.* 2008; 30:823–828. [PubMed: 18989159]
46. Knopf JM, Hornung RW, Slap GB, et al. Views of treatment decision making from adolescents with chronic illnesses and their parents: A pilot study. *Health Expect.* 2008; 11:343–354. [PubMed: 19076663]
47. Beresford BA, Sloper P. Chronically ill adolescents' experiences of communicating with doctors: A qualitative study. *J Adolesc Health.* 2003; 33:172–179. [PubMed: 12944007]
48. Young B, Dixon-Woods M, Windridge K, et al. Managing communication with young people who have a potentially life threatening chronic illness: Qualitative study of patients and parents. *Br Med J.* 2003; 326:305. [PubMed: 12574042]
49. Oosterhuis BE, Goodwin T, Kiernan M, et al. Concerns about infertility risks among pediatric oncology patients and their parents. *Pediatr Blood Cancer.* 2008; 50:85–89. [PubMed: 17514741]
50. Edge B, Holmes D, Makin G. Sperm banking in adolescent cancer patients. *Arch Dis Child.* 2006; 91:149–152. [PubMed: 16174641]
51. Burns KC, Boudreau C, Panepinto JA. Attitudes regarding fertility preservation in female adolescent cancer patients. 2006:350–354.
52. Ginsberg JP, Ogle SK, Tuchman LK, et al. Sperm banking for adolescent and young adult cancer patients: Sperm quality, patient, and parent perspectives. *Pediatr Blood Cancer.* 2008; 50:594–598. [PubMed: 17514733]
53. Chapple A, Salinas M, Ziebland S, et al. Fertility issues: The perceptions and experiences of young men recently diagnosed and treated for cancer. *J Adolesc Health.* 2007; 40:69–75. [PubMed: 17185208]
54. Schover LR, Brey K, Lichtin A, et al. Oncologists' attitudes and practices regarding banking sperm before cancer treatment. *J Clin Oncol.* 2002; 20:1890–1897. [PubMed: 11919249]
55. Vadaparampil ST, Clayton H, Quinn GP, et al. Pediatric oncology nurses' attitudes related to discussing fertility preservation with pediatric cancer patients and their families. *J Pediatr Oncol Nurs.* 2007; 24:255–263. [PubMed: 17827491]
56. Anderson RA, Weddell A, Spoudeas HA, et al. Do doctors discuss fertility issues before they treat young patients with cancer? *Hum Reprod.* 2008; 23:2246–2251. [PubMed: 18614615]
57. Achille MA, Rosberger Z, Robitaille R, et al. Facilitators and obstacles to sperm banking in young men receiving gonadotoxic chemotherapy for cancer: The perspective of survivors and health care professionals. *Hum Reprod.* 2006; 21:3206–3216. [PubMed: 16887922]
58. Goodwin T, Oosterhuis BE, Kiernan M, et al. Attitudes and practices of pediatric oncology providers regarding fertility issues. *Pediatr Blood Cancer.* 2007; 48:80–85. [PubMed: 16572406]
59. Odigwe C. Children say they are not involved enough in their treatment. *BMJ.* 2004; 328:600. [PubMed: 15016670]
60. Ellis R, Leventhal B. Information needs and decision-making preferences of children with cancer. *Psychooncology.* 1993; 2:277–284.
61. Pyke-Grimm KA, Degner L, Small A, et al. Preferences for participation in treatment decision making and information needs of parents of children with cancer: A pilot study. *J Pediatr Oncol Nurs.* 1999; 16:13–24. [PubMed: 9989013]
62. Boylan, P. Children's voices project: Feedback from children and young people about their experience and expectations of healthcare. Commission for Health Improvement; United Kingdom: 2004.
63. Diiorio C, Kelley M, Hockenberry-Eaton M. Communication about sexual issues: Mothers, fathers, and friends. *J Adolesc Health.* 1999; 24:181–189. [PubMed: 10195801]
64. Koopman HM, Baars RM, Chaplin J, et al. Illness through the eyes of the child: The development of children's understanding of the causes of illness. *Patient Educ Couns.* 2004; 55:363–370. [PubMed: 15582342]

65. Deatrick J, Dickey S, Wright R, et al. Correlates of children's competence to make healthcare decisions. *J Clin Ethics*. 2003; 14:152–163. [PubMed: 14740350]
66. Bahadur G, Whelan J, Ralph D, et al. Gaining consent to freeze spermatozoa from adolescents with cancer: Legal, ethical and practical aspects. *Hum Reprod*. January 1.2001 16:188–193. 2001. [PubMed: 11139561]
67. Clayman M, Galvin K, Arntson P. Shared decision making: Fertility and pediatric cancers: Oncofertility fertility preservation for cancer survivors. *Cancer Treat Res*. 2007; 138:149–160. [PubMed: 18080663]
68. Sharma M, Meert KL, Sarnaik AP. What influences parents' decisions to limit or withdraw life support? *Pediatr Res*. 2004; 56:672.
69. Stewart JL, Pyke-Grimm KA, Kelly KP. Parental treatment decision making in pediatric oncology. *Semin Oncol Nurs*. 2005; 21:89–97. [PubMed: 15991660]
70. Spinetta JJ. Adjustment in children with cancer. *J Pediatr Psychol*. 1977; 2:49–51.
71. Manne SL, Jacobsen PB, Gorfinkle K, et al. Treatment adherence difficulties among children with cancer: The role of parenting style. *J Pediatr Psychol*. 1993; 18:47–62. [PubMed: 8463933]
72. Bergus GR, Levin IP, Elstein AS. Presenting risks and benefits to patients: The effect of information order on decision making. *J Gen Intern Med*. 2002; 17:612–617. [PubMed: 12213142]
73. Susman EJ, Hersh SP, Nannis ED, et al. Conceptions of cancer: The perspectives of child and adolescent patients and their families. *J Pediatr Psychol*. 1982; 7:253–261. [PubMed: 7131213]
74. *Pierce v. Society of Sisters*. 268 U.S. 510 1925. p. 534-35.
75. Litton P. Non-Beneficial Pediatric Research and the Best Interests Standard: A Legal and Ethical Reconciliation. *Yale J Health Policy Law Ethics*. 2008; 8:359–420. [PubMed: 19049018]
76. Dolin G, Roberts DE, Rodriguez LM, Woodruff TK. Medical hope, legal pitfalls: potential legal issues in the emerging field of oncofertility. *Cancer Treat Res*. 2010; 156:111–34. [PubMed: 20811829]
77. Griffith DB. The best interests standard: a comparison of the state's parens patriae authority and judicial oversight in best interests determinations for children and incompetent patients. *Issues Law Med*. 1991; 7:283–338. [PubMed: 1765451]
78. Schlam L, Wood J. Informed consent to the medical treatment of minors: Law and practice. *Health Matrix*. 2000; 10:141–174. [PubMed: 14655526]
79. Cohen CB. Ethical issues regarding fertility preservation in adolescents and children. *Pediatr Blood Cancer*. 2009; 53:249–253. [PubMed: 19340857]
80. Savedra M, Tesler M. Coping strategies of hospitalized school-age children. *West J Nurs Res*. 1981; 3:371–384. [PubMed: 6916567]

Table 1**Adolescent Decision Making in Oncology (n=10)**

Study	Purpose	Method	Sample	Results
Decker et al., 2004 [40]	This study describes how adolescents with cancer rate the importance of specific cancer-related information, analyzes gender and age group differences, and compares 2 different adolescents with cancer groups	Questionnaire	N=201 aged 11-21	Adolescents with cancer rated the need for information as high. There were no significant differences by age, but females had significantly higher total scores for both groups and for many item means in the newly diagnosed group. Qualitative analysis of the additional write-in items generated 4 themes: treatment/side effects, uncertainty, social issues, and personal/emotional issues
Geller et al., 2003 [38]	To better understand the process by which families at increased risk of disease would decide to enroll their children in genetic susceptibility research	Face-to-face, audio-taped, semi-structured interviews	N=74 Parents and children 10-17	In general, the older/more mature the child, the less risky the research and the more open the communication style, the greater the likelihood that decisions would be made jointly. Most children wanted some parental input, but still thought the final decision should be theirs.
Hokkanen et al., 2004 [35]	To understand what adolescents with cancer think about their life situation, the support they get, and the information they receive about their illness	Focus groups (3)	N=60 aged 13-18 Three focus groups, N=20 participants each	Adolescents made very little, if any, conscious effort to plan ahead for the future. As for the adolescents' chances to take part in making decisions about their care and life, the analysis yielded 6 categories: joint decision making, inadequate chances for decision making, independent decision making, illusion of decision making, reluctant to make decisions, and excluded from decision making.
Hui, 2008 [33]	This paper reports two cases in Hong Kong involving two native Chinese adolescent cancer patients (APs) who were denied their rights to consent to necessary treatments refused by their parents, resulting in harm	Observational	N=2 aged 15-17	Specifically for adolescent cancer patients, we suggest that parental authority and family integrity should be re-interpreted in terms of parental responsibility and the enhancement of children's interests respectively.
Miller, 2009 [34]	To present a case study on parent refusal of treatment for their child if it is deemed as the court to be in the	Secondary data analysis	N=1 aged 13	Judge ordered that family cannot refuse chemotherapy for their 13 year old son and

Study	Purpose	Method	Sample	Results
	child's best interest			mother was arrested for evading police.
Palmer et al., 2007 [31]	This study reports the outcomes of a focus group conducted to explore issues of unmet physical, psychological, and social needs of adolescents with cancer	Focus group	N=6 mean age 17	The participants identified a number of concerns relating to three broad categories: information provision, treatment processes, and survivorship. This indicates gaps in current cancer services provided to adolescents.
Pousset et al., 2009 [36]	To investigate the attitudes of adolescent cancer survivors toward end-of-life decisions with life shortening effects, including decisions for non-treatment, alleviation of pain, euthanasia, and how illness influences these decisions	Questionnaire	N=83 aged 11 to 18	In terminal situations, 70% to 90% found requests for non-treatment acceptable, 84% requests for APS, and 57% to 64% requests for euthanasia. Frequently cited reasons for holding back physicians from administering a lethal drug to a child were the child not being well informed about his or her condition (92%) and the parents' opinion not being asked (92%). Cancer survivors were more accepting toward requests for non-treatment decisions and alleviation of pain and symptoms in terminal situations.
Stegenga & Ward-Smith, 2008 [39]	Few research studies have obtained data directly from the adolescent with cancer regarding decision making. The purpose of this pilot, or exploratory, study was to determine whether the research question is appropriate and whether participants between 13 and 15 years of age were capable of providing rigorous data	Semi-structured interviews	N=3 aged 13-15	The results suggest that treatment decision making was not a concern for these patients and that they were capable and appropriate study participants. Infusing information and education into the health care plan from time of diagnosis may allow the adolescent to become knowledgeable about the disease process. This may enable him or her to participate in health care treatment decisions, when appropriate.
Tyc et al., 2001 [39]	To examine sociodemographic and health perception variables as predictors of health-protective behaviors in pediatric cancer survivors	Cross sectional survey	N=46 aged 10--18	Results indicated that the practice of health-protective behaviors was best predicted by patient's age and socioeconomic status (SES); younger adolescents and patients from higher SES more frequently engaged in healthy behaviors. Survivors perceived themselves to be vulnerable to health problems.
Zwaanswijk et		Cross sectional survey	N=36 aged 8–17	

Study	Purpose	Method	Sample	Results
al., 2007 [32]	To investigate the communication preferences of childhood cancer patients, parents, and survivors of childhood cancer			They agreed on the importance of interpersonal and informational aspects of communication, such as honesty, support, and the need to be fully informed. Participants generally preferred a collaborative role in medical decision making. Patients differed in their satisfaction with their parents' role as managers of the communication.

Table 2

Adolescent Decision Making with Chronic Illnesses (n=10)

Study	Purpose	Method	Sample	Results
Beresford & Sloper, 2003 [47]	To explore the experiences of chronically ill adolescents in communicating with health professionals, including the identification of factors which hinder or facilitate their use of health professionals as an information source	Individual interviews and focus groups	N=63 aged 11-16	Factors affecting the openness and effectiveness of communication included: duration and frequency of contact; gender; perceived attitudes towards adolescents; the communication skills of the adolescent and health professional; and the presence of parents and medical students/trainee doctors.
Broome & Richards, 1999 [21]	To describe how relationships with adults influenced pediatric patients' involvement in clinical research	Semi-structured interviews	N=34 aged 8-22	Were shown to articulate their ability and right to make decisions about involvement in research, along with their parent(s).
Coyne, 2009 [42]	To explore the views of children, parents, and nurses to determine desired engagement in decision making within a healthcare setting	In-person interviews, questionnaires and observation	N= 11 children and adolescents aged 9-15 N= 10 parents N= 12 nurses	Adolescents expressed the need to be consulted so that they could understand their illness, be involved in their care, and prepare themselves for procedures. However, children's views were underused and had varying experiences of being consulted about their care and treatment.
Garth et al., 2009 [43]	To explore the interaction between physicians, parents, and pediatric patients when the child is a participant in decision-making	Semi-structured interviews	N=33 9 pediatricians, 14 parents, and 10 children aged 8-12 with cerebral palsy	The child was not perceived to be an 'equal' or 'regular' partner as there were reports of variability in the involvement between children, as well as variability in the progressive involvement of each child.
Knopf et al., 2008 [46]	To describe the decision-making preferences of adolescents with chronic illnesses and their parents, and the extent to which they agree	Survey	N= 82 adolescents mean age 15 years, 62 parents	Nearly equal percentages of adolescents (37%) and parents (36%) preferred shared decision making. Overall, the largest proportion of adolescents (46%) and parents (53%) preferred passive decision making compared to active or shared decision making. Across five response choices, 33% of pairs agreed. Physicians need to inquire directly about patient and parent preferences
Lyon et al., 2004 [41]	To explore whether differences exist between chronically ill and healthy adolescents with regard to their attitudes about end-of-life	Survey	N=25 adolescents with chronic illness, 25 healthy adolescents aged 13-21	Ninety-six percent of chronically ill and 88% of healthy teens want to share decision-making if they are very ill; 76% prefer to discuss their wishes earlier in the course of a life-

Study	Purpose	Method	Sample	Results
	(EOL) issues			threatening illness. Half of all adolescents believed that they would be allowed to change their minds once they had made a decision. Eighty percent of chronically ill and 68% of healthy teens believed that their doctor would respect their wishes, while many were uncertain.
Runeson et al., 2002 [44]	The aim of this study was: (1) to identify everyday situations in medical and nursing care that illustrate children's participation in decision making; (2) to identify various levels of participation, actual and optimal; and (3) to compare actual and optimal participation. A further aim was: (4) to describe the way in which children, parents and staff interact during the decision-making process	Observation	N=24 aged 5 months to 18	Children and adolescents are not always allowed to participate in decision making to the extent that is considered optimal. This showed that parents do not always support their children in difficult situations and that health care staff often inform children about what is going to happen without presenting alternatives or asking for their views. It is of great importance that children are looked upon as potentially autonomous individuals and that staff members realize that one of their core duties is to facilitate children's participation in decision making concerning their health care.
Young et al., 2003 [48]	To examine young people's and parents' accounts of communication about cancer in childhood	Semi-structured interviews	N=32 aged 8-17	The patients identified elements of their parents' role that both facilitated and constrained their communication, and while they welcomed their parents' involvement, some expressed unease with the constraining aspects of their parents' role.
Varma et al., 2008 [45]	To know how children and parents make decisions regarding pediatric research	Survey	N=117 children aged 7-14	In the research group, 90.5% of pediatric respondents believed they should be involved in making research enrollment decisions, whereas only 61.5% of the adult respondents believed children should be involved in making these decisions.

Table 3**Adolescent Decision Making in Fertility (n=9)**

Study	Purpose	Method	Sample	Results
Anderson et al., 2008 [56]	To assess UK practice relating to information provision about the effects of cancer treatment on fertility and options for fertility preservation	Survey	N=1030 aged 1-22	The majority (83%) of post-pubertal boys assessed as high/medium risk of infertility were referred for semen cryopreservation. This rate fell to 39% of those in early puberty. Only 1% (n=4) of girls were referred to an assisted conception unit.
Burns et al., 2006 [51]	To determine whether female adolescents with a diagnosis of cancer and their parents were interested in trying to preserve fertility	Cross-sectional survey	N= 50 39 parent/adolescent pairs, 3 parent-only responses, and 8 adolescent-only responses aged 10-21	Adolescents and parents had thought about the future and were interested in research treatments to help preserve fertility, but not willing to postpone cancer therapy. Achieving a state of good health was most important to the adolescent group.
Chapple et al., 2006 [53]	To explore fertility concerns for adolescent males treated for cancer and identify communication barriers of these issues	Narrative interviews	N=21 aged 16-21	Communication about sperm storage was sometimes difficult and embarrassing. Patients wanted the opportunity to bank their sperm but decisions were often rushed. Some indicated a desire for counseling as they were unprepared to undergo the process of sperm banking. Uncertainty about fertility status caused worries for the future.
Crawshaw & Sloper, 2006 [30]	To assess the experiences of adolescents and young adults faced with possible infertility following cancer treatment	In-depth interviews	N=38 aged 13-21	All desired to have fertility information prior to treatment.
Edge et al., 2006 [50]	To identify the barriers to successful sperm banking in a group of adolescent and young adult patients	Questionnaires	N=55 aged 13-21	Those who had been unsuccessful were younger and described higher levels of anxiety at diagnosis and greater difficulty in talking about fertility. They also described less understanding of sperm banking at the time of diagnosis. Younger patients may be helped by the provision of high quality information and more open discussion of the technique.
Ginsberg et al., 2008 [52]	To evaluate the feasibility of offering newly diagnosed patients the opportunity to sperm bank and to determine the decision-making processes of patients and their parents who considered sperm banking	Survey	N=68 aged 12-24	Parent and patient made the decision together to bank 80% of the time. All sons who attempted to bank and their parents felt they had made the right decision, including those who attempted but failed.

Study	Purpose	Method	Sample	Results
Oosterhuis et al., 2008 [49]	To ascertain concerns about infertility risks among pediatric oncology patients and their parents	Survey	N=97 aged 14-25	Only 29.9% of parents were satisfied with the amount of information received on fertility.
Schover et al., 2002 [54]	To survey oncologists in three different practice settings to determine their knowledge, attitudes, and practices regarding referring patients to bank sperm before cancer treatment	Postal survey	N=718 physicians regarding post-pubertal teen boys	52% of physicians thought any teen producing sperm should be eligible to bank and 86% believed males under age 18 should be able to bank. Parental consent is required before offering sperm banking to a teen patient (37% agree) or to allow a teen to view erotic materials in the collection room (61% agree). Seventy-eight percent believed that both parents and teen should be in the room when sperm banking is discussed, 11% would start with the parents only, and 9% preferred to talk to the teen alone.
Vadaparampil et al., 2008 [55]	To examine factors that may influence the discussion with pediatric oncologists, pediatric cancer patients, and their families	In-depth interviews	N=24 pediatric oncologists who treat patients aged 0-21, mean age 9	Physicians perceived girls were more open to discussing issues of fertility than boys due to embarrassment, however girls showed more characteristics of distress than boys regarding the potential for infertility.