Published in final edited form as:

Tetrahedron Asymmetry. 2008 May 1; 19(8): 1020–1026. doi:10.1016/j.tetasy.2008.04.005.

Asymmetric multi-component reactions: convenient access to acyclic stereocenters and functionalized cyclopentenoids

Arun K. Ghosh*, Sarang S. Kulkarni, Chun-Xiao Xu, and Khriesto Shurrush

Department of Chemistry and Medicinal Chemistry, Purdue University, 560 Oval Drive, West Lafayette, IN 47907, USA

Abstract

Asymmetric multi-component reactions of optically active phenyl dihydrofuran, keto ester or *N*-tosyl imino ester, and allylsilane provided functionalized phenyl tetrahydrofurans with multiple stereogenic centers diastereoselectively. Cleavage of the resulting substituted tetrahydrofurans readily provided acyclic derivatives with three contiguous asymmetric centers via an acyloxycarbenium ion intermediate. Ring closing olefin metathesis, using Grubbs catalyst, afforded functionalized cyclopentene derivatives in optically active form. A one pot tandem tetrahydrofuran ring cleavage followed by ring closing olefin metathesis also provided functionalized cyclopentenones in good yield.

1. Introduction

Recently, we reported the syntheses of a variety of substituted tetrahydrofurans and tetrahydropyrans with the creation of multiple stereogenic centers in a single operation by novel multicomponent coupling reactions.¹ The overall protocol is practical, efficient and provided rapid access to functionalized heterocycles. Subsequently, we have shown that multicomponent reactions of optically active phenyl dihydrofuran, *N*-tosyl imino ester, and carbon nucleophiles can provide stereocontrolled synthesis of unnatural optically active heterocyclic amino acids with multiple stereocenters.² To broaden the scope and utility of this reaction further, we envisioned the cleavage of such functionalized phenyl tetrahydrofuran derivatives to provide access to acyclic derivatives with multiple stereocenters in optically active form. Also, for all previous investigation, we synthesized optically active 2-phenyl-2,5-dihydrofuran using the Hayashi-Heck³ reaction of aryl triflate and dihydrofuran in 80–88% ee. While the methodology provided enantioenriched phenyldihydrofuran in a single step, operational complexity, low enantiomeric excess and lack of generality led us to explore alternative synthesis of optically active dihydrofuran. Herein we report an alternative synthesis of optically active phenyl dihydrofuran, its conversion to functionalized tetrahydrofurans in high diastereomeric purity and a Lewis acid-catalyzed cleavage of the substituted tetrahydrofurans via an acyloxycarbenium ion intermediate to provide acyclic derivatives with three contiguous asymmetric centers. The acyclic precursors can be converted to functionalized cyclopentene derivatives in optically active form by a ring closing metathesis using Grubbs catalyst. We have also demonstrated a one-pot tandem Lewis acid catalyzed cleavage followed by ring closing olefin metathesis to

© 2008 Elsevier Science. All rights reserved.

*Corresponding author. Tel.: +1-765-494-5323; fax: +1-765-496-1612; akghosh@purdue.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

afford functionalized cyclopentene derivatives in good yield. Such functionalized cyclopentenoids are often structural features of a variety of bioactive natural products.⁴

2. Results and Discussion

The synthesis of optically active phenyldihydrofuran was carried out as outlined in Scheme 1. The known γ -lactone **1** was prepared in multigram quantity using an enantioselective CBS-reduction as the key step.^{5,6} The reduction of **1** by DIBAL-H provided the corresponding lactol. Reaction of the resulting lactol with mesyl chloride and triethylamine at $-50\text{ }^{\circ}\text{C}$ provided the mesylate. Heating of the mesylate at $42\text{ }^{\circ}\text{C}$ furnished phenyldihydrofuran **2** in 60% yield. The protocol is convenient and both enantiomers of phenyldihydrofuran **2** can be prepared in high enantiomeric purity.⁷

An asymmetric multi-component reaction was then carried out with this optically active phenyldihydrofuran **2**. As described, the reaction of **2** (1.2 equiv) and ethyl pyruvate **3a** ($R = \text{Me}$, 1 equiv) in the presence of TiCl_4 (1 M solution in CH_2Cl_2 , 1.1 equiv) at $-78\text{ }^{\circ}\text{C}$ for 1 h, followed by the addition of trimethylsilane (3equiv) at $-78\text{ }^{\circ}\text{C}$ and warming up the reaction to $23\text{ }^{\circ}\text{C}$ for 1 h. The reaction was quenched with saturated NaHCO_3 solution, standard work up and flash chromatography over silica gel provided the single condensation product **4a** in 75% yield as a single diastereomer (by ^1H - and ^{13}C - NMR analysis). We have also carried out multicomponent reaction with ethyl 4-methoxy-2-oxobutanoate which was prepared according to published procedure.⁸ Multicomponent reaction of **2** with ethyl 4-methoxy-2-oxobutanoate **3b** ($R = \text{CH}_2\text{CH}_2\text{OCH}_3$) as described above, furnished product **4b** as a 3:1 inseparable mixture of diastereomers in 67% yield. These multicomponent reactions afforded highly functionalized tetrahydrofuranyl derivatives **4a** and **4b** diastereoselectively with efficient construction of three contiguous asymmetric centers including a quaternary carbon center. Conceivably, the appropriate cleavage of the tetrahydrofuran ring would provide access to stereocontrolled acyclic derivatives in optically active form. As it turned out, the 2-phenyltetrahydrofuran is suitably positioned to be cleaved by an acyloxycarbenium ion mediated ring opening reaction. Unsubstituted phenyltetrahydrofuran was known to react with excess of metal halide in acetic anhydride to provide 4-acetoxy-1-phenylbut-1-ene.⁹ Cleavage of a more substituted phenyltetrahydrofuran has been carried out by us recently using a catalytic amount of ZnCl_2 in acetic anhydride at $120\text{ }^{\circ}\text{C}$. We elected to utilize this protocol. Thus, 2-phenyl tetrahydrofuran derivative **4a** was exposed to the above reaction conditions for 4 h. This has afforded the styryl derivative **6a** in 25% yield. The reaction proceeded first by Lewis acid catalyzed formation of the acetate (monitored by TLC and and by ^1H -NMR) followed by formation of a possible acyloxycarbenium ion intermediate **5**, then stable benzylic carbonium ion and loss of a proton to provide **6a**.¹⁰ To further improve the reaction yield, we have examined a number of other Lewis acids under a variety of reaction conditions. The results are shown in Table 1. Entry 8 provided the optimum result.

Thus, the reaction of **4a** with a catalytic amount of $\text{Cu}(\text{OTf})_2$ in a mixture (10:1) of toluene and acetic anhydride at reflux provided acyclic derivative **6a** in 85% yield as a single diastereomer. Similarly, phenyltetrahydrofuran **4b** afforded acyclic derivative **6b** in 83% yield as a 3:1 inseparable mixture of diastereomers. Thus, the asymmetric multicomponent reaction followed by acyloxycarbenium ion mediated ring opening afforded **6a** and **6b** (major product) containing multiple stereocenters in an optically active form. Both these derivatives are also suitable precursors for the synthesis of substituted cyclopentene derivatives by a ring closing olefin metathesis.¹¹ Thus, exposure of **6a** to Grubbs II catalyst (3 mol%) in CH_2Cl_2 at $23\text{ }^{\circ}\text{C}$ for 4 h afforded cyclopentene derivative **7a** in optically active form (Scheme 2).¹² We then attempted to carry out Lewis acid catalyzed ring opening and ring closing metathesis in one operation. Indeed, treatment of **4a** with a catalytic amount (20

mol%) of $\text{Zn}(\text{OTf})_2$ in the presence of acetic anhydride in toluene at 110 °C for 12h followed by exposure of the resulting ring opening product to 3% Grubbs catalyst at 23 °C for 4 h afforded cyclopentene **7a** in 58% yield. Treatment of **4a** with 20 mol% $\text{Zn}(\text{OTf})_2$ and 5 mol% Grubbs II catalyst at 110 °C for 12 h afforded only a trace amount of **7a** with starting material mostly unchanged.

Exposure of **6b** and its diastereomer to Grubbs II catalyst (3 mol%) in CH_2Cl_2 at 23 °C furnished cyclopentenones **7b** and **7c**. The isomers were separated by flash column chromatography. The stereochemistry of the cyclopentenones was determined based on the NOESY studies on the corresponding bicyclic lactones **8b** and **8c**.

We have previously reported that the multicomponent reaction of *N*-tosyl imino ester^{2a} with 2-phenyldihydrofuran **2** in the presence of TiCl_4 with allyltrimethylsilane as a nucleophile provided the single diastereomer **10** in 72% yield (Scheme 3). To determine the cyclic stereochemistry in an acyclic form, we attempted acyloxycarbenium ion mediated ring opening of tosylamine derivative **9** as described above for phenyltetrahydrofuran derivatives **4a** and **4b**. However, the above reaction conditions failed to provide any ring opening product, even in presence of three equivalents of Lewis acid. The starting material remained mostly unchanged. The formation of the acyloxycarbenium ion was possibly retarded because of sequestering of the Lewis acid by the tosylamide functionality.¹⁴ Therefore, ethyl ester **10** was reduced using NaBH_4 in the presence of CaCl_2 and resulting alcohol was treated with triphosgene in pyridine to provide the oxazolidinone **11** in 61% yield (over two steps).¹⁵ The oxazolidinone **11** was then treated with a catalytic amount (5 mol%) of $\text{Cu}(\text{OTf})_2$ which gave the expected ring opening product **12** in 67% yield, increasing the catalyst loading to 10 mol% resulted in improvement of isolated yield to 73%. Exposure of oxazolidinone **11** to a catalytic amount (20 mol%) of $\text{Zn}(\text{OTf})_2$ in presence of excess acetic anhydride in toluene however, provided the ring opening product **12** in 81% isolated yield. All three asymmetric centers in acyclic derivative **12** were constructed diastereoselectively during the multicomponent reaction. Diene **12** is set to form a functionalized cyclopentene derivative **13**. Thus, treatment of **12** with Grubbs catalyst (3 mol%) in CH_2Cl_2 at 23 °C for 4 h afforded cyclopentene in optically active form. The enantiomeric excess of compound **13** was determined by reduction of this compound with Dibal-H in CH_2Cl_2 at -78 °C followed by conversion of the alcohol to the Mosher ester. The ^{19}F -NMR analysis of the Mosher ester established enantiomeric excess of **10** as 87% ee.¹⁶

3. Conclusions

In conclusion, the asymmetric multicomponent reaction diastereoselectively constructed functionalized tetrahydrofuran derivatives containing multiple stereocenters. Lewis acid catalyzed cleavage of phenyltetrahydrofuran ring via acyloxycarbenium ion intermediate provided useful access to acyclic derivatives with three contiguous asymmetric centers. Acyclic derivatives were converted to functionalized cyclopentene derivatives in optically active form by ring closing olefin metathesis.

4. Experimental

4.1 General

All melting points were recorded on a melting point apparatus and are uncorrected. Anhydrous solvents were obtained as follows: THF and diethyl ether by distillation from sodium and benzophenone; pyridine and dichloromethane from CaH_2 . All other solvents were reagent grade. All moisture sensitive reactions were carried out in flame dried flask under nitrogen atmosphere. Column chromatography was performed with 240–400 mesh silica gel under low pressure of 3–5 psi. TLC was carried out with silica gel 60-F-254 plates.

The title α -*N*-tosyl imino ester was made from ethyl glyoxylate and *N*-toluenesulfonylisocyanate by Weinreb's procedure.¹³ Ethyl 4-methoxy-2-oxobutanoate **3b** was prepared according to the reported procedure.⁸

4.2.1. (*R*)-2-Phenyl-2,3-dihydrofuran 2—To a stirred solution of lactone **1** (0.5 g, 3 mmol) in CH₂Cl₂ (10 mL) at -78°C , was added a solution of DIBAL-H (1 M in CH₂Cl₂, 3.7 mmol). After stirred at -78°C for 1 h, the reaction was quenched with saturated aqueous potassium sodium tartrate. The mixture was diluted with CH₂Cl₂ and vigorously stirred at room temperature for 2 h. The aqueous layer was extracted twice with CH₂Cl₂. The combined organic layers were dried over anhydrous Na₂SO₄ and concentrated in vacuo to provide the crude lactol as a colorless oil, which was immediately used in the next reaction without further purification.

To a stirred solution of the above lactol in CH₂Cl₂ (25 mL) at -50°C was added triethylamine (1.2 mL, 12.2 mmol) followed by methanesulfonyl chloride (0.3 mL, 3.9 mmol). The solution was stirred for 1 h at -50°C and then the reaction mixture was warmed to 23°C and heated at reflux for 12 h. The reaction mixture was then cooled to 23°C , solvents evaporated and purified by column chromatography to afford phenyl dihydrofuran **2** (272 mg, 60% yield over 2 steps) as clear oil $R_f = 0.3$ (pentane: CH₂Cl₂ = 3:1), $[\alpha]_D^{23} = -66$ (c 1.05, CHCl₃); ¹H NMR (400 MHz, CDCl₃): δ 2.63 (qt, $J = 2, 8$ Hz, 1 H), 3.06–3.13 (m, 1H), 4.98 (dd, $J = 2.5, 6.4$ Hz, 1H), 5.53 (dd, $J = 8.5, 13.3$ Hz, 1H), 6.47 (dd, $J = 2.4, 4.7$ Hz, 1H), 7.28–7.45 (m, 5 H); ¹³C NMR (100 MHz, CDCl₃): 37.8, 82.3, 98.9, 125.5, 127.6, 128.5, 142.9, 145.2. IR (film, NaCl) 2928, 1619, 1135, 1051 cm⁻¹.

4.2.2. (*R*)-Ethyl2-((2*R*,3*R*,5*R*)-2-allyl-5-phenyltetrahydrofuran-3-yl)-2-hydroxypropanoate 4a—To a mixture of ethyl pyruvate (232 mg, 2 mmol) and (*R*)-2,3-dihydro-2-phenylfuran **1** (350 mg, 2.4 mmol) in dry CH₂Cl₂ (20 mL) at -78°C was added a solution of TiCl₄ (1.0 M in CH₂Cl₂, 2.2 mmol) and the resulting orange solution was stirred at -78°C for 1 h. Allyltrimethylsilane (686 mg, 6 mmol) was added and the mixture was warmed from -78°C to 23°C over 1 h. Then the reaction mixture was quenched carefully with 15 mL saturated sodium bicarbonate solution and the aqueous layer was extracted with dichloromethane (3 x). The combined organic extracts were dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The residue was purified further by flash column chromatography on silica gel to afford the tertiary alcohol **4a** (454 mg, 75% yield) as a yellow oil, $R_f = 0.48$ (25% EtOAc in hexanes), $[\alpha]_D^{23} = +15.7$ (c 1.3, CHCl₃); ¹H NMR (500 MHz, CDCl₃): δ 1.27 (t, $J = 7$ Hz, 3H), 1.42 (s, 3H), 1.78–1.84 (m, 1H), 2.03–2.07 (m, 1H), 2.42–2.48 (m, 2H), 2.61–2.66 (m, 1H), 3.51 (s, 1H), 4.20–4.28 (m, 3H), 4.89 (dd, $J = 10, 6.5$ Hz, 1H), 5.15–5.21 (m, 2H), 5.98–6.07 (m, 1H), 7.23–7.37 (m, 5H); ¹³C NMR (125 MHz, CDCl₃) δ 14.2, 24.8, 38.2, 40.9, 49.8, 62.3, 74.4, 75.3, 78.9, 80.6, 117.4, 125.8, 127.3, 128.2, 134.7, 142.2, 176.8; FT-IR (film, NaCl) 3462, 2979, 1784 cm⁻¹. m/z (CI) 305.05 (M + H)⁺.

4.2.3. ((*S*)-Ethyl2-((2*R*,3*S*,5*R*)-2-allyl-5-phenyltetrahydrofuran-3-yl)-2-hydroxy-4-methoxybutanoate 4b—The procedure described for **3a** was used for **3b**. Accordingly, ethyl-4-methoxy-2-oxobutanoate **1** (160 mg, 1 mmol), (*R*)-2,3-dihydro-2-phenylfuran **1** (175 mg, 1.2 mmol), TiCl₄ (1 M in CH₂Cl₂, 1.2 mmol) and allyltrimethylsilane (345 mg, 3 mmol) afforded furan **4b** (230 mg, 66% yield) as a 3:1 mixture of inseparable diastereomers, $R_f = 0.36$ (hex : EtOAc = 4:1), $[\alpha]_D^{23} = -15.3$ (c 1.2, CHCl₃); ¹H NMR (500 MHz, CDCl₃): δ 1.29 (t, $J = 7$ Hz, 3H, major), 1.34 (t, $J = 7.5$ Hz, minor), 1.76–1.97 (m, 2H), 1.98–2.11 (m, 2H), 2.17–2.23 (m, minor), 2.39–2.47 (m, 2H), 2.58–2.63 (m, 1H), 3.26 (s, minor), 3.27 (s, 3H, major), 3.41–3.46 (m, 1H), 3.49–3.56 (m, 1H), 3.75 (s, minor), 3.91 (s, 1H, major), 4.01–4.05 (m, minor), 4.16–4.30 (m, 3H), 4.87–4.92 (m, 1H), 5.09–5.20 (m, 2H), 5.92–6.04 (m, 1H), 7.21–7.36 (m, 5H); ¹³C NMR (125

MHz, CDCl₃): δ 14.1 (minor), 14.2 (major), 31.2 (minor), 37.0 (major), 37.2 (minor), 37.7 (minor), 37.9 (major), 39.5 (minor), 41.2 (major), 50.3 (major), 50.5 (minor), 58.9 (major), 61.2 (major), 68.4 (minor), 68.5 (major), 117.3 (minor), 117.4 (major), 126 (major), 126.1 (minor), 127.3 (major), 128.2 (major), 128.5 (minor), 134.7 (minor), 134.8 (major), 141.9 (major), 142.4 (minor), 175.6 (major), 175.9 (minor); FT-IR (film, NaCl) 3493, 2362, 1724 cm⁻¹. HRMS (CI) [M+H]⁺ calcd for C₂₀H₂₈O₅ 349.2015, found 349.2018.

4.2.4. (2*R*,3*R*,5*R*)-1-Ethoxy-2-methyl-1-oxo-3-styrylhept-6-ene-2,4-diyl

diacetate 6a—A mixture of tertiary alcohol **4** (350 mg, 1.15 mmol), Cu(OTf)₂ (21 mg, 0.058 mmol) and acetic anhydride (5.5 mL, 57.5 mmol) was heated to 110 °C in toluene (50 mL) with stirring. After 2 h, the reaction mixture was cooled to room temperature and the reaction mixture was washed with saturated aqueous NaHCO₃ (2 x). The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The residue was purified further by flash column chromatography on silica gel to afford the styrene **6a** (355 mg, 80% yield) as a yellow oil, *R*_f = 0.48 (20% EtOAc in hexanes), ¹H NMR (500 MHz, CDCl₃): δ 1.23 (t, *J* = 7 Hz, 3H), 1.67 (s, 3H), 2.04 (s, 3H), 2.05 (s, 3H), 2.23–2.27 (m, 1H), 2.34–2.38 (m, 1H), 2.77 (d, *J* = 10 Hz, 1H), 4.07–4.20 (m, 2H), 5.07–5.11 (m, 2H), 5.38 (t, *J* = 5 Hz, 1H), 5.76–5.88 (m, 1H), 6.24 (dd, *J* = 16, 10 Hz, 1H), 6.46 (d, *J* = 16 Hz, 1H), 7.26–7.30 (m, 1H), 7.35 (t, *J* = 7 Hz, 2H), 7.43 (d, *J* = 8.5 Hz, 2H); ¹³C NMR (125 MHz, CDCl₃) δ 13.9, 20.8, 21.2, 21.3, 37.9, 52.7, 61.4, 70.3, 81.8, 118.5, 122.7, 126.4, 127.7, 128.6, 133.1, 135.7, 136.6, 169.7, 169.9, 170.6; FT-IR (film, NaCl), 2981, 1742, 1642 cm⁻¹. HRMS (CI) [M+H]⁺ calcd for C₂₂H₂₈O₆ 389.1964, found 389.1968.

4.2.5. (3*R*,4*S*,5*R*)-3-(Ethoxycarbonyl)-1-methoxy-4-styryloct-7-ene-3,5-diyl

diacetate 6b—The procedure described for **5a** was used for **5b**. Accordingly furan **4b** (230 mg, 0.66 mmol), Cu(OTf)₂ (12 mg, 0.03 mmol), acetic anhydride (3.1 mL, 33 mmol) were heated in toluene (30 mL) to give styrene **6b** (216 mg, 83% yield) as a 3:1 mixture of inseparable diastereomers, *R*_f = 0.42 (hex : EtOAc = 4:1), [α]_D²³ = +44.9 (*c* 1.0, CHCl₃); ¹H NMR (400 MHz, CDCl₃): δ 1.27 (t, *J* = 7 Hz, 3H, major), 1.99 (s, 3H, major), 2.01 (s, minor), 2.02 (s, minor), 2.08 (s, 3H, major), 2.23–2.32 (m, 1H), 2.33–2.41 (m, 3H), 2.52–2.60 (m, 1H), 3.09 (d, *J* = 10 Hz, minor), 3.18 (d, *J* = 10 Hz, 1H), 3.26 (s, 3H), 3.27 (s, minor), 3.41–3.49 (m, 3H), 4.05–4.29 (m, 3H), 5.04–5.11 (m, 3H), 5.33 (t, *J* = 7.5 Hz, minor), 5.45 (t, *J* = 7 Hz, 1H), 5.65–5.79 (m, 1H), 6.17 (dd, *J* = 16, 10 Hz, 1H), 6.25 (dd, *J* = 16, 10 Hz, minor), 6.48 (d, *J* = 16 Hz, 1H), 7.13–7.19 (m, 2H), 7.34 (t, *J* = 7.5 Hz, 3H), 7.69 (d, *J* = 7.5 Hz, 2H); ¹³C NMR (100 MHz, CDCl₃): δ 13.9 (major), 21.0 (minor), 21.1 (major), 21.2 (major), 31.8 (minor), 32.1 (major), 37.9 (major), 38.0 (minor), 50.2 (major), 50.7 (minor), 58.4 (minor), 58.5 (major), 61.1 (minor), 61.3 (major), 67.7 (major), 67.9 (minor), 70.4 (major), 70.7 (minor), 82.3 (major), 82.6 (minor), 118.0 (major), 118.1 (minor), 126.3 (minor), 126.4 (major), 127.5 (minor), 127.7 (major), 128.5 (major), 133.2 (minor), 133.4 (major), 135.7 (minor), 136.0 (major), 136.8 (major), 137.1 (minor), 169.6 (major), 169.7 (minor), 169.8 (major), 169.9 (major), 170.0 (minor); FT-IR (film, NaCl) 2361, 1742 1370, 972 cm⁻¹. HRMS (CI) [M+Na]⁺ calcd for C₂₄H₃₂O₇Na 455.2046, found 455.2050.

4.2.6. (*R*)-Ethyl 2-acetoxy-2-((1*S*,5*R*)-5-acetoxycyclopent-2-enyl)propanoate 7a

—To a stirring solution of **6a** (230 mg, 0.59 mmol) in CH₂Cl₂ (120 mL) was added a second generation Grubbs catalyst (15 mg, 0.018 mmol) under argon. The reaction was allowed to stir at 23 °C for 4 h. After this period the reaction mixture was concentrated under reduced pressure and purified by flash column chromatography on silica gel to afford the cyclopentene **6a** (147 mg, 88% yield) as a colorless oil, *R*_f = 0.4 (20% EtOAc in hexanes), [α]_D²³ = −3.5 (*c* 1, CHCl₃); ¹H NMR (500 MHz, CDCl₃): δ 1.28 (t, *J* = 7 Hz, 3H), 1.69 (s, 3H), 2.03 (s, 3H), 2.07 (s, 3H), 2.23–2.37 (m, 1H), 2.69–2.74 (m, 1H), 3.40–3.42 (m, 1H),

4.14–4.22 (m, 2H), 5.51 (dt, $J = 6.5$, 3 Hz 1H), 5.38 (t, $J = 5$ Hz, 1H), 5.76–5.88 (m, 1H), 6.24 (dd, $J = 16$, 10 Hz, 1H), 6.46 (d, $J = 16$ Hz, 1H), 7.26–7.30 (m, 1H), 7.35 (t, $J = 7$ Hz, 2H), 7.43 (d, $J = 8.5$ Hz, 2H); ^{13}C NMR (125 MHz, CDCl_3) δ 13.9, 20.8, 21.2, 21.3, 37.9, 52.7, 61.4, 70.3, 81.8, 118.5, 122.7, 126.4, 127.7, 128.6, 133.1, 135.7, 136.6, 169.7, 169.9, 170.6; FT-IR (film, NaCl), 2981, 1742, 1642 cm^{-1} . HRMS (ESI) $[\text{M}+\text{Na}]^+$ calcd for $\text{C}_{14}\text{H}_{20}\text{O}_6\text{Na}$ 307.1158, found 307.1155.

4.2.7. (*R*)-Ethyl-2-acetoxy-2-((1*S*,5*R*)-5-cetoxycyclopent-2-enyl)-4-

methoxybutanoate 7b—The procedure described for **7a** was used for **7b**. Accordingly, styrene **6b** (210 mg, 0.16 mmol), second generation Grubbs catalyst (12 mg, 0.015 mmol) in CH_2Cl_2 (30 mL) afforded cyclopentenones **7b** (92 mg, 58%) and **7c** (25 mg, 16%).

7b $[\alpha]_{\text{D}}^{23} = -3.2$ (c 1, CHCl_3); ^1H NMR (500 MHz, CDCl_3) : δ 1.30 (t, $J = 7$ Hz, 3H), 2.02 (s, 3H), 2.10 (s, 3H), 2.52 (dt, $J = 14.8$, 6.5 Hz, 1H), 2.60 (dt, $J = 14.8$, 6.5 Hz, 1H), 2.65–2.71 (m, 1H), 3.30 (s, 3H), 3.47–3.54 (m, 2H), 3.78–3.80 (m, 1H), 4.16–4.26 (m, 2H), 5.56 (dt, $J = 6.5$, 1.7 Hz, 1H), 5.82–5.85 (m, 1H), 5.87–5.93 (m, 1H); ^{13}C NMR (125 MHz, CDCl_3) δ 14.0, 21.0, 21.5, 34.2, 39.9, 54.5, 61.4, 68.1, 73.0, 82.7, 128.8, 129.5, 169.9, 170.1, 170.4; FT-IR (film, NaCl) 1739, 1370, 1236 cm^{-1} . HRMS (CI) $[\text{M}+\text{H}]^+$ calcd for $\text{C}_{16}\text{H}_{25}\text{O}_7$ 329.1600, found 329.1607.

7c $[\alpha]_{\text{D}}^{23} = -5.5$ (c 1, CHCl_3); ^1H NMR (500 MHz, CDCl_3) : δ 1.30 (t, $J = 7$ Hz, 3H), 2.04 (s, 3H), 2.09 (s, 3H), 2.31–2.37 (m, 2H), 2.43–2.47 (m, 1H), 2.65–2.71 (m, 1H), 3.32 (s, 3H), 3.49–3.54 (m, 2H), 3.80 (br s, 1H), 4.14–4.28 (m, 2H), 5.56 (t, $J = 7$ Hz, 1H), 5.81–5.87 (m, 1H), 5.99–6.04 (m, 1H); ^{13}C NMR (125 MHz, CDCl_3) δ 13.9, 21.1, 21.2, 33.2, 40.1, 53.7, 58.6, 61.4, 74.0, 81.8, 128.3, 130.4, 170.2, 170.4, 170.5; FT-IR (film, NaCl) 1738, 1371, 1233 cm^{-1} . HRMS (CI) $[\text{M}+\text{H}]^+$ calcd for $\text{C}_{16}\text{H}_{25}\text{O}_7$ 329.1600, found 329.1603.

4.2.8. (3*S*,3*aS*,6*aR*)-3-(2-Methoxyethyl)-2-oxo-3,3*a*,6,6*a*-tetrahydro-2H-

cyclopenta[b]furan-3-yl acetate 8b—To a stirred solution of **6b** (90 mg, 0.27 mmol) in a mixture (3:1:1) of THF:EtOH:H₂O (3 mL) at 0 °C was added solid LiOH (35 mg, 0.82 mmol) and the mixture was stirred for 6 h at 23 °C. After this period, the solvent was evaporated and aqueous layer was then acidified with 1 M HCl to pH 3 and extracted with EtOAc (3 x). The combined organic layer was dried over Na_2SO_4 and concentrated under reduced pressure and purified by flash column chromatography on silica gel to provide the lactone (34 mg, 64% yield) as colorless oil.

To the above lactone (34 mg, 0.17 mmol) in dry CH_2Cl_2 (3 mL) at 0 °C was added DMAP (103 mg, 0.84 mmol) and acetic anhydride (160 μL , 1.7 mmol). The resulting solution was stirred at 23 °C for 12 h. After this time the reaction mixture was diluted with CH_2Cl_2 and washed with 1N HCl, water and brine. The organic layer was concentrated under reduced pressure and the residue purified by column chromatography on silica gel to provide **8b** (37 mg, 92% yield) as colorless oil. ^1H NMR (500 MHz, CDCl_3) : δ 2.07–2.11 (m, 2H), 2.13 (s, 3H), 2.66 (d, $J = 18$ Hz, 1H), 2.73–2.79 (m, 1H), 3.36 (s, 3H), 3.51–3.54 (m, 1H), 3.67–3.75 (m, 2H), 5.35 (t, $J = 6.7$ Hz, 1H), 5.76–5.79 (m, 1H), 5.84–5.88 (m, 1H); ^{13}C NMR (125 MHz, CDCl_3) δ 20.7, 34.3, 38.7, 56.5, 58.8, 67.379.6, 81.4, 127.1, 131.8, 174.2.

4.2.9. (3*S*,3*aS*,6*aR*)-3-(2-Methoxyethyl)-2-oxo-3,3*a*,6,6*a*-tetrahydro-2H-

cyclopenta[b]furan-3-yl acetate 8c—The procedure described for **8b** was used for **8c**. Accordingly, cyclopentene **7c** (25 mg, 0.08 mmol), LiOH (10 mg, 0.23 mmol) afforded lactone (13 mg, 85% yield) as colorless oil. The above lactone (12 mg, 0.06 mmol), DMAP (37 mg, 0.30 mmol) and acetic anhydride (57 μL , 0.6 mmol) in CH_2Cl_2 (2 mL) afforded **8c** (11 mg, 70% yield) as a colorless oil. ^1H NMR (500 MHz, CDCl_3) : δ 2.16 (s, 3H), 2.23–

2.28 (m, 1H), 2.58–2.64 (m, 1H), 2.78–2.79 (m, 2H), 3.34 (s, 3H), 3.52–3.3.57 (m, 1H), 3.62–3.66 (m, 1H), 3.95–3.97 (m, 1H), 5.06–5.09 (m, 1H), 5.40–5.42 (m, 1H), 5.86–5.88 (m, 1H); ^{13}C NMR (125 MHz, CDCl_3) δ 21.2, 35.2, 40.0, 56.1, 58.7, 67.3, 80.3, 81.8, 126.7, 131.8, 169.8, 173.3.

4.2.10. (*R*)-Ethyl2-((2*R*,3*R*,5*R*)-2-allyl-tetrahydro-5-phenylfuran-3-yl)-2-(tosylamino)acetate (10**)**—To a mixture of freshly distilled α -*N*-tosyl imino ester **9** (255 mg, 1 mmol) and (*R*)-2,3-dihydro-2-phenylfuran **1** (175 mg, 1.2 mmol) in dry CH_2Cl_2 (10 mL) at -78°C was added a solution of TiCl_4 (1.0 M in CH_2Cl_2 , 1.1 mmol) and the resulting orange solution was stirred at -78°C for 50 min. Dry acetonitrile (205 mg, 5 mmol) was added at -78°C and the mixture was stirred at -78°C for 10 min. Allyltrimethylsilane (342 mg, 3 mmol) was added and the mixture was stirred at -78°C for 1 h. Then the mixture was warmed to -20°C and stirred at -20°C for 2 h. Then the mixture was cooled to -30°C and the reaction was quenched carefully with 15 mL ice-cooled saturated sodium bicarbonate solution. After the mixture was warmed to room temperature, the aqueous layer was extracted with dichloromethane (3 x). The combined organic extracts were dried over anhydrous Na_2SO_4 and concentrated under reduced pressure. The residue was further purified by flash column chromatography on silica gel to afford the α -*N*-tosylamino ester **10** (320 mg, 72% yield) as a yellow oil, $R_f = 0.38$ (hex : EtOAc 3:2), $[\alpha]_D^{23} = -11.8$ (c 0.9, CHCl_3); ^1H NMR (500 MHz, CDCl_3) : δ 1.05 (t, $J = 7.1$ Hz, 3H), 1.7–1.80 (m, 1H), 2.21–2.29 (m, 1H), 2.31–2.33 (m, 1H), 2.38–2.22 (m, 5H), 3.87–3.91 (m, 3H), 4.04–4.06 (m, 1H), 4.89 (t, $J = 7.7$ Hz, 1H), 5.11–5.17 (m, 2H), 5.39 (d, $J = 9.9$ Hz, 1H), 5.87–5.93 (m, 1H), 7.33–7.23 (m, 7H), 7.73 (d, $J = 8.3$ Hz, 2H); ^{13}C NMR (125 MHz, CDCl_3) δ 13.8, 21.5, 36.1, 38.6, 45.9, 55.9, 62.0, 79.5, 79.7, 117.7, 125.7, 127.4, 128.3, 129.7, 134.1, 136.3, 142.1, 144.0, 171.0; FT-IR (film, NaCl), 2980, 1746, 1349.2, 1163 cm^{-1} ; m/z (ESI) 444.1 ($\text{M} + \text{H}$) $^+$, 466.3 ($\text{M} + \text{Na}$) $^+$.

4.2.11. (*S*)-4-((2*R*,3*R*,5*R*)-2-allyl-5-phenyltetrahydrofuran-3-yl)-3-tosyloxazolidin-2-one (11**)**—To a stirring mixture of furan **10** (220 mg, 0.5 mmol) in THF:ethanol (2 mL:3 mL) at 0°C was added CaCl_2 (116 mg, 1.05 mmol) followed by addition of NaBH_4 (80 mg, 2.1 mmol) and the resulting solution was allowed to warm to room temperature over 3 h. After this time the reaction mixture was cooled to 0°C and quenched carefully with 5% citric acid solution. This mixture was then concentrated under vacuum and the aqueous layer was extracted with ethyl acetate (3 x). The combined organic extracts were dried over anhydrous Na_2SO_4 and concentrated under reduced pressure. The residue was further purified by flash column chromatography on silica gel to afford the primary alcohol (153 mg, 76% yield) as a colorless solid, $R_f = 0.2$ (25% EtOAc in hexanes), $[\alpha]_D^{23} = +4.7$ (c 1, CHCl_3); ^1H NMR (500 MHz, CDCl_3) : δ 1.78–1.82 (m, 1H), 2.08–2.18 (m, 2H), 2.31 (t, $J = 7$ Hz, 2H), 2.43 (s, 1H), 3.31–3.35 (m, 1H), 3.53 (d, $J = 4$ Hz, 2H), 3.76 (q, $J = 6$ Hz, 1H), 4.68 (dd, $J = 8.5, 6.5$ Hz, 2H), 5.07–5.11 (m, 2H) 5.24 (d, $J = 9.5$ Hz, 1H), 5.78–5.83 (m, 1H), 7.19–7.34 (m, 7H), 7.79 (d, $J = 8.3$ Hz, 2H); ^{13}C NMR (125 MHz, CDCl_3) δ 21.5, 29.6, 36.5, 39.4, 44.7, 63.4, 79.6, 80.5, 117.6, 125.5, 127.1, 127.2, 128.2, 129.9, 134.4, 137.3, 142.2, 143.9; FT-IR (film, NaCl), 3546, 3263, 2844 cm^{-1} .

To a stirring mixture of above amino alcohol (133 mg, 0.33 mmol) in dry CH_2Cl_2 (3 mL) was added pyridine (158 mg, 2 mmol) and triphosgene (49 mg, .17 mmol) at 0°C and reaction was allowed to warm to room temperature over 3 h. The reaction mixture was diluted with CH_2Cl_2 and was washed with water (3 x) and saturated aqueous CuSO_4 (5 x) followed by brine. The organic layer was dried over MgSO_4 and concentrated under reduced pressure. The residue was further purified by flash column chromatography on silica gel to afford the oxazolidinone **11** (115 mg, 81% yield) as a yellow oil, $R_f = 0.5$ (25% EtOAc in hexanes), $[\alpha]_D^{23} = -32.5^\circ$ (c 1.6, CHCl_3); ^1H NMR (500 MHz, CDCl_3) : δ 2.03–2.08 (m, 1H), 2.24–2.29 (m, 1H), 2.37–2.44 (m, 2H), 2.45 (s, 3H), 2.58–2.62 (m, 1H) 4.06 (q, $J = 6.5$

Hz, 1H), 4.24 (dd, $J = 2, 9$ Hz, 1H), 4.38 (t, $J = 8.5$ Hz, 1H), 4.65–4.68 (m, 1H), 4.96 (t, $J = 7.5$ Hz, 1H), 5.13–5.19 (m, 2H), 5.85–5.97 (m, 1H), 7.27–7.38 (m, 7H), 7.96 (d, $J = 8.5$ Hz, 2H); ^{13}C NMR (125 MHz, CDCl_3) δ 21.8, 36.5, 39.6, 47.4, 57.8, 67.3, 79.7, 79.8, 118.2, 125.6, 127.6, 128.5, 129.9, 134.0, 134.9, 142.3, 146.0, 152.6; ; FT-IR (film, NaCl), 2924, 1783 cm^{-1} .

4.2.12. (3*R*,4*R*)-3-(2-oxo-3-tosyloxazolidin-4-yl)-1-phenylhepta-1,6-diene-4-yl acetate (12)—A mixture of oxazolidinone **11** (80 mg, 0.19 mmol), $\text{Zn}(\text{OTf})_2$ (14.0 mg, 0.04 mmol) and acetic anhydride (890 μL , 9.4 mmol) were heated to 110 $^\circ\text{C}$ in toluene (15 mL) with stirring. After 2h the reaction mixture was cooled to room temperature and the reaction mixture was washed with saturated aqueous NaHCO_3 (2×5 mL). The organic layer was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure. The residue was further purified by flash column chromatography on silica gel to afford acetate **12** (71 mg, 81% yield) as a yellow oil, $R_f = 0.4$ (25% EtOAc in hexanes), $[\alpha]_D^{23} = +41.4$ (c 1.05, CHCl_3); ^1H NMR (500 MHz, CDCl_3) : δ 2.1 (s, 3H), 2.22–2.25 (m, 1H), 2.32–2.38 (m, 1H), 2.40 (s, 3H), 2.92–2.95 (m, 1H) 4.31 (t, $J = 7$ Hz, 1H), 4.47 (dd, $J = 2, 9$ Hz, 1H), 4.60 (dt, $J = 2, 8.5$ Hz, 1H), 5.03–5.08 (m, 3H), 5.50–5.59 (m, 1H), 6.20 (dd, $J = 9.5, 16$ Hz, 1H), 6.57 (d, $J = 16$, 1H), 7.22–7.40 (m, 7H), 7.96 (d, $J = 8.5$ Hz, 2H); ^{13}C NMR (125 MHz, CDCl_3) δ 21.1, 21.7, 37.6, 48.5, 59.1, 66.3, 70.4, 118.8, 122.06, 126.6, 128.2, 128.5, 128.7, 129.7, 132.7, 135.1, 136.2, 137.4, 145.6, 152.4, 170.3; ; FT-IR (film, NaCl), 2921, 1786, 1596 cm^{-1} ; HRMS (EI) $[\text{M}+\text{H}]^+$ calcd for $\text{C}_{25}\text{H}_{28}\text{NO}_6\text{S}$ 470.1637, found 470.1648.

4.2.13. (1*R*,2*R*)-2-((*R*)-2-Oxo-3-tosyloxazolidin-4-yl)cyclopent-3-enyl acetate 13—The procedure described for **7a** was used for **13**. Accordingly, styrene **12** (71 mg, 0.15 mmol), second generation Grubb's catalyst (6 mg, 0.008 mmol) in CH_2Cl_2 (45 mL) afforded cyclopentene **13** (48 mg, 86% yield) as a white solid, $R_f = 0.3$ (20% EtOAc in hexanes), $[\alpha]_D^{23} = -87.4$ (c 0.85, CHCl_3); ^1H NMR (500 MHz, CDCl_3) : δ 1.60 (s, 3H), 2.04–2.09 (m, 1H), 2.26 (s, 3H), 2.66–2.74 (m, 1H), 3.58 (s, 1H), 4.00 (dd, $J = 4, 9$ Hz, 1H), 4.15 (t, $J = 9$ Hz, 1H), 4.57 (quintet, $J = 4$ Hz, 1H), 5.00 (dt, $J = 7.7, 4$ Hz, 1H), 5.34–5.36 (m, 1H), 5.73–5.77 (m, 1H), 7.17 (d, $J = 8$ Hz, 2H), 7.76 (d, $J = 8$ Hz, 2H); ^{13}C NMR (125 MHz, CDCl_3) δ 20.5, 21.7, 40.9, 48.6, 56.1, 64.9, 73.6, 127.5, 128.1, 129.8, 132.6, 134.9, 146.7, 152.6, 169.9; FT-IR (film, NaCl) 1783, 1728, 1370, 1172 cm^{-1} ; HRMS (ESI) $[\text{M}+\text{Na}]^+$ calcd for $\text{C}_{17}\text{H}_{19}\text{NO}_6\text{SNa}$ 388.0831, found 388.0833.

Acknowledgments

Financial support by the National Institute of Health (GM 53386) is gratefully acknowledged.

References

- (a) Ghosh AK, Kawahama R, Wink D. *Tetrahedron Lett.* 2000; 41:8425. (b) Ghosh AK, Kawahama R. *Tetrahedron Lett.* 1999; 40:1083. (c) Ghosh AK, Kawahama R. *Tetrahedron Lett.* 1999; 40:4751. (d) Ghosh AK, Kawahama R. *J Org Chem.* 2000; 65:5433. [PubMed: 10993380]
- (a) Ghosh AK, Xu CX, Kulkarni SS, Wink D. *Org Lett.* 2005; 7:7. [PubMed: 15624964] (b) Ghosh AK, Kulkarni S, Xu C-X, Fanwick EP. *Org Lett.* 2006; 8:4509. [PubMed: 16986937]
- Ozawa F, Kubo A, Hayashi T. *J Am Chem Soc.* 1991; 113:1417.
- (a) Mayo P, Tam W. *Tetrahedron.* 2002; 58:9513. (b) Ito H, Hasegawa M, Takenaka Y, Kobayashi T, Iguchi K. *J Am Chem Soc.* 2004; 126:4520. [PubMed: 15070361] (c) Beale MH, Ward JL. *Nat Prod Rep.* 1998; 15:533. [PubMed: 10036771] (d) Noyori R, Suzuki M. *Science.* 1993; 259:44. and references cited therein. [PubMed: 8418493]
- Corey EJ, Bakshi RK, Shibata S, Chen C, Singh VK. *J Am Chem Soc.* 1987; 109:7925.
- Ghosh AK, Swanson L. *J Org Chem.* 2003; 68:9823. [PubMed: 14656116]
- Optical purity of phenyldihydrofuran was determined to be >95% ee.

8. Tavecchai P, Gantili P, Kruz M, Sottani C, Bonfichi R, Selva E, Lociuro S, Restelli E, Ciabatti R. *Tetrahedron*. 1995; 51:4867.
9. Ispriyan RM, Belen'kii LI, Gol'dfarb YL. *Bull Acad Sci USSR, Div Chem Sci*. 1967:2391.
10. (a) Ishihara K, Kubota M, Kurihara H, Yamamoto H. *J Am Chem Soc*. 1995; 117:4413. (b) Ishihara K, Kubota M, Kurihara H, Yamamoto H. *J Org Chem*. 1996; 61:4560. [PubMed: 11667380] (c) Chakraborti AK, Gulhane R. *Tetrahedron Lett*. 2003; 44:6749.
11. (a) Ghosh S, Sinha S, Drew MGB. *Org Lett*. 2006; 8:3781. [PubMed: 16898816] (b) Prunet J, Funel JA. *J Org Chem*. 2004; 69:4555. [PubMed: 15202921] (c) Hale KJ, Domostoj MM, Tocher DA, Irving E, Scheinmann F. *Org Lett*. 2003; 5:2927. [PubMed: 12889910]
12. Scholl M, Ding S, Lee C, Grubbs RH. *Org Lett*. 1999; 1:953. [PubMed: 10823227]
13. Tschaen DH, Turos E, Wienreb SM. *J Org Chem*. 1984; 49:5058.
14. (a) Furstner A, Langermann K. *J Am Chem Soc*. 1997; 119:9130. (b) Ghosh AK, Cappiello J, Shin D. *Tetrahedron Lett*. 1998; 8:4651.
15. Blakemore PR, Browder CC, Hong J, Lincoln CM, Nagorny PA, Robarge LA, Wardrop DJ, White JD. *J Org Chem*. 2005; 70:5449. [PubMed: 15989326]
16. The Mosher ester was formed by reaction of Mosher acid chloride and alcohol in presence of DMAP. The ^{19}F NMR analysis of Mosher ester revealed an enantiomeric purity of 87%. For procedure see Dale JA, Dull DL, Mosher HS. *J Org Chem*. 1969; 34:2543.

Scheme 1.

Scheme 2.

Scheme 3.

Table 1

Optimization of conditions of ring opening reaction

Entry	Lewis Acid	Equivalents	Solvent ^a	Yield % ^b
1	ZnCl ₂	0.06	Ac ₂ O	25
2	ZnCl ₂	0.20	Ac ₂ O/Toluene	--
3 ^b	Zn(OTf) ₂	1.00	Ac ₂ O/Toluene	79
4	Zn(OTf) ₂	0.20	Ac ₂ O/Toluene	75
5	Zn(OTf) ₂	0.10	Ac ₂ O/Toluene	75
6	Sc(OTf) ₃	0.20	Ac ₂ O/Toluene	73
7	Cu(OTf) ₂	0.20	Ac ₂ O/Toluene	85
8	Cu(OTf) ₂	0.05	Ac ₂ O/Toluene	82

^aReaction temperature of 110°C.^bIsolated yield after silica gel chromatography.