KpnI RFLP in the 3' flanking region of the C1 inhibitor gene

A.R.McPhaden, P.E.Carter¹, G.D.Birnie² and K.Whaley

Department of Pathology, Western Infirmary, Glasgow G11 6NT, ¹Department of Biochemistry, Aberdeen University, Aberdeen and ²Beatson Institute, Bearsden, Glasgow, UK

SOURCE AND DESCRIPTION OF CLONE : The 527bp EcoRI fragment (Ex 8) from the 3' end of the "full-length" C1 inhibitor cDNA in pUC 13 (1).

POLYMORPHISM : KpnI identifies two alleles represented by 30kb (A) and 13kb (B) hybridising bands.

FREQUENCY : Studied in 25 unrelated Caucasians.

A : 0.2

B : 0.8

NOT POLYMORPHIC FOR : ApaI, AvaI, BamHI, BglI, BglIII, BstEII, ClaI, DdeI, DraI, EcoRI, HaeII, HaeIII, HhaI, HincII, HindIII, HinfI, HpaI, HpaII, MspI, PstI, PvuI, PvuII, RsaI, SalI, Sau3AI, ScaI, SmaI, SstI, SstII, StuI, StyI, TaqI, XbaI, XhoI, XmnI.

CHROMOSOMAL LOCALISATION : The C1 inhibitor gene has been mapped to 11q11 - q 13.1 by in situ hybridisation (2).

MENDELIAN INHERITANCE : Co-dominant segregation has been demonstrated in one three generation family (21 individuals).

PROBE AVAILABILITY : Contact A.R.M.

OTHER COMMENTS : The polymorphism has been shown to arise due to a mutation involving the first KpnI site which lies approximately 10kb downstream of the C1 inhibitor gene. This polymorphism may be useful as an indirect marker of the defective C1 inhibitor gene in certain families with inherited C1 inhibitor deficiency. No relationship has been demonstrated between the polymorphism and plasma C1 inhibitor protein levels in normal individuals.

REFERENCES : 1. Bock, S.C. et al., *Biochemistry* **25** : 4292 - 4301 (1986)
2. Theriault, A. et al. (in preparation).

ACKNOWLEDGEMENTS : The full length C1 inhibitor cDNA clone was a generous gift from Dr S.C. Bock, Rockefeller University, New York. Support for this work came from the MRC (London) as Training Award Grant G106/320.