NICE news in the annals

Recent NICE Guidance of interest to surgeons

HEATHER STEGENGA and BRUCE CAMPBELL

Interventional Procedures

Single-incision laparoscopic cholecystectomy (May 2010)

Laparoscopic cholecystectomy usually leaves three small scars from closure of the port sites, but there is now a move to improve cosmesis still further by single-incision laparoscopic cholecystectomy. In this procedure, both the laparoscope and the instruments are introduced at a single site, commonly using a special port at the umbilicus where any scar is easily concealed.

The evidence presented to the Committee included 472 patients from one randomised (90 patients) and three non-randomised trials (130, 51 and 40 patients) all of which compared single incision against standard laparoscopic cholecystectomy; and eight small case series (total 161 patients). The techniques used in the studies varied substantially, particularly with relation to the use of proprietary multiport system solutions or more than one trocar. Extra ports were required in between 6% and 40% of patients in these studies and one patient required open surgery. Few or no follow-up data were presented but two of the case series reported 'barely' or 'hardly' visible scars in all patients (total 25) at 6-week or 6-month follow-up.

With respect to safety, gallbladder perforation was reported in 30% of patients in one case series. Mesenteric injury treated conservatively, injury to the right hepatic duct requiring placement of an endoscopic retrograde biliary drainage stent, cystic artery haemorrhage, bilioma from cystic duct leak due to clip slippage, intra-abdominal free-fluid collection caused by 'bleeding from the liver' which resolved spontaneously, cystic duct stump leak requiring biliary stenting and sphincterotomy, and bleeding requiring transfusion were each reported in one patient in the studies.

Since the main potential advantage of this procedure is cosmetic, the Committee felt that good evidence on safety was particularly important and that this was currently inadequate. They, therefore, recommended that single incision laparoscopic cholecystectomy should only be used with 'special arrangements' for consent and for audit or research. They recommended that future research should include the thorough reporting of complications and a comparison of outcomes with the standard laparoscopic approach.

Mini-incision surgery for total knee replacement (May 2010)

Total knee replacement usually involves patellar eversion (or replacement) and the replacement of joint surfaces of the femur and tibia with prostheses. Mini-incision surgery has been introduced with the aims of improving cosmesis, blood loss and recovery time. The procedure involves an incision usually less than 14 cm long and avoids both patellar inversion and dislocation of the tibiofemoral joint, thereby minimising damage to tendons and muscles. Specially designed instruments are required but the prostheses are the same as those used in standard total knee replacement.

NICE recently reviewed 'special arrangements' guidance published in 2005 which was based on 368 patients from one non-randomised controlled study (70 patients) and four case series with most follow-up less than 1 year. The 2005 guidance encouraged further research on long-term outcomes. The updated evidence presented to the Committee included approximately 2600 patients from three randomised and four non-randomised studies, and two case series with a follow-up intervals around 2 years. There was no difference at 6 months between patients treated by mini-incision and standard incision surgery in the randomised controlled trial, in terms of freedom from pain, but one of the non-randomised studies reported a significantly better mean pain score in those treated with the mini-incision procedure.

The evidence in 2005 included very little information about adverse events. The updated evidence reported a number of complications, including isolated cases of deep wound infection requiring revision surgery, haemarthrosis requiring evacuation, tibial component failure requiring revision, periprosthetic femur fracture after a fall, and patella tendon rupture requiring surgical repair. One non-randomised study reported that patella tendon shortening greater than 5% occurred significantly more often in those treated with conventional surgery but another non-randomised study reported that 'patella clunk' occurred significantly more often in those treated with mini-incision procedure.

The Committee considered that the evidence on both safety and efficacy of the procedure was now adequate to remove the need for any 'special arrangements'. The guidance stipulates that mini-incision total knee replacement

should only be carried out by surgeons with specific training in the procedure and that results should be submitted to the National Joint Register.

Endoscopic saphenous vein harvest for coronary artery bypass grafting (May 2010)

Harvesting of the long saphenous vein for coronary artery bypass is traditionally done by a long incision over the vein. These incisions on the leg may be complicated by delayed healing, wound infection, dehiscence, local loss of sensation and pain. Endoscopic vein harvesting has been introduced with a view to reducing these problems. It is interesting to observe that cardiac surgeons have continued to use a single, long incision for vein harvesting while peripheral vascular surgeons almost universally use multiple smaller incisions with a view to reducing wound complications.

NICE had produced 'normal arrangements' guidance for this procedure in 2007 but was stimulated to review that guidance following publication in the *New England Journal of Medicine* of a non-randomised trial of 3000 patients which suggested a higher rate of death, myocardial infarction and the need for further revascularisation in patients whose veins had been harvested endoscopically, compared with open vein harvest.¹

The evidence presented to the Committee in 2007 was based on 4416 patients from one meta-analysis (1156 patients), one randomised (112 patients) and four non-randomised studies (all short-term follow-up with maximum 42 days except for the randomised controlled trial, which observed all 112 patients to 5 years) and two letters. The updated evidence included approximately 16,855 patients from two meta-analyses (9462 and 3689 patients), five randomised (200, 144, 132, 120 and 112 patients) and three non-randomised studies, and a number of other reports.

Both graft patency and 5-year event-free survival (free from death, myocardial infarction, recurrent angina or congestive heart failure) were reported to be similar between those treated with the open and endoscopic techniques in three randomised controlled trials. A meta-analysis of eight studies reported a similar mortality rate between those treated with the endoscopic and open techniques. However, the recent *NEJM* non-randomised comparative study of 3000 patients reported both a significantly higher vein graft failure at 12.6-month follow-up and also significantly more incidences of death and myocardial infarction in those treated with the endoscopic technique than those treated with the open technique.

The inconsistency in the literature led the Committee to change their recommendations so that the procedure must only be used with 'special arrangements' for clinical governance, consent and audit or research. The new guidance also recommends that patients should be entered into the Central Cardiac Audit Database (CCAD). It suggests that

further research should include both retrospective data analyses and prospective studies including graft imaging to demonstrate patency rates and long-term clinical outcomes – to try to resolve the uncertainty about this procedure.

Stapled transanal rectal resection for obstructed defaecation syndrome (June 2010)

Obstructed defaecation syndrome (ODS) is a complex, multifactorial condition. Conservative treatments such as diet, biofeedback, laxatives and pelvic floor retraining are used as first-line treatments. ODS is often associated with structural abnormalities and surgical treatment to correct these may be required when conservative treatments fail. Prolapse or rectocele are commonly responsible for ODS and stapled transanal rectal resection (STARR) has been introduced as a treatment. The procedure is similar to stapled haemorrhoidopexy where the prolapsing rectum is stapled circumferentially, either with two circular staples or a specially-designed stapling device.

NICE recently reviewed 'special arrangements' guidance published in 2006 which was based on approximately 274 patients from one randomised controlled trial (50 patients) and five case series with a maximum mean follow-up of 22 months. The updated evidence presented to the Committee included approximately 3530 patients from two randomised (119 and 50 patients) and one non-randomised studies (73 patients), eight case series (one including 2838 patients) and one case report. Both randomised controlled studies reported that STARR had a better outcome (less symptoms or obstructed defaecation syndrome score) than stapled transanal prolapsectomy (STAPL) or biofeedback. A case series of 2838 patients (based on a register) reported a significant reduction in ODS score after 12 months.

With respect to safety, shock and death as a result of necrotising pelvic fasciitis (1), rectal necrosis (1), rectovaginal fistula (2), dyspareunia (3) were reported. 'Septic events' were reported in 4% of patients in one study, and postoperative faecal incontinence in 9% and defaecatory urgency in 6% in another study. Early stenosis rates ranged from 1–2%. Comparative studies against STAPL reported that postoperative faecal incontinence and defaecatory urgency both occurred less often with STARR (8% vs 3% and 16% vs 4%).

In the context of a condition which can significantly affect quality of life, the Committee considered that the evidence on safety and efficacy was adequate to support the use of the procedure without any 'special arrangements'. However, the guidance stipulated that the procedure should only be carried out in units specialising in the investigation and management of pelvic floor disorders and that patient selection and management should involve a multidisciplinary team including a colorectal surgeon experienced in the procedure.

Other Guidance

Between May and June 2010, NICE published further guidance (or reviews of existing guidance) of relevance to surgeons (listed below) which could not be summarised here. Please refer to the NICE website (<www.nice.org.uk>) for details.

Interventional Procedures Guidance

- Epithelial radiofrequency ablation for Barrett's oesophagus
- Laparoscopic radical hysterectomy for early stage cervical cancer
- Limited macular translocation for wet age-related macular degeneration
- Macular translocation with 360° retinotomy for wet age-related macular degeneration
- Prosthetic intervertebral disc replacement in the cervical spine
- Therapeutic hypothermia with intracorporeal temperature monitoring for hypoxic perinatal brain injury
- Haemorrhoidal artery ligation

- Extracranial to intracranial bypass for cerebrovascular disease
- Photodynamic therapy for Barrett's oesophagus
- Percutaneous occlusion of the left atrial appendage in non-valvular atrial fibrillation for the prevention of thrombo-embolism

Clinical Guidelines

- Lower urinary tract symptoms
- Constipation in children and young people
- Chronic obstructive pulmonary disease

Cancer Service Guidance

- Skin tumours including melanoma (cancer service guidance)

Reference

1. Lopes RD, Hafley GE, Allen KB, Ferguson TB, Peterson ED, Harrington RA *et al.* Endoscopic versus open vein-graft harvesting in coronary-artery bypass surgery. *N Engl J Med* 2009; **361**: 235–44.

NHS Evidence – promoting efficiency and quality through evidence-based decisions

GILLIAN LENG* and CARMEL THOMASON**

**Chief Operating Officer for NHS Evidence, **Senior Publications Manager*

NHS Evidence was launched last year to manage the synthesis and spread of knowledge in the NHS.¹ Its introduction has ensured that everyone working in health and social care has free access to the quality-assured, best-practice information required to inform evidence-based decision making, quickly and easily.

An information revolution

NHS Evidence was developed to address the frustration health and social care staff reported at trying to find the best evidence to support decision making.² It was not that there was insufficient information, but rather too much in too many different places.

The full range of evidence relating to health and social care was available only to those who had been granted access and had the skills and knowledge to search for it.

This is no longer the case. Now, by using NHS Evidence, staff only need to do a single search to find out what best practice looks like as well as how to deliver it. NHS Evidence does the rest, sifting through more than 150 quality sources of information simultaneously and prioritising

the results. Sources include the National Institute for Health and Clinical Excellence, The Cochrane Library, British National Formulary and The Royal College of Surgeons of England.

From the homepage there is also a link to NHS Athens Resources, from which you can use your NHS Athens login to view e-books, journals and healthcare databases purchased for NHS staff by the SHAs and local trusts.

In the first year, over 9 million searches were carried out on NHS Evidence. Feedback highlighted that users wanted to save and share search results. To accommodate this, a personalisation service, My Evidence, was developed. Registering allows users to create a personal space, which allows search results to be saved and sends out automatic updates when new evidence has been added.

Other developments to improve access have included a downloadable search bar, which any organisation can add to its intranet or website and a mobile version which allows access to evidence on the move.

By putting health and social care practitioners in control of information, NHS Evidence has set in train an information revolution which is helping to make clinical excellence a reality.

Specialist resources

In addition to providing web, medical database, journal and e-book searches, NHS Evidence is also home to more than 30 digital Specialist Collections, which provide Annual Evidence Updates on key areas, giving a straight-forward and succinct overview of what new research and evidence has been published so that you do not need to spend the time searching.

NHS Evidence – surgery, anaesthesia, peri-operative and critical care is run by a specialist team based at The Royal College of Surgeons of England library. Under the guidance of four clinical leads, it provides quick and easy access to current, quality assured, evidence-based resources to support surgeons, anaesthetists, intensivists and all members of the theatre and critical care team.

Regular evidence updates from the collection highlight any new evidence that might challenge current practice, as described in the most recent, accredited national guidance, and provide a commentary on its likely impact. Recent updates published by NHS Evidence – surgery, anaesthesia, peri-operative and critical care have focused on the surgical management of faecal incontinence and critical illness rehabilitation. The updates included a list of relevant national policy, guidelines and systematic reviews, as well as other significant primary research in these fields.

QIPP – Quality, Innovation, Productivity and Prevention

QIPP has been designed to help the NHS meet the current challenges of addressing inefficiencies while also improving patient care.⁵

The QIPP collection on NHS Evidence brings together practical examples of how NHS organisations are improving quality whilst making efficiency savings. It allows colleagues across the NHS to share best practice and deliver successful improvements without having to re-invent the wheel.

Examples on the QIPP collection range from one-off studies from single organisations to robustly tested large-scale changes that have been replicated in multiple organisations with results published in academic journals.

For example, the productive operating theatre programme is estimated to produce savings of £1.7 million for an average-sized trust per annum with an improvement in quality of care in multiple areas. The programme, developed by the NHS Institute for Innovation and Improvement, has been designed to produce a long-term cultural change in the performance of operating theatres.

The aim of the QIPP collection is to build a comprehensive base of best-practice examples that others can learn from. It also recognises the good work being done in many NHS

organisations and NHS Trusts across the country by identifying them as beacons of best practice.⁴

Raising standards

NHS Evidence not only encourages the use of evidence-based medicine but is also working to raise the standards of available evidence.

A key achievement for NHS Evidence has been the development of the first evidence accreditation scheme. Guidance producers, for example, those producing clinical guidelines, referral guidelines and best practice statements, are invited to apply for accreditation. Processes for producing guidance are then assessed against a strict set of criteria and passed to an independent advisory committee for approval.

If the application is successful, the guidance producer is awarded the NHS Evidence Accreditation Mark, a quality mark which helps users sift out the best guidance in a given area.

The NICE Interventional Procedures Programme is one of 13 organisations recognised under the scheme, being formally accredited in February this year. The NICE Interventional Procedures Programme assesses the efficacy and safety of interventional procedures, with the aim of protecting patients and helping clinicians, healthcare organisations and the NHS to introduce procedures appropriately. Most of the procedures that the Interventional Procedures Programme examines are not well established in clinical practice; however, the Programme also scrutinises established procedures if there is uncertainty about their efficacy and/or safety. The award confirms the high quality of the scientific work involved in producing the guidance and the clarity of the advice given.

The Accreditation Mark provides added confidence to any users looking for guidance when making clinical decisions. It is a quick way of identifying that guidance has been produced to exacting standards and can be relied upon.

Further information on NHS Evidence is available from <www.evidence.nhs.uk>.

References

1. Leng GC. NHS Evidence – better decisions and improved patient care. *Ann R Coll Surg Engl* 2009; **91**: 526–31.
2. Masters K. For what purpose and reasons do doctors use the Internet: a systematic review. *Int J Med Inform* 2006; **77**: 4–16.
3. Department of Health. *The NHS quality, innovation, productivity and prevention challenge: an introduction for clinicians*. London: DH, 2010 <www.dh.gov.uk/en/Publicationsandstatistics/Publications/PublicationsPolicyAndGuidance/DH_113806>.
4. Department of Health. *National Innovation Procurement Plan*. Department of Health. London: DH, 2009 <www.dh.gov.uk/en/Publicationsandstatistics/Publications/PublicationsPolicyAndGuidance/DH_109969>.