arthroscopy port to ensure that no irrigation fluid is lost as back-flow out of the port. This allows the surgeon to concentrate on adequate wash-out of the entire joint. It also enables accurate assessment of the volume of fluid used and minimises saline spillage onto the floor of the operating theatre. This is a useful technique which helps to simplify the procedure and also improves theatre safety.

An alternative technique for mesh repair of an incisional hernia

AZ PANTLING, H ALDEAN, D NEHRA
Department of General Surgery, Epsom and St Helier Hospitals, Carshalton, UK

CORRESPONDENCE TO

Anna Z Pantling, E: anna.pantling@esth.nhs.uk

BACKGROUND

Incisional hernias are reported in up to 13% of patients who undergo primary abdominal incisions¹ and high recurrence rates of 25–90% three years following primary repair.^{1,2} Due to high recurrence rates many methods are described, commonly using a mesh that is placed in an on-lay, preperitoneal, or intraperitoneal position. Many patients find incisional hernias uncomfortable and aesthetically displeasing and, like any hernia, may lead to bowel obstruction. Meshes placed intraperitoneally despite being dual layered have a significant incidence of adhesion formation and fistulation into the bowel.³

TECHNIQUE

We suggest an on-lay mesh repair technique in an incisional hernia where it is possible to close the defect. The defect is closed primarily with a continuous 1 loop PDS suture incorporating a lightweight mesh (Ultrapro®; Ethicon, PO Box 1988, Livingston EH54 0AB, UK) within the suture line. This should reduce the

possibility of the suture cutting through tissues and, therefore, add strength to the repair at an early stage. Tacks (Versatek® Covidien, 154 Fareham Road, Gosport PO13 0AS, UK) are placed around the perimeter of the mesh to hold it flat (Fig. 1).

DISCUSSION

This method is similar to the on-lay mesh method; however, it incorporates the mesh into the repair of the defect. Twenty-five cases have been performed and followed up for 2 years with only one recurrence reported so far. We continue to follow-up these patients to gain long-term results. This method could be extended to close any laparotomy where an incisional hernia is deemed likely to occur such as in obese patients.

References

1. The Society for Surgery of the Alimentary Tract. *Surgical Repair of Incisional Hernias*. Patient Care Guidelines, 2005.
2. Khaira HS, Lall P, Hunter B, Brown JH. Repair of incisional hernias. *J R Coll Surg Edinb* 2001; **46**: 39–43.
3. Leber GE, Garb JL, Alexander AI, Reed WP. Long-term complications associated with prosthetic repair of incisional hernias. *Arch Surg* 1998; **133**: 378–82.

A safe technique for olecranon osteotomy

MARTIN MARSH, NEIL PATEL, DAVID LIMB
Department of Trauma and Orthopaedics, Leeds General Infirmary, Leeds, UK

CORRESPONDENCE TO

Martin Marsh, E: martin.marsh@doctors.org.uk

BACKGROUND

Olecranon osteotomy is a well-established technique, providing access to the distal humerus for reduction and fixation of intra-articular fractures.¹ The sigmoid notch of the proximal ulna articulates with the trochlea of the distal humerus and has two articular facets separated by the bare area. Ideally, the osteotomy should enter the joint at the bare area while ensuring the saw itself does not enter the joint, where it can damage the distal humerus.

TECHNIQUE

The fascia and muscle are elevated from the proximal ulna and the capsule of the elbow joint is opened both laterally and medially. The ulna nerve is identified and protected. A swab is pushed through, which is very easy when the distal humerus is fractured (Fig. 1). The swab then indicates the narrowest part of the ulna and, therefore, locates the bare area, allowing for a safe osteotomy with protection of the articular surface (Fig. 2). The osteotomy is completed with the use of an osteotome.

DISCUSSION

As the bare area lies at a relatively constant location,² at the narrowest part of the ulna, we advocate the use of a swab placed through the

Figure 1 Diagrammatic of on-lay mesh showing tacks and sutures.