Published in final edited form as:

Spine (Phila Pa 1976). 2012 March 15; 37(6): 508–514. doi:10.1097/BRS.0b013e318221e83f.

Back Pain among Farmers in A Northern Area of China

Xiaotong Liu¹, Limin Wang, MD, MPH², Lorann Stallones, PhD³, Krista K. Wheeler, MS¹, Weiyan Zhao, MD, PhD¹, Gary A. Smith, MD, DrPH¹, and Huiyun Xiang, MD, PhD¹

¹Center for Injury Research and Policy, The Research Institute at Nationwide Children's Hospital, The Ohio State University College of Medicine, Columbus, Ohio, USA

²Division of Disease Surveillance, National Center for Chronic Disease Control and Prevention, Chinese Center for Disease Control and Prevention, Beijing, People's Republic of China

³Colorado Injury Control Research Center, Department of Psychology, Colorado State University, Fort Collins, Colorado, USA

Abstract

Study Design—This was a population-based survey conducted in 2008 in a northern area of China.

Objective—To investigate back pain prevalence and to examine the associations between potential risk factors and back pain among Chinese farmers.

Summary of Background Data—Few studies have investigated back pain and its associated risk factors among farmers in low-income and developing nations.

Methods—Farmers ages 15 years and older were chosen from 800 families in Heilongjiang Province of People's Republic of China using cluster sampling methods. Complete survey data were obtained from 2045 farmers. The prevalence of self-reported back pain during the previous three-months was reported. Associations between back pain and potential risk factors were examined in logistic regression models which included age, gender, education levels, perceived stress, main farm activities, smoking, and drinking status.

Results—A total of 786 (38.4%) farmers reported back pain. Two-thirds of those with back pain (66.0%) reported that back pain affected work quantity and quality. The adjusted odds ratios of reporting back pain increased with advancing age. Females and farmers who experienced stress regularly were also more likely to report back pain.

Conclusion—Back pain is a common problem among Chinese farmers and is reported more frequently by females. Significant positive associations of gender, age, and perceived stress with back pain warrant additional study.

Keywords

back pain; stress; farmers; agricultural work; China; epidemiology

Corresponding author Huiyun Xiang, Center for Injury Research and Policy, The Research Institute at Nationwide Children's Hospital, Columbus, Ohio, 43205. Huiyun.Xiang@nationwidechildrens.org Telephone: 614-722-2400, Fax: 614-722-2448.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

INTRODUCTION

Among working adults, low back pain (LBP) is one of the principal causes of disability in industrialized nations.¹ Occupational risk factors contribute to 37% of LBP worldwide.² Moreover, back pain leads to substantial losses in financial revenue and to reductions in quality of life. Farming has become increasingly recognized as an occupation that poses a high risk for back pain. The risk of back and joint pain has been found to be higher among farmers when compared with workers from other occupational sectors, including manual workers.³

A number of contributing factors have been associated with back pain among farmers. Potential risk factors include physical occupational demands such as manual handling, repetitive motion, and whole body vibration and mechanical shock from farming equipment.⁴⁻⁶ Additionally, several studies in industrialized nations have shown that people with mental distress and people who smoke are more likely to suffer from back pain.⁷⁻⁹

While there has been research in the United States^{4,10} and other developed countries,^{5,11} less is known about the factors associated with back pain among farmers in low-income and developing nations. In the past three decades, the still-developing People's Republic of China (PRC) has emerged as a prevailing force on the world market. Producing 30% of the world's rice, 20% of the world's corn, 25% of the world's cotton, an estimated 37% of the world's fruit and vegetables, and half of the world's pork, China has become the largest agricultural producer and consumer.¹² Not only does China export to the rest of the world, it must also sustain its own population of 1.3 billion. It is thus imperative to investigate and to address the issue of back pain among China's vast agricultural workforce.

The objectives of this study were to: 1) describe the prevalence of back pain among Chinese farmers by selected demographic characteristics and 2) examine the associations between potential risk factors (perceived stress, smoking and alcohol drinking) and back pain among Chinese farmers.

MATERIALS and METHODS

Data Source

We analyzed data from a population-based survey conducted in 2008 that was a collaborative effort between the Center for Injury Research and Policy (CIRP) of the Research Institute at Nationwide Children's Hospital, the Colorado Injury Control Research Center (CICRC) of the Psychology Department at Colorado State University, and a research team from the School of Public Health of Qiqihar Medical University. These data were previously used by Wang et al. in a study investigating the association between alcohol consumption and work-related injuries among Chinese farmers. The survey procedures and study methods were described in detail in their study.¹³

Study Design and Sampling

Qiqihar is a population center located in China's northernmost province, Heilongjiang Province. Approximately 59% of Qiqihar's population lives in surrounding agricultural areas. Information about residents was obtained from Qiqihar's mandatory registration of residents. The following procedures were utilized to select the sample for this study. First, 800 families were selected through systematic sampling from 9 villages with similar soil and farming techniques. Using cluster survey sampling methods, rural residents 15 years of age and older were surveyed.

With help from CIRP and CICRC, the research team at the School of Public Health of Qiqihar Medical University developed the survey questionnaire. After obtaining approval from the Colorado State University Institutional Review Board and the Scientific Research Committee of the Qiqihar Medical University, the study was officially launched.

Trained in May 2008 by the principal investigator, 25 students from the School of Public Health of Qiqihar Medical University served as interviewers. Student interviewers filled in all questionnaires by conducting face-to-face interviews. Data collection began on May 7 and was completed by May 25 in 2008. The principal investigator at the university randomly chose 25 questionnaires following completion of the initial survey and met with the respondents to verify the consistency of the data collected. Data agreement was greater than 90%, and so the previously collected data were considered reliable.

Definition of Back Pain

A specific question about back pain was asked in the survey: “During the past three months, that is, February to April, 2008, did you have back pain that lasted for a full day or more?” In this study, a positive response (yes) to this question was defined as back pain. To those who self-reported back pain, another question was asked: “Did any of the back pain you had during the past three months affect your work quality or quantity?” Respondents who answered “yes” were defined as having back pain that affected work.

Potential Risk Factors

The potential risk factors examined in this study were gender, age, education, main agricultural activities, perceived life stress, cigarette use, and alcohol consumption. Subjects were asked to evaluate the overall level of stress they felt in relation to external factors, as well as whether they perceived stress in their family and neighbor relationships. Subjects were asked whether they smoked or drank, as well as a detailed history of their use.

Statistical Analysis

EpiData 3.02 was used to collect and store the data.¹³ Both SPSS 13.0 and SAS 9.1 were used in the data analysis. Our analysis was completed in phases. First, the prevalence of self-reported back pain was compared across gender, age, education, and main agricultural activities using Pearson Chi-square tests. Similarly, comparisons in back pain prevalence were considered across stress, smoking, and alcohol consumption variables. Finally, logistic regression models were fitted with back pain and back pain that affected work as the outcome variables and gender, age, education, main farm activities, perceived stress, smoking, and alcohol use status as independent variables. Odds ratios (ORs) with 95% confidence intervals (CIs) were reported. We also evaluated the use of the Generalized Estimating Equation (GEE) to account for non-independence because more than one adult could participate from the each family (using the SAS GENMOD procedure). The original survey did not include a cluster variable, but we could identify clusters using phone numbers for just over 43% of our sample. On average, 2 individuals from each family (i.e. with the same phone number) participated in our survey. Our statistical results from the standard logistic regression analysis and the GEE that accounted for clustering effects were similar, and the conclusions were the same. Therefore, we choose to present the results of logistic regression models, so we could utilize the full dataset.

RESULTS

Of the 2264 respondents initially selected, only 15 (0.7%) declined to participate in the survey, and another 51 (2.3%) provided incomplete information. After data collection, we found that 148 persons actually spent little time performing farm work so they were

excluded from the study. Therefore, 2045 farmers were included in the final analyses. There were 786 (38.4%) farmers who reported having experienced general back pain and 518 (25.4%) who reported having experienced back pain that affected work within the 3 months preceding the survey. Two-thirds of those with back pain (66.0%) reported that back pain affected work quantity and quality. Because patterns and potential risk factors were similar for back pain and back pain that affected work, the results and discussion in this article focus primarily on general back pain.

Demographic Variables

As shown in Table 1, 40.7% of females reported back pain compared with 36.3% of males ($P=0.041$). Additionally, farmers ages 25 and older reported significantly higher rates of back pain relative to younger farmers (15-24 year olds) with those over age 35 reporting back pain over 40% more often than the younger farmers. Farmers whose main farm activity was crop cultivation were more likely to report back pain than farmers whose main activity was breeding livestock, although this difference was not statistically significant (42.1% vs. 37.9%, $P=0.092$).

Perceived Stress, Smoking and Alcohol Consumption

Of the 584 farmers who reported experiencing stress “regularly,” 53.9% reported back pain compared with 31.3% of farmers who reported back pain among those that said they never experienced stress ($P<0.001$) (Table 2). Overall, less than 2% of respondents reported stress in their familial and neighbor relationships, but those with familial stress reported back pain more often than those without that stress ($P=0.003$). Back pain was self reported more often among smokers than among non-smokers ($P=0.008$). Current and former drinkers reported back pain more often than those abstaining from alcohol.

Multivariate Analyses

Gender, age, and stress were statistically significant in the multivariate model (Table 3). The associations of smoking and drinking with back pain were not statistically significant after controlling for other variables in the multivariate model. Females were more likely to experience back pain than males (OR 1.22; 95% CI: 1.09-1.36). The odds ratios of back pain increased with increasing age among farmers. Farmers with regularly perceived stress had an increased odds ratio of reported back pain (OR 2.51; 95% CI: 2.00-3.16) compared with farmers who “never or rarely” experienced stress.

DISCUSSION

The overall prevalence of self-reported back pain in the preceding 3 months among Chinese farmers in Qiqihar was 38.4%; back pain affected the quantity and quality of work for over a quarter of the farmers. Though comparisons to other studies face limitations because of differences in reference periods, study population, and back pain definitions, the results of a U.S. study and several other Chinese studies provide context for our results. In the United States, a study using a large nationally representative dataset, reported a previous 3 month prevalence of LBP (with and without neck pain) of 26.3%.¹⁴ Studies of LBP among farmers, mostly males in the U.S. have sometimes reported higher levels of prevalence.^{4,15} Several studies have described self-reported LBP prevalence rates in Chinese populations. Nearly 67% of farmers in eastern China self-reported back pain in the previous 12 months, and the prevalence rate among women was consistently higher than that among men until age 60.¹⁶ Jin, et al. reported an annual LBP prevalence rate of 50% for 3 types of workers in Shanghai: teachers, battery/ kiln factory workers, and garment workers.¹⁷ A cross-sectional study of villages in Tibet found a point prevalence rate of LBP of 34.1% and a 12-month prevalence of 41.9%.¹⁸ In our multivariate modeling, sex, age, and perceived stress were

statistically associated with back pain while controlling for education, farm activity, smoking, and alcohol use, but gender had only a weak association in the multivariate logistic model. Greater back pain reported among women has been seen in other large population based studies.^{14,19} Of the 4 identified studies about back pain among Chinese general populations or workers,^{16-18,20} only one study of a rural population in Tibet reported no gender-related difference in prevalence of back pain.¹⁸ The three other previous studies and our study suggest that females have slightly higher prevalence of back pain.^{16-18,20} This higher prevalence of back pain among Chinese females may be due to reporting bias (i.e., females may be more likely to report back pain than males) or pain among females due to menstruation.^{16,17}

In the multivariate modeling, advancing age and perceived stress were more strongly associated with back pain than gender. Close to half of Chinese farmers older than 45 years of age reported back pain, and those farmers with regular stress reported back pain more often than those who never or rarely perceived stress. There are a number of literature reviews regarding back pain and psychosocial factors at work^{21,22} and in private life.^{23,24} More recently, a review of review studies has been published.²⁵ While there appears to be value in including psychosocial factors in studies of back pain,^{25,26} associations should be viewed cautiously.²² It is not possible to determine the direction of causality between pain and stress in this cross-sectional survey.

In the univariate analysis, we observed that smokers were significantly more likely to have self-reported general back pain, and individuals with 20 or more years of smoking cigarettes had a significantly higher prevalence of general back pain. In addition, former drinkers were significantly more likely to report general back pain and back pain that affected work quality and quantity. The prevalence of self-reported back pain that affected work also increased with years of alcohol drinking. Our results further suggested that years of alcohol drinking has a larger impact on back pain that affected work than on general back pain. These results could be caused by the confounding effects of age. We found a strong association between age and both general back pain and back pain that affected work in this study. Older individuals are more likely to be former alcohol drinkers and smokers than young individuals in China. Cigarette smoking and alcohol drinking were not associated with back pain in the multivariate model controlling for confounding effects of age. These results suggest that association between years of cigarette smoking/alcohol drinking and self-reported back pain could be explained by the confounding effects of age. According to a systematic literature review of original research reports published between 1987 and 1995, there is no positive association between alcohol consumption and low back pain.²⁷ However, one recent prospective study suggests that moderate alcohol consumption had a protective effect on back pain.²⁸ A meta-analysis of 40 studies published prior to February 2009 indicates that the association between smoking and low back pain was fairly modest.²⁹

This study has several limitations. First, because of the variability of estimates and their high sensitivity to various factors, such as inquiry phrasing, setting and cultural differences, comparison of prevalence rates of back pain among various populations and countries can be problematic.^{3,30} In future studies, standard definitions of back pain for use in epidemiological studies developed and agreed upon by a group of international experts should be used.³¹ Their use will facilitate comparisons among back pain studies internationally.³² Second, results were based on self-reports of back pain and the potential risk factors and may have been affected by recall bias. Also, due to cultural stigmatization regarding alcohol consumption and cigarette usage, it is possible that participants underreported actual usage. Third, although the study questionnaire asked a general question about whether subjects “perceived life stresses,” the more detailed questions focused only on familial and neighbor relationships, not allowing assessment of other sources of stress, such

as job, economic and other life pressures. In addition, the psychometric properties of the stress questions used in the study have not been validated. Finally, because our study was cross-sectional, causality cannot be evaluated.

Conclusions

This study found a high prevalence of self-reported back pain among farmers in this northern area of China. Because two-thirds of those with back pain (66.0%) reported that back pain affected work quantity and quality, back pain is an important public health issue among agricultural workers in China. Future research should address remaining important questions regarding prevalence of back pain, risk factors, and effective interventions.

Acknowledgments

Dr Limin Wang was a trainee of the USA-China Agricultural Injury Research Training project. Grant sponsor: National Institutes of Health Fogarty International Center (PIs: L Stallones and H Xiang; Grant #: 1D43TW007257-01A2). The findings and conclusions in this report are those of authors and do not necessarily represent the views of the funding agency.

We thank researchers and students at the School of Public Health, Qiqihar Medical University for data collection.

REFERENCES

1. Mngoma N, Corbiere M, Stevenson J. Pain profiles and psychosocial distress symptoms in workers with low back pain. *Physiother Can.* 2008; 60:239–45. [PubMed: 20145756]
2. Punnett L, Pruss-Ustun A, Nelson DI, et al. Estimating the global burden of low back pain attributable to combined occupational exposures. *Am J Ind Med.* 2005; 48:459–69. [PubMed: 16299708]
3. Leino-Arjas P, Hanninen K, Puska P. Socioeconomic variation in back and joint pain in Finland. *Eur J Epidemiol.* 1998; 14:79–87. [PubMed: 9517877]
4. Rosecrance J, Rodgers G, Merlino L. Low back pain and musculoskeletal symptoms among Kansas farmers. *Am J Ind Med.* 2006; 49:547–56. [PubMed: 16685722]
5. Milosavljevic S, Bergman F, Rehn B, et al. All-terrain vehicle use in agriculture: exposure to whole body vibration and mechanical shock. *Appl Ergon.* 2010; 41:530–5. [PubMed: 19944407]
6. Walker-Bone K, Palmer KT. Musculoskeletal disorders in farmers and farm workers. *Occup Med (Lond).* 2002; 52:441–50. [PubMed: 12488514]
7. Lepine JP, Briley M. The epidemiology of pain in depression. *Hum Psychopharmacol.* 2004; 19(Suppl 1):S3–7. [PubMed: 15378670]
8. Leijon O, Mulder M. Prevalence of low back pain and concurrent psychological distress over a 16-year period. *Occup Environ Med.* 2009; 66:137–9. [PubMed: 19028803]
9. Zvolensky MJ, McMillan KA, Gonzalez A, et al. Chronic musculoskeletal pain and cigarette smoking among a representative sample of Canadian adolescents and adults. *Addict Behav.* 2010
10. Xiang H, Stallones L, Keefe TJ. Back pain and agricultural work among farmers: an analysis of the Colorado Farm Family Health and Hazard Surveillance Survey. *Am J Ind Med.* 1999; 35:310–6. [PubMed: 9987565]
11. Manninen P, Riihimaki H, Heliovaara M. Incidence and risk factors of low-back pain in middle-aged farmers. *Occup Med (Lond).* 1995; 45:141–6. [PubMed: 7605976]
12. Lohmar B, Gale F. Who Will China Feed? *Amber Waves.* 2008; 6:10–5.
13. Wang L, Wheeler K, Bai L, et al. Alcohol consumption and work-related injuries among farmers in Heilongjiang Province, People's Republic of China. *Am J Ind Med.* 2010
14. Strine TW, Hootman JM. US national prevalence and correlates of low back and neck pain among adults. *Arthritis Rheum.* 2007; 57:656–65. [PubMed: 17471542]
15. Park H, Sprince NL, Whitten PS, et al. Risk factors for back pain among male farmers: analysis of Iowa Farm Family Health and Hazard Surveillance Study. *Am J Ind Med.* 2001; 40:646–54. [PubMed: 11757041]

16. Barrero LH, Hsu YH, Terwedow H, et al. Prevalence and physical determinants of low back pain in a rural Chinese population. *Spine (Phila Pa 1976)*. 2006; 31:2728–34. [PubMed: 17077743]
17. Jin K, Sorock GS, Courtney TK. Prevalence of low back pain in three occupational groups in Shanghai, People's Republic of China. *J Safety Res*. 2004; 35:23–8. [PubMed: 14992843]
18. Hoy D, Toole MJ, Morgan D, et al. Low back pain in rural Tibet. *Lancet*. 2003; 361:225–6. [PubMed: 12547548]
19. Leboeuf-Yde C, Nielsen J, Kyvik KO, et al. Pain in the lumbar, thoracic or cervical regions: do age and gender matter? A population-based study of 34,902 Danish twins 20–71 years of age. *BMC Musculoskelet Disord*. 2009; 10:39. [PubMed: 19379477]
20. Wigley RD, Zhang NZ, Zeng QY, et al. Rheumatic diseases in China: ILAR-China study comparing the prevalence of rheumatic symptoms in northern and southern rural populations. *J Rheumatol*. 1994; 21:1484–90. [PubMed: 7983651]
21. Linton SJ. Occupational psychological factors increase the risk for back pain: a systematic review. *J Occup Rehabil*. 2001; 11:53–66. [PubMed: 11706777]
22. Hartvigsen J, Lings S, Leboeuf-Yde C, et al. Psychosocial factors at work in relation to low back pain and consequences of low back pain; a systematic, critical review of prospective cohort studies. *Occup Environ Med*. 2004; 61:e2. [PubMed: 14691283]
23. Hoogendoorn WE, van Poppel MN, Bongers PM, et al. Systematic review of psychosocial factors at work and private life as risk factors for back pain. *Spine (Phila Pa 1976)*. 2000; 25:2114–25. [PubMed: 10954644]
24. Linton SJ. A review of psychological risk factors in back and neck pain. *Spine (Phila Pa 1976)*. 2000; 25:1148–56. [PubMed: 10788861]
25. Macfarlane GJ, Pallewatte N, Paudyal P, et al. Evaluation of work-related psychosocial factors and regional musculoskeletal pain: results from a EULAR Task Force. *Ann Rheum Dis*. 2009; 68:885–91. [PubMed: 18723563]
26. Linton SJ. Do psychological factors increase the risk for back pain in the general population in both a cross-sectional and prospective analysis? *Eur J Pain*. 2005; 9:355–61. [PubMed: 15979015]
27. Leboeuf-Yde C. Alcohol and low-back pain: a systematic literature review. *J Manipulative Physiol Ther*. 2000; 23:343–6. [PubMed: 10863254]
28. Skillgate E, Vingard E, Josephson M, et al. Is smoking and alcohol consumption associated with long-term sick leave due to unspecific back or neck pain among employees in the public sector? Results of a three-year follow-up cohort study. *J Rehabil Med*. 2009; 41:550–6. [PubMed: 19543666]
29. Shiri R, Karppinen J, Leino-Arjas P, et al. The association between smoking and low back pain: a meta-analysis. *Am J Med*. 2010; 123:87 e7–35. [PubMed: 20102998]
30. Hoy D, March L, Brooks P, et al. Measuring the global burden of low back pain. *Best Pract Res Clin Rheumatol*. 2010; 24:155–65. [PubMed: 20227638]
31. Dionne CE, Dunn KM, Croft PR, et al. A consensus approach toward the standardization of back pain definitions for use in prevalence studies. *Spine (Phila Pa 1976)*. 2008; 33:95–103. [PubMed: 18165754]
32. Walker BF. The prevalence of low back pain: a systematic review of the literature from 1966 to 1998. *J Spinal Disord*. 2000; 13:205–17. [PubMed: 10872758]

Key Points

- The overall prevalence of self-reported back pain in the preceding 3 months among Chinese farmers in Qiqihar was 38.4%.
- Back pain affected the quantity and quality of work for over a quarter of the farmers.
- In multivariate logistic models, gender, age, and perceived stress were associated with back pain while controlling for education, farm activities, cigarette and alcohol use.
- Females and older farmers were more likely to report back pain.
- Farmers under regular stress reported back pain significantly more often than those without perceived stress.

Mini Abstract

We investigated back pain and back pain that affected work among Chinese farmers in a northern area of China. Both were self-reported in face-to-face interviews in this population-based survey. High prevalence of back pain was found among farmers, especially among females, older farmers, and those reporting perceived stress.

Table 1
Demographics and Perceived Stress of Chinese Farmers with Self-reported Back Pain and Back Pain that Affected Work

	Back Pain			Back Pain that Affected Work		
	Total	n	%	P	n	P
Total	2045	786	38.4		519	25.4
Sex						
Male	1073	390	36.3	Ref.	252	23.5 Ref.
Female	972	396	40.7	0.041	267	27.5 0.042
Age (Yrs)						
15-24	195	31	15.9	Ref.	17	8.7 Ref.
25-34	438	130	29.7	<0.001	63	14.4 0.048
35-44	515	213	41.4	<0.001	135	26.2 <0.001
45-54	426	203	47.7	<0.001	154	36.2 <0.001
55-64	326	145	44.5	<0.001	102	31.3 <0.001
65-74	105	49	46.7	<0.001	37	35.2 <0.001
75-84	29	14	48.3	<0.001	11	37.9 <0.001
Education						
Elementary or Below	828	348	42.0	Ref.	248	30.0 Ref.
Middle School	1049	374	35.7	0.005	220	21.0 <0.001
High School and Above	168	64	38.1	0.345	51	30.4 0.807
Main Farm Activities						
Livestock Breeding	1271	482	37.9	Ref.	311	24.5 Ref.
Crop Cultivation	573	241	42.1	0.092	167	29.1 0.034
Other	201	63	31.3	0.073	41	20.4 <0.001

* Row percentages; Ref. - Reference Group

Table 2
Perceived stress, Cigarette and Alcohol Use in Chinese Farmers Self-reporting Back Pain and Back Pain that Affected Work

	Total	Back Pain		Back Pain that Affected Work	
		n	%*	n	%*
Perceived Stress					
Never or Rarely	785	246	31.3%	152	19.4%
Sometimes	667	223	33.4%	144	21.6%
Regularly	584	315	53.9%	222	38.0%
Familial Stress					
No	2003	763	38.1%	501	25.0%
Yes	33	21	63.6%	17	51.5%
Neighbor Stress					
No	2005	769	38.4%	506	25.2%
Yes	31	15	48.4%	12	38.7%
Smoker					
No	1214	438	36.1%	297	24.5%
Yes	831	348	41.9%	222	26.7%
Years Smoking [†]					
Less than 10	152	50	32.9%	31	20.4%
10 to 19	201	77	38.3%	40	19.9%
20 or More	478	221	46.2%	151	31.6%
Drinker					
Never (Abstainer)	1358	495	36.5%	333	24.5%
Former (Quit)	76	39	51.3%	27	35.5%
Current	611	252	41.2%	159	26.0%
Years Drinking [†]					
Less than 5	56	21	37.5%	7	12.5%
5 to 9	89	31	34.8%	16	18.0%
10 to 14	149	55	36.9%	36	24.2%
15 or More	391	182	46.6%	126	32.2%

* Row percentages; Ref.- Reference Group;

† Approximate figures from self-reports

Table 3

Logistic Regression Model of Self-reported Back Pain and Back Pain that Affected Work among Chinese Farmers

	<u>Back Pain</u>		<u>Back Pain that Affected Work</u>	
	OR	95% CI *	OR	95% CI *
Sex				
Male	1.00		1.00	
Female	1.22	1.09-1.36	1.16	1.03-1.32
Education				
Elementary or Below	1.00	0.69-1.43	0.82	0.56-1.21
Middle School	1.07	0.75-1.53	0.77	0.52-1.12
High School and Above	1.00		1.00	
Age (Yrs)				
15-24	1.00		1.00	
25-34	2.12	1.36-3.30	1.72	0.97-3.04
35-44	3.29	2.14-5.07	3.31	1.92-5.70
45-54	4.16	2.67-6.48	5.01	2.89-8.69
55-64	4.00	2.52-6.35	4.46	2.53-7.87
65-74	4.97	2.82-8.75	5.96	3.08-11.55
75-84	5.43	2.32-12.73	6.73	2.66-16.99
Perceived Stress				
Never or Rarely	1.00		1.00	
Sometimes	1.16	0.92-1.46	1.22	0.94-1.59
Regularly	2.51	2.00-3.16	2.52	1.96-3.25
Main Farm Activities				
Livestock Breeding	1.00		1.00	
Crop Cultivation	1.11	0.90-1.37	1.18	0.93-1.49
Other	0.78	0.56-1.09	0.82	0.56-1.20
Smoker				
No	1.00		1.00	
Yes	1.19	0.97-1.47	1.04	0.83-1.31
Drinker				
Never (Abstainer)	1.00		1.00	
Current	1.26	0.99-1.61	1.11	0.85-1.46
Former (Quit)	1.64	1.00-2.68	1.49	0.88-2.51

*
CI: confidence interval