Genome Sequence of *Weissella thailandensis* fsh4-2

Nabil Benomar,¹ Hikmate Abriouel,¹ Heejae Lee,² Gyu-Sung Cho,³ Melanie Huch,³
Rubén Pérez Pulido,¹ Wilhelm H. Holzapfel,² Antonio Gálvez,¹
and Charles M. A. P. Franz^{3*}

Departamento de Ciencias de la Salud, Área de Microbiología, Facultad de Ciencias Experimentales, Universidad de Jaén, 23071 Jaén, Spain¹; Handong Global University, School of Life Sciences, Pohang, Gyeongbuk 791-708, Republic of Korea²; and Max Rubner-Institut, Federal Research Institute for Nutrition and Food, Department of Safety and Quality of Fruit and Vegetables, Haid-und-Neu-Straße 9, D-76131 Karlsruhe, Germany³

Received 27 July 2011/Accepted 1 August 2011

***Weissella thailandensis* fsh4-2 is a heterofermentative lactic acid bacterium isolated from the Korean fermented seafood condiment jeotkal. Here we report the draft genome sequence of *W. thailandensis* fsh4-2 (1,651 genes, 1,436 encoding known proteins, 183 encoding unknown proteins, 32 RNA genes), which consists of 50 large contigs of >100 bp.**

Weissella thailandensis, originally isolated from fermented fish (4), was reported to be able to grow at 10% NaCl. *W. thailandensis* strain fsh4-2 was isolated from jeotkal, a Korean fermented fish condiment which is based on fish, shrimp, and shellfish and contains up to 20% salt. *W. thailandensis* fsh4-2 grew at 10% NaCl *in vitro* using de Man, Rogosa, and Sharpe broth at 30°C. The only other fermented food product from which this species has also been isolated was Mexican Cotija cheese. These isolates grew in medium containing 7% NaCl (3). *W. thailandensis* thus appears to be a particularly salt-tolerant lactic acid bacterial species associated with high-salt and protein-rich food fermentations.

The genomic DNA of *W. thailandensis* fsh4-2 was isolated using the High Pure PCR template preparation kit (Roche). The genome was sequenced by a whole-genome shotgun strategy using Roche 454 GS FLX Titanium pyrosequencing (Lifesequencing S.L.) (386,107 reads totaling 1,968,992 bp [~1.97 Mbp], 76.11-fold coverage of the genome). Quality filtered reads were assembled *in silico* using 454 Life Science Newbler Assembler 2.3, giving 50 contigs (>100 bp in size). Annotation transfer was performed based on open reading frame (ORF) prediction and BlastN gene searching of public databases (UniProt and GenBank). Functional annotation of predicted protein-coding sequences was done using Gene Ontology terms (1) and InterPro motifs compared to the different *Weissella* and *Leuconostoc* proteomes.

The GC content was 40.0 mol%. There were 1,651 possible ORFs in 50 contigs, with sizes ranging from 64 to 1,711 bp. Most were under 1,000 bp, and only a few ORFs ($n = 10$) were >1,000 bp. Predicted copies of the 5S, 16S, and 23S rRNA genes (5 genes); 24 putative tRNA genes; and 3 genes encoding other RNA molecules were also detected. ORFs encoding the enzymes associated with obligately heterofermentative sugar metabolism via the phosphogluconate/phosphoketolase

pathway (2) were present, as well as genes for polysaccharide breakdown and transport and 9 aminopeptidases.

In accordance with the association of *W. thailandensis* with high-salt fermented foods, stress response genes encoding resilience to osmotic stress and changing redox and oxygen levels were identified. In addition to GroES and GroEL chaperonin, HrcA-GrpE-DnaK-DnaJ heat shock protein and an HSP33 heat shock protein were present. *W. thailandensis* also encodes a CspA cold shock protein and two alkaline shock proteins, as well as 4 systems to cope with osmotic shock, including the proline/betaine transporter ProP6, two glycine-betaine/carnitine/choline ATP binding cassette transporters, and one ABC-type proline/glycine-betaine transporter system. Although the genome of *W. thailandensis* does not encode a superoxide dismutase or a catalase, other gene products thought to contribute to intracellular redox balance include three glutathione reductases, one glutathione peroxidase, one iron-dependent peroxidase, one organic hydroperoxide resistance protein, thioredoxin, a possible thioredoxin disulfide reductase, and a thioredoxin reductase.

Nucleotide sequence accession numbers. The draft genome sequence of *W. thailandensis* fsh4-2 is available in the EMBL nucleotide sequence database (project 68571) under accession numbers HE575133 to HE575182.

This work was supported by the Junta de Andalucía (PAI Research Group AGR230) and the University of Jaén (Campus Excelencia Internacional Agroalimentario CEIA3). H. Abriouel was financed by the Ramón y Cajal Research Program.

REFERENCES

1. Ashburner, M., et al. 2000. Gene Ontology: tool for the unification of biology. *Nat. Genet.* **25**:25–29.
2. Axelsson, L. 2004. Lactic acid bacteria: classification and physiology, p. 1–50. In S. von Salminen, A. Wright, and A. Ouwehand (ed.), *Lactic acid bacteria*. Marcel Dekker, Inc., New York, NY.
3. Morales, F., J. I. Morales, C. H. Hernández, and H. Hernández-Sánchez. 2011. Isolation and partial characterization of halotolerant lactic acid bacteria from two Mexican cheeses. *Appl. Biochem. Biotechnol.* **164**:889–905.
4. Tanasupawat, S., O. Shida, S. Okada, and K. Komagata. 2000. *Lactobacillus acidipiscis* sp. nov. and *Weissella thailandensis* sp. nov., isolated from fermented fish in Thailand. *Int. J. Syst. Evol. Microbiol.* **50**:1479–1485.

* Corresponding author. Mailing address: Haid-und-Neu Straße 9, D-76131 Karlsruhe, Germany. Phone: 49 721 6625 225. Fax: 49 721 6625 453. E-mail: Charles.Franz@mri.bund.de.