Published in final edited form as:

J Med Chem. 2011 October 13; 54(19): 6969–6983. doi:10.1021/jm200980u.

Cyclopentane-1,3-dione: A Novel Isostere for the Carboxylic Acid Functional Group. Application to the Design of Potent Thromboxane (A₂)-Receptor Antagonists

Carlo Ballatore^{a,b,*}, James Soper^b, Francesco Piscitelli^a, Michael James^b, Longchuan Huang^a, Onur Atasoylu^a, Donna M. Huryn^a, John Q. Trojanowski^b, Virginia M.-Y. Lee^b, Kurt R. Brunden^b, and Amos B. Smith III^a

^aDepartment of Chemistry, School of Arts and Sciences, University of Pennsylvania, 231 South 34th St., Philadelphia, PA 19104-6323

^bCenter for Neurodegenerative Diseases Research, Department of Pathology and Laboratory Medicine, Institute on Aging, University of Pennsylvania, 3600 Spruce Street, Philadelphia, PA 19104-6323

Abstract

Cyclopentane-1,3-diones are known to exhibit pK_a values typically in the range of carboxylic acids. To explore the potential of the cyclopentane-1,3-dione unit as a carboxylic acid isostere, the physical-chemical properties of representative congeners were examined and compared with similar derivatives bearing carboxylic acid or tetrazole residues. These studies suggested that cyclopentane-1,3-diones may effectively substitute for the carboxylic acid functional group. To demonstrate the use of the cyclopentane-1,3-dione isostere in drug-design, derivatives of a known thromboxane-A₂ prostanoid (TP) receptor antagonist, 3-(3-(2-(4-chlorophenylsulfonamido)ethyl)-phenyl)propanoic acid (**12**), were synthesized and evaluated in both functional and radioligand-binding assays. A series of mono- and di-substituted cyclopentane-1,3-dione derivatives (**41–45**) were identified that exhibit nM IC₅₀ and K_d values similar to **12**. Collectively, these studies demonstrate that the cyclopentane-1,3-dione moiety comprises a novel isostere of the carboxylic acid functional group. Given the combination of the relatively strong acidity, tunable lipophilicity, and versatility of the structure, the cyclopentane-1,3-dione moiety may constitute a valuable addition to the palette of carboxylic acid isosteres.

Keywords

isostere; bio-isostere; TP receptor; antagonist; cyclopentane-1,3-dione

Introduction

Replacement of a specific atom, or group of atoms, with surrogates that exhibit similar physicochemical properties comprises a strategy known as isosteric replacement, adopted broadly by the medicinal chemistry community to improve the properties of a wide variety

*Corresponding author: Department of Chemistry, School of Arts and Sciences, University of Pennsylvania, 231 South 34th St., Philadelphia, PA 19104-6323; bcarlo@sas.upenn.edu; T (215) 898-4891.

Supporting Information Available: Experimental procedures for the synthesis of compounds **12**, **17–19**, **60–62**; X-ray crystal structures of **7–9**, and 2-(cyclopropylmethyl)-3-hydroxycyclopent-2-enone; ¹H-NMR spectra of the complex **1**-benzamidinium used for Job Plot analysis; determination of solubility in acetonitrile for compounds **7–11**; Details of computational studies; Analytical reports for pK_a determinations for compounds **1** and **2**, and pK_a /logP determinations for compounds **7–9**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

of biologically active compounds, such as potency, selectivity, metabolic/chemical stability, and ADME-PK properties.^{1, 2} A typical example of such a strategy is the replacement of a carboxylic acid moiety with a surrogate structure. Indeed, the presence of a carboxylic acid residue in the structure of a drug or drug-candidate is often responsible for significant drawbacks, including limited permeability, toxicity and metabolic instability.² At the same time, however, the acidity of the carboxylic acid residue may be required for biological activity; this is often the case when the carboxylic acid group is directly involved in relatively strong ionic interactions with the biological target (*e.g.*, salt-bridge within the active site of an enzyme). Under such circumstances, replacement of the carboxylic acid moiety with a surrogate structure may be beneficial. The outcome of any isosteric replacement, however, cannot be readily predicted, and thus multiple isosteres are typically required for evaluation. Among the most commonly employed isosteres of the carboxylic acid group, 1-*H*-tetrazoles and 3-hydroxycyclobutene-1,2-diones (Figure 1A), proved to be useful in a number of cases.^{2, 3} Surprisingly, however, cyclopentanepolyones have not been evaluated as carboxylic acid surrogates, although the acidic properties of such compounds have been known for more than 50 years.⁴ For example, cyclopentane-1,3-dione (CPD) and the 2-methyl congener (**1** and **2**, Figure 1) have been reported to have pK_a values of 4.4 and 4.7, respectively.⁴ Equally important, nuclear magnetic resonance (NMR),⁵ infrared (IR) absorption,⁵ and X-ray crystal structure^{6, 7} studies demonstrate that such compounds exist almost exclusively as fast exchanging enol-ketone tautomers (see Figure 1B), which, like carboxylic acids, can establish “head-to-tail” intermolecular hydrogen bonds. In light of these interesting properties, we reasoned that CPDs may hold considerable promise as non-classical isosteres of the carboxylic acid moiety.

To test this hypothesis, we designed, synthesized and evaluated a focused set of model compounds to compare the physical chemical properties of similar structures bearing a carboxylic acid, tetrazole or CPD residue. Electrospray ionization (ESI) mass spectrometry (MS), ¹H-NMR and computational techniques were employed to evaluate the ability of the congeners to form salts with benzamidine. To test the potential of CPD isosteres in drug design, we also designed, synthesized and compared the biological properties of a series of CPD derivatives based on a representative thromboxane-A₂ prostanoid (TP) receptor antagonist (**12**, Figure 2)⁸ to the corresponding analogues bearing other carboxylic acid isosteres such as the tetrazole and amino squaric acid.

Design and Synthesis of Model Compounds

The CPD fragment presents two non-equivalent points of attachment (*i.e.*, the C2 and one of the prochiral C4/C5 positions, Figure 1B) for analogue development. Thus, to evaluate the physical chemical properties of CPDs relative to carboxylic acids and tetrazoles, representative model 2- and 5-monosubstituted (**9** and **7**, respectively, Scheme 1), as well as 2,5-disubstituted (**8**, Scheme 1) CPD derivatives were designed and synthesized. In each case, a 4-bromobenzyl moiety was used as a substituent to ensure appropriate solubility in organic solvents, a prerequisite to enable ESI-MS studies (*vide infra*).

Compounds **7** and **8** were prepared by reacting the appropriate lithiated 3-isobutoxycyclopent-2-enone (*i.e.*, **3** or **4**) with 4-bromobenzylbromide, under the reaction conditions developed by Koreeda and co-workers⁹ to achieve regioselective alkylation at the C5 position, to obtain monoalkylated products **5** and **6** as racemic mixtures. Hydrolysis of the enol-ether under acidic conditions then furnished **7** and **8**. For the synthesis of the 2-substituted CPD **9**, we employed the reductive alkylation protocol reported by Ramachary and Kishor;¹⁰ specifically, 4-bromobenzaldehyde was reacted with CPD in the presence of diethyl 1,4-dihydro-2,6-dimethyl-3,5-pyridinedicarboxylate (Hantzsch ester) to furnish **9**. Construction of tetrazole **11** was achieved as previously reported,¹¹ while the 4-bromophenylpropionic acid **10** was purchased.

Design and Synthesis of CPD-Containing TP-Receptor Antagonists

With respect to TP-receptor antagonist design, depending on the point of attachment on the CPD moiety and the length of the aliphatic spacer, two alternative structures were envisioned, which exhibit a promising degree of overlap with the carboxylic acid moiety of the parent compound **12** (Figure 2).

We therefore synthesized a series of both 2- and 5-substituted CPDs, as well as different 2,5-disubstituted CPD derivatives (Scheme 2). Tetrazole and amino squarate derivatives of **12** (respectively **61** and **62**, Table 2) were also constructed for comparison.

The synthesis of the 5-substituted CPD derivatives **41** and 2,5-disubstituted congeners **42–45** (Scheme 2) employed similar strategies as employed for **7** and **8**. In all cases, alkylation of the CPD was achieved by reacting the appropriate lithiated 3-isobutoxycyclopent-2-enone⁹ (*i.e.*, **3**, **4**, **17–19**) with benzyl iodide **16**, obtained in three steps from aldehyde **13**, to furnish **20–24** (Scheme 2). Removal of the silyl-protecting group then furnished alcohols **25–29**. For installation of the phenylsulfonamide moiety, we initially employed a reaction sequence consisting of: (a) conversion of alcohol **25** to the corresponding alkyl iodide **30**; (b) displacement of the iodide with sodium azide (**31**); and (c) reduction of the latter to the corresponding amine, followed by *in situ* sulfonylation of the amine with 4-chlorobenzenesulfonyl chloride to yield **32** (Scheme 2). Similar overall transformations could be achieved in comparatively fewer steps and in higher overall yield by treating alcohols **26–29** with *N*-Boc-protected phenyl-sulfonylamide under Mitsunobu conditions¹² to furnish **33–36**. Finally, treatment of **32** with 2N hydrochloric acid in acetone provided the desired compound **41**, whereas **33–36** were subjected to stepwise deprotection of the Boc-carbamate and enol-ether, respectively, to furnish **37–40** and **42–45** (Scheme 2).

Construction of the 2-substituted CPD derivatives, **50** and **59** (Scheme 3), began with aldehydes **14** and **54**, respectively. Silyl-protected aldehyde **54** was obtained from di-acid **51** via lithium aluminum hydride reduction, followed in turn by protection as silyl ether of one of the hydroxyl moieties (**53**) and pyridinium dichlorochromate-mediated oxidation of the free hydroxyl. The aldehydes (**14**, **54**) were then reacted with CPD under the previously described reductive alkylation conditions¹⁰ to furnish the 2-alkylated CPD derivatives **46** and **55**. These intermediates were then directly converted to alcohols **47** and **56** by treatment with *i*-butanol/benzene in the presence of a catalytic amount of *p*-toluenesulfonic acid at reflux. Installation of the *N*-Boc-protected 4-chlorophenylsulfonamide moiety via Mitsunobu reaction¹² and removal of Boc and *i*-butyl ether protecting groups then furnished the desired 2-substituted CPD derivatives **50** and **59**.

Synthesis of carboxylic acid **12** (Figure 2), the corresponding alcohol **60**, tetrazole **61**, and amino squaric acid **62** (Table 1), are detailed in the Supporting Information Section.

Physical-Chemical Property Evaluation

The lipophilicity and/or acidity of compounds **1**, **2**, **7–11** were determined experimentally, followed by an examination of their ability to form salts with benzamidine, the latter investigated by ESI-MS and NMR. Finally, computational studies were carried out to estimate the structure and stability of the complexes of CPD derivatives with benzamidine.

Determination of pK_a and logP values—pK_a values of compounds **1**, **2**, **7–11**, as well as the logP and logD_{pH 7.4} values of compounds **7–11**, were determined experimentally by Sirius Analytical (UK) via UV-metric or potentiometric (pH-metric) methods. The results are summarized in Table 1 (see Supporting Information for full reports).

Evaluation of the ability of model CPD-derivatives to form salts with benzamidine

The structure and stability of the amidine complexes with carboxylate and tetrazolate anions have been investigated with a combination of ESI-MS and NMR techniques,^{13, 14} suggesting that this experimental model could be useful to evaluate and compare carboxylic acid isosteres. Thus, we employed similar strategies to evaluate the ability of CPDs **7–9** to form salts with amidines. For ESI-MS studies, a 10 μ M solution of each test compound in acetonitrile was mixed with an equal volume of a 10 μ M solution of benzamidine in acetonitrile. The solubility of **7–11** in acetonitrile had been pre-determined and each of the compounds was found to be fully soluble at a concentration of 10 μ M (see Supporting Information Section). After a 5 min incubation period, the resulting mixtures were analyzed by ESI-MS to detect the mass of the [CPD-benzamidine] complex. As illustrated in Figure 3, in all cases the test compound-benzamidine ion⁺ was readily detectable by ESI-MS.

Competition-binding studies were conducted by comparing the intensity of the signal corresponding to the [**8**-benzamidine] complex with and without co-addition of a competing acid/acid surrogate (*i.e.*, either **7**, **9–11**; see Figure 4). In these experiments, benzamidine, compound **8** and the competing acid/acid surrogate were mixed in a 1:1:1 ratio in acetonitrile and, after a 5 min incubation time, the mixtures were analyzed by ESI-MS to monitor for the presence of the [**8**-benzamidine] complex. The relative intensity of the [**8**-benzamidine] complex in the presence or absence of competing acid/acid surrogate provided qualitative information on the binding affinity of CPD derivatives for amidines, compared to acid **10** or tetrazole **11** (Figure 4). As shown in Figure 4, co-incubation of equimolar amounts of benzamidine, **8** and **9** resulted in a ~50% reduction in the peak intensity assigned to the [**8**-benzamidine] complex, while analogous co-incubations of benzamidine and **8** with either **7**, **10** or **11** resulted in a comparatively smaller reduction (*i.e.*, <50%) of the [**8**-benzamidine] signal. Thus, under the experimental conditions employed in the competition studies, all CPD derivatives appeared to have greater affinity for benzamidine than either the corresponding carboxylic acid (**10**) or tetrazole (**11**) compound (Figure 4).

The interaction of the CPD-benzamidine complex was also investigated by ¹H-NMR. The observed upfield shift of the signal corresponding to the proton in position 2 of CPD **1** at different molar ratios with benzamidine was used to conduct Job plot analysis (see Supporting Information).¹⁵ Based on such an analysis, the minimum of the plot corresponds to the stoichiometry of the CPD-benzamidine interaction. These studies confirmed that CPD forms a 1:1 complex with benzamidine (Figure 5).

Computational studies—CPDs possess two tautomeric forms that can potentially generate different salt bridge geometries. Initially, quantum mechanical (QM) calculations were performed both to explore the ground state geometries and to define the energy differences of the two tautomeric forms of a representative CPD (*i.e.*, **7** and **8**). These studies revealed that regardless of solvent media (*i.e.*, no solvent/vacuum, water or chloroform), the enol-ketone tautomer is comparatively more stable than the diketone tautomer (*e.g.*, for **8** the energy difference is 0.91 kcal/mol; B3LYP 6-31G+(d,p) in the gas phase and, on average, 0.74 kcal/mol in water and chloroform, see Supporting Information). Next, the geometry and stability of [**8**-benzamidine] complexes was investigated. Optimization of possible salt geometries revealed two possible scenarios (A and B), originating from the initial tautomers (*i.e.*, di-ketone and enol-ketone): (a) a bi-coordinated “stacked” geometry (Figure 6A) characterized by two intermolecular interactions between the di-ketone tautomer and the amidine moiety; and (b) a mono-coordinated “linear” geometry, wherein a single interaction between the amidine nitrogen and the enolate oxygen is observed (Figure 6B). Quantum mechanical calculations (gas phase) and with a water solvation model suggest that the latter

salt bridge geometry (6B) is more stable than the former. Disruption of the π system in geometry 6A is believed to be responsible for the energy difference. Calculated energies for the mono-coordinated CPD salt (6B) were found to be similar to those corresponding to the bi-coordinated salt bridges of carboxylic acids with amidine bases (see Supporting Information).

Biological Evaluation

To evaluate whether CPD isosteres could mimic the activity of biologically active compounds, CPD analogues **41–45** and **59** were evaluated for their ability to act as antagonists of the human and mouse TP-receptor by a functional inositol monophosphate (IP₁) assay that measures receptor activation. The activity of test compounds was compared with known antagonist **12**,⁸ as well as with tetrazole **61** and amino squaric acid **62**. Analogues **37–39** and **50** were also examined for comparison. Briefly, the IP₁ assay depends on a homogeneous time-resolved fluorescence method that permits the measurement of IP₁, which is a stable metabolite of the intercellular signal transduction molecule, inositol triphosphate (IP₃). Activation of the TP receptor is known to result in increased release of IP₃,¹⁶ thus an antagonist to this receptor will inhibit agonist-induced increase of IP₁ in QBI293 cells that have been stably-transfected with the human or mouse TP receptor. As illustrated in Table 2, a range of CPD analogues were evaluated and derivatives **42–45** exhibited activity comparable to the known antagonist **12** as well as the tetrazole and amino squarate derivatives (**61** and **62**, respectively).

Finally, the binding affinity of a representative subset of active and inactive compounds against the human and mouse TP-receptor were evaluated via radioligand scintillation proximity assay. In this assay, membrane preparations from cells expressing the human or mouse TP receptor associate with scintillant beads that emit measurable photons upon binding of the radiolabeled TP antagonist, [³H]SQ 29548, to receptors within the membrane. Thus, a ligand that competes for binding of the radiolabeled compound to the TP receptor will cause a concentration-dependent decrease in signal that can be quantified to yield a binding constant for the competitive molecule.¹⁷ Consistent with the functional assay results, these radioligand binding studies reveal that CPD-derivatives **41** and **42** exhibit binding affinity values that are comparable to the known antagonist **12**, as well as the tetrazole (**61**) and the amino squarate (**62**) derivatives (Table 3).

Docking Studies

A TP-receptor model was developed via homology modeling as previously described.¹⁸ Docking studies were then carried out using Autodock software package, initially employing the natural ligand, thromboxane A₂ (TXA₂). Since previous reports suggest that Arg-295 and Ser-201 residues in the intracellular region are important constituents of the binding pocket within the TP-receptor,¹⁹ docking of TXA₂ was conducted by setting a salt bridge between the carboxylic acid residue of TXA₂ and the side-chain of Arg-295 as a constraint. In agreement with previous docking studies by Yamamoto and coworkers,²⁰ our studies indicate that the polycarbon side-chain of TXA₂ extends into the intracellular hydrophobic segment between the III, V and VI helices, with a favorable hydrogen bond to the Ser-201 residue (Figure 7A). Next, the binding modes of compound **12**, and of a representative CPD derivative (**42**), were evaluated and compared (Figure 7B and 7C, respectively). As shown in Figure 7B–C and in Figure 8, compounds **12** and **42** were found to share similar hydrophobic interactions between the central phenyl ring and the pocket formed by Leu-261, Leu-294, Ala-297, and Met-112, as well as between the phenyl-sulfonamide moiety and the pocket between helices III and IV (Trp-258 and Gly-116, see Figure 8). Interestingly, while the carboxylic acid moiety of **12** interacted with Arg-295 via the expected bi-coordinated salt bridge, the corresponding CPD isostere in **42** was found to establish a mono-coordinated

salt bridge with the guanidinium moiety of Arg-295 as well as an hydrogen bond with the backbone nitrogen of Thr-298 (Figure 8). Equally interesting, the 2-position of the CPD fragment appears to be directed through the groove of helix VII, suggesting that an increase in steric hindrance at this position may be well tolerated. This observation is consistent with functional and receptor binding data (*cf.*, Table 2 and 3).

Discussion

Although CPDs share important similarities with the carboxylic acid moiety, including comparable pK_a values and the ability to establish *H*-bonds, there are no reports of this isostere employed as a surrogate of the carboxylic acid moiety in drug design. To evaluate CPDs as potential carboxylic acid isosteres, we compared the properties of representative mono- and di-substituted CPD analogues with similar compounds bearing carboxylic acid or tetrazole moieties. These studies confirmed the intrinsic acidity of the CPD fragment and revealed that compounds **7–9**, depending on the substitution pattern, can be somewhat more acidic than the corresponding carboxylic acid and tetrazole counterparts. Furthermore, the lipophilicity of these compounds is also either equal or higher than acid **10** and tetrazole **11** (Table 1). Importantly, the ability of CPD derivatives **7–9** to form complexes with benzamidine was demonstrated by ESI-MS and NMR. Of particular note, our studies suggest that CPDs can reversibly form 1:1 complexes with benzamidine regardless of the substitution pattern (*cf.*, Figure 3 and 5). Computational studies indicated that the most likely salt bridge geometry involves mono-coordinated structures between the amidine nitrogen and the enolate oxygen of the CPD. Although our studies focused exclusively on the CPD-benzamidine complexes, similar salt geometries may also exist between CPDs and other amines, including the amino groups on lysine and histidine side-chains. Interestingly, results from competition-binding studies indicate that, under the experimental conditions, the CPD-benzamidine complex may be more stable than the corresponding tetrazolate-benzamidine or carboxylate-benzamidine complexes (Figure 4). These results do not appear to correlate exactly with either the pK_a values or calculated salt bridge formation energies; however, the lack of correlation observed may be due to the experimental conditions employed in the ESI-MS studies, particularly the fact that these experiments are performed in organic solvent. Taken together, these results strongly suggested that the CPD fragment holds great promise as an effective surrogate for the carboxylic acid moiety.

To verify this hypothesis in the context of a biologically active system, analogues of the known TP-receptor antagonist **12**⁸ were designed by replacing the carboxylic acid moiety of this compound with a CPD fragment linked at either the C2 or C5 position (Figure 2). The TP-receptor is an important G-protein coupled receptor (GPCR) implicated in several pathophysiological processes, such as vasoconstriction, aggregation of platelets,²¹ and more recently, Alzheimer's disease.²² A number of TP-receptor antagonists have been reported and many of these, like **12**, comprise an aromatic ring connected both to a phenylsulfonamide and a carboxylic acid moiety via intervening aliphatic spacers, typically of 2–3 carbons in length.²¹ Previous studies suggest that the carboxylic acid moiety of TP-receptor antagonists are involved in the formation of a salt-bridge with Arg-295 of the receptor.²⁰ Thus, the presence of a carboxylic acid moiety may be a general pre-requisite for receptor binding and activity. This requirement is clearly the case for compound **12**, as highlighted by the dramatic loss in activity in both the functional and radioligand-binding assays observed when the corresponding alcohol derivative **60** is employed (Table 2 and 3). Conversely, replacement of the carboxylic acid moiety with either tetrazole (**61**) or squaric acid (**62**) led to derivatives with activity and binding affinity generally comparable to **12**. Among the CPD derivatives, interesting differences emerged depending on where the CPD moiety is attached (*i.e.*, C2 or C5). While C2 substituted derivatives **50** and **59** were essentially devoid of any activity in the functional assay, all C5 substituted compounds (**41–**

45) exhibited IC₅₀ values in the same range of carboxylic acid **12** (Table 2). These observations were confirmed by radioligand-binding data (Table 3), demonstrating that representative examples of C2 and C5 linked CPD derivatives exhibited drastically different binding affinities. Given that ESI-MS studies with model compounds **7–9** did not reveal significant differences between the C2 and C5 linked CPD congeners in their ability to generate complexes with benzamidine, the results from the functional assay suggest that the position and/or orientation of the enol-ketone moiety may prove critical in the relatively confined environment of the receptor. Also of note, the steric hindrance of the C2 substituents of **41–45** did not appear to impact the activity of the analogues, suggesting that effective electrostatic interactions with the TP-receptor may take place if the CPD moiety is linked at C5, even in the presence of relatively large substituents linked at C2. Consistent with these results, docking studies revealed that the 2-position of the CPD can be adjacent to the groove of helix VII of the TP receptor, indicating that steric hindrance at the C2 would be relatively well tolerated. This characteristic may be exploited to design analogues with increased complementarity with the receptor and/or to modify the physical-chemical properties (*e.g.*, lipophilicity) of the compound.

Conclusions

Collectively, the studies presented here demonstrate that the CPD fragment comprises a viable new surrogate for the carboxylic acid moiety with potential applications in drug design. In addition to the significant intrinsic acidity and the ability to form salt-bridges, our results demonstrate that the CPD moiety may permit substantial structural differentiation of the acidic residue. This characteristic may be particularly desirable when attempting to modulate drug-target/off-target interactions and/or physical-chemical properties of biologically active compounds. As a result, the CPD would appear to be a valuable addition to the existing palette of carboxylic acid isosteres.

Experimental Section

Materials and Methods

All solvents were reagent grade. All reagents were purchased from Aldrich or Acros and used as received. Thin layer chromatography (TLC) was performed with 0.25 mm E. Merck pre-coated silica gel plates. Flash chromatography was performed with silica gel 60 (particle size 0.040 – 0.062 mm) supplied by Silicycle and Sorbent Technologies. Spots were detected by viewing under a UV light. Yields refer to chromatographically and spectroscopically pure compounds. Infrared spectra were recorded on a Jasco Model FT/IR-480 Plus spectrometer. All melting points were obtained on a Thomas-Hoover apparatus. Proton (¹H) and carbon (¹³C) NMR spectra were recorded on a Bruker AMX-500 spectrometer. Chemical shifts were reported relative to solvents (CDCl₃ 7.27 ppm; CD₃OD 3.35 ppm; DMSO-d₆ 2.5 ppm, Acetone-d₆ 2.05 ppm). *As previously observed,¹⁰ CPDs are mixtures of rapidly interconverting tautomers which can complicate the interpretation of ¹³C-NMR spectra.* High-resolution mass spectra were measured at the University of Pennsylvania Mass Spectrometry Service on either a VG Micromass 70/70H or VG ZAB-E spectrometer. Single-crystal X-ray structure determinations were performed at the University of Pennsylvania with an Enraf Nonius CAD-4 automated diffractometer. Analytical reversed-phased (Sunfire™ C18; 4.6×50 mm, 5 mL) high-performance liquid chromatography (HPLC) was performed with a Water binary gradient module 2525 equipped with Waters 2996 PDA and Water micromass ZQ. All samples were analyzed employing a linear gradient from 10% to 90% of acetonitrile in water over 8 minutes and flow rate of 1 mL/min, and unless otherwise stated, the purity level was >95%. Preparative reverse phase HPLC purification was performed employing Waters SunFire™ prep C₁₈ OBD™ columns (5 μm 19×50 or 19×100 mm). All samples were purified employing a linear

gradient from 10% to 90% of acetonitrile in water over 15 minutes and flow rate of 20 mL/min. The preparative HPLC system was equipped with Gilson 333 pumps, a 215 Liquid Handler, 845Z injection module, and UV detector. Unless otherwise stated, all final compounds were found to be >95% as determined by HPLC/MS and NMR.

5-(4-Bromobenzyl)-3-isobutoxycyclopent-2-enone (5)—3-Isobutoxycyclopent-2-enone (**3**, 0.130 g, 0.84 mmol) in anhydrous tetrahydrofuran (2.0 mL) was cooled to -78°C and a freshly prepared 1M solution of lithium diisopropylamide (1.0 mL, 1.0 mmol) was added dropwise. The mixture was stirred at -78°C for 45 min and a solution of 1-bromo-4-(bromomethyl)benzene (0.210 g, 0.84 mmol) in anhydrous tetrahydrofuran (3.0 mL) was added dropwise. The resulting mixture was stirred for 1 h allowing the temperature to rise to rt. The reaction was quenched with a saturated aqueous solution of ammonium chloride (3.0 mL). The aqueous phase was extracted with ethyl acetate (2×10 mL) and combined organic layers were washed with brine, dried over magnesium sulfate, filtered and concentrated *in vacuo*. Purification by silica gel column chromatography using a gradient of ethyl acetate in hexanes as eluant provided **5** as a white solid (40% yield).

^1H NMR (CDCl_3): δ 0.94 (d, $J = 6.7$ Hz, 6H), 2.00 (m, 1H), 2.27 (dd, $J = 17.7, 2.2$ Hz, 1H), 2.55–2.61 (m, 2H), 2.71–2.74 (m, 1H), 3.12 (dd, $J = 14.0, 4.2$ Hz, 1H), 3.67 (d, $J = 6.4$ Hz, 2H), 5.20 (s, 1H), 7.04 (d, $J = 8.3$ Hz, 2H), 7.35–7.36 (m, 2H) ppm.

^{13}C NMR (CDCl_3): δ 19.05, 19.07, 27.9, 34.1, 36.4, 46.4, 78.1, 104.0, 120.3, 130.8, 131.6, 138.5, 189.2, 206.7 ppm.

IR (film): ν 2962, 2931, 2876, 1692, 1593 cm^{-1} .

5-(4-Bromobenzyl)-3-isobutoxy-2-methylcyclopent-2-enone (6)—Prepared as **5** from **4**. Yield 20%.

^1H NMR (CDCl_3): δ 0.97 (d, $J = 6.5$ Hz, 6H), 1.65 (s, 3H), 1.65–2.00 (m, 1H), 2.21–2.25 (m, 1H), 2.52–2.57 (m, 1H), 2.60–2.65 (m, 1H), 2.73–2.76 (m, 1H), 3.22 (dd, $J = 16.5, 4.5$ Hz, 1H), 3.83 (dd, $J = 1.5, 0.5$ Hz, 2H), 7.08 (d, $J = 8.5$ Hz, 2H), 7.42 (d, $J = 8.5$ Hz, 2H) ppm.

^{13}C NMR (CDCl_3): δ 6.2, 18.9, 28.8, 31.0, 36.7, 45.8, 75.7, 115.4, 120.2, 130.7, 131.6, 138.6, 183.2, 206.1 ppm.

IR (film): ν 3384, 3283, 1725 cm^{-1} .

MS (ESI^+): calculated for $\text{C}_{17}\text{H}_{22}\text{BrO}_2^+$ 337.08; found 337.01.

5-(4-Bromobenzyl)-3-hydroxycyclopent-2-enone (7)—To a mixture of **5** (0.110 g, 0.34 mmol) in acetone (4.2 mL) 2N hydrochloric acid (1.7 mL) was added at rt and the mixture was stirred for 16 h. The reaction mixture was then concentrated under reduced pressure and the residue is purified by preparative reverse phase HPLC providing **7** as a white solid (46% yield).

mp: 206–208 $^{\circ}\text{C}$ (from methanol)

^1H NMR (MeOD): δ 2.20 (dd, $J = 18.1, 2.5$ Hz, 1H), 2.52 (dd, $J = 18.1, 6.9$ Hz, 1H), 2.63 (dd, $J = 13.8, 9.3$ Hz, 1H), 2.93 (m, 1H), 3.08 (dd, $J = 13.8, 4.3$ Hz, 1H), 7.13 (d, $J = 8.3$ Hz, 2H), 7.40 (d, $J = 8.4$ Hz, 2H) ppm.

^{13}C NMR (MeOD): δ 37.47, 37.61, 46.0, 105.9, 121.3, 132.2, 132.6, 139.7, 200.1, 204.9 ppm.

IR (film): ν 2920, 2680, 2562, 1713, 1555 cm^{-1} .

HRMS [ESI] $^{-}$: calculated for $\text{C}_{12}\text{H}_{10}\text{O}_2\text{Br}^{-}$ 264.9864; found 264.9869.

5-(4-Bromobenzyl)-3-hydroxy-2-methylcyclopent-2-enone (8)—Prepared as **7** from **6**. Yield 84%.

mp: 180–181 $^{\circ}\text{C}$ (from methanol).

^1H NMR (CD_3OD): δ 1.55 (s, 3H), 2.16 (d, J = 17.5 Hz, 1H), 2.49 (d, J = 17.5 Hz, 1H), 2.58 (dd, J = 13.5, 8.5 Hz, 1H), 2.84–2.88 (m, 1H), 3.11 (dd, J = 14.0, 4.0 Hz, 1H), 7.13 (d, J = 8.5 Hz, 2H), 7.40 (d, J = 8.5 Hz, 2H) ppm.

^{13}C NMR (CD_3OD): δ 5.7, 36.0, 37.8, 45.2, 114.1, 121.3, 132.2, 132.6, 139.8 ppm.

IR (film): ν 2922, 2639, 1570 cm^{-1} .

MS (ESI $^{+}$): calculated for $\text{C}_{13}\text{H}_{14}\text{BrO}_2^{+}$ 281.02; found 281.13.

2-(4-Bromobenzyl)-3-hydroxycyclopent-2-enone (9)—To a solution of cyclopentane-1,3-dione (0.200 g, 2.0 mmol), diethyl 2,6-dimethyl-1,4-dihydropyridine-3,5-dicarboxylate (0.520 g, 2.0 mmol) and *L*-proline (0.012 g) in dichloromethane (6.6 mL) 4-bromobenzaldehyde (1.130 g; 6.0 mmol) was added and the resulting mixture was allowed to stir at rt for 30 min. Purification by silica gel column chromatography using a gradient of ethyl acetate in hexanes as eluant provided **9** (75% yield).

mp: 199–201 $^{\circ}\text{C}$ (from methanol).

^1H NMR (CD_3OD): δ 2.51 (s, 4H), 3.38 (s, 2H), 7.12 (d, J = 8.2 Hz, 2H), 7.33 (d, J = 8.3 Hz, 2H) ppm.

^{13}C NMR (MeOD): δ 27.2, 31.5, 117.6, 120.5, 131.5, 132.3, 141.0 ppm.

IR (film): ν 2911, 2532, 1565 cm^{-1} .

HRMS (ESI $^{-}$): calculated for $\text{C}_{12}\text{H}_{10}\text{BrO}_2^{-}$ 264.9864; found 264.9872.

3-(2-Hydroxyethyl)benzaldehyde (13)—A solution 2-(3-bromophenyl)ethanol (1.00 g, 0.67 mL, 4.9 mmol) and *N,N,N',N'*-tetramethylethylenediamine (1.5 mL) in anhydrous diethyl ether (10 mL) was cooled to -78°C in dry-ice acetone bath. *n*-Butyl lithium (2.4 M solution in hexanes, 4.0 mL, 9.8 mmol) was added dropwise and the resulting mixture was allowed to warm to -20°C (over 1 h) and then cooled again to -78°C . Anhydrous *N,N*-dimethylformamide (5 mL) was added and the resulting mixture was warmed to rt and stirred for an additional 1 h. The reaction was quenched with a saturated solution of ammonium chloride and extracted with ethyl acetate. The combined extracts were washed with brine, dried over magnesium sulfate, filtered and concentrated *in vacuo*. Purification by silica gel column chromatography using ethyl acetate-hexanes 2:3 as eluant furnished **13** as colorless oil (88% yield).

^1H NMR (CDCl_3): δ 1.69 (broad s, 1H), 2.96 (t, J = 6.5 Hz, 2H), 3.91 (t, J = 6.5 Hz, 2H), 7.47–7.53 (m, 2H), 7.74–7.76 (m, 2H), 10.00 (s, 1H) ppm.

^{13}C NMR (CDCl_3): δ 39.0, 63.4, 128.4, 129.4, 130.1, 135.5, 136.9, 140.1, 192.6 ppm.

IR (film): ν 3381, 1696 cm^{-1} .

3-(2-((Triisopropylsilyl)oxy)ethyl)benzaldehyde (14)—Chlorotriisopropylsilane (0.640 g, 0.71 mL, 3.3 mmol) was added dropwise to a solution of **13** (0.450 g, 3.0 mmol) and 1*H*-imidazole (0.410 g, 6.0 mmol) in anhydrous *N,N*-dimethylformamide (10 mL) at 0 °C and the resulting mixture was stirred at 0 °C for 3 h. The reaction mixture was then diluted with water and extracted with diethyl ether. The combined extracts were washed with brine, dried over magnesium sulfate, filtered and concentrated *in vacuo*. Purification by silica gel column chromatography using ethyl acetate-hexanes 1:9 as eluant furnished **14** as colorless oil (96% yield).

^1H NMR (CDCl_3): δ 1.00–1.10 (m, 21H), 2.93 (t, J = 6.6 Hz, 2H), 3.93 (t, J = 6.6 Hz, 2H), 7.45 (t, J = 7.5 Hz, 1H), 7.52 (d, J = 7.6 Hz, 1H), 7.73 (d, J = 7.5 Hz, 1H), 7.76 (s, 1H), 10.00 (s, 1H) ppm.

^{13}C NMR (CDCl_3): δ 12.1, 18.1, 39.5, 64.4, 127.9, 129.0, 130.6, 135.7, 136.7, 140.9, 192.7 ppm.

IR (film): ν 1703 cm^{-1} .

MS $[\text{ESI}]^+$: calculated for $\text{C}_{18}\text{H}_{31}\text{O}_2\text{Si}^+$ 307.20; found 307.20.

3-(2-((Triisopropylsilyl)oxy)ethyl)phenyl)methanol (15)—To a solution of **14** (0.500 g, 1.63 mmol) in tetrahydrofuran (4 mL) and water (0.25 mL), sodium borohydride (0.062 g, 1.63 mmol) was added. The reaction mixture was heated to reflux for 2 h. After cooling, water was added and the resultant mixture was extracted with ethyl acetate. The combined extracts were washed with brine, dried over magnesium sulfate, filtered and concentrated *in vacuo*. The residue was dried under high vacuum to afford **15** as colorless oil (99% yield).

^1H NMR (CDCl_3): δ 0.91–1.17 (m, 21H), 1.62 (broad t, J = 5.9 Hz, 1H), 2.88 (t, J = 7.1 Hz, 2H), 3.89 (d, J = 7.1 Hz, 2H), 4.68 (d, J = 5.7 Hz, 2H), 7.17 (d, J = 7.4 Hz, 1H), 7.21 (d, J = 7.5 Hz, 1H), 7.24 (s, 1H), 7.28 (t, J = 7.5 Hz, 1H) ppm.

^{13}C NMR (CDCl_3): δ 12.2, 18.2, 39.9, 64.9, 65.7, 125.0, 128.1, 128.7, 128.8, 139.9, 141.0 ppm.

IR (film): ν 3325 cm^{-1} .

3-(Iodomethyl)phenethoxy)triisopropylsilane (16)—A mixture of **15** (0.500 g, 1.63 mmol), triphenylphosphine (1.280 g, 4.89 mmol), imidazole (0.360 g, 5.22 mmol) and iodine (1.320 g, 5.22 mmol) in diethyl ether (20 mL) and acetonitrile (6.0 mL) was stirred at 0 °C for 2 h. The reaction mixture was diluted with ether and washed with water and a 20 wt % solution of thiosulfate in water. The organic layer was dried over magnesium sulfate, filtered and concentrated *in vacuo*. Purification by silica gel column chromatography using ethyl acetate-hexanes 1:9 as eluant provided **16** as colorless oil (64% yield).

^1H NMR (CDCl_3): δ 1.00–1.11 (m, 21H), 2.83 (t, J = 6.9 Hz, 2H), 3.89 (t, J = 6.9 Hz, 2H), 4.4 (s, 2H), 7.11 (d, J = 6.9 Hz, 1H), 7.20–7.27 (m, 3H) ppm.

^{13}C NMR (CDCl_3): δ 6.1, 12.2, 18.2, 39.7, 64.7, 126.8, 128.9, 129.0, 129.8, 139.2, 140.3 ppm.

MS [ESI]⁺: calculated for C₁₈H₃₂IOSi⁺ 419.12; found 419.10.

3-Isobutoxy-5-(3-(2-((triisopropylsilyl)oxy)ethyl)benzyl)cyclopent-2-enone (20)

—A solution of diisopropylamine (0.040 g, 0.05 mL, 0.38 mmol) in anhydrous tetrahydrofuran (0.3 mL) was cooled to −78 °C and *n*-Butyl lithium (2.3 M solution in hexanes, 0.15 mL, 0.35 mmol) was added dropwise. After stirring at −78 °C for 1 h, a solution of **3**⁹ (0.050 g, 0.32 mmol) in anhydrous tetrahydrofuran (2 mL) was added dropwise and the reaction mixture is stirred at −78 °C for 45 min. A solution of **16** (0.150 g, 0.35 mmol) in anhydrous tetrahydrofuran (1.5 mL) was then added dropwise and the reaction mixture was allowed to warm to rt. The reaction was quenched with saturated solution of ammonium chloride and the aqueous phase was extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over magnesium sulfate, filtered and concentrated *in vacuo*. Purification by silica gel column chromatography using ethyl acetate-hexanes 1:3 as eluant furnished **20** as colorless oil (49% yield).

¹H NMR (CDCl₃): δ 0.95 (d, *J* = 6.5 Hz, 6H), 1.00–1.10 (m, 21H), 1.20–2.10 (m, 1H), 2.32–2.37 (m, 1H), 2.49–2.61 (m, 2H), 2.76–2.80 (m, 1H), 2.82 (t, *J* = 7.1 Hz, 2H), 3.26 (dd, *J* = 13.9, 4.0 Hz, 1H), 3.71 (d, *J* = 7.1 Hz, 2H), 3.87 (t, *J* = 7.1 Hz, 2H), 5.25 (s, 1H), 7.03–7.07 (m, 3H), 7.19 (t, *J* = 8.1 Hz, 1H) ppm.

¹³C NMR (CDCl₃): δ 12.1, 18.1, 19.1, 28.0, 34.5, 37.3, 39.9, 46.9, 65.0, 78.1, 103.8, 126.7, 127.3, 128.5, 129.9, 139.6, 139.7, 189.4, 207.3 ppm.

IR (film): ν 1696, 1595 cm^{−1}.

MS [ESI]⁺: calculated for C₂₇H₄₄O₃NaSi⁺ 467.30; found 467.30.

3-Isobutoxy-2-methyl-5-(3-(2-((triisopropylsilyl)oxy)ethyl)benzyl)cyclopent-2-enone (21)—Prepared as **20** from **16** and **4**. Yield: 56%.

¹H NMR (CDCl₃): δ 0.95–1.08 (m, 27H), 1.65 (s, 3H), 1.93–1.99 (m, 1H), 2.25–2.30 (m, 1H), 2.45 (dd, *J* = 14.0, 10.8 Hz, 1H), 2.58–2.63 (m, 1H), 2.76–2.78 (m, 1H), 2.83 (t, *J* = 7.1 Hz, 2H), 3.30 (dd, *J* = 14.1, 3.9 Hz, 1H), 3.82 (d, *J* = 6.6 Hz, 2H), 3.87 (t, *J* = 7.3 Hz, 2H), 7.03–7.08 (m, 3H), 7.20 (t, *J* = 7.5 Hz, 1H) ppm.

¹³C NMR (CDCl₃): δ 6.2, 12.1, 18.1, 18.9, 28.8, 31.3, 37.5, 39.8, 46.2, 64.9, 75.6, 115.3, 126.6, 127.3, 128.4, 129.8, 139.6, 139.8, 183.3, 206.6 ppm.

IR (film): ν 1736, 1652, 1634 cm^{−1}.

MS [ESI]⁺: calculated for C₂₈H₄₇O₃Si⁺ 459.32; found 459.28.

2-Ethyl-3-isobutoxy-5-(3-(2-((triisopropylsilyl)oxy)ethyl)benzyl)cyclopent-2-enone (22)—Prepared as **20** from **16** and **17**. Yield: 48%.

¹H NMR (CDCl₃): δ 0.96 (dd, *J* = 6.7, 1.6 Hz, 6H), 0.99–1.07 (m, 24H), 1.97 (dt, *J* = 13.3, 6.7 Hz, 1H), 2.16 (q, *J* = 7.6 Hz, 2H), 2.27 (dt, *J* = 17.5, 1.0 Hz, 1H), 2.48 (dd, *J* = 14.0, 10.7 Hz, 1H), 2.60 (dd, *J* = 17.4, 6.8 Hz, 1H), 2.74–2.77 (m, 1H), 2.83 (t, *J* = 7.2 Hz, 2H), 3.29 (dd, *J* = 14.1, 4.0 Hz, 1H), 3.80 (d, *J* = 6.5 Hz, 2H), 3.86–3.89 (m, 2H), 7.06 (dt, *J* = 12.4, 6.7 Hz, 3H), 7.20 (t, *J* = 7.5 Hz, 1H) ppm.

¹³C NMR (CDCl₃): δ 12.2, 12.8, 14.8, 18.2, 19.0, 28.9, 31.0, 37.5, 39.9, 46.2, 65.0, 75.5, 121.4, 126.8, 127.3, 128.5, 129.9, 139.65, 139.80, 183.3, 206.2 ppm.

3-Isobutoxy-2-isopropyl-5-(3-(2-((triisopropylsilyl)oxy)ethyl)benzyl)cyclopent-2-enone (23)—Prepared as **20** from **16** and **18**. Yield: 68%.

^1H NMR (CDCl_3): δ 0.96 (dd, $J = 6.7, 0.9$ Hz, 6H), 1.02–1.08 (m, 21H), 1.09–1.14 (m, 6H), 1.96 (dd, $J = 13.3, 6.6$ Hz, 1H), 2.25 (dd, $J = 17.4, 2.3$ Hz, 1H), 2.49 (dd, $J = 14.0, 10.5$ Hz, 1H), 2.58 (dd, $J = 17.4, 6.9$ Hz, 1H), 2.72 (td, $J = 7.3, 3.3$ Hz, 1H), 2.77 (t, $J = 7.1$ Hz, 1H), 2.83 (t, $J = 7.1$ Hz, 2H), 3.25 (dd, $J = 14.0, 4.0$ Hz, 1H), 3.78 (d, $J = 6.0$ Hz, 2H), 3.87 (t, $J = 7.2$ Hz, 2H), 7.05 (dt, $J = 13.4, 7.0$ Hz, 3H), 7.19 (t, $J = 7.5$ Hz, 1H) ppm.

^{13}C NMR (CDCl_3): δ 12.2, 18.2, 19.1, 20.5, 23.0, 28.9, 30.7, 37.5, 39.9, 46.0, 65.0, 75.5, 124.8, 126.8, 127.3, 128.4, 129.9, 139.59, 139.72, 183.0, 205.7 ppm.

2-(Cyclopropylmethyl)-3-isobutoxy-5-(3-(2-((triisopropylsilyl)oxy)ethyl)benzyl)cyclopent-2-enone (24)—Prepared as **20** from **16** and **19**. Yield: 44%.

^1H NMR (CDCl_3): δ 0.10–0.13 (m, 2H), 0.33–0.36 (m, 2H), 0.87 (m, 1H), 0.95 (dd, $J = 6.7, 0.6$ Hz, 6H), 1.02–1.07 (m, 21H), 1.97 (m, 1H), 2.06 (d, $J = 6.8$ Hz, 2H), 2.27–2.31 (m, 1H), 2.49 (dd, $J = 14.1, 10.6$ Hz, 1H), 2.63 (dd, $J = 17.4, 6.9$ Hz, 1H), 2.78 (m, 1H), 2.82 (t, $J = 7.1$ Hz, 2H), 3.28 (dd, $J = 14.1, 4.0$ Hz, 1H), 3.80 (d, $J = 6.5$ Hz, 2H), 3.86 (t, $J = 7.2$ Hz, 2H), 7.05 (m, 3H), 7.18 (t, $J = 7.9$ Hz, 1H) ppm.

^{13}C NMR (CDCl_3): δ 4.6, 10.0, 12.1, 18.1, 19.0, 26.0, 28.8, 31.0, 37.4, 39.9, 46.2, 64.9, 75.5, 119.6, 126.7, 127.3, 128.4, 129.8, 139.6, 139.7, 183.6, 206.1 ppm.

5-(3-(2-Hydroxyethyl)benzyl)-3-isobutoxycyclopent-2-enone (25)—A solution of tetrabutylammonium fluoride (1M in tetrahydrofuran, 0.47 mL, 0.47 mmol) was added dropwise to a solution of **20** (0.071 g, 0.16 mmol) in anhydrous tetrahydrofuran (1 mL) at 0 °C. The reaction mixture was stirred at 0 °C for 3 h and then quenched with saturated solution of ammonium chloride. The aqueous layer was extracted with diethyl ether and the combined organic layers were washed with brine, dried over magnesium sulfate, filtered and concentrated *in vacuo*. Purification by silica gel column chromatography using dichloromethane-methanol 95:5 as eluant provided **25** as colorless oil (82% yield).

^1H NMR (CDCl_3): δ 0.96 (d, $J = 6.7$ Hz, 6H), 1.98–2.16 (m, 1H), 2.34 (dd, $J = 17.7, 2.3$ Hz, 1H), 2.54–2.63 (m, 2H), 2.75–2.80 (m, 1H), 2.84 (t, $J = 6.7$ Hz, 2H), 3.22 (dd, $J = 13.9, 4.1$ Hz, 2H), 3.70 (d, $J = 5.8$ Hz, 2H), 3.83 (t, $J = 6.6$ Hz, 2H), 5.24 (s, 1H), 7.05–7.08 (m, 3H), 7.21 (t, $J = 8.1$ Hz, 1H) ppm.

^{13}C NMR (CDCl_3): δ 19.1, 28.0, 34.4, 37.2, 39.3, 46.7, 63.7, 78.2, 103.9, 127.1, 127.2, 128.8, 129.8, 139.0, 139.9, 189.5, 207.5 ppm.

IR (film): ν 3399, 1685, 1590 cm^{-1} .

MS $[\text{ESI}]^+$: calculated for $\text{C}_{18}\text{H}_{25}\text{O}_3^+$ 289.18; found 289.21.

5-(3-(2-Hydroxyethyl)benzyl)-3-isobutoxy-2-methylcyclopent-2-enone (26)—Prepared as **25** from **21**. Yield: 78%.

^1H NMR (CDCl_3): δ 0.95–0.96 (m, 6H), 1.64 (s, 3H), 1.92–2.00 (m, 2H), 2.28 (d, $J = 17.4$ Hz, 1H), 2.52 (dd, $J = 14$ and 10.4 Hz, 1H), 2.60–2.65 (m, 1H), 2.75–2.80 (m, 1H), 2.85 (t, $J = 7.5$ Hz, 2H), 3.26 (dd, $J = 14.0$ and 4.0 Hz, 1H), 3.80–3.86 (m, 4H), 7.05–7.09 (m, 3H), 7.23 (t, $J = 7.8$ Hz, 1H) ppm.

^{13}C NMR (CDCl_3): δ 6.2, 19.0, 28.9, 31.3, 37.5, 39.3, 46.1, 63.8, 75.7, 115.4, 127.1, 127.2, 128.8, 129.8, 139.0, 140.1, 183.3, 206.7 ppm.

IR (film): ν 3404, 1680, 1620 cm^{-1} .

MS $[\text{ESI}]^+$: calculated for $\text{C}_{19}\text{H}_{27}\text{O}_3^+$ 303.19; found 303.20.

2-Ethyl-5-(3-(2-hydroxyethyl)benzyl)-3-isobutoxycyclopent-2-enone (27)—

Prepared as **25** from **22**. Yield: 86%.

^1H NMR (Acetone- d_6): δ 0.93–0.96 (m, 9H), 1.94 (dt, J = 13.3, 6.7 Hz, 1H), 2.08 (q, J = 7.6 Hz, 2H), 2.42 (dt, J = 17.3, 1.1 Hz, 1H), 2.54 (dd, J = 13.7, 9.7 Hz, 1H), 2.67 (ddt, J = 9.6, 4.4, 2.3 Hz, 1H), 2.72–2.79 (m, 3H), 3.10 (dd, J = 13.7, 4.0 Hz, 1H), 3.70 (bs, 1H), 3.73 (t, J = 7.1 Hz, 2H), 3.92 (dd, J = 6.5, 3.1 Hz, 2H), 7.07 (t, J = 6.4 Hz, 2H), 7.12 (s, 1H), 7.17 (t, J = 7.5 Hz, 1H) ppm.

^{13}C NMR (Acetone- d_6): δ 13.5, 15.7, 19.5, 29.9, 31.6, 38.1, 40.8, 47.2, 64.4, 76.3, 121.4, 127.9, 128.1, 129.5, 131.0, 140.96, 140.98, 184.4, 205.8 ppm.

5-(3-(2-Hydroxyethyl)benzyl)-3-isobutoxy-2-isopropylcyclopent-2-enone (28)—

Prepared as **25** from **23**. Yield: 90%.

^1H NMR (Acetone- d_6): δ 1.07 (d, J = 6.7 Hz, 6H), 1.22 (t, J = 7.6 Hz, 6H), 2.02–2.09 (m, 1H), 2.43–2.48 (m, 1H), 2.69–2.84 (m, 4H), 2.89 (t, J = 7.0 Hz, 2H), 3.00 (s, 1H), 3.18 (dd, J = 13.1, 3.1 Hz, 1H), 3.83 (t, J = 7.0 Hz, 2H), 3.97 (dd, J = 6.4, 1.1 Hz, 2H), 7.17–7.21 (m, 3H), 7.33 (t, J = 7.5 Hz, 1H) ppm.

^{13}C NMR (Acetone- d_6): δ 19.7, 21.3, 21.3, 24.4, 30.1, 31.6, 38.3, 40.7, 47.1, 64.5, 76.8, 118.9, 125.2, 128.4, 128.5, 129.8, 131.5, 141.1, 185.2, 206.7 ppm.

2-(Cyclopropylmethyl)-5-(3-(2-hydroxyethyl)benzyl)-3-isobutoxycyclopent-2-enone (29)—Prepared as **25** from **24**. Yield: 99%.

^1H NMR (CDCl_3): δ 0.09–0.12 (m, 2H), 0.33–0.37 (m, 2H), 0.87–0.91 (m, 1H), 0.96 (d, J = 6.7 Hz, 6H), 1.80 (bs, 1H), 1.97 (dt, J = 13.3, 6.7 Hz, 1H), 2.06 (d, J = 6.8 Hz, 2H), 2.28–2.32 (m, 1H), 2.55 (dd, J = 14.1, 10.3 Hz, 1H), 2.66 (dd, J = 17.4, 6.8 Hz, 1H), 2.77–2.81 (m, 1H), 2.85 (t, J = 6.6 Hz, 2H), 3.26 (dd, J = 14.1, 4.2 Hz, 1H), 3.81 (d, J = 6.4 Hz, 2H), 3.85 (t, J = 6.6 Hz, 2H), 7.07–7.09 (m, 3H), 7.23 (t, J = 7.7 Hz, 1H) ppm.

^{13}C NMR (CDCl_3): δ 4.6, 10.0, 19.0, 26.0, 28.9, 31.1, 37.4, 39.4, 46.1, 63.8, 75.6, 119.7, 127.2, 128.8, 129.8, 139.0, 140.1, 183.9, 206.3 ppm.

HRMS (ESI^+): calculated for $\text{C}_{22}\text{H}_{31}\text{O}_3^+$ 343.2267; found 343.2268.

5-(3-(2-Iodoethyl)benzyl)-3-isobutoxycyclopent-2-enone (30)—Prepared as **16** from **25**. Yield: 88%.

^1H NMR (CDCl_3): δ 0.96 (d, J = 6.7 Hz, 6H), 1.99–2.07 (m, 1H), 2.35 (dd, J = 17.8, 2.3 Hz, 1H), 2.59–2.63 (m, 2H), 2.77–2.81 (m, 1H), 3.14 (t, J = 7.7 Hz, 2H), 3.23 (dd, J = 13.9, 4.1 Hz, 1H), 3.35 (t, J = 7.3 Hz, 2H), 3.70 (d, J = 6.5 Hz, 2H), 5.25 (s, 1H), 7.02–7.04 (m, 2H), 7.10 (d, J = 7.6 Hz, 1H), 7.22 (t, J = 7.7 Hz, 1H) ppm.

^{13}C NMR (CDCl_3): δ 6.0, 19.1, 28.0, 34.3, 37.1, 40.3, 46.7, 78.1, 104.0, 126.5, 127.6, 128.9, 129.2, 140.0, 140.9, 189.4, 207.2 ppm.

IR: ν 1691, 1592 cm^{-1} .

MS [ESI]⁺: calculated for $\text{C}_{18}\text{H}_{24}\text{IO}_2^+$ 398.08 found 398.07.

5-(3-(2-Azidoethyl)benzyl)-3-isobutoxycyclopent-2-enone (31)—A mixture of **30** (0.044 g, 0.11 mmol) and sodium azide (0.007 g, 0.11 mmol) in dry *N,N*-dimethylformamide (0.5 mL) was heated at 50 °C for 45 min. The reaction mixture was then diluted with water and the resulting mixture was extracted with ethyl acetate. The organic extracts were washed with brine, dried over magnesium sulfate, filtered and concentrated *in vacuo*. The residue so obtained was dried under high vacuum to furnish **31** (76% yield).

¹H NMR (CDCl_3): δ 0.97 (d, J = 6.7 Hz, 6H), 2.00–2.08 (m, 1H), 2.34 (dd, J = 17.8, 2.2 Hz, 1H), 2.57–2.63 (m, 2H), 2.77–2.81 (m, 1H), 2.87 (t, J = 7.2 Hz, 2H), 3.24 (dd, J = 13.9, 4.1 Hz, 1H), 3.49 (t, J = 7.2 Hz, 2H), 3.70 (d, J = 6.5 Hz, 2H), 5.25 (s, 1H), 7.06–7.09 (m, 3H), 7.23 (t, J = 7.9 Hz, 1H) ppm.

¹³C NMR (CDCl_3): δ 19.1, 27.9, 34.3, 35.4, 37.1, 46.7, 52.6, 78.1, 103.9, 126.9, 127.5, 128.9, 129.5, 138.4, 140.1, 189.4, 207.2 ppm.

IR (film): ν 2097, 1692, 1592 cm^{-1} .

MS [ESI]⁺: calculated for $\text{C}_{18}\text{H}_{24}\text{N}_3\text{O}_2^+$ 314.18; found 314.07.

4-Chloro-*N*-(3-((4-isobutoxy-2-oxocyclopent-3-en-1-yl)methyl)phenethyl)benzenesulfonamide (32)—A mixture of **31** (0.059 g, 0.19 mmol) and Pd on C (10 wt%, 0.006 g) in methanol (3 mL) was stirred at rt for 1 h under 1 atmosphere of hydrogen. The reaction mixture was filtered to remove the catalyst. Water (0.5 mL) and 2N sodium hydroxide (0.27 mL) were added and the resultant mixture was cooled to 0 °C. 4-Chlorophenylsulfonyl chloride (0.082 g, 0.39 mmol) was added; the reaction mixture was stirred at 0 °C for 4 h and then diluted with ethyl acetate. The organic layers were collected, washed with brine, dried over magnesium sulfate, filtered and concentrated *in vacuo*. Purification by silica gel column chromatography, using ethyl acetate-hexanes 1:1 as eluant provided **32** (22% yield).

¹H NMR (CDCl_3): δ 0.98 (d, J = 6.5 Hz, 6H), 1.99–2.10 (m, 1H), 2.32 (dd, J = 17.5, 2.7 Hz, 1H), 2.61–2.65 (m, 2H), 2.75 (t, J = 6.6 Hz, 2H), 3.15 (dd, J = 13.9, 4.5 Hz, 1H), 3.22–3.26 (m, 2H), 3.71–3.73 (m, 3H), 4.47 (broad t, J = 6.2 Hz, 1H), 5.26 (s, 1H), 6.92–6.94 (m, 2H), 7.08 (d, J = 7.6 Hz, 1H), 7.21 (t, J = 7.8 Hz, 1H), 7.47 (d, J = 8.6 Hz, 2H), 7.75 (d, J = 8.6 Hz, 2H) ppm.

MS [ESI]⁺: calculated for $\text{C}_{24}\text{H}_{29}\text{ClNO}_4\text{S}^+$ 462.15; found 462.21.

***tert*-Butyl (4-chlorophenyl)sulfonyl(3-((2-isobutoxy-3-methyl-4-oxocyclopent-2-en-1-yl)methyl)-phenethyl)carbamate (33)**—Diethyl diazodicarboxylate (40 wt% in toluene, 0.08 mL, 0.28 mmol) was added dropwise to a solution of **26** (0.021 g, 0.07 mmol), *N*-*boc*-4-chlorobenzenesulfonamide (0.032 g, 0.11 mmol), and triphenylphosphine (0.058 g, 0.22 mmol) in anhydrous tetrahydrofuran (3 mL) at 0 °C under Ar atmosphere and was stirred at rt for 4 h. The reaction was quenched with water and the resulting mixture was extracted with diethyl ether. The combined organic layers were washed with brine, dried over magnesium sulfate, filtered and concentrated *in vacuo*. Purification by silica gel column chromatography using ethyl acetate-hexanes 1:3 as eluant furnished **33** (96% yield).

^1H NMR (CDCl_3): δ 0.94 (d, J = 6.8 Hz, 6H), 1.34 (s, 9H), 1.65 (s, 3H), 1.91–1.99 (m, 1H), 2.29 (d, J = 17.5 Hz, 1H), 2.45–2.49 (m, 1H), 2.60 (dd, J = 17.5, 6.7 Hz, 1H), 2.26–2.28 (m, 1H), 3.03 (t, J = 7.4 Hz, 2H), 3.31 (dd, J = 14.0, 3.4 Hz, 1H), 3.79–3.84 (m, 2H), 4.04 (t, J = 5.85 Hz, 2H), 7.09–7.12 (m, 3H), 7.24 (t, J = 7.5 Hz, 1H), 7.43 (d, J = 8.3 Hz, 2H), 7.69 (d, J = 8.3 Hz, 2H) ppm.

^{13}C NMR (CDCl_3): δ 6.2, 19.0, 28.1, 28.9, 31.4, 36.6, 37.5, 46.2, 48.6, 75.7, 84.8, 115.3, 127.4, 129.0, 129.1, 129.6, 129.9, 138.4, 138.9, 139.9, 140.5, 150.8, 183.5, 206.5 ppm.

IR (film): ν 1729, 1688, 1631 cm^{-1} .

MS $[\text{ESI}]^+$: calculated for $\text{C}_{30}\text{H}_{39}\text{ClNO}_6\text{S}^+$ 576.22; found 576.06.

***tert*-Butyl (4-chlorophenyl)sulfonyl(3-((3-ethyl-4-isobutoxy-2-oxocyclopent-3-en-1-yl)methyl)-phenethyl)carbamate (34)**—Prepared as **33** from **27**.

Yield: 100%.

^1H NMR (Acetone- d_6): δ 1.00 (dd, J = 6.7, 1.3 Hz, 6H), 1.04 (t, J = 7.5 Hz, 3H), 1.39 (s, 9H), 1.99 (dt, J = 13.4, 6.7 Hz, 1H), 2.16 (q, J = 7.5 Hz, 2H), 2.43–2.39 (m, 1H), 2.63 (dd, J = 13.8, 9.7 Hz, 1H), 2.72 (dd, J = 17.3, 6.8 Hz, 1H), 2.77 (m, 1H), 3.07 (t, J = 7.3 Hz, 2H), 3.18 (dd, J = 13.8, 4.0 Hz, 1H), 3.93 (d, J = 6.6 Hz, 2H), 4.12 (t, J = 7.3 Hz, 2H), 7.19 (m, 3H), 7.32 (t, J = 7.5 Hz, 1H), 7.63 (d, J = 8.6 Hz, 2H), 7.82 (d, J = 8.6 Hz, 2H) ppm.

***tert*-Butyl (4-chlorophenyl)sulfonyl(3-((4-isobutoxy-3-isopropyl-2-oxocyclopent-3-en-1-yl)methyl)-phenethyl)carbamate (35)**—Prepared as **33** from **28**.

Yield: 80%.

^1H NMR (CD_3OD): δ 0.98 (d, J = 6.7 Hz, 6H), 1.15 (dd, J = 10.3, 7.1 Hz, 6H), 1.30–1.42 (m, 9H), 1.96 (dt, J = 13.3, 6.6 Hz, 1H), 2.42 (d, J = 15.9 Hz, 1H), 2.63 (dd, J = 13.7, 9.1 Hz, 1H), 2.71–2.78 (m, 3H), 3.03 (t, J = 7.1 Hz, 2H), 3.16 (dd, J = 13.7, 3.8 Hz, 1H), 3.92 (d, J = 6.3 Hz, 2H), 4.12 (t, J = 7.1 Hz, 2H), 7.14 (m, 3H), 7.26 (d, J = 7.4 Hz, 1H), 7.59 (d, J = 8.7 Hz, 2H), 7.78 (d, J = 8.7 Hz, 2H) ppm.

^{13}C NMR (CD_3OD): δ 19.3, 20.85, 20.88, 24.1, 28.3, 30.0, 31.7, 37.4, 38.0, 47.0, 77.1, 85.7, 125.3, 128.63, 128.67, 129.9, 130.3, 131.0, 131.3, 139.8, 140.4, 140.85, 140.94, 152.2, 187.4, 208.8 ppm.

***tert*-Butyl (4-chlorophenyl)sulfonyl(3-((3-(cyclopropylmethyl)-4-isobutoxy-2-oxocyclopent-3-en-1-yl)methyl)phenethyl)carbamate (36)**—Prepared as **33** from **29**.

Yield: 78%.

^1H NMR (CDCl_3): δ 0.09–0.12 (m, 2H), 0.33–0.36 (m, 2H), 0.87–0.90 (m, 1H), 0.93 (dd, J = 6.7, 1.8 Hz, 6H), 1.33 (s, 9H), 1.95 (dt, J = 13.3, 6.7 Hz, 1H), 2.06 (d, J = 6.8 Hz, 2H), 2.31 (dd, J = 17.6, 1.9 Hz, 1H), 2.51 (dd, J = 14.1, 10.7 Hz, 1H), 2.63 (dd, J = 17.6, 6.8 Hz, 1H), 2.78–2.82 (m, 1H), 3.02 (t, J = 7.5 Hz, 2H), 3.30 (dd, J = 14.1, 4.0 Hz, 1H), 3.80 (dd, J = 6.5, 3.3 Hz, 2H), 4.03 (t, J = 7.5 Hz, 2H), 7.10 (q, J = 8.8 Hz, 3H), 7.23 (t, J = 7.8 Hz, 1H), 7.43 (d, J = 8.7 Hz, 2H), 7.69 (d, J = 8.7 Hz, 2H) ppm.

4-Chloro-*N*-(3-((2-hydroxy-4-oxocyclopent-2-en-1-yl)methyl)phenethyl)benzenesulfonamide (41)

—A mixture of **32** (0.018 g, 0.04 mmol) in acetone (0.5 mL) and 2N hydrochloric acid (0.2 mL) was stirred at rt for 6 h. The reaction mixture was diluted with water and the acetone was evaporated under reduced pressure. The aqueous phase was extracted with ethyl acetate and the combined organic layers were washed with brine, dried over magnesium sulfate, filtered and concentrated *in vacuo*. Purification by reverse phase preparative HPLC provided **41** as white solid. Yield: 88%.

¹H NMR (DMSO-*d*₆): δ 2.05–2.09 (m, 1H), 2.37–2.39 (m, 1H), 2.46 (dd, *J* = 13.7, 10.2 Hz, 1H), 2.64 (t, *J* = 7.3 Hz, 2H), 2.74–2.78 (m, 1H), 2.95–3.01 (m, 3H), 5.06 (s, 1H), 6.97 (d, *J* = 7.6 Hz, 1H), 6.99 (s, 1H), 7.04 (d, *J* = 7.6 Hz, 1H), 7.16 (t, *J* = 7.5 Hz, 1H), 7.64 (d, *J* = 8.6 Hz, 2H), 7.76 (d, *J* = 8.6 Hz, 2H), 7.82 (t, *J* = 5.7 Hz, 1H) ppm.

¹³C NMR (DMSO-*d*₆): δ 22.1, 29.0, 35.2, 36.7, 44.0, 104.5, 126.5, 126.8, 128.3, 128.4, 129.1, 129.6, 137.2, 138.5, 139.4, 139.6 ppm.

IR (film): ν 3402, 1730, 1647, 1580 cm^{−1}.

HRMS [ESI]⁺: calculated for C₂₀H₂₀ClNNaO₄S⁺ 428.0699; found 428.0714.

4-Chloro-*N*-(3-((4-isobutoxy-3-methyl-2-oxocyclopent-3-en-1-yl)methyl)phenethyl)-benzenesulfonamide (37)

—To a solution of **33** (0.100 g, 0.173 mmol) in dichloromethane (5 mL) 2,2,2-trifluoroacetic acid (1.5 mL) was added. The resultant mixture was stirred at rt for 1 h. The reaction mixture was concentrated *in vacuo* and purified by preparative reverse phase HPLC to obtain **37** (73% yield).

¹H NMR (CDCl₃): δ 0.97 (dd, *J* = 6.7, 1.3 Hz, 6H), 1.65 (t, *J* = 1.6 Hz, 3H), 1.98 (dt, *J* = 13.4, 6.7 Hz, 1H), 2.28 (dt, *J* = 17.4, 1.9 Hz, 1H), 2.53 (dd, *J* = 14.0, 10.2 Hz, 1H), 2.65 (ddd, *J* = 17.4, 6.8, 1.6 Hz, 1H), 2.75–2.79 (m, 3H), 3.21–3.26 (m, 3H), 3.85 (d, *J* = 6.6 Hz, 2H), 4.45 (t, *J* = 6.2 Hz, 1H), 6.95 (d, *J* = 8.7 Hz, 2H), 7.09 (d, *J* = 7.6 Hz, 1H), 7.22 (t, *J* = 7.4 Hz, 1H), 7.48 (d, *J* = 8.7 Hz, 2H), 7.75 (d, *J* = 8.7 Hz, 2H) ppm.

¹³C NMR (CDCl₃): δ 5.3, 18.3, 28.4, 30.9, 35.7, 36.9, 44.1, 45.7, 75.7, 114.9, 126.65, 126.85, 128.04, 128.13, 128.5, 129.09, 138.3, 138.59, 138.72, 139.2, 185.5, 208.1 ppm.

IR (film): ν 2926, 1679, 1618 cm^{−1}.

HRMS [ESI]⁺: calculated for C₂₅H₃₁ClNO₄S⁺ 476.1662; found 476.1656.

4-Chloro-*N*-(3-((3-ethyl-4-isobutoxy-2-oxocyclopent-3-en-1-yl)methyl)phenethyl)-benzenesulfonamide (38)

—A solution of **34** (0.122 g, 0.207 mmol) in dichloromethane (5 mL) and 2,2,2-trifluoroacetic acid (3 mL) was stirred at rt for 5 h. The reaction mixture was concentrated *in vacuo* and purified by reverse phase preparative HPLC to provide **38** as a colorless oil (80% yield).

¹H NMR (CDCl₃): δ 0.95 (d, *J* = 6.7 Hz, 6H), 0.98 (t, *J* = 7.5 Hz, 3H), 1.96 (dt, *J* = 13.3, 6.7 Hz, 1H), 2.14 (q, *J* = 7.5 Hz, 2H), 2.26 (dd, *J* = 17.4, 0.9 Hz, 1H), 2.53 (dd, *J* = 14.0, 10.0 Hz, 1H), 2.64 (dd, *J* = 17.4, 6.8 Hz, 1H), 2.71–2.78 (m, 3H), 3.17–3.24 (m, 3H), 3.82 (d, *J* = 6.5 Hz, 2H), 4.87 (t, *J* = 6.1 Hz, 1H), 6.94–6.96 (m, 2H), 7.06 (d, *J* = 7.5 Hz, 1H), 7.18 (d, *J* = 7.5 Hz, 1H), 7.46 (d, *J* = 8.6 Hz, 2H), 7.74 (d, *J* = 8.6 Hz, 2H) ppm.

¹³C NMR (CDCl₃): δ 12.8, 14.8, 19.0, 28.9, 31.1, 36.0, 37.4, 44.4, 46.0, 75.7, 121.4, 126.9, 127.6, 128.7, 129.0, 129.56, 129.62, 138.0, 138.8, 139.2, 140.4, 183.5, 206.1 ppm.

IR (film): ν 2960, 2920, 2852, 1613 cm^{-1} .

HRMS [ESI]⁺: calculated for $\text{C}_{26}\text{H}_{32}\text{NO}_4\text{NaCl}^+$ 512.1638; found 512.1627.

4-Chloro-*N*-(3-((4-isobutoxy-3-isopropyl-2-oxocyclopent-3-en-1-yl)methyl)phenethyl)-benzenesulfonamide (39)—Prepared as **38** from **35**. Yield: 99%.

¹H NMR (CDCl_3): δ 0.87 (d, J = 6.7 Hz, 6H), 1.02 (dd, J = 15.4, 7.0 Hz, 6H), 1.85 (m, 1H), 2.29–2.33 (m, 1H), 2.55–2.69 (m, 6H), 2.98 (dd, J = 13.4, 3.2 Hz, 1H), 3.05 (t, J = 7.2 Hz, 2H), 3.82 (d, J = 6.3 Hz, 2H), 6.90–6.93 (m, 2H), 6.97 (d, J = 7.6 Hz, 1H), 7.08 (d, J = 7.5 Hz, 1H), 7.46 (d, J = 8.5 Hz, 2H), 7.71 (d, J = 8.5 Hz, 2H) ppm.

¹³C NMR (CDCl_3): δ 19.3, 20.78, 20.81, 24.1, 29.9, 31.4, 37.0, 37.9, 45.6, 46.9, 77.0, 125.4, 128.0, 128.4, 129.64, 129.71, 130.5, 130.8, 139.7, 140.0, 140.4, 140.9, 187.5, 208.9 ppm.

IR (film): ν 3268, 2963, 2930, 2874, 1671, 1608, 1470, 1383, 1335 cm^{-1} .

HRMS (ESI[−]): calculated for $\text{C}_{27}\text{H}_{33}\text{ClNO}_4\text{S}^-$ 502.1819; found 502.1802.

4-Chloro-*N*-(3-((3-(cyclopropylmethyl)-4-isobutoxy-2-oxocyclopent-3-en-1-yl)methyl)phenethyl)-benzenesulfonamide (40)—Prepared as **37** from **36**. Yield: 29%.

¹H NMR (CD_3OD): δ 0.05–0.06 (m, 2H), 0.30–0.32 (m, 2H), 0.84 (m, 1H), 0.98 (d, J = 7.0 Hz, 6H), 1.95 (m, 1H), 2.03 (d, J = 6.8 Hz, 2H), 2.45 (m, 1H), 2.65–2.69 (m, 1H), 2.74 (t, J = 7.2 Hz, 2H), 2.80–2.83 (m, 2H), 3.09–3.15 (m, 3H), 3.95 (dd, J = 6.4, 0.7 Hz, 2H), 7.00–7.04 (m, 2H), 7.08 (d, J = 7.6 Hz, 1H), 7.19 (t, J = 7.6 Hz, 1H), 7.57 (d, J = 8.7 Hz, 2H), 7.78–7.80 (m, 2H) ppm.

¹³C NMR (CD_3OD): δ 5.0, 10.8, 19.2, 26.7, 30.0, 31.7, 37.0, 37.9, 45.6, 47.2, 77.2, 120.3, 128.1, 128.4, 129.73, 129.74, 130.5, 130.8, 139.8, 140.2, 140.5, 141.0, 188.4, 209.7 ppm.

HRMS (ESI[−]): calculated for $\text{C}_{28}\text{H}_{33}\text{ClNO}_4\text{S}^-$ 514.1819; found 514.1809.

4-Chloro-*N*-(3-(2-hydroxy-3-methyl-4-oxocyclopent-2-en-1-yl)methyl)phenethyl)-benzenesulfonamide (42)—Prepared as **43** from **37**. Yield 54%.

¹H NMR ($\text{DMSO}-d_6$): δ 1.43 (s, 3H), 1.98 (d, J = 17.4 Hz, 1H), 2.30 (dd, J = 17.3, 7.0 Hz, 1H), 2.37 (dd, J = 13.5, 10.2 Hz, 1H), 2.62–2.65 (m, 3H), 2.94–2.98 (m, 2H), 3.03 (dd, J = 13.5, 3.6 Hz, 1H), 6.94–6.97 (m, 2H), 7.02 (d, J = 7.6 Hz, 1H), 7.14 (t, J = 7.5 Hz, 1H), 7.64 (d, J = 8.6 Hz, 2H), 7.76 (d, J = 8.6 Hz, 2H), 7.84 (broad t, J = 5.6 Hz, 1H) ppm.

¹³C NMR ($\text{DMSO}-d_6$): δ 5.9, 22.0, 35.3, 37.0, 43.6, 44.0, 110.8, 126.4, 126.8, 128.2, 128.4, 129.2, 129.4, 137.2, 138.4, 139.3, 139.9, 212.2, 214.7 ppm.

IR (film): ν 3279, 2925, 1608, cm^{-1} .

HRMS [ESI]⁺: calculated for $\text{C}_{21}\text{H}_{22}\text{ClNNaO}_4\text{S}^+$ 442.0856; found 442.0847.

4-Chloro-*N*-(3-((3-ethyl-4-hydroxy-2-oxocyclopent-3-en-1-yl)methyl)phenethyl)-benzenesulfonamide (43)—To a mixture of **38** (0.05 g, 0.11 mmol) in acetone (5 mL) 2N hydrochloric acid (5 mL) was added. The reaction mixture was stirred at rt for 13 h and

then concentrated *in vacuo*, extracted with ethyl acetate (10 mL $\times$ 2), and washed with brine (5 mL $\times$ 2). The organic layer was dried with anhydrous sodium sulfate, filtered and concentrated *in vacuo*. Purification by reverse phase preparative HPLC provided the desired compound (75% yield).

^1H NMR (CD_3OD): δ 0.93–0.96 (m, 3H), 2.05–2.17 (m, 3H), 2.45 (dd, J = 17.7, 6.8 Hz, 1H), 2.59 (dd, J = 13.7, 9.2 Hz, 1H), 2.69–2.74 (m, 2H), 2.82 (m, 1H), 3.11 (m, 3H), 6.99 (t, J = 2.0 Hz, 2H), 7.07 (s, 1H), 7.16 (d, J = 7.8 Hz, 1H), 7.57–7.58 (m, 2H), 7.79–7.81 (m, 2H) ppm.

^{13}C NMR (CDCl_3): δ 12.8, 14.5, 29.4, 29.9, 32.0, 35.9, 37.8, 44.4, 53.9, 120.1, 127.0, 127.8, 128.7, 129.1, 129.7, 130.1, 137.9, 138.5, 139.5, 139.6 ppm.

IR (film): ν 3388, 2922, 2354, 1608 cm^{-1} .

HRMS (ESI^-): calculated for $\text{C}_{22}\text{H}_{23}\text{NO}_4\text{SCl}^-$ 432.1036; found 432.1036.

4-Chloro-*N*-(3-((4-hydroxy-3-isopropyl-2-oxocyclopent-3-en-1-yl)methyl)phenethyl)benzenesulfonamide (44)—Prepared as **43** from **39**.

Yield: 50%.

^1H NMR (CDCl_3): δ 1.10 (d, J = 6.9 Hz, 6H), 2.17 (bs, 1H), 2.45 (dd, J = 17.4, 5.3 Hz, 1H), 2.59–2.78 (m, 5H), 3.04–3.11 (m, 3H), 6.97 (d, J = 9.7 Hz, 2H), 7.03 (d, J = 5.9 Hz, 1H), 7.15 (s, 1H), 7.55 (d, J = 7.3 Hz, 2H), 7.78 (d, J = 7.7 Hz, 2H) ppm.

^{13}C NMR (CDCl_3): δ 20.70, 20.72, 24.1, 35.84, 37.2, 38.3, 45.3, 45.7, 123.6, 128.0, 128.5, 129.64, 129.78, 130.5, 130.8, 139.8, 140.1, 140.5, 141.0 ppm.

IR (film): ν 3268, 2962, 2930, 2873, 1672, 1609 cm^{-1} .

HRMS (ESI^+): calculated for $\text{C}_{23}\text{H}_{26}\text{ClNO}_4\text{SNa}^+$ 470.1169; found 470.1165.

4-Chloro-*N*-(3-((3-(cyclopropylmethyl)-4-hydroxy-2-oxocyclopent-3-en-1-yl)methyl)phenethyl)benzenesulfonamide (45)—Prepared as **43** from **40**. Yield 56%.

^1H NMR (CD_3OD): δ 0.04–0.05 (m, 2H), 0.27–0.30 (m, 2H), 0.81–0.85 (m, 1H), 2.00 (d, J = 6.7 Hz, 2H), 2.17 (dd, J = 17.8, 1.8 Hz, 1H), 2.46 (d, J = 6.8 Hz, 1H), 2.58 (dd, J = 13.8, 9.1 Hz, 1H), 2.68 (t, J = 7.4 Hz, 2H), 2.80–2.88 (m, 1H), 3.07 (m, 3H), 6.95–6.96 (m, 2H), 7.02 (d, J = 7.7 Hz, 1H), 7.14 (t, J = 7.8 Hz, 1H), 7.53 (d, J = 8.7 Hz, 2H), 7.76 (d, J = 8.7 Hz, 2H) ppm.

^{13}C NMR (CD_3OD): δ 4.85, 4.87, 10.9, 26.2, 30.9, 37.2, 38.3, 45.8, 118.4, 128.0, 128.5, 129.72, 129.81, 130.6, 130.8, 139.8, 140.1, 140.6, 141.0 ppm.

IR (film): ν 3279, 2921, 2852, 1729, 1584 cm^{-1} .

HRMS (ESI^+): calculated for $\text{C}_{24}\text{H}_{25}\text{ClNO}_4\text{S}^+$ 458.1193; found 458.1190.

3-Hydroxy-2-(3-(2-((triisopropylsilyl)oxy)ethyl)benzyl)cyclopent-2-enone (46)—To a solution of **14** (0.920 g, 3.0 mmol), cyclopentane-1,3-dione (0.098 g, 1.0 mmol), *L*-proline (0.005 g) in dichloromethane (5 mL) diethyl 2,6-dimethyl-1,4-dihydropyridine-3,5-dicarboxylate (0.253 g, 1.0 mmol) was added and the resulting mixture was allowed to stir at rt for 12 h. Purification by silica gel column chromatography using a gradient of ethyl acetate in hexanes as eluant provided **46** (76% yield).

^1H NMR (CD_3OD): δ 1.01–1.04 (m, 21H), 2.47 (s, 4H), 2.75 (t, J = 6.8 Hz, 2H), 3.40 (s, 2H), 3.85 (t, J = 6.8 Hz, 2H), 6.95–6.97 (m, 1H), 7.03 (d, J = 7.6 Hz, 1H), 7.08 (t, J = 7.4 Hz, 2H) ppm.

^{13}C NMR (CD_3OD): δ 13.3, 18.6, 27.8, 31.5, 40.9, 66.2, 118.4, 127.4, 127.8, 129.1, 130.4, 140.4, 141.5 ppm.

2-(3-(2-Hydroxyethyl)benzyl)-3-isobutoxycyclopent-2-enone (47)—A mixture containing **46** (0.294 g, 0.757 mmol), p-toluenesulfonic acid (0.005 g), *i*-butanol (1.0 mL) and benzene (4.0 mL) was heated to reflux for 12 h. The reaction mixture was then concentrated *in vacuo* and the residue was purified by silica gel column chromatography using 5% of methanol in dichloromethane as eluant, providing **47** (81% yield).

^1H NMR (CDCl_3): δ 0.96 (d, J = 6.7 Hz, 6H), 2.00 (m, 1H), 2.39 (dt, J = 5.0, 2.4 Hz, 2H), 2.61 (t, J = 4.6 Hz, 3H), 2.77 (t, J = 6.9 Hz, 2H), 3.41 (s, 2H), 3.76 (t, J = 6.9 Hz, 2H), 3.88 (d, J = 6.4 Hz, 2H), 6.97 (d, J = 7.3 Hz, 1H), 7.11 (m, 3H) ppm.

^{13}C NMR (CDCl_3): 19.0, 25.2, 27.4, 28.8, 33.6, 39.4, 63.7, 75.7, 119.9, 126.6, 126.7, 128.4, 129.4, 138.7, 140.6, 185.3, 204.5 ppm.

IR (film): ν 3399, 1685, 1590 cm^{-1} .

MS (ESI^+): calculated for $\text{C}_{18}\text{H}_{25}\text{O}_3^+$ 289.18; found 289.16.

tert-Butyl (4-chlorophenyl)sulfonyl(3-((2-isobutoxy-5-oxocyclopent-1-en-1-yl)methyl)phenethyl)carbamate (48)—Prepared as **33** from **47**. Yield: 92%.

^1H NMR (CDCl_3): δ 0.99 (d, J = 6.7 Hz, 6H), 1.37 (s, 9H), 2.04 (td, J = 13.1, 6.4 Hz, 1H), 2.44 (m, 2H), 2.65 (m, 2H), 2.98 (t, J = 7.8 Hz, 2H), 3.46 (s, 2H), 3.91 (d, J = 6.5 Hz, 2H), 4.00 (dd, J = 8.6, 7.0 Hz, 2H), 7.02 (m, 1H), 7.17–7.18 (m, 3H), 7.44 (d, J = 8.8 Hz, 2H), 7.69 (d, J = 8.7 Hz, 2H) ppm.

^{13}C NMR (CDCl_3): δ 19.1, 25.3, 27.5, 28.1, 28.9, 33.7, 36.6, 48.6, 75.8, 84.7, 119.9, 126.8, 127.2, 128.7, 129.1, 129.53, 129.63, 138.1, 138.9, 139.8, 140.9, 150.8, 185.2, 204.4 ppm.

MS (ESI^+): calculated for $\text{C}_{29}\text{H}_{37}\text{ClNO}_6\text{S}^+$ 562.20; found 561.83.

4-Chloro-N-(3-((2-isobutoxy-5-oxocyclopent-1-en-1-yl)methyl)phenethyl)benzenesulfonamide (49)—To a solution of **48** (0.120 g, 0.214 mmol) in dichloromethane (5.0 mL) 2,2,2-trifluoroacetic acid (1.5 mL) was added and the resulting mixture was stirred at rt for 0.5 h. The reaction mixture was concentrated *in vacuo* and purified by reverse phase preparative HPLC to furnish **49** as white solid (51% yield).

^1H NMR (CDCl_3): δ 0.99 (d, J = 6.7 Hz, 6H), 2.00–2.06 (m, 1H), 2.45 (m, 2H), 2.66 (m, 2H), 2.71 (t, J = 6.8 Hz, 2H), 3.19 (q, J = 6.6 Hz, 2H), 3.42 (s, 2H), 3.92 (d, J = 6.5 Hz, 2H), 4.71 (t, J = 6.1 Hz, 1H), 6.84–6.86 (m, 1H), 7.02 (s, 1H), 7.13 (m, 2H), 7.45 (d, J = 8.6 Hz, 2H), 7.71 (d, J = 8.6 Hz, 2H) ppm.

^{13}C NMR (CDCl_3): δ 19.1, 25.3, 27.5, 28.9, 33.7, 35.9, 44.4, 75.8, 119.9, 126.3, 127.4, 128.71, 128.89, 129.3, 129.5, 137.6, 138.8, 139.2, 141.1, 185.2, 204.5 ppm.

IR (film): ν 3270, 3202, 2962, 2929, 2875, 1678, 1616 cm^{-1} .

HRMS (ESI^-): calculated for $\text{C}_{24}\text{H}_{27}\text{ClNO}_4\text{S}^-$ 460.1349; found 460.1348.

4-Chloro-*N*-(3-((2-hydroxy-5-oxocyclopent-1-en-1-yl)methyl)phenethyl)benzenesulfonamide (50)—Prepared as **43** from **49**.

mp: 176–178 °C (from ethyl acetate).

¹H NMR (DMSO-*d*₆): δ 2.37 (s, 4H), 2.60 (t, *J* = 7.5 Hz, 2H), 2.91–2.95 (m, 2H), 3.27 (s, 2H), 6.89 (d, *J* = 7.5 Hz, 1H), 6.94–7.01 (m, 2H), 7.09 (t, *J* = 7.5 Hz, 1H), 7.65 (d, *J* = 7.8 Hz, 2H), 7.76 (d, *J* = 8.0 Hz, 2H), 7.82 (broad t, *J* = 5.7 Hz, 1H), 11.9 (broad s, 1H) ppm.

¹³C NMR (DMSO-*d*₆): δ 26.4, 35.3, 44.0, 115.2, 125.7, 126.1, 128.0, 128.4, 128.5, 129.3, 137.1, 138.1, 139.3, 140.7 ppm.

IR (film): ν 3090, 1610 cm^{−1}.

HRMS [ESI]⁺: calculated for C₂₀H₂₀ClNNaO₄S⁺ 428.0699; found 428.0713.

2,2'-(1,3-Phenylene)diethanol (52)—To a solution of 2,2'-(1,3-phenylene)diacetic acid (1.000 g, 5.15 mmol) in anhydrous tetrahydrofuran (15 mL) lithium aluminum hydride (0.5860 g, 15.45 mmol) was added at 0 °C. The reaction mixture was stirred at 0 °C for 3 h, and finally quenched by addition of hydrochloric acid (1N) until pH = 7. The solution was filtered through filter paper, and extracted with ethyl acetate (15 mL × 2). The organic layer was then washed with brine (15 mL × 2), dried over sodium sulfate, filtered and concentrated *in vacuo*. Purification by silica gel column chromatography eluting with a gradient of ethyl acetate in hexanes (50%–80%) provided **52** (61% yield).

¹H NMR (MeOD): δ 2.79 (t, *J* = 7.1 Hz, 4H), 3.74 (t, *J* = 7.1 Hz, 4H), 7.05 (dd, *J* = 7.5, 1.5 Hz, 2H), 7.09 (s, 1H), 7.19 (t, *J* = 7.5 Hz, 1H) ppm.

¹³C NMR (MeOD): δ 40.3, 64.3, 127.9, 129.5, 130.8, 140.4 ppm.

2-(3-(2-((Triisopropylsilyl)oxy)ethyl)phenyl)ethanol (53)—To a solution of 2,2'-(1,3-Phenylene)diethanol (0.430 g, 2.59 mmol) and imidazole (0.176 g, 2.59 mmol) in anhydrous *N,N*-dimethylformamide (15 mL) chlorotriisopropylsilane (0.22 mL, 1.03 mmol) was slowly added at 0 °C and the resulting mixture was stirred at 0 °C for 3 h. The reaction was quenched by addition of water (5 mL), extracted with ethyl acetate (15 mL × 2), and washed with brine (10 mL × 2). The organic layer was dried over anhydrous sodium sulfate, filtered, and concentrated *in vacuo*. Purification by column chromatography using a mixture of 1:10 ethyl acetate/hexanes as eluant provided **53** as colorless oil (35% yield).

¹H NMR (CDCl₃): δ 1.09–1.05 (m, 21H), 1.85 (s, 1H), 2.86 (q, *J* = 6.7 Hz, 4H), 3.85 (t, *J* = 6.7 Hz, 2H), 3.91 (t, *J* = 7.1 Hz, 2H), 7.09 (d, *J* = 7.6 Hz, 1H), 7.11–7.12 (m, 2H), 7.24 (t, *J* = 7.8 Hz, 1H) ppm.

¹³C NMR (CDCl₃): δ 12.2, 18.1, 39.4, 39.9, 63.8, 65.0, 126.9, 127.5, 128.6, 130.1, 138.5, 139.8 ppm.

2-(3-(2-((Triisopropylsilyl)oxy)ethyl)phenyl)acetaldehyde (54)—To a solution of **53** (0.170 g, 0.527 mmol) in dichloromethane (4 mL) Dess-Martin periodinane (0.270 g, 0.632 mmol) was added and stirred at rt for 4 h. The reaction mixture was filtered through a pad of celite and concentrated *in vacuo*. Purification by silica gel column chromatography, using ethyl acetate in hexanes as eluant, provided **54** (77% yield).

^1H NMR (CDCl_3): δ 1.01–1.11 (m, 21H), 2.87 (t, J = 7.0 Hz, 2H), 3.65 (d, J = 2.5 Hz, 2H), 3.92 (t, J = 7.0 Hz, 2H), 7.08 (d, J = 7.5 Hz, 1H), 7.12 (s, 1H), 7.19 (d, J = 7.5 Hz, 1H), 7.30 (t, J = 7.5 Hz, 1H), 9.74 (t, J = 2.5 Hz, 1H) ppm.

^{13}C NMR (CDCl_3): δ 12.1, 18.1, 39.7, 50.7, 64.8, 127.5, 128.4, 129.0, 130.7, 131.8, 140.5, 199.5 ppm.

3-Hydroxy-2-(3-(2-((triisopropylsilyl)oxy)ethyl)phenethyl)cyclopent-2-enone (55)—Prepared as **46** from **54**. Yield: 74%.

^1H NMR (CD_3OD): δ 0.98 (m, 21H), 2.33 (q, J = 7.7 Hz, 2H), 2.40 (d, J = 7.4 Hz, 4H), 2.60 (q, J = 7.5 Hz, 2H), 2.73 (q, J = 6.9 Hz, 2H), 3.82 (q, J = 7.0 Hz, 2H), 6.97 (m, 3H), 7.07 (t, J = 7.4 Hz, 1H) ppm.

^{13}C NMR (CD_3OD): δ 13.3, 18.6, 24.2, 31.4, 34.9, 40.9, 66.2, 118.2, 127.3, 127.8, 129.2, 130.5, 140.44, 140.45, 143.3 ppm.

2-(3-(2-Hydroxyethyl)phenethyl)-3-isobutoxycyclopent-2-enone (56)—Prepared as **47** from **55**. Yield: 66%.

^1H NMR (CDCl_3): δ 0.98 (d, J = 6.5 Hz, 6H), 1.98–2.03 (m, 1H), 2.41–2.47 (m, 4H), 2.60–2.62 (m, 2H), 2.73 (t, J = 7.5 Hz, 2H), 2.84 (t, J = 6.5 Hz, 2H), 3.84–3.87 (m, 4H), 7.02–7.08 (m, 3H), 7.20 (t, J = 7.5 Hz, 1H) ppm.

tert-Butyl (4-chlorophenyl)sulfonyl(3-(2-(2-isobutoxy-5-oxocyclopent-1-en-1-yl)ethyl)phenethyl)-carbamate (57)—Prepared as **33** from **56**. Yield: 43%.

^1H NMR (CDCl_3): δ 0.98 (d, J = 6.5 Hz, 6H), 1.59 (s, 9H), 1.98–2.03 (m, 1H), 2.43–2.45 (m, 4H), 2.62 (t, J = 4.5 Hz, 2H), 2.73 (t, J = 8.0 Hz, 2H), 2.99–3.03 (m, 2H), 3.83–3.86 (m, 2H), 3.98–4.01 (m, 1H), 4.19–4.24 (m, 1H), 7.05–7.11 (m, 3H), 7.21 (t, J = 7.5 Hz, 1H), 7.76 (d, J = 8.5 Hz, 2H), 7.77 (d, J = 8.5 Hz, 2H) ppm.

^{13}C NMR (CDCl_3): δ 19.1, 23.5, 25.2, 28.1, 28.9, 33.8, 33.9, 36.7, 48.8, 75.6, 84.8, 119.9, 126.7, 127.0, 128.7, 129.1, 129.4, 129.7, 138.0, 139.0, 139.9, 142.9, 150.9, 184.9, 204.9 ppm.

4-Chloro-N-(3-(2-(2-isobutoxy-5-oxocyclopent-1-en-1-yl)ethyl)phenethyl)benzenesulfonamide (58)—Prepared as **42** from **57**. Yield 28%

^1H NMR (CDCl_3): δ 1.00 (d, J = 6.7 Hz, 6H), 2.06–1.98 (m, 1H), 2.45–2.39 (m, 4H), 2.61 (m, 2H), 2.69 (t, J = 7.7 Hz, 2H), 2.73 (t, J = 6.7 Hz, 2H), 3.22 (q, J = 6.3 Hz, 2H), 3.88 (d, J = 6.5 Hz, 2H), 5.03–5.04 (bs, 1H), 6.86 (d, J = 7.5 Hz, 1H), 6.95 (s, 1H), 6.99 (d, J = 7.6 Hz, 1H), 7.13 (t, J = 7.5 Hz, 1H), 7.44 (d, J = 8.6 Hz, 2H), 7.75 (d, J = 8.6 Hz, 2H) ppm.

^{13}C NMR (CDCl_3): δ 19.1, 23.2, 25.2, 28.9, 33.6, 33.8, 36.0, 44.5, 75.8, 119.8, 126.4, 127.2, 128.67, 128.72, 129.2, 129.5, 137.7, 139.00, 139.08, 142.6, 185.9, 205.5 ppm.

HRMS [ESI] $^+$: calculated for $\text{C}_{25}\text{H}_{30}\text{ClNO}_4\text{SNa}^+$ 498.1482 found 498.1476.

4-Chloro-N-(3-(2-(2-hydroxy-5-oxocyclopent-1-en-1-yl)ethyl)phenethyl)benzenesulfonamide (59)—Prepared as **43** from **58**. Yield 84%

^1H NMR ($\text{DMSO}-d_6$): δ 2.26 (dd, J = 9.5, 6.8 Hz, 2H), 2.34 (bs, 4H), 2.57 (dd, J = 9.4, 6.7 Hz, 2H), 2.66 (t, J = 7.5 Hz, 2H), 2.98 (q, J = 6.7 Hz, 2H), 6.95 (d, J = 6.6 Hz, 2H), 7.02 (d,

$J = 7.6$ Hz, 1H), 7.16 (t, $J = 7.9$ Hz, 1H), 7.64 (d, $J = 8.7$ Hz, 2H), 7.77 (d, $J = 8.6$ Hz, 2H), 7.83 (t, $J = 5.7$ Hz, 1H), 11.55 (s, 1H) ppm.

^{13}C NMR (DMSO- d_6): δ 22.8, 33.3, 35.3, 44.1, 115.5, 125.98, 126.09, 128.2, 128.41, 128.51, 129.3, 137.1, 138.3, 139.4, 142.0 ppm.

IR: ν 3253, 2924, 1709, 1587 cm^{-1} .

HRMS [ESI] $^-$: calculated for $\text{C}_{21}\text{H}_{21}\text{ClNO}_4\text{S}^-$ 418.0880 found 418.0878.

CPD-benzamidine Salt Detection with ESI-MS

CPD-benzamidine salts were detected using ESI-MS with an Acquity TQMS controlled by MassLynx software (Waters Corporation, Milford, MA, USA). Source gas flow rates, temperatures and voltages were optimized for the detection of intact salt ions.

CPD:benzamidine mixtures (10 μM :10 μM) in acetonitrile were vortexed at rt for five min and then infused using the detector's syringe pump at 30 $\mu\text{L}/\text{min}$. Mass spectra were acquired in positive ion mode with a 0.5 second scan rate over 30 seconds. Scans one through fifty were combined and analyzed.

CPD-benzamidine Relative Affinity by Competition

Compound **8** and benzamidine were mixed with a competing acid at equimolar amounts (10 μM :10 μM :10 μM) in acetonitrile. Mixtures were vortexed at rt for 10 min then infused into the MS at 30 $\mu\text{L}/\text{min}$. After stabilization of the infusion flow, mass spectra were acquired in positive ion mode with a 0.2 second scan rate over 30 seconds. Scans 1–100 were combined and the intensity of the **8**-benzamidine salt ion, [**8**(^{79}Br)•benzamidine+H] $^+$ (m/z 401), was recorded. Affinities between acids and the benzamidine base were determined relative to **8** as the average ($N=3$) reduction of intensity of the **8**-benzamidine salt ion in the presence of a competing acid.

IP₁ Functional Assay

The activity of the TP receptor was measured by quantifying cellular levels of the IP₃ metabolite, IP₁, using a homogenous time-resolved fluorescence (HTRF) assay kit (IP-One Tb, Cisbio, Bedford, MA, USA). QBI-HEK 293A cells that were stably-transfected with human or mouse TP receptor cDNA were plated into 384-well plates at 10,000 cells/well in DMEM containing 4.5 g/L glucose, 10% fetal bovine serum, *L*-glutamine, and penicillin/streptomycin. The cells were incubated for 16 h at 37 °C with 5% CO₂, after which culture media was removed and the cells were then incubated for 15 min at 37 °C with 5% CO₂ in 10 mM Hepes, 1 mM CaCl₂, 0.4 mM MgCl₂, 4.2 mM KCl, 146 mM NaCl, 5.5 mM glucose, 50 mM LiCl, pH 7.4 containing varying concentrations of test antagonist. The TP receptor agonist, I-BOP ([15-(1 α ,2 β (5Z),3 α -(1E,3S),4 α)]-7-[3-hydroxy-4-(p-iodophenoxy)-1-butenyl]-7-oxabicycloheptenoic acid), was added at 1.6 nM and incubated for 1 h at 37 °C with 5% CO₂. Tb-labeled Anti-IP₁ cryptate and D2-labeled IP₁ were subsequently added in lysis buffer and incubated for 1 hour at 25 °C according to the manufacturer's instructions. Plates were read on a Spectramax M5 microplate reader, with data expressed as the ratio of 665 nm/620 nm fluorescence.

Scintillation Proximity Binding Assay

QBI-HEK 293A cells expressing hTP or mTP receptor were grown as described above and harvested in phosphate-buffered saline with 1 mM EDTA. The cell pellet underwent homogenization in 20 mM Hepes, 1 mM EGTA, and 0.5 mM DTT with protease inhibitor cocktail, followed by centrifugation at 1000 $\times g$ for 10 min at 8 °C to remove cell debris. The resulting supernatant was centrifuged at 21,000 rpm for 30 min at 4 °C, with the pellet

resuspended in 20 mM Hepes, 1 mM EGTA, 100 mM NaCl. Membrane preparations were normalized to protein level as determined with a bicinchoninic acid assay and stored at -80°C . Test antagonists were incubated at 10 different concentrations with 100 μg PVT-WGA SPA beads (PerkinElmer, Waltham, MA, USA), 62.5 μg membrane and 20 nM ^3H -SQ29,548 (PerkinElmer, Waltham, MA, USA) in 50 mM Tris, 4 mM CaCl_2 , 0.1% ascorbic acid pH 7.5 for 2 h at 25°C in 384-well polystyrene plates. Plates were sealed and read on a scintillation counter. The percent total binding was determined, with total binding calculated from a minimum of three wells containing membrane, beads, and ^3H -SQ29,548 without antagonist. Non-specific binding was determined by incubating ^3H -SQ29,548 at multiple concentrations in the presence of 100 μM cold SQ29,548. Binding constants were determined from the determined IC_{50} values that were used in the Cheng-Prusoff equation.²³

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

ABS dedicates this article to Professor William C. Agosta, Emeritus (Rockefeller University), outstanding scientist/scholar, author and mentor, who first brought to our attention the remarkable acidity of the cyclopentane-1,3-dione class of molecules. Financial support for this work has been provided by the NIH/NIA (AG034140), and the NSF (CHE-0840438, X-ray facility).

List of Abbreviations

CPD	cyclopentane-1,3-dione
TP	thromboxane A_2 prostanoid
IP_1	inositol monophosphate

References

1. Patani GA, LaVoie EJ. Bioisosterism: A Rational Approach in Drug Design. *Chem Rev.* 1996; 96:3147–3176. [PubMed: 11848856]
2. Meanwell NA. Synopsis of Some Recent Tactical Application of Bioisosteres in Drug Design. *J Med Chem.* 2011; 54:2529–2591. [PubMed: 21413808]
3. Herr RJ. 5-Substituted-1H-tetrazoles as carboxylic acid isosteres: medicinal chemistry and synthetic methods. *Bioorg Med Chem.* 2002; 10:3379–3393. [PubMed: 12213451]
4. Boothe J, Wilkinson R, Kushner S, Williams J. Synthesis of Aureomycin Degradation Products. II. *J Am Chem Soc.* 1953; 75:1732–1733.
5. Hiraga K. Structures of cyclopentanepolyones. *Chem Pharm Bull (Tokyo).* 1965; 13:1300. [PubMed: 5864720]
6. Katrusiak A. Structure of 1, 3-cyclopentanedione. *Acta Crystallogr C.* 1990; 46:1289–1293.
7. Katrusiak A. Structure of 2-methyl-1, 3-cyclopentanedione. *Acta Crystallogr C.* 1989; 45:1897–1899.
8. Dickinson RP, Dack KN, Long CJ, Steele J. Thromboxane modulating agents. 3. 1H-imidazol-1-ylalkyl- and 3-pyridinylalkyl-substituted 3-[2-[(arylsulfonyl)amino]ethyl]benzenepropanoic acid derivatives as dual thromboxane synthase inhibitor/thromboxane receptor antagonists. *J Med Chem.* 1997; 40:3442–3452. [PubMed: 9341919]
9. Koreeda M, Liang Y, Akagi H. Easy generation of the dianions of 3-isobutoxycyclopent-2-en-1-ones and their reactions. *J Chem Soc Chem Comm.* 1979:449–450.
10. Ramachary DB, Kishor M. Direct amino acid-catalyzed cascade biomimetic reductive alkylations: application to the asymmetric synthesis of Hajos-Parrish ketone analogues. *Org Biomol Chem.* 2008; 6:4176–4187. [PubMed: 18972048]

11. Tilley JW, Danho W, Lovey K, Wagner R, Swistok J, Makofske R, Michalewsky J, Triscari J, Nelson D, Weatherford S. Carboxylic acids and tetrazoles as isosteric replacements for sulfate in cholecystokinin analogs. *J Med Chem.* 1991; 34:1125–1136. [PubMed: 2002454]
12. Henry JR, Marcin LR, McIntosh MC, Scola PM, Davis Harris G, Weinreb SM. Mitsunobu reactions of N-alkyl and N-acyl sulfonamides—an efficient route to protected amines. *Tet Lett.* 1989; 30:5709–5712.
13. Peters L, Frohlich R, Boyd ASF, Kraft A. Noncovalent Interactions between Tetrazole and an N,N'-Diethyl-Substituted Benzamidine. *J Org Chem.* 2001; 66:3291–3298. [PubMed: 11348109]
14. Tominey AF, Docherty PH, Rosair GM, Quenardelle R, Kraft A. Unusually Weak Binding Interactions in Tetrazole-Amidine Complexes. *Org Lett.* 2006; 8:1279–1282. [PubMed: 16562871]
15. Job P. Formation and stability of inorganic complexes in solution. *Ann Chim.* 1925; 9:113–125.
16. Shenker A, Goldsmith P, Unson CG, Spiegel AM. The G protein coupled to the thromboxane A2 receptor in human platelets is a member of the novel Gq family. *J Biol Chem.* 1991; 266:9309–9313. [PubMed: 1851174]
17. Bylund D, Toews M. Radioligand binding methods for membrane preparations and intact cells. *Methods Mol Biol (Clifton, NJ).* 2011; 746:135.
18. Ruan KH, Wu J, So SP, Jenkins LA, Ruan CH. NMR structure of the thromboxane A2 receptor ligand recognition pocket. *Eur J Biochem.* 2004; 271:3006–3016. [PubMed: 15233797]
19. Hirata M, Hayashi Y, Ushikubi F, Yokota Y, Kageyama R, Nakanishi S, Narumiya S. Cloning and expression of cDNA for a human thromboxane A2 receptor. *Nature.* 1991; 349:617–620. [PubMed: 1825698]
20. Yamamoto Y, Kamiya K, Terao S. Modeling of human thromboxane A2 receptor and analysis of the receptor-ligand interaction. *J Med Chem.* 1993; 36:820–825. [PubMed: 8464036]
21. Dogne J, Hanson J, Leval X, Pratico D, Pace-Asciak C, Drion P, Pirotte B, Ruan K. From the design to the clinical application of thromboxane modulators. *Curr Pharm Des.* 2006; 12:903–923. [PubMed: 16533159]
22. Shineman DW, Zhang B, Leight SN, Pratico D, Lee VMY. Thromboxane Receptor Activation Mediates Isoprostane-Induced Increases in Amyloid Pathology in Tg2576 Mice. *J Neurosci.* 2008; 28:4785–4794. [PubMed: 18448655]
23. Yung-Chi C, Prusoff WH. Relationship between the inhibition constant (KI) and the concentration of inhibitor which causes 50 per cent inhibition (I50) of an enzymatic reaction. *Biochem Pharmacol.* 1973; 22:3099–3108. [PubMed: 4202581]

Figure 1. Selected examples of known (A) and proposed (B) isosteres of the carboxylic acid moiety.

Figure 2.

The structure of known TP-receptor antagonist **12** (top) and overlap of the carboxylic acid moiety of **12** with CPD derivatives attached either at C2 or C4/C5 (bottom, A or B, respectively).

Figure 3.
[CPD-benzamidine] complexes detected by ESI-MS.

Figure 4. Relative intensity of the $[8\cdot\text{benzamidinium}]^+$ signal in a 1:1 mixture of **8** and benzamidinium (100%) or in 1:1:1 mixtures of **8**, benzamidinium and competing acid/acid surrogate (*i.e.*, **7**, **9**–**11**).

Figure 5. ^1H -NMR Job plot analysis of [**1**-benzamidine] complex showing 1:1 stoichiometry. The plot is derived from the chemical shift of the C2- H of **1** at different molar ratios with benzamidine. The total concentration is kept constant (*i.e.*, 0.125 M) in each experiment. The mole fraction x is defined as $[\textbf{1}]/([\textbf{1}] + [\text{benzamidine}])$; $\Delta\theta = \theta_{\text{complex}} - \theta_{\text{free}}$.

Figure 6. Calculated bi-coordinated “stacked” (A), and mono-coordinated “linear” (B) geometries of the [8-benzamidine] complex in gas phase.

Figure 7.
Docking of thromboxane A₂ (A), **12** (B) and **42** (C) to the TP-receptor.

Figure 8.
Schematic representation of key interactions of **12** and **42** with TP-receptor.

Scheme 1. Reagents and reaction conditions

a) (i) lithium diisopropylamide, -78°C (ii) 4-bromobenzylbromide, -78°C to rt over 1 h; *b*) 2N hydrochloric acid, acetone; *c*) 4-bromobenzaldehyde, Hantzsch ester, *L*-proline, dichloromethane, rt, 2h.

Scheme 2. Reagents and reaction conditions

a) $(i\text{-Pr})_3\text{Si-Cl}$, imidazole, N,N -dimethylformamide, 0°C , 3 h; b) NaBH_4 , H_2O , tetrahydrofuran, 70°C , 2 h; c) PPh_3 , I_2 , imidazole, Et_2O , acetonitrile, 0°C , 2 h; d) appropriate i -butyl-protected cyclopentane-1,3-dione, lithium diisopropylamide, tetrahydrofuran, from -78°C to rt; e) tetra- n -butylammonium fluoride, tetrahydrofuran, 0°C , 3 h; f) NaN_3 , N,N -dimethylformamide, 50°C , 45 min; g) PPh_3 , diethyl azodicarboxylate, $tert$ -butyl (4-chlorophenyl)sulfonylcarbamate, tetrahydrofuran, rt, 4 h; h) (i) H_2 , Pd-C, methanol, rt, 16 h, (ii) 4-chlorobenzenesulfonyl chloride, 2N NaOH, 0°C , 3 h; i) 2N HCl, acetone, rt, 6–12 h; j) 2,2,2-trifluoroacetic acid, dichloromethane, rt, 2–5 h.

Scheme 3. Reagents and reaction conditions

a) CPD (1), Hantzsch ester, *L*-proline, dichloromethane, rt, 12 h; *b*) p-toluenesulfonic acid (cat.), *i*-butanol/benzene, reflux, 16 h; *c*) *tert*-butyl (4-chlorophenyl)sulfonylcarbamate, PPh₃, diethyl azodicarboxylate, tetrahydrofuran, rt, 4 h; *d*) 2,2,2-trifluoroacetic acid, dichloromethane, rt, 2 h; *e*) 2N hydrochloric acid, acetone, rt, 12 h; *f*) LiAlH₄, diethyl ether, -78 °C, 3 h; *g*) (*i*-Pr)₃Si-Cl, imidazole, *N,N*-dimethylformamide, 0 °C, 3 h; *h*) pyridinium chlorochromate, dichloromethane, rt, 2 h.

Table 1

pK_a , logP and logD_{pH 7.4} values. All determinations were carried out via UV-metric or potentiometric (pH-metric) method.

Compound #	pK_a	LogP	LogD _{pH 7.4}
1	4.20 ± 0.01 (4.4) [§]	ND [‡]	ND [‡]
2	4.47 ± 0.01 (4.7) [§]	ND [‡]	ND [‡]
7	3.96 ± 0.01	3.02 ± 0.01	-0.42
8	4.24 ± 0.01	3.28 ± 0.01	0.11
9	4.20 ± 0.02	3.01 ± 0.01	-0.19
10	4.33 ± 0.01	3.00 ± 0.01	0.01
11	4.62 ± 0.01	2.21 ± 0.01	-0.42

[‡]Not determined;

[§]Literature value.⁵

Table 2

TP-receptor antagonist activity of test compounds

Cpd#	X	TP Receptor Antagonist Activity	
		IP ₁ Assay IC ₅₀ (nM)	
		Human TP	Mouse TP
12		61 (+/- 21)	27 (+/- 11)
60		15117 (+/- 7994)	1627 (+/- 721)
61		64 (+/- 41)	8.0 (+/- 6)
62		377 (+/- 203)	91 (+/- 33)
50		>10,000	>10,000
59		>10,000	1229 (+/-462)
41		250 (+/- 33)	52 (+/- 17)
42		131 (+/- 70)	3.1 (+/- 3.4)

	TP Receptor Antagonist Activity	
		IP ₁ Assay IC ₅₀ (nM)	
Cpd#	X	Human TP	Mouse TP
37		>10,000	1786 (+/- 1212)
43		171 (+/- 35)	9.4 (+/- 8.0)
38		3801 (+/-443)	463 (+/-250)
44		198 (+/- 97)	18 (+/- 21)
39		5303 (+/-1078)	259 (+/- 90)
45		41 (+/- 32)	5.8 (+/- 2.5)
40		21853 (+/-2151)	2524 (+/-233)

Table 3

TP-receptor binding affinity of selected compounds, determined by radioligand-binding assay.

Cpd#	X	TP Receptor Binding	
		Scintillation Proximity Assay Kd (nM)	
		Human TP	Mouse TP
12		141 (+/- 31)	26 (+/- 9.6)
60		>10,000	>10,000
61		331 (+/- 109)	19 (+/- 9.0)
62		359 (+/- 60)	17 (+/- 7.5)
50		11400 (+/- 3960)	1196 (+/- 1334)
41		203 (+/- 87)	11 (+/- 9.5)
42		328 (+/- 87)	26 (+/- 5.7)