Published in final edited form as:

Biol Psychiatry. 2011 November 1; 70(9): 859–865. doi:10.1016/j.biopsych.2011.05.033.

Decreased FMRP Expression Underlies Amygdala Dysfunction in Carriers of the Fragile X Premutation

David Hessel^{1,2}, John M. Wang⁵, Andrea Schneider^{1,2}, Kami Koldewyn^{6,7}, Lien Le⁵, Christine Iwahashi³, Katherine Cheung³, Flora Tassone^{1,3}, Paul J. Hagerman^{1,3}, and Susan M. Rivera^{1,4,5}

¹ Medical Investigation of Neurodevelopmental Disorders (M.I.N.D.) Institute, University of California-Davis, Medical Center, Sacramento, CA

² Departments of Psychiatry and Behavioral Sciences, University of California-Davis, Medical Center, Sacramento, CA

³ Department of Biochemistry and Molecular Medicine, University of California-Davis, School of Medicine, Davis, CA

⁴ Department of Psychology, University of California-Davis, Davis, CA

⁵ Center for Mind and Brain, University of California-Davis, Davis, CA

⁶ Department of Brain and Cognitive Sciences, Massachusetts Institute of Technology, Cambridge, MA

⁷ McGovern Institute for Brain Research, Massachusetts Institute of Technology, Cambridge, MA

Abstract

Background—The fragile X premutation provides a unique opportunity for the study of genetic and brain mechanisms of behavior and cognition in the context of neurodevelopment and neurodegeneration. Although the neurodegenerative phenotype, fragile X-associated tremor/ataxia syndrome (FXTAS), is well described, evidence of a causal link between the premutation and psychiatric disorder earlier in life, clear delineation of a behavioral/cognitive phenotype, and characterization of the physiological basis of observed symptoms have been elusive.

Methods—We completed functional magnetic resonance imaging (fMRI) targeting the amygdala with an emotion-matching task and concurrent infra-red eyetracking, *fMRI* molecular genetic testing, and neuropsychological assessment in 23 men with the premutation (mean age = 32.9 years) and 25 male controls (mean age = 30.1).

Results—Premutation carriers had significantly smaller left and right amygdala volume and reduced right amygdala activation during the task relative to controls. Although both elevated

© 2011 Society of Biological Psychiatry. Published by Elsevier Inc. All rights reserved.

Address correspondence to: David Hessel, Ph.D., Associate Professor, M.I.N.D. Institute, University of California, Davis Medical Center, 2825 50th Street, Sacramento, CA 95817, Ph: 916-703-0249, Fax: 916-703-0244, david.hessel@ucdmc.ucdavis.edu.

Financial Disclosures

Dr. Hessel reports financial support from Hoffman LaRoche, Novartis, and Seaside Therapeutics for clinical trials of patients with fragile X syndrome. Drs. P Hagerman and Tassone report a patent for screening/detection of expanded CGG-repeat *FMR1* alleles. Drs. Rivera, Koldewyn, Tassone, and Schneider, and Ms. Cheung, Ms. Iwahashi, Mr. Wang, and Ms. Le report no biomedical financial interests or potential conflicts of interest.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

FMRI mRNA and reduced *FMRI* protein (FMRP) were associated with the reduced activation, multiple regression analysis suggested that reduced FMRP is the primary factor. Premutation carriers also had higher ratings of autism spectrum symptoms than controls that were associated with the reduced amygdala response.

Conclusions—Although prior studies have emphasized a toxic gain-of-function effect of elevated mRNA associated with the premutation, the current results point to the role of reduced FMRP in alterations of brain activity and behavior.

Keywords

FMRI; fragile X; premutation; amygdala; insula; FXTAS

Introduction

Abnormal trinucleotide (CGG)-repeat expansions of the fragile X mental retardation 1 (*FMRI*) gene confer significant risk for a surprisingly wide range of health problems, including premature ovarian insufficiency (1, 2), neurodegenerative disease (fragile X-associated tremor/ataxia syndrome or FXTAS (3), and, of course, fragile X syndrome (FXS), the most frequent known inherited cause of intellectual disability and autism (4). While these three clinical outcomes are well-defined and substantiated by empirical data, other apparent manifestations of *FMRI* mutations, especially those associated with the premutation, appear to have either a variable penetrance (perhaps due to secondary genetic or environmental factors) or a more subtle or sub-clinical phenotype. Experimenter bias and sampling methods, in some studies, may also lead to discovery of deficits or symptoms mistakenly attributed to the premutation. For example, while studies have suggested higher rates of psychiatric or neurodevelopmental disorders among premutation carriers, these disorders are common in the general population, making the clear attribution of increased risk related to the premutation challenging, especially when studies are small or inadequately controlled. The fragile X premutation occurs in 1 in 113–259 females and 1 in 251–813 males (5–8) and therefore it is important to understand its potential influence on brain structure and function, as well as on cognition and behavior.

Neuroimaging techniques such as structural and functional magnetic brain imaging (MRI), in conjunction with molecular genetic measures specific to the *FMRI* gene, can provide objective methods for investigation of the premutation phenotype and insight into gene-brain-behavior associations explaining a physiological basis of clinical observations. For example, Moore and colleagues (9) reported reduced grey matter volume in the amygdalo-hippocampal complex that was associated with reduced percentage of blood lymphocytes expressing the *FMRI* protein (FMRP) among a group of men with the fragile X premutation. This observation led us to hypothesize that *FMRI*-mediated altered limbic system function may partially explain neuropsychiatric and neurodevelopmental disorders, including autism spectrum disorders (10–13) and anxiety and depression (14–17) that are known to occur in at least a subgroup of premutation carriers.

We conducted a preliminary structural and functional brain MRI (1.5 Tesla) study, focused on the amygdala, in a convenience sample of 12 primarily clinic-referred men with the premutation (mean age 42.9 years) and 13 men without the premutation of similar age, level of education, and intellectual functioning (18). Relative to controls, men with the premutation had significantly reduced amygdala activation while passively viewing fearful vs. scrambled faces. The reduced amygdala activation was significantly associated with abnormal elevation of *FMRI* mRNA in the premutation group, suggesting a dose-dependent pattern in which fold-increase in mRNA was associated with decreased amygdala

responsiveness. The clinical relevance of reduced amygdala activation in the premutation carriers was demonstrated by significant correlation with self-reported current psychological symptoms. In light of reports of autism spectrum disorders in young males with the premutation (10–13), we suggested that either reduced FMRP or elevated *FMR1* mRNA or a combination of both could contribute to these symptoms via aberrant limbic system function.

The purpose of the present study was to examine amygdala volume and function in a larger sample of non-referred participants with the premutation relative to controls; to determine the relative effects of reduced FMRP and elevated *FMR1* mRNA on these measures of interest; and to study the clinical relevance of amygdala morphology and function, especially with regard to autism spectrum symptomology.

Methods

Participants

Participants included 23 men with the *FMR1* premutation (mean age 32.9 years) and 25 matched controls (mean age 30.1 years). *FMR1* DNA testing was used to confirm allele status for all participants. None of the participants with the premutation were mosaic for either repeat size or methylation – all had between 55 and 200 CGG repeats with no methylation. The control group was matched for age, IQ, level of education, handedness, psychoactive medication use, and ethnicity. Group demographic statistics are shown in Table 1. Males with the premutation were selected from fragile X pedigrees either because a) they were known to carry the premutation (and confirmed by *FMR1* DNA testing) or b) were untested sons of females with the premutation who were later confirmed through study testing. None was ascertained or referred to the clinic due to clinical symptoms. Neurological examinations including assessment of possible FXTAS were normal for all participants. Exclusion criteria included any acute medical condition such as renal, liver, or cardiac disease, migraine headache, history of head trauma, history of alcoholism or drug abuse, and use of current medications that effect cerebral blood flow. Two participants with the premutation were included in a previous study (16).

Neuropsychological Assessment

Cognitive ability was based on Full Scale IQ using the Wechsler Adult Intelligence Scale, Third Edition (WAIS-III;19). Autism phenotype features were assessed with the Autism Diagnostic Observation Schedule, Module 4 (ADOS; 20) and the Social Responsiveness Scale (SRS), adult version (21).

Molecular Genetic Measures

CGG repeat size—Genomic DNA was isolated from peripheral blood lymphocytes using standard methods (Qiagen, Valencia, CA). Repeat size and methylation status were determined using both PCR and Southern Blot analysis using an Alpha Innotech FluorChem 8800 Image Detection System (San Leandro, CA) as previously described (22, 23).

***FMR1* mRNA**—All quantifications of *FMR1* mRNA were performed using a 7900 Sequence detector (PE Biosystems) as previously described (24).

FMRP levels (ELISA)—FMRP was quantified in lymphocytes utilizing a recently described sandwich Enzyme Linked ImmunoSorbent Assay (ELISA) for FMRP, as described previously (25). The FMRP ELISA assay differs from the commonly used immunocytochemistry (IHC) method in that the ELISA approach provides a quantitative

measure of FMRP level, whereas the IHC method does not measure protein level, only the proportion of cells with detectable staining.

MRI Acquisition

Images were acquired on a 3.0 T Siemens scanner with Echospeed gradients and a Siemens 8-channel whole head coil. Functional MRI (fMRI) was performed using a single-shot gradient recalled echo planar imaging sequence that is auto corrected for motion and magnetic field distortions utilizing a point spread function mapping approach (26). Thirty-eight slices (3.4 mm thick), aligned 30 clockwise from parallel to the anterior and posterior commissure plane and covering the whole brain, were imaged with a temporal resolution of 2 sec using a T2* weighted gradient echo planar pulse sequence with TE 13 ms, flip angle 84, FoV 220 mm, and base resolution 64. For the purpose of manual segmentation and to aid in localization, co-registration, and normalization of functional data, a high-resolution T1-weighted magnetization prepared rapid acquisition gradient echo (MPRAGE) sequence of 192 slices with TR 2170 ms, TE 4.82 ms, FoV 256 mm, and 1.0 mm slice thickness was acquired.

Image Preprocessing

Pre-processing of the imaging data was completed using statistical parametric mapping software (SPM5; Wellcome Department of Imaging Neuroscience, University College London, UK). Images were corrected for movement using least squares minimization without higher-order corrections for spin history and normalized to stereotaxic MNI (Montreal Neurological Institute), then resampled every 2 mm using 4th Degree B-spline interpolation and smoothed with a 5 mm Gaussian kernel to decrease spatial noise.

Amygdala Volume

Amygdala volumes were quantified by operator-guided manual segmentations using Mayo BIR's Analyze 8.5 and 9.0 (27–29). These guidelines were developed from the anatomical analysis of post-mortem human brains using histological sections of tissue cut perpendicular to the hippocampus axis. For a detailed description of this protocol, see Schumann et al. (30). Amygdala volumes were corrected for total cerebral volume. The reliability between tracers on 10 cases was 94.7% for the left amygdala and 96.9% for the right amygdala.

Total Cerebral Volume

To obtain a measure of total cerebral volume, each series of images was edited manually using Mayo BIR's Analyze 8.5 and 9.0 (27–29) to remove non-brain structures, the brainstem, and the cerebellum. Using a Gaussian cluster multispectral thresholding tool, the ventricles were defined and excluded.

fMRI Emotion Processing Task

The emotion processing task was modeled after a task used by Brown and colleagues (31). Participants viewed three faces expressing either fear or anger, where the emotion presented at the bottom of the screen was identical to one of the two emotions presented above. Participants were asked to press a button to indicate which top emotion matched the sample while accuracy and reaction time were recorded. Faces were taken from two standard sets of pictures of facial affect (32, 33) and presented in grayscale with hair and other non-face features cropped. Fearful and angry expressions are perceived as threatening and robustly engage the amygdala (34, 35). Control stimuli consisted of three simple geometric shapes (squares, circles, and ovals), colored in grayscale gradients matching the face stimuli in average luminance and arranged in the same configuration. Subjects matched the stimuli similarly to the emotion trials. Each set of three images, whether emotions or objects, was

presented for three seconds in a pseudo-random order (see Figure 1). Intervals between stimuli were jittered between 1 and 5 seconds. A total of 104 face trials and 104 control trials were presented over 2 runs.

Eye Tracking

Eye tracking data was collected concurrent with the fMRI emotion-matching task using a long-range infrared eye tracking system (Applied Science Laboratories, Bedford, MA) and processed using Eyeanal 6.0 (Eyetracker, North Sydney, Australia). Areas of interests (AOI) were rectangles covering all three face or shape ovals individually.

fMRI analysis

Statistical analysis was performed on individual and group data using the general linear model and the theory of Gaussian random fields as implemented in SPM5 (36) with one predictor (convolved with a standard canonical hemodynamic response function) for each condition. For both within-group and between-group comparisons, significant clusters were defined as those that exceeded a threshold value equivalent to a one-tailed t -test $p < 0.01$ for a higher level of quality control, corrected False Discovery Rate (FDR) on the voxel level (37). Once thresholded, activation foci were superposed on normalized and averaged high-resolution MPRAGE images, their locations were identified both manually using an atlas with known neuroanatomical landmarks (38) and automatically using xjView (<http://www.alivelearn.net/xjview8/>).

A within-subjects procedure was used to model all the effects of interest for each subject by contrasting experimental and control trials. This model estimates the error variance for each condition of interest across participants rather than across scans (39). These contrast images generated previously on the individual level were analyzed using a general linear model to determine voxel-wise t -statistics and generating one contrast image per participant, per effect of interest.

Within-group analyses of each contrast were performed to identify brain regions showing similar response modulations across participants in both the premutation and control groups for each contrast. Between group analyses were then performed to determine differences in average activation responses to each contrast between the two groups.

Region of interest (ROI) analyses for the amygdala and insula were carried out using MarsBar toolbox for SPM (40). Contrasts defined remained the same as previously described. Each contrast of interest was analyzed only in voxels that fell either within the MNI template (41) of the area of interest (e.g. insula) on normalized images or within boundaries defined by operator-guided manual segmentations of the individual specific amygdala, as described above, applied to individual unnormalized functional images. A t -statistic termed “contrast value” was then calculated as the average of all voxels’ contrast values that fell within the defined ROI. The contrast value in these analyses is comparable to the Z-score reported in the whole-brain analyses tables. The contrast values were then used in between group independent sample T-tests and Spearman’s correlation and regression analyses in conjunction with biological measures and neuropsychological data. In all analyses utilizing ROI-derived data, outlier values of greater than 1.5 interquartiles from the median of the sample were excluded from analysis.

Results

Participant *FMRI* genetic, demographic and clinical descriptive statistics are shown in Table 1. The FMRP distribution was positively skewed and therefore log-transformed to achieve normality. FMRP was reduced by 12% in the premutation group relative to controls, $t(38) =$

2.19, $p < 0.034$. Premutation carriers had significantly higher ADOS total scores than controls, with mean level (5.87) somewhat below the autism spectrum cut-off (7). Two controls and 6 premutation carriers had ADOS scores beyond the autism spectrum cut-off. On the SRS, reported by a spouse or significant other, premutation carriers had elevated scores but this difference did not reach significance.

Total cerebral and amygdala volume

Independent samples t-tests revealed that premutation carriers had significantly smaller total amygdala volume than controls for both raw [$t(46) = 2.328$, $p = 0.024$] and corrected [$t(46) = 2.129$, $p = 0.039$] volumes (Supplement 1 and Figure 2). This amygdala volume reduction in the corrected data was stronger on the right [$t(46) = 2.073$, $p = 0.044$] but was still marginally significant on the left [$t(46) = 1.984$, $p = 0.053$]. Both raw values for right [$t(46) = 2.033$, $p = 0.048$] and left [$t(46) = 2.267$, $p = 0.028$] amygdala were significantly smaller in the premutation group. Total cerebral volume did not differ between the two groups [$t(46) = -0.423$, $p = 0.674$].

In the premutation carrier and control groups, all correlations between *fMRI* molecular measures and corrected amygdala volumes were non-significant.

Emotion matching task performance

There were no between group differences for percentage correct in response accuracy [premutation mean = 81.4%, control mean = 78.8%, $t(44) = -0.952$, $p = 0.346$] or reaction time [premutation mean = 2.18 s, control mean = 2.20 s, $t(45) = 0.210$, $p = 0.835$] during the emotion matching task.

Eye-tracking results

There were no group differences in the average looking time per trial for the defined AOIs covering the faces or proportion of looking time at faces (premutation means 1.40 s and 75.75%, control means 1.40 s and 71.74%, respectively) or shapes (premutation means 1.14 s and 70.09%, control means 1.08 s and 63.72%, respectively), all $p > 0.17$.

Amygdala and associated brain activation

Within-Group analysis—When matching emotional faces compared to matching shapes (emotion faces > shapes), the premutation group showed similar brain activation patterns compared with controls at a thresholding of $p < 0.01$ with FDR correction (Supplement 1). Both groups showed robust activation bilaterally in the insula, mid cingulate gyrus, fusiform gyrus, parietal lobe, superior frontal gyrus and middle frontal gyrus. Both controls and premutation carriers activated the amygdala bilaterally which was evident in the normalized whole brain analysis (see Figure 3).

Between-group analysis—A direct comparison between premutation and control groups for the emotion faces > shapes contrast across the whole brain showed no significant activation differences in either direction (premutation > control or control > premutation) using a FDR correction at $p < 0.01$. Region of interest (ROI) analyses, where activity specific to each participant was extracted from manually-traced amygdala ROIs, revealed that left amygdala activation was significantly less for the premutation group [$t(44) = 2.161$, $p = 0.036$] (supplement 1). The group difference in the right amygdala was non-significant, [$t(44) = 1.43$, $p = 0.160$]. No significant group differences were observed in the insula (all $p > 0.80$).

Molecular and Functional MRI Correlation Analysis

As expected in the premutation group, CGG repeat number was strongly and positively correlated with *FMR1* mRNA expression ($r = 0.842$, $p < 0.001$). There was also a negative trend for FMRP with increasing *FMR1* mRNA concentration ($\rho = -0.482$, $p = 0.069$), as expected for reduced efficiency of translational initiation with increasing repeat size in RNA (42).

FMR1 mRNA was correlated negatively with right amygdala activation (ROI analysis; $\rho = -0.625$, $p = 0.002$) and with right insula activity ($\rho = -0.610$, $p = 0.003$) in the emotion faces > control contrast. FMRP correlated positively with both right ($\rho = 0.697$, $p = 0.003$) and left ($\rho = 0.771$, $p < 0.0001$) amygdala activation (Figure 4) and with right insula activation ($\rho = 0.524$, $p = 0.037$). No significant correlations between activation of these brain regions and the genetic measures were observed in the control group.

Because we observed strong associations between reduced amygdala activation and both elevated *FMR1* mRNA and decreased FMRP in premutation carriers, we sought to determine the relative and independent effects of these factors in multiple regression analyses ($n = 16$). Using right and left amygdala activation as the dependent variables, and FMRP and *FMR1* mRNA as independent variables, we found that FMRP was positively correlated with activation (left, $\beta = 0.681$, $p = 0.009$; right, $\beta = 0.572$, $p = 0.023$) whereas *FMR1* mRNA showed no significant association (left, $\beta = 0.024$, $p = 0.915$; right, $\beta = -0.327$, $p = 0.158$). Analogous regression analyses with left and right insula activation demonstrated no significant associations with *FMR1* mRNA, and an association between FMRP and right insula activation that approached significance ($\beta = -0.490$, $p = 0.055$).

Neuropsychological Correlates of Amygdala Activation

The planned correlations between amygdala activation for the emotion > shapes contrast and SRS total score was significant in the premutation group only for the right amygdala ROI ($\rho = -0.469$, $p = 0.032$). Post-hoc exploratory correlations using the subscales of the SRS showed that reduced right amygdala activation was most strongly associated with problems in social information processing ($\rho = -0.588$, $p = 0.005$) and reciprocal social communication ($\rho = -0.497$, $p = 0.022$), although similar correlations were observed for the left amygdala. No other significant correlations between amygdala activation and clinical measures were observed including those with IQ and ADOS scores.

Discussion

The results of this study demonstrate that relative to controls, men with an *FMR1* premutation allele had smaller amygdala volumes and reduced amygdala activation during an emotion-matching task. In contrast to our prior study (18), in this investigation we were able to determine that aberrant amygdala function was more strongly associated with reduced FMRP (the cause of FXS) than abnormal elevation of *FMR1* mRNA (the hypothesized toxic mRNA gain-of-function model underlying FXTAS). To our knowledge this the first evidence of the impact of reduced FMRP without methylation in *FMR1* premutation carriers. Standardized clinical assessment indicated that men with the premutation had more significant communication and reciprocal social behavior symptoms of autism spectrum disorder compared to controls; however, on a group level the severity of these symptoms did not reach clinical significance. Exploratory analyses showed that the FMRP-mediated blunted amygdala response may be correlated with deficits in social information processing, a component of the broader autism phenotype. Since no evidence of transcriptional silencing was observed in any individual with a premutation included in this study, who do show levels of *FMR1* mRNA substantially higher than normal levels (24, 43,

44), the mild reductions in FMRP with increasing CGG repeat length (45–48) are due primarily to reduced efficiency of translational initiation (42). Although the full range of FMRP levels in individuals with the full mutation, premutation and general population controls has not been fully elucidated, it is important to emphasize that the relatively lower FMRP levels observed in some premutation carriers in this study are likely to be quite modest compared to most males with the full mutation and FXS who express essentially no or very little FMRP and who demonstrate more substantial behavioral and cognitive deficits.

Overall, the study indicates that neurodevelopmental and psychiatric symptoms previously reported among younger men and boys with the premutation may be due to the same molecular genetic mechanism leading to FXS, but to a milder degree. Previous findings, including our own (16, 18) highlighting the potential negative impact of elevated mRNA on these symptoms suggests that both mRNA- and protein-level effects may be operating in the premutation range, with protein decrements perhaps playing a larger role, at least in amygdala-mediated functions.

In the current cohort of participants, we found reduced amygdala volumes in premutation carriers, whereas in our prior study of a different group of participants (18), we did not find this effect. A number of factors might explain this difference. For example, the more powerful 3T scanner used here provided higher signal-to-noise ratio, enabling higher levels of image quality, speed and resolution, perhaps making the group effect easier to detect. Also, somewhat larger sample sizes in the present study provided additional data and greater power to detect differences.

Prior fMRI studies in FXS using social stimuli (human faces showing direct or averted gaze) (49) have documented greater sensitization in the left amygdala with successive exposure to direct gaze in a sample of adolescent boys. Given that patients with FXS and some individuals with the premutation have reduced FMRP (albeit to a different degree), we might expect premutation carriers to have a similarly increased amygdala response. The explanation for this inconsistency is unclear but could be related to the nature of the tasks, the added complexity of mildly reduced FMRP and elevated message in premutation carriers, or due to developmental differences. Nevertheless, this research highlights the critical role of amygdala function in perception of and response to human social stimuli and in social behavior in individuals affected by *FMR1* mutations.

In our previous report, we suggested that the altered limbic system function observed in men with the premutation could reflect early prodromal brain changes associated with later-onset FXTAS. However, given that the primary molecular genetic mechanism underlying FXTAS neuropathology is thought to involve abnormal elevation of the expanded-CGG-repeat *FMR1* mRNA, our current results suggest that lowered FMRP contributes to amygdala dysfunction, and that the latter effect could be both neurodevelopmental in origin and independent of FXTAS progression.

The reduced amygdala activation in premutation carriers observed here cannot be explained by decreased visual fixation on stimuli, as we observed similar fixation patterns between groups. Recruitment biases are unlikely to have affected our findings because participants were selected at random from pedigrees and were not clinic referred. The blunted amygdala response cannot be explained by reduced amygdala volume as the ROI method controls for individual differences in amygdala size and, additionally, we did not find a significant correlation between volume and activation (data not shown). Interestingly, although differences in brain activation were observed during the emotion-matching task, we found no difference in performance, indicating that brain activation patterns may be more sensitive to effects of the premutation than observable behavior.

There were important limitations to the study. Although we took several steps to minimize recruitment bias, this form of bias cannot be completely eliminated because individuals make decisions whether to participate based on many factors, some of which may be associated with effects of the premutation being studied here. Although men with the premutation were rated as having higher rates of autism symptoms on clinical exam, it is not possible to rule out effects of experimenter bias. Many participants were assessed blind to group; however, blinding was not possible in all cases due to personal information revealed during the interview portions of the ADOS. Approximately one-third of participants in each group were taking psychoactive medication which may have influenced our results, and somewhat detracts from the generalizability of the findings. It is important to emphasize that the *FMR1* molecular genetic measures were ascertained from blood samples and may not necessarily reflect what would be found in brain tissue. Finally, we were unable to derive valid FMRP values for all participants in the study; as such the results pertaining to the association between FMRP level and brain function must be considered preliminary.

Our findings may have treatment implications for some individuals with the premutation who present with significant symptoms of FXS or ASD. The “mGluR theory” of FXS states that mental impairment and phenotypic behaviors associated with FXS arise, at least in part, from constitutive activation of translational pathways normally controlled by group 1 metabotropic glutamate receptor (mGluR1 and mGluR5) activity (50). Accordingly, down regulation of mGluR5 activity, both pharmacologically and genetically, has rescued many phenotypes examined in the animal models of FXS (51). Phase II clinical trials of mGluR5 antagonists in patients with FXS are now ongoing, and it is intriguing to consider whether premutation carriers with significantly lowered FMRP would benefit from these treatments as well.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We are grateful to the research participants and their families; Diana Selmeczy and Jessica Burris for manual tracings of the amygdala; Jose Fon for cerebrum tracings; Alyssa Chavez for data collection and management; Louise Gane for assistance with recruitment and genetic counseling to participants; and Randi Hagerman for team leadership, mentoring, and manuscript review. Funding from the National Institutes of Health grants MH078041 (D.H. and S.M.R.), MH77554 (D.H.), UL1 DE019583 (P.J.H.), RL1AG032119 (P.J.H.) and HD02274 (F.T.) supported this work.

References

1. Sherman SL. Premature ovarian failure in the fragile X syndrome. *Am J Med Genet.* 2000; 97:189–194. [PubMed: 11449487]
2. Allingham-Hawkins DJ, Babul-Hirji R, Chitayat D, Holden JJ, Yang KT, Lee C, et al. Fragile X premutation is a significant risk factor for premature ovarian failure: the International Collaborative POF in Fragile X study--preliminary data. *Am J Med Genet.* 1999; 83:322–325. [PubMed: 10208170]
3. Hagerman PJ, Hagerman RJ. Fragile X-associated tremor/ataxia syndrome--an older face of the fragile X gene. *Nat Clin Pract Neurol.* 2007; 3:107–112. [PubMed: 17279084]
4. Schneider A, Hagerman RJ, Hessl D. Fragile X syndrome -- from genes to cognition. *Dev Disabil Res Rev.* 2009; 15:333–342. [PubMed: 20014363]
5. Fernandez-Carvajal I, Walichiewicz P, Xiaosen X, Pan R, Hagerman PJ, Tassone F. Screening for expanded alleles of the *FMR1* gene in blood spots from newborn males in a Spanish population. *J Mol Diagn.* 2009; 11:324–329. [PubMed: 19460941]

6. Dombrowski C, Levesque S, Morel ML, Rouillard P, Morgan K, Rousseau F. Premutation and intermediate-size FMR1 alleles in 10572 males from the general population: loss of an AGG interruption is a late event in the generation of fragile X syndrome alleles. *Hum Mol Genet.* 2002; 11:371–378. [PubMed: 11854169]
7. Rousseau F, Rouillard P, Morel ML, Khandjian EW, Morgan K. Prevalence of carriers of premutation-size alleles of the FMRI gene--and implications for the population genetics of the fragile X syndrome. *Am J Hum Genet.* 1995; 57:1006–1018. [PubMed: 7485149]
8. Toledano-Alhadeff H, Basel-Vanagaite L, Magal N, Davidov B, Ehrlich S, Drasinover V, et al. Fragile-X carrier screening and the prevalence of premutation and full-mutation carriers in Israel. *Am J Hum Genet.* 2001; 69:351–360. [PubMed: 11443541]
9. Moore CJ, Daly EM, Tassone F, Tysoe C, Schmitz N, Ng V, et al. The effect of pre-mutation of X chromosome CGG trinucleotide repeats on brain anatomy. *Brain.* 2004; 127:2672–2681. [PubMed: 15483045]
10. Aziz M, Stathopulu E, Callias M, Taylor C, Turk J, Oostra B, et al. Clinical features of boys with fragile X premutations and intermediate alleles. *Am J Med Genet.* 2003; 121B:119–127. [PubMed: 12898586]
11. Farzin F, Perry H, Hessl D, Loesch D, Cohen J, Bacalman S, et al. Autism Spectrum Disorders and Attention-Deficit/Hyperactivity Disorder in Boys with the Fragile X Premutation. *J Dev Behav Pediatr.* 2006; 27:S137–S144. [PubMed: 16685180]
12. Goodlin-Jones BL, Tassone F, Gane LW, Hagerman RJ. Autistic spectrum disorder and the fragile X premutation. *J Dev Behav Pediatr.* 2004; 25:392–398. [PubMed: 15613987]
13. Borghgraef, M.; Steyaert, J.; Deroo, S.; Maes, B.; Fryns, JP. Preliminary findings in boys with fragile X premutation: Is there a distinct behavioral phenotype?. *International Fragile X Conference*; Washington, D.C.. 2004.
14. Bourgeois J, Seritan A, Casillas EM, Hessl D, Schneider A, Yang Y, et al. Lifetime prevalence of mood and anxiety disorders in fragile X premutation carriers. *Journal of Clinical Psychiatry.* (In press).
15. Bourgeois JA, Coffey SM, Rivera SM, Hessl D, Gane LW, Tassone F, et al. A review of fragile X premutation disorders: expanding the psychiatric perspective. *J Clin Psychiatry.* 2009; 70:852–862. [PubMed: 19422761]
16. Hessl D, Tassone F, Loesch DZ, Berry-Kravis E, Leehey MA, Gane LW, et al. Abnormal elevation of FMR1 mRNA is associated with psychological symptoms in individuals with the fragile X premutation. *Am J Med Genet B Neuropsychiatr Genet.* 2005; 139:115–121. [PubMed: 16184602]
17. Roberts JE, Bailey DB Jr, Mankowski J, Ford A, Sideris J, Weisenfeld LA, et al. Mood and anxiety disorders in females with the FMR1 premutation. *Am J Med Genet B Neuropsychiatr Genet.* 2009; 150B:130–139. [PubMed: 18553360]
18. Hessl D, Rivera S, Koldewyn K, Cordeiro L, Adams J, Tassone F, et al. Amygdala dysfunction in men with the fragile X premutation. *Brain.* 2007; 130:404–416. [PubMed: 17166860]
19. Wechsler, D. *Manual.* 3. San Antonio: The Psychological Corporation; 1997. Wechsler Adult Intelligence Scale.
20. Lord, CM.; Rutter, M. *Autism Diagnostic Observation Schedule - WPS Edition.* Los Angeles, CA: Western Psychological Services; 1989.
21. Constantino JN, Todd RD. Intergenerational transmission of subthreshold autistic traits in the general population. *Biol Psychiatry.* 2005; 57:655–660. [PubMed: 15780853]
22. Tassone F, Pan R, Amiri K, Taylor AK, Hagerman PJ. A rapid polymerase chain reaction-based screening method for identification of all expanded alleles of the fragile X (FMR1) gene in newborn and high-risk populations. *J Mol Diagn.* 2008; 10:43–49. [PubMed: 18165273]
23. Filipovic-Sadic S, Sah S, Chen L, Krosting J, Sekinger E, Zhang W, et al. A novel FMR1 PCR method for the routine detection of low abundance expanded alleles and full mutations in fragile X syndrome. *Clin Chem.* 56:399–408. [PubMed: 20056738]
24. Tassone F, Hagerman RJ, Taylor AK, Gane LW, Godfrey TE, Hagerman PJ. Elevated levels of FMR1 mRNA in carrier males: a new mechanism of involvement in the fragile-X syndrome. *Am J Hum Genet.* 2000; 66:6–15. [PubMed: 10631132]

25. Iwahashi C, Tassone F, Hagerman RJ, Yasui D, Parrott G, Nguyen D, et al. A quantitative ELISA assay for the fragile x mental retardation 1 protein. *J Mol Diagn.* 2009; 11:281–289. [PubMed: 19460937]
26. Zeng H, Constable RT. Image distortion correction in EPI: comparison of field mapping with point spread function mapping. *Magn Reson Med.* 2002; 48:137–146. [PubMed: 12111941]
27. Robb RA. The biomedical imaging resource at Mayo Clinic. *IEEE Trans Med Imaging.* 2001; 20:854–867. [PubMed: 11585203]
28. Robb RA, Barillot C. Interactive display and analysis of 3-D medical images. *IEEE Trans Med Imaging.* 1989; 8:217–226. [PubMed: 18230519]
29. Robb RA, Hanson DP, Karwoski RA, Larson AG, Workman EL, Stacy MC. Analyze: a comprehensive, operator-interactive software package for multidimensional medical image display and analysis. *Comput Med Imaging Graph.* 1989; 13:433–454. [PubMed: 2688869]
30. Schumann CM, Hamstra J, Goodlin-Jones BL, Lotspeich LJ, Kwon H, Buonocore MH, et al. The amygdala is enlarged in children but not adolescents with autism; the hippocampus is enlarged at all ages. *J Neurosci.* 2004; 24:6392–6401. [PubMed: 15254095]
31. Brown SM, Peet E, Manuck SB, Williamson DE, Dahl RE, Ferrell RE, et al. A regulatory variant of the human tryptophan hydroxylase-2 gene biases amygdala reactivity. *Mol Psychiatry.* 2005; 10:805.
32. Ekman, F.; Friesen, W. *Pictures of Facial Affect.* Palo Alto, CA: Psychologists Press; 1976.
33. Tottenham N, Tanaka JW, Leon AC, McCarry T, Nurse M, Hare TA, et al. The NimStim set of facial expressions: judgments from untrained research participants. *Psychiatry Res.* 2009; 168:242–249. [PubMed: 19564050]
34. Bertolino A, Arciero G, Rubino V, Latorre V, De Candia M, Mazzola V, et al. Variation of human amygdala response during threatening stimuli as a function of 5'HTTLPR genotype and personality style. *Biol Psychiatry.* 2005; 57:1517–1525. [PubMed: 15953488]
35. Hariri AR, Mattay VS, Tessitore A, Kolachana B, Fera F, Goldman D, et al. Serotonin transporter genetic variation and the response of the human amygdala. *Science.* 2002; 297:400–403. [PubMed: 12130784]
36. Friston KJ, Holmes AP, Worsley J-P, Poline CD, Frith CD, Frackowiak RSJ. Statistical parametric maps in functional imaging: A general linear approach. *Human Brain Mapping.* 1995; 2:189–210.
37. Genovese CR, Lazar NA, Nichols T. Thresholding of statistical maps in functional neuroimaging using the false discovery rate. *Neuroimage.* 2002; 15:870–878. [PubMed: 11906227]
38. Duvernoy, HM.; Bourgouin, P. *The Human Brain: Surface, Three-Dimensional sectional Anatomy with MRI, and Blood Supply.* New York: Springer; 1999.
39. Holmes AP, Friston KJ. Generalisability, random effects, and population inference. *Human Brain Mapping.* 1998; 7:S754.
40. Brett M, Anton JL, Valabregue R, Pauline JB. Region of interest analysis using an spm toolbox. *Neuroimage.* 2002; 16:S497.
41. Tzourio-Mazoyer N, Landeau B, Papathanassiou D, Crivello F, Etard O, Delcroix N, et al. Automated anatomical labeling of activations in SPM using a macroscopic anatomical parcellation of the MNI MRI single-subject brain. *Neuroimage.* 2002; 15:273–289. [PubMed: 11771995]
42. Ludwig AL, Hershey JW, Hagerman PJ. Initiation of translation of the FMR1 mRNA Occurs predominantly through 5'-end-dependent ribosomal scanning. *J Mol Biol.* 407:21–34. [PubMed: 21237174]
43. Allen EG, He W, Yadav-Shah M, Sherman SL. A study of the distributional characteristics of FMR1 transcript levels in 238 individuals. *Hum Genet.* 2004; 114:439–447. [PubMed: 14758538]
44. Tassone F, Beilina A, Carosi C, Albertosi S, Bagni C, Li L, et al. Elevated FMR1 mRNA in premutation carriers is due to increased transcription. *RNA.* 2007; 13:555–562. [PubMed: 17283214]
45. Kenneson A, Zhang F, Hagedorn CH, Warren ST. Reduced FMRP and increased FMR1 transcription is proportionally associated with CGG repeat number in intermediate-length and premutation carriers. *Hum Mol Genet.* 2001; 10:1449–1454. [PubMed: 11448936]

46. Tassone F, Hagerman RJ, Taylor AK, Hagerman PJ. A majority of fragile X males with methylated, full mutation alleles have significant levels of FMR1 messenger RNA. *J Med Genet.* 2001; 38:453–456. [PubMed: 11432964]
47. Chen LS, Tassone F, Sahota P, Hagerman PJ. The (CGG)_n repeat element within the 5' untranslated region of the FMR1 message provides both positive and negative cis effects on in vivo translation of a downstream reporter. *Hum Mol Genet.* 2003; 12:3067–3074. [PubMed: 14519687]
48. Peprah E, He W, Allen E, Oliver T, Boyne A, Sherman SL. Examination of FMR1 transcript and protein levels among 74 premutation carriers. *J Hum Genet.* 55:66–68. [PubMed: 19927162]
49. Watson C, Hoeft F, Garrett AS, Hall SS, Reiss AL. Aberrant brain activation during gaze processing in boys with fragile X syndrome. *Arch Gen Psychiatry.* 2008; 65:1315–1323. [PubMed: 18981343]
50. Bear MF. Therapeutic implications of the mGluR theory of fragile X mental retardation. *Genes Brain Behav.* 2005; 4:393–398. [PubMed: 16098137]
51. Dolen G, Osterweil E, Rao BS, Smith GB, Auerbach BD, Chattarji S, et al. Correction of fragile X syndrome in mice. *Neuron.* 2007; 56:955–962. [PubMed: 18093519]

Figure 1. Sample trials of the emotion matching task (33). Participants matched the bottom emotion to either the top left or top right emotion. For the control trials, participants matched the shapes.

Figure 2. Distributions of left and right amygdala volume (corrected for total cerebral volume) in premutation carriers and controls.

Figure 3.

Whole brain within group analysis in the emotion > control condition for controls and premutation carriers. Significant clusters were thresholded on the voxel level at $p < 0.01$ after FDR correction. Both groups significantly activated the bilateral amygdala during this task. Bars at left represent degree of activation using the t-statistic.

Figure 4. Scatterplots showing the association between right (top) and left (bottom) amygdala activation (contrast value extracted from ROI analyses) and FMRP in premutation carriers.

Table 1

Molecular genetic, demographic and clinical descriptive statistics by group.

	Control (n=25)	Premutation (n=23)	Significance
CGG repeats	28.32 (3.56)	106.91 (40.0)	$\underline{P} < 0.001$
Range	19 – 33	55 – 199	
<i>FMR1</i> mRNA	1.46 (0.28)	3.65 (1.71)	$\underline{P} < 0.001$
Range	1.01 – 1.93	2.16 – 8.33	
FMRP [§]	112.81 (95.36)	66.29 (51.95)	$\underline{P} = 0.034$
Range	13.40 – 473.60	14.50 – 182.60	
Handedness (RH)	80.0%	95.5%	ns
Psychoactive medication	33.3%	28.6%	ns
Age	30.12 (7.75)	32.95 (8.89)	ns
Caucasian	68.5%	82.6%	ns
Years of education	15.39 (3.60)	15.47 (2.97)	ns
Full scale IQ	125.52 (17.77)	118.00 (16.05)	ns
SRS total	27.96 (23.75)	39.61 (31.95)	$\underline{P} = 0.16$
ADOS total	2.36 (2.74)	5.87 (6.42)	$\underline{P} = 0.01$

[§]FMRP raw data is shown. Log transformation was used to normalize distributions for t-test of group differences.