Published in final edited form as:

Microbes Infect. 2010 November ; 12(12-13): 978–989. doi:10.1016/j.micinf.2010.06.007.

Topical flagellin protects the injured corneas from *Pseudomonas aeruginosa* infection

Ashok Kumar¹, Nan Gao¹, Theodore J Standiford², Richard L Gallo³, and Fu-Shin X. Yu^{1,*}

¹ Kresge Eye Institute/Departments of Ophthalmology and Anatomy & Cell Biology, Wayne State University School of Medicine, Detroit, MI 48201

² Division of Pulmonary and Critical Care Medicine, Department of Internal Medicine, University of Michigan Medical Center, Ann Arbor, Michigan, USA

³ Division of Dermatology, Department of Medicine, University of California-San Diego, San Diego, California

Abstract

Among bacterial pathogens, *Pseudomonas (P.) aeruginosa* infection is the most sight threatening. The corneal innate immune responses are key mediators of the host's defense to *P. aeruginosa*. Using a mouse model of *Pseudomonas* keratitis, we evaluated the protective effects of topical application of flagellin, a ligand for Toll-Like receptor 5 (TLR5), on the development of *Pseudomonas* keratitis and elucidated the underlying mechanisms. Topical application of purified flagellin 6 and 24 h prior to *P. aeruginosa* inoculation on injured mouse corneas significantly attenuated clinical symptoms of *P. aeruginosa* keratitis, decreased bacterial burden, and suppressed infection induced inflammation in the B6 mouse cornea. Topical application of flagellin on wounded cornea induced PMN infiltration and markedly up-regulated cathelicidin-related antimicrobial peptide (CRAMP) expression. In PMN depleted mice, flagellin promoted bacterial clearance in the cornea compared to that of the PBS treated mice, but was unable to prevent corneal perforation and systemic bacterial dissemination and sepsis. Deletion of CRAMP increased corneal susceptibility to *P. aeruginosa* and abolished flagellin induced protection in B6 mice. Our findings illustrate the profound protective effect of flagellin on the cornea innate defense, a response that can be exploited for prophylactic purposes to prevent contact lens associated *Pseudomonas* keratitis.

Keywords

Innate immunity; Toll-like receptors (TLRs); Flagellin; Antimicrobial Peptides; (AMPs); Cathelicidin; Keratitis

1. Introduction

Bacterial keratitis is a leading cause of visual morbidity and a significant public health problem. *Pseudomonas (P.) aeruginosa* is a major, potentially devastating ocular pathogen

*Corresponding Author: Kresge Eye Institute, Wayne State University School of Medicine, 4717 St. Antoine, Detroit, MI 48201, Tel: (313) 577-1657, Fax: (313) 577-7781, fyu@med.wayne.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

and is the most common cause of ulcerative keratitis associated with contact lens wear [1]. Therapy for such infections must be instituted rapidly following the onset of symptoms to minimize corneal damage [2]. The current treatment regimen includes the frequent administration of topical ocular antibacterial agents. The practice of using multiple antibiotics simultaneously and with frequent dosing may result in corneal toxicity [3,4] and selection of antibiotic resistant bacterial strains [5,6]. Thus, in the light of the number of contact lens wearers, currently 34 million in the United States and 140 million worldwide [7], and an increase in antibiotic-resistant bacteria including *P. aeruginosa* [8], the development of new adjunctive prophylactic/therapeutic modalities for preventing bacterial keratitis is warranted.

In the cornea, bacterial infection occurs only when the epithelial barrier function is compromised [9], for example, due to the chronic mild injury from contact lens wearing [10]. The opportunist pathogens such as *P. aeruginosa* gain access to the cellular layers of the epithelial and/or the stroma and become infectious or pathogenic [11]. As a first line of defense, corneal epithelium possesses the ability to recognize and respond to the presence of pathogens [9]. The studies in recent years have shown that TLR5 is the major pathogen recognizing receptor at the ocular surface [12], similar to that observed in many other mucosal surfaces including airway [13], intestinal [14] and urinary tract [15]. It is now generally believed that the epithelium, via the action of TLRs, is an active participant in mucosal innate defense against microbial infection by producing pro-inflammatory cytokines that recruit PMNs to the infected site, and by secreting defense molecules that directly kill bacteria [16,17]. Indeed, in humans a common stop codon polymorphism in the ligand-binding domain of TLR5 (TLR5392STOP) that abrogates flagellin signaling is associated with susceptibility to pneumonia caused by *Legionella pneumophila* [18,19] and recurrent urinary tract infection [20]. Mice lacking TLR5 are more susceptible to *Escherichia coli* urinary tract infection [21]. Thus, TLR5 plays an important role in the protection of mucosal surfaces against bacterial infection.

Flagellin is the sole ligand of TLR5, compared to other TLR ligands, such as LPS, flagellin is a potent stimulator of epithelial cells, but not of macrophages [19,22]. We recently reported that subconjunctival and intraperitoneal injection of flagellin was highly effective in attenuating the development of *P. aeruginosa* keratitis in C57BL/6 (B6) mice [23]. Similarly, systemic injection of flagellin was found to protect mice from orally administered pathogens (*Salmonella typhimurium* and Rhesus rotavirus) and flagellin induced activation of epithelial cells and the elevation of non-pathologic cytokines were proposed as a mechanism for observed protection and innate defense against infection [24]. Interestingly, systematic application of flagellin also had profound cytoprotective effects on radiation; flagellin protected mice [24,25] as well as monkeys from lethal radiation [24,25]. The underlying mechanisms for flagellin induced cytoprotection remain elusive. Our *in vitro* studies revealed that flagellin pretreatment dampened the pathogen-induced inflammatory response but dramatically induced the expression of antimicrobial peptides (AMPs), specifically LL-37 in corneal epithelial cells [26]. Based on these prior studies we suggested that induced expression of AMPs as a result of flagellin treatment is one of the protective mechanisms.

In this study, we investigated whether topical administration of flagellin can evoke protective effect in B6 mice and can be used as a potential prophylactic agent to prevent keratitis. We also elucidated the role of flagellin-induced expression of Camp gene (murine homolog of human LL-37) in corneal innate immunity using Camp^{-/-} knockout mice. Our results showed that topical application of flagellin on wounded cornea is very effective in preventing the development of keratitis and that antimicrobial peptide CRAMP and neutrophil plays an important role in innate defense of the cornea.

2. Material and Methods

2.1. Bacterial strains and flagellin purification

A cytotoxic strain of *P. aeruginosa* (ATCC 19660) which provides a reproducible inflammatory response in the B6 mouse cornea [23,27] was used in this study. Flagellin was prepared from *P. aeruginosa* strain PAO1 by ammonium sulfate precipitation as described earlier [12]. The amount of LPS in the flagellin preparation was 0.0027 endotoxin unit/ μ g protein, as determined with a quantitative *Limulus* amoebocyte lysate kit. In addition, up to 1 μ g/ml purified flagellin failed to elicit activation of HeLa cells expressing functional TLR4, CD14, and MD2 [28] that were fully activated by 1 ng/ml purified *P. aeruginosa* LPS (Sigma).

2.2. Mice

Wild type C57BL/6 (B6) mice (age, 8 weeks; weight, 20 to 24 g) were purchased from the Jackson Laboratory (Bar Harbor, Me). *Camp*^{-/-} (CRAMP deficient) mice on B6 background [29] were obtained from Dr. R. L. Gallo (University of California San Diego, San Diego, CA). Animals were treated in compliance with the ARVO Statement for the Use of Animals in Ophthalmic and Vision Research. The Institutional Animal Care and Use Committee of Wayne State University approved all animal procedures.

2.3. Flagellin pretreatment and infection procedure

Mice were anesthetized with ketamine/xylazine and placed beneath a stereoscopic microscope for needle scratching with three 1-mm incisions using a 26-gauge needle. Purified flagellin (500 ng in 5 μ l of PBS) or PBS (the control) was topically applied on the needle-injured corneas. At the indicated time (6 or 24 h) post pretreatments, the corneas were scarified again and inoculated with 5 μ l bacterial suspension containing 1×10^4 CFU of *P. aeruginosa* strain ATCC 19660. Eyes were examined daily to monitor the disease progression with a dissection microscope equipped with a digital camera.

2.4. Clinical examination

For the assessment of clinical scores, mice were color coded and examined in a blinded fashion by two independent observers at 1, 3, and 5 days post infection (dpi) to visually grade the disease severity. The ocular disease was graded, and clinical scores were assigned using the following previously [23] described scale: 0, clear; 1, slight opacity partially or fully covering the pupil; 2, slight opacity fully covering the anterior segment; 3, dense opacity partially or fully covering the pupil; and 4, dense opacity covering the anterior segment and corneal perforation. At 5 dpi, all infected corneas were photographed with a microscope to illustrate the disease response.

2.5. Histopathology

For histopathological examination, eyes from euthanized mice were enucleated at 5 dpi and fixed in 4% paraformaldehyde. After paraffin embedding, anterior sections of the corneal epithelium were prepared and stained by hematoxylin and eosin at the pathology research core facility. Sections were analyzed and micrographs prepared.

2.6. Cytokine ELISA

To measure mouse monocyte-inhibitory protein-2 (MIP-2, or CXCL2) and interleukin (IL)-1 β , corneal extracts were prepared by homogenization in 1.0 ml phosphate-buffered saline (PBS) with a glass micro tissue grinder, followed by centrifugation at $14,000 \times g$ for 10 min. Protein concentration of the corneal lysate was determined by Micro BCA™

protein assay kit (Pierce) and an equal amounts of protein was used to perform ELISA according to the manufacturer's instructions (R & D Systems).

2.7. PMN infiltration assay

Measurement of myeloperoxidase (MPO) activity was employed to determine PMN infiltration in the cornea as previously described [23]. The corneas were excised from enucleated eyes at 5 dpi and homogenized in 1 ml of hexadecyltrimethylammonium bromide (HTAB) buffer (0.5% HTAB in 50 mM phosphate buffer, pH 6.0). The samples were then subjected to three freeze-thaw cycles, followed by centrifugation at $16,000 \times g$ for 20 min. Each supernatant was mixed with 50 mM phosphate buffer (pH 6.0) containing 16.7 mg/ml *O,O*-dianisidine hydrochloride and 0.0005% hydrogen peroxide at a 1:30 ratio to obtain a total volume of 3 ml. The change in absorbance at 460 nm was monitored continuously for 5 min. The results were expressed in units of MPO activity/cornea. One unit of MPO activity corresponded to approximately 2.0×10^5 PMN [30].

2.8. Bacterial load in the cornea

Corneas (n=5/group) from *P. aeruginosa*-infected mice were collected, and the numbers of viable bacteria were determined by plate count method. Individual corneas were homogenized in sterile PBS, and aliquots (100 μ l) of serial dilutions were plated onto *Pseudomonas* isolation agar (BD biosciences) plates in triplicate. The plates were incubated overnight at 37°C and bacterial colonies were counted. The results were expressed as the mean number of cfu/cornea $\pm$ standard error of the mean.

2.9. PMN depletion, flagellin treatment and immunohistochemistry

Mice made neutropenic by IP injections of 100–200 μ g of RB6-8C5 mAb in 0.2 ml PBS, 48 h prior to flagellin or *P. aeruginosa* challenge. The control mice received the same dose of rat IgG. To assess the efficacy of the RB6-8C5 antibody, corneas of neutropenic and control mice were wounded by needle scratching and treated by topical application of flagellin or PBS. After 6 h of flagellin treatment eyes were either snapped frozen for staining with antibody (NIMP-R15) specific for mouse PMNs or inoculated with 1×10^4 CFU of *P. aeruginosa*.

2.10. Statistical analysis

An unpaired, two-tailed Student *t* test was used to determine statistical significance for data from bacterial count, cytokine ELISA and MPO assay. A nonparametric Mann-Whitney U test was performed for clinical score. Mean differences were considered significant at a *P* value of <0.05 . Experiments were repeated at least twice to ensure reproducibility, each time with 5 mice in each group.

3. Results

3.1. Topical application of flagellin attenuates clinical symptoms of *P. aeruginosa* keratitis in B6 mouse

We previously showed that TLR5 was expressed in the internal layers of stratified corneal epithelium [12], indicating that the topical application of TLR ligands without injury would not be effective. Indeed, we found topical application of flagellin to the intact, uninjured corneas 6 or 24 h prior to bacterial inoculation had no effects on the progress of corneal disease monitored for up to five day (data not shown), by this time most *P. aeruginosa*-infected corneas were perforated. To allow flagellin to function, we slightly injured corneas with a 26-gauge needle and topically applied solution containing flagellin or PBS (vehicle control) immediately after the injury. To determine whether topical application of flagellin

induces protective mechanisms we applied 500 ng of flagellin or PBS topically (in 5 μ l volume) prior to inoculation of 1.0×10^4 CFU of *P. aeruginosa* (strain ATCC 19660) to needle re-injured corneas. The disease response was graded on various days post infection (dpi) using the previously described scale [23]. As shown in Fig. 1A, a significant difference was observed in the clinical scores of the control vs 6 or 24 h flagellin pretreated mice ($p > 0.05$) at 1 dpi. At 3 dpi, the average clinical scores, 0.2 (6 h pretreatment) and 0.4 (24 h pretreatment) for the flagellin-pretreated mice, were significantly lower than their respective PBS controls with an average clinical score of 1.8 (6 h) and 2.4 (24 h). A similar trend was observed at 5 dpi, where control eyes had a mean clinical score of 3.6 (6 h) and 3.8 (24 h), and the flagellin pretreated were scored 0.2 (6 h) or 0.4 (24 h). Overall, clinical scores of all flagellin treated mice were significantly (ANOVA, $p < 0.001$) lower than the control at 1, 3 and 5 dpi.

Fig.1B shows representing micrographs of infected corneas at 5 dpi. The PBS-pretreated eyes exhibited dense opacity and perforation. In contrast, the eyes pre-exposed to flagellin were mostly clear with no sign of infection while a few eyes were slightly opaque, indicating residual infiltration and inflammation. The infected eyes were also subjected to histopathological analysis (Fig. 1C). In the PBS pretreated eyes, *P. aeruginosa* infection caused heavy infiltrate in the cornea and the anterior chamber along with severe corneal edema, and perforation. On the other hand, flagellin pretreatment preserved the morphological integrity of the cornea with little infiltration detected in the anterior chamber. Thus, flagellin administration prior to bacterial challenge attenuates the development of *P. aeruginosa* keratitis and protects the cornea from damages caused by bacteria -and/or host inflammation.

3.2. Topical flagellin application suppressed infection induced inflammation in the B6 mouse cornea

Having shown a strong protection induced by topical flagellin, we next determined whether flagellin pretreatment reduces the production of inflammatory cytokines and the infiltration of PMNs in the infected cornea. Compared to the mice pretreated with PBS, the levels of IL-1 β in the flagellin pretreated (6 and 24 h) corneas at 5 dpi were dramatically reduced (Fig. 2A). Moreover, the MPO activity was much higher in the PBS pretreated, needle-injured corneas than that of flagellin pretreated ones (Fig. 2B) at 5 dpi. Thus, flagellin pretreatment reduced inflammation in the B6 mouse cornea in response to *P. aeruginosa* infection.

3.3. Flagellin induces cytokines expression and antimicrobial peptide production in the cornea

Having shown that flagellin pretreatment attenuated the development of *P. aeruginosa* keratitis and decreased inflammation in B6 cornea, we next investigated the underlying mechanisms for flagellin induced protection by assessing the effect of topical flagellin on the corneal innate response. First we found that topical application of flagellin or PBS to uninjured corneas did not induce cytokine expression (data not shown). Fig. 3 shows the expression of pro-inflammatory cytokines and the antimicrobial peptide, CRAMP (a 17-kDa precursor protein) after topical application of flagellin on injured mouse corneas. Compare to PBS treatment, topical flagellin induced a rapid and transient expression of MIP-2 and IL-1 β in injured corneas, detectable at 1 h, peaked at 3 h for both cytokines. At 6 h post wounding, the levels of MIP2 declined to a level similar to the baseline while IL-1 β remained relatively high. It should be noted that the levels of IL-1 β and MIP2 (pg/mg tissue extract) in flagellin pretreated corneas were much lower than that of the infected corneas (ng/mg tissue extract, Fig. 2 and Fig. 6). The expression of CRAMP was detected by Western blotting. While wounding caused detectable expression of CRAMP, topical

application of flagellin markedly increased its levels both at 3 and 6 h (Fig. 3B) and, in other experiments, CRAMP levels remained higher than that in PBS treated corneas 24 h post injury (data not shown).

3.4. Flagellin-mediated PMN recruitment plays a key role in protection

Rapid induction of chemokines such as MIP2 suggests recruitment of PMNs to the site of injury and flagellin application. We next assessed if PMNs are involved in enhanced bacterial clearance in flagellin pretreated cornea. We first used immunohistochemistry (IHC) to assess PMN infiltration in flagellin treated B6 mouse corneas. While in corneas without injury, there were no detectable PMNs, injury caused detectable PMN infiltration evidenced by sporadically distributed NIMP-R14 positive cells in the stroma of corneas 6 h post-needle injury. Compared to the PBS treated cornea, flagellin treatment greatly increased the number of PMNs recruited to the cornea in 6 h post injury with more NIMP-R14 positive cells at peripheral region, consistent with the migration of PMNs from the limbal region towards the central cornea. At 24 h, persistent strong PMN staining remained near the site of injury and flagellin application (Fig. 4A). We also performed MPO activity to quantitate PMN recruitment. Clearly, flagellin treated cornea exhibited significantly higher MPO activity compared to normal and injury controls (Fig. 4B). Thus, flagellin pretreatment induced PMN recruitment to the cornea.

To assess whether flagellin-mediated PMNs infiltration also contribute to the observed protection in B6 cornea, we made neutropenic mice by intraperitoneal injection of MAb RB6-8C5. The specificity and efficacy of RB6-8C5 have been well established [31,32] and the approach has been used to study the role of neutrophils in keratitis by several investigators [33–36]. To confirm that PMN were depleted, we used flagellin induced PMN recruitment to the cornea as a parameter and found that while abundant PMNs were detected in the control and flagellin treated corneas, no positively NIMP-R14 stained cells were observed in the corneas of RB6-8C5 treated mice challenged with or without topical flagellin, indicating depletion of PMNs from mice (Supplement Fig.1). PMN depletion was also confirmed by PMNs blood smear staining with Wright Stain (data not shown).

To determine the protective role of flagellin-mediated PMN infiltration, corneas of both normal and RB6-8C5-treated mice were pretreated with flagellin or PBS for 6 h and then infected with *P. aeruginosa*. At 1 dpi *P. aeruginosa* infection caused opacity (clinical score 1) in normal PBS pretreated mice, whereas flagellin pretreated mice showed no sign of inflammation or infection (Figure 5A, upper panels). In the corneas of RB6-8C5-treated mice, there was less opacity at 1 dpi compared to that of PBS treated normal mice and no difference in appearance of the corneas pretreated with flagellin or PBS (Fig. 5A, lower panels). Although the corneas of RB6-8C5-treated mice appeared clear, they showed sign of meting and perforation, especially at 2 d.p.i. Most RB6-8C5-treated mice died overnight between 1 and 2 dpi. The experiment was therefore terminated and the bacterial loadings in the kidney, liver, spleen, and corneas were determined. While no bacteria were detected in the kidney, liver, spleen of normal mice, heavy bacterial loads were detected in these tissues of RB6-8C5-treated mice (data not shown), suggesting severe sepsis caused by the bacterial spread from the infected cornea in these mice. In the cornea, the bacterial load was much higher in RB6-8C5-treated mice than that in non-PMN depleted corneas and, more importantly, flagellin pretreatment significantly reduced the bacterial load in both neutropenic and normal mice when compared to PBS pretreated eyes (Fig. 5B).

3.5. Flagellin pretreatment does not protect the *Camp*^{-/-} mice from *P. aeruginosa*

The observation that topical flagellin treatment induced the expression of CRAMP in B6 cornea suggests CRAMP may contribute to the enhanced innate immunity in the cornea.

Previous work by Huang et al [37] showed that cathelicidin-knockout markedly increased the susceptibility of a *P. aeruginosa* resistant strain, 129/SVJ, to *P. aeruginosa* keratitis. To test the role of CRAMP in corneal innate immunity, we used Camp^{-/-} knockout mice with B6 background [29]. We first investigated the effects of CRAMP deficiency on the susceptibility of B6 mice by assessing the number of bacteria required to cause corneal infection in anesthetized B6 mice. For the wild type B6 mice, 10⁴ bacteria were required to cause 90% infection while 10³ bacteria were sufficient to cause 100% infection in needle-injured corneas of Camp^{-/-} mice. When 10⁴ bacteria were used, the disease progressed much faster in Camp^{-/-} knockout mice (Fig. 6A). At 1 dpi the corneas of PBS pretreated WT mice showed mild opacity and anterior chamber inflammatory reaction (mean clinical score = 0.9), whereas the corneas of Camp^{-/-} exhibited severe inflammatory response and dense opacity (mean clinical score = 2.0). Most corneas were perforated at 3 dpi for of Camp^{-/-} and 5 dpi for the wild type. We next investigated the role of CRAMP in flagellin-induced protection against *P. aeruginosa* keratitis using the knockout mice. At 1 dpi the disease severity was less in both WT (mean clinical score = 0.2) and Camp^{-/-} (mean clinical score = 0.6) mice pretreated with flagellin compared to their respective PBS controls. However, no significant difference was observed in the mean clinical scores of Camp^{-/-} mice pretreated with flagellin or PBS at 3 & 5 dpi (3.6 vs 3.8.0 at 3 dpi and 3.8 vs 4.0 at 5dpi) (Fig. 6A and 6B).

The effects of flagellin pretreatment on bacterial load, inflammatory cytokine and PMN infiltration in WT and Camp^{-/-} mice at 5 dpi were also investigated (Fig. 6C–E). The Camp^{-/-} mice showed significant higher (17 fold) bacterial load compared to WT at 5 dpi. The effects of flagellin pretreatment on bacterial clearance in Camp^{-/-} mice were greatly reduced compared to that in wild type B6 mice. Although Flagellin pretreatment resulted in reduced bacterial loads in Camp^{-/-} mice, this difference did not reach statistical significance ($P > 0.5$, Student's *t*-test) (Fig. 6C). A similar trend was observed for cytokines expression and PMN infiltration in the infected WT and Camp^{-/-} mice. Flagellin pretreatment dramatically reduced the expression of MIP-2 and IL-1 β in WT mice compared to their PBS controls, whereas no significant difference was observed in Camp^{-/-} mice with or without flagellin pretreatment (Fig. 6D). Finally, flagellin pretreatment significantly reduced the MPO activity in the corneal homogenate of *P. aeruginosa* infected WT mice but not in Camp^{-/-} mice and MPO levels in the corneas of Camp^{-/-} mice were much higher than that of the wild type with or without flagellin pretreatment (Fig. 6E).

4. Discussion

In the present report, we used topical application of flagellin as a stimulus of corneal innate immunity and demonstrated that the treatment induced rapid and transient production of cytokine and chemokines, greatly increased the expression of antimicrobial peptide CRAMP, and enhanced infiltration of PMN to the injured cornea. Importantly, topical application of flagellin protected the injured corneas of from *P. aeruginosa* infection. This is clinical relevance since chronic mild injury is thought to be a prerequisite for contact lens wearing associated microbial keratitis. Moreover, using antibody mediated neutrophil depletion; we showed that PMNs is a key player in cornea innate immunity by clearing bacteria at the site of infection and by limiting the systemic dissemination of the pathogens from the eye. Finally, using Camp^{-/-} mice, we demonstrated that cathelicidin is important for cornea defense and bacterial clearance, confirming a previous study reporting CRAMP as a determinant factor for corneal susceptibility to PA keratitis [38]. Taken together, our studies suggest that strong innate defense mechanisms exist in the cornea and these mechanisms can be induced readily when there is a need to prevent sight threatening corneal infections.

In our previous study, we administrated flagellin through subconjunctival and intraperitoneal injection of a total of 225 ng flagellin per mouse 24 h prior to bacterial infection and found that pre-exposure of the corneas to flagellin through these routes was effective in preventing *P. aeruginosa* keratitis [23]. Although that approach was likely to induce maximal protection, a necessary first step for the study, subconjunctival and intraperitoneal injection in particular is not amendable for clinical applications. In this study, we demonstrated that one time topical application of flagellin was sufficient to protect the mouse corneas from *P. aeruginosa* keratitis. More recently, we observed that this treatment was also effective in preventing fungal infection of the cornea although TLR5 is not a fungal recognizing receptor (Gao and Yu, unpublished results). We showed that topical application of flagellin to intact cornea had no detectable effects on cytokine and CRAMP expression, suggesting it is safe to use flagellin for short time durations such as eye drop for contact lens wearers. In such cases, if the epithelial barrier is intact, flagellin should have minimal effects on the cornea. On the other hand, if barrier is disrupted, flagellin may trigger activation of corneal innate immunity, resulting in greatly enhanced bacterial clearance and hence preventing potential infection from occurring. Taken together, our results suggest that incorporation of low dosage of flagellin in a contact lens solution or eye drops may be used prophylactically for contact lens wearer or as a post-ocular surgery remedy to reduce the incidence of bacterial keratitis. Further studies to test the safety and efficacy of repeated topical treatments are warranted.

Among different TLR ligands, flagellin is very unique in inducing protection mechanisms against infection, radiation and several other adverse environmental challenges [24,39]. To date, the mechanism by which flagellin activates cyto- and immuno-protection responses has not been fully understood [19]. In the cornea, the full protection was inducible by flagellin in 6 hrs post administration, suggesting primarily the stimulation of innate immunity. It is interesting to note that at this time point, the levels of MIP2 and IL-1 β were still somewhat elevated and infection might exacerbate inflammatory response in the cornea. However, our results showed significant decrease in the severity of the diseases as early as 1 dpi (Fig. 1A). This may be due to flagellin induced tolerance (hyporesponsiveness) of corneal epithelial cells to bacterial challenge observed as early as 4 h *in vitro* [26]. At the cytokine levels, we showed that flagellin induced rapid and transient production of chemokine MIP2, a human IL-8 homolog and a potent PMN chemoattractant, resulting in the influx of PMN towards the site of injury and flagellin application. Interestingly, while the expression of MIP2 subsided rapidly, the infiltrated PMN remained in the cornea. Using antibody-mediated PMN depletion we showed that PMN is critical for corneal innate defense against *P. aeruginosa* infection and the lack of them resulted in greatly increased bacterial burden in the cornea and dissemination to other organs such as the kidney, liver and lungs. In another study using a lethal murine *P. aeruginosa* pneumonia model, we showed that the effects of flagellin were primarily mediated by TLR5, as evidenced by a substantial loss of flagellin-mediated protection in TLR5^{-/-} mice [40]. Hence, we propose that topically applied flagellin activates TLR5 and triggers controlled innate immune responses including infiltration of neutrophils and production of antimicrobial peptides in the cornea, combination of which form a biological shield at the ocular surface against invading pathogens.

The present study revealed that topical flagellin is capable to significantly decrease bacterial loading, suppress pro-inflammatory cytokine such as IL-1 β expression, and reduce PMN infiltration at 5 dpi, resulting in the preservation of corneal function in B6 mice. We showed the suppression of IL-1 β in flagellin pretreated corneas since persistent production of the cytokine is known detrimental to the corneas and inhibition of IL-1 β production by Caspase-1 inhibitor reduced severity of *Pseudomonas aeruginosa* keratitis in mice [41]. The low levels of pro-cytokines and infiltrated cells in flagellin pretreated corneas at 5 dpi are

likely the result of much reduced bacterial burden (Figure 6) and flagellin induced abundant expression of AMPs such as CRAMP is likely to play a role in bacterial clearance in the cornea. The cathelicidin have been detected in many different cell types and organs, including innate immune cells such as neutrophils [42], monocytes [43], mast cells [44], and epithelial cells [42]. An increasing number of studies have established a clear link between TLR activation and the expression of AMPs [45]. Gordon et al first reported LL-37 expression in ocular surface epithelia and demonstrated its antimicrobial activity against ocular pathogens [46]. We also showed that Pam3Cys [47] and natural SA lipopeptide [48] treatment induced human β -defensin-2 and LL-37 expression in human corneal epithelial cells *in-vitro*. In this study, we provided evidence that cathelicidin gene products are a determinant factor for the corneal innate defense and susceptibility to microbial infection. Consistent with our *in vitro* studies showing that flagellin pretreatment dampened the production of proinflammatory cytokines and yet enhanced the expression of antimicrobial peptides [26], we showed in the present study that while cytokine expression is transient, the expression of CRAMP in flagellin treated corneas is persistent at least for 24 h. Hence, the flagellin-induced reprogramming also occurs at the ocular surface *in vivo*.

Cathelicidin exhibits antimicrobial activity against a variety of bacterial strains, including *Escherichia coli*, *P. aeruginosa*, *Klebsiella pneumoniae*, and *Staphylococcus aureus* [49,50]. Its bactericidal activity involves membrane disruption (pore formation, change of lipid packing and organization) following interaction with negatively charged bacterial molecules and insertion into the membrane. Additionally, at a very low concentration (0.5 μ g/ml), LL-37 inhibits the formation of *P. aeruginosa* biofilms by decreasing the attachment of bacterial cells and stimulating their twitching motility [51]. LL-37 was shown to be effective in killing *P. aeruginosa* in tears [52]. Hence, the presence of abundant levels of CRAMP at the ocular surface, alone or acting in synergy with other AMPs such as β -defensins [53] and surfactant D [54], may result in rapid bacterial clearance. Using CRAMP deficient mice, our study revealed that up-regulation of cathelicidin gene is important for flagellin induced protection in the corneas. Overall, flagellin was unable to induce protection against *P. aeruginosa* in the cornea of Camp^{-/-} mice. Similar results were also observed in the lung study where flagellin-induced protective effects on bacterial clearance and survival were significantly diminished in Camp^{-/-} mice relative to wild-type mice [40]. Interestingly, although flagellin pretreatment does not protect Camp^{-/-} mice effectively compared to WT mice, we observed a significant difference in clinical scores of PBS vs flagellin pretreated Camp^{-/-} mice at 1 dpi (Fig. 6A), suggesting that in addition to CRAMP, flagellin induce other protective mechanism(s) including other AMPs such as defensins in early stage on infection. However, the lack of sufficient bacterial killing in flagellin pretreated Camp^{-/-} mice resulted in a rapid increase in bacterial burden that overwhelms the innate immune system, resulting in a rapid progression of infection and total destruction of the cornea. Compared to Camp^{-/-} mice, the low levels of pro-cytokines and infiltrated cells in flagellin pretreated corneas of wild type mice observed at the later stage of infection-5 dpi-were likely the result of much reduced bacterial burden. Hence, our results indicate that CRAMP is a dominant mediator of immune protection induced by flagellin in response to infection.

In addition to its antimicrobial activity, human cathelicidin LL-37 was also shown to exhibit immunomodulatory effects, such as inhibition of pro-inflammatory responses to bacterial LPS in human monocytic cells [55] and promotes epithelial wound healing [56]. It is possible that the presence of abundant CRAMP may promote epithelial wound healing and modulate the inflammatory response of infiltrated cells in *P. aeruginosa* infected corneas, thus reducing tissue destruction resulted from inflammatory cells. Indeed, we recently showed that Topical flagellin blocked *C. albicans*-induced epithelial apoptosis in the B6

mouse cornea, suggesting strong cytoprotective effects induced by topical flagellin in the cornea (Yu and Gao, unpublished results).

Several studies have shown that flagellin induces dendritic cell (DC) activation, eventually leading to CD4 T cell activation [57,58]. In our case, topical flagellin on injured cornea may induce the activation and maturation of residential DCs [59]; which in turn expands and activates innate lymphocytes (natural killer and gammadelta T cells) and initiates adaptive T cell immunity [60]. Gammadelta T cells [61] and natural killer cells [62,63] have recently been shown to play an important role in corneal homeostasis and defense and are likely to contribute to flagellin mediated protection. Further study to define the involvement of adaptive immunity in flagellin induced protection is warranted.

In summary, our results showed that strong innate defense mechanisms exist in the cornea and flagellin is a unique stimulator of corneal innate immunity. Activation of innate immunity prior to infection is an effective way to prevent infection from occurring in vulnerable organs such as the cornea. Thus, flagellin can be exploited as a prophylactic modality to prevent and/or treat bacterial infections of the cornea and other tissues such as the respiratory and gastrointestinal tracts.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by in part by NIH Grants EY017960 and EY010869 (F.S.Y.), Fight for Sight grant-in-aid (A.K.) and Midwest Eye-Banks (A.K.) and an unrestricted Grant from the Research to Prevent Blindness to the Department of Ophthalmology, Wayne State University School of Medicine.

References

1. Hazlett LD. Bacterial infections of the cornea (*Pseudomonas aeruginosa*). *Chem Immunol Allergy*. 2007; 92:185–194. [PubMed: 17264494]
2. Fleiszig SM, Evans DJ. The pathogenesis of bacterial keratitis: studies with *Pseudomonas aeruginosa*. *Clin Exp Optom*. 2002; 85:271–278. [PubMed: 12366347]
3. Thompson AM. Ocular toxicity of fluoroquinolones. *Clin Experiment Ophthalmol*. 2007; 35:566–577. [PubMed: 17760640]
4. Smith A, Pennefather PM, Kaye SB, Hart CA. Fluoroquinolones: place in ocular therapy. *Drugs*. 2001; 61:747–761. [PubMed: 11398907]
5. Moshirfar M, Mirzaian G, Feiz V, Kang PC. Fourth-generation fluoroquinolone-resistant bacterial keratitis after refractive surgery. *J Cataract Refract Surg*. 2006; 32:515–518. [PubMed: 16631067]
6. Goldstein MH, Kowalski RP, Gordon YJ. Emerging fluoroquinolone resistance in bacterial keratitis: a 5-year review. *Ophthalmology*. 1999; 106:1313–1318. [PubMed: 10406613]
7. Stapleton F, Keay L, Edwards K, Naduvilath T, Brian G, Jacobs R. Studies of contact lens-related microbial keratitis in Australia and New Zealand: new learnings. *Eye Contact Lens*. 2007; 33:354–357. discussion 362–353. [PubMed: 17975419]
8. Jhanji V, Sharma N, Satpathy G, Titiyal J. Fourth-generation fluoroquinolone-resistant bacterial keratitis. *J Cataract Refract Surg*. 2007; 33:1488–1489. [PubMed: 17662450]
9. Kumar A, Yu FS. Toll-like receptors and corneal innate immunity. *Curr Mol Med*. 2006; 6:327–337. [PubMed: 16712478]
10. Fleiszig SM, Evans DJ. Contact lens infections: can they ever be eradicated? *Eye Contact Lens*. 2003; 29:S67–71. discussion S83–64, S192–194. [PubMed: 12772735]
11. Willcox MD. *Pseudomonas aeruginosa* infection and inflammation during contact lens wear: a review. *Optom Vis Sci*. 2007; 84:273–278. [PubMed: 17435510]

12. Zhang J, Xu K, Ambati B, Yu FS. Toll-like receptor 5-mediated corneal epithelial inflammatory responses to *Pseudomonas aeruginosa* flagellin. *Invest Ophthalmol Vis Sci*. 2003; 44:4247–4254. [PubMed: 14507868]
13. Zhang Z, Louboutin J-P, Weiner DJ, Goldberg JB, Wilson JM. Human Airway Epithelial Cells Sense *Pseudomonas aeruginosa* Infection via Recognition of Flagellin by Toll-Like Receptor 5. *Infect Immun*. 2005; 73:7151–7160. [PubMed: 16239509]
14. Vijay-Kumar M, Sanders CJ, Taylor RT, Kumar A, Aitken JD, Sitaraman SV, Neish AS, Uematsu S, Akira S, Williams IR, Gewirtz AT. Deletion of TLR5 results in spontaneous colitis in mice. *J Clin Invest*. 2007; 117:3909–3921. [PubMed: 18008007]
15. Andersen-Nissen E, Hawn TR, Smith KD, Nachman A, Lampano AE, Uematsu S, Akira S, Aderem A. Cutting Edge: Tlr5^{-/-} Mice Are More Susceptible to *Escherichia coli* Urinary Tract Infection. *J Immunol*. 2007; 178:4717–4720. [PubMed: 17404249]
16. Yu FS, Hazlett LD. Toll-like Receptors and the Eye. *Invest Ophthalmol Vis Sci*. 2006; 47:1255–1263. [PubMed: 16565355]
17. Johnson A, Pearlman E. Toll-like Receptors in the Cornea. *Ocul Surf*. 2005; 3:S187–189. [PubMed: 17216116]
18. Hawn TR, Berrington WR, Smith IA, Uematsu S, Akira S, Aderem A, Smith KD, Skerrett SJ. Altered inflammatory responses in TLR5-deficient mice infected with *Legionella pneumophila*. *J Immunol*. 2007; 179:6981–6987. [PubMed: 17982089]
19. Vijay-Kumar M, Gewirtz AT. Flagellin: key target of mucosal innate immunity. *Mucosal Immunol*. 2009; 2:197–205. [PubMed: 19242410]
20. Hawn TR, Scholes D, Li SS, Wang H, Yang Y, Roberts PL, Stapleton AE, Janer M, Aderem A, Stamm WE, Zhao LP, Hooton TM. Toll-like receptor polymorphisms and susceptibility to urinary tract infections in adult women. *PLoS One*. 2009; 4:e5990. [PubMed: 19543401]
21. Andersen-Nissen E, Hawn TR, Smith KD, Nachman A, Lampano AE, Uematsu S, Akira S, Aderem A. Cutting Edge: Tlr5^{-/-} Mice Are More Susceptible to *Escherichia coli* Urinary Tract Infection. *J Immunol*. 2007; 178:4717–4720. [PubMed: 17404249]
22. Franchi L, Amer A, Body-Malapel M, Kanneganti TD, Ozoren N, Jagirdar R, Inohara N, Vandenabeele P, Bertin J, Coyle A, Grant EP, Nunez G. Cytosolic flagellin requires Ipaf for activation of caspase-1 and interleukin 1beta in salmonella-infected macrophages. *Nat Immunol*. 2006; 7:576–582. [PubMed: 16648852]
23. Kumar A, Hazlett LD, Yu FS. Flagellin suppresses the inflammatory response and enhances bacterial clearance in a murine model of *Pseudomonas aeruginosa* keratitis. *Infect Immun*. 2008; 76:89–96. [PubMed: 17938214]
24. Vijay-Kumar M, Aitken JD, Sanders CJ, Frias A, Sloane VM, Xu J, Neish AS, Rojas M, Gewirtz AT. Flagellin treatment protects against chemicals, bacteria, viruses, and radiation. *J Immunol*. 2008; 180:8280–8285. [PubMed: 18523294]
25. Burdelya LG, Krivokrysenko VI, Tallant TC, Strom E, Gleiberman AS, Gupta D, Kurnasov OV, Fort FL, Osterman AL, DiDonato JA, Feinstein E, Gudkov AV. An Agonist of Toll-Like Receptor 5 Has Radioprotective Activity in Mouse and Primate Models. *Science*. 2008; 320:226–230. [PubMed: 18403709]
26. Kumar A, Yin J, Zhang J, Yu FS. Modulation of corneal epithelial innate immune response to *pseudomonas* infection by flagellin pretreatment. *Invest Ophthalmol Vis Sci*. 2007; 48:4664–4670. [PubMed: 17898290]
27. Hazlett LD, McClellan S, Kwon B, Barrett R. Increased severity of *Pseudomonas aeruginosa* corneal infection in strains of mice designated as Th1 versus Th2 responsive. *Invest Ophthalmol Vis Sci*. 2000; 41:805–810. [PubMed: 10711697]
28. Re F, Strominger JL. Monomeric recombinant MD-2 binds toll-like receptor 4 tightly and confers lipopolysaccharide responsiveness. *J Biol Chem*. 2002; 277:23427–23432. [PubMed: 11976338]
29. Buchau AS, Morizane S, Trowbridge J, Schaubert J, Kotol P, Bui JD, Gallo RL. The Host Defense Peptide Cathelicidin Is Required for NK Cell-Mediated Suppression of Tumor Growth. *J Immunol*. 2009 jimmunol.0902110.

30. Williams RN, Paterson CA, Eakins KE, Bhattacharjee P. Quantification of ocular inflammation: evaluation of polymorphonuclear leucocyte infiltration by measuring myeloperoxidase activity. *Curr Eye Res.* 1982; 2:465–470. [PubMed: 6303695]
31. Leendertse M, Willems RJ, Giebelen IA, Roelofs JJ, Bonten MJ, van der Poll T. Neutrophils are essential for rapid clearance of *Enterococcus faecium* in mice. *Infect Immun.* 2009; 77:485–491. [PubMed: 19001080]
32. Tvinnereim AR, Hamilton SE, Harty JT. Neutrophil involvement in cross-priming CD8+ T cell responses to bacterial antigens. *J Immunol.* 2004; 173:1994–2002. [PubMed: 15265934]
33. Zaidi TS, Zaidi T, Pier GB. Role of neutrophils, MyD88-mediated neutrophil recruitment and complement in antibody-mediated defense against *Pseudomonas aeruginosa* keratitis. *Invest Ophthalmol Vis Sci.* 2009
34. Brown CR, Blaho VA, Loiacono CM. Treatment of mice with the neutrophil-depleting antibody RB6-8C5 results in early development of experimental lyme arthritis via the recruitment of Gr-1 polymorphonuclear leukocyte-like cells. *Infect Immun.* 2004; 72:4956–4965. [PubMed: 15321987]
35. Khatri S, Lass JH, Heinzel FP, Petroll WM, Gomez J, Diaconu E, Kalsow CM, Pearlman E. Regulation of endotoxin-induced keratitis by PECAM-1, MIP-2, and toll-like receptor 4. *Invest Ophthalmol Vis Sci.* 2002; 43:2278–2284. [PubMed: 12091428]
36. Zhou Z, Barrett RP, McClellan SA, Zhang Y, Szliter EA, van Rooijen N, Hazlett LD. Substance P delays apoptosis, enhancing keratitis after *Pseudomonas aeruginosa* infection. *Invest Ophthalmol Vis Sci.* 2008; 49:4458–4467. [PubMed: 18566468]
37. Huang LC, Reins RY, Gallo RL, McDermott AM. Cathelicidin-Deficient (Cnlp /) Mice Show Increased Susceptibility to *Pseudomonas aeruginosa* Keratitis. *Invest Ophthalmol Vis Sci.* 2007; 48:4498–4508. [PubMed: 17898271]
38. Huang LC, Reins RY, Gallo RL, McDermott AM. Cathelicidin-deficient (Cnlp $-/-$) mice show increased susceptibility to *Pseudomonas aeruginosa* keratitis. *Invest Ophthalmol Vis Sci.* 2007; 48:4498–4508. [PubMed: 17898271]
39. Burdelya LG, Krivokrysenko VI, Tallant TC, Strom E, Gleiberman AS, Gupta D, Kurnasov OV, Fort FL, Osterman AL, Didonato JA, Feinstein E, Gudkov AV. An agonist of toll-like receptor 5 has radioprotective activity in mouse and primate models. *Science.* 2008; 320:226–230. [PubMed: 18403709]
40. Yu FS, Cornicelli MD, Kovach MA, Newstead MW, Zeng X, Kumar A, Gao N, Yoon SG, Gallo RL, S TJ. Flagellin Stimulates Protective Lung Mucosal Immunity: Role of Cathelicidin Related Antimicrobial Peptide (CRAMP). *Journal of Immunology.* 2010 in press.
41. Thakur A, Barrett RP, Hobden JA, Hazlett LD. Caspase-1 inhibitor reduces severity of *pseudomonas aeruginosa* keratitis in mice. *Invest Ophthalmol Vis Sci.* 2004; 45:3177–3184. [PubMed: 15326138]
42. Zanetti M. Cathelicidins, multifunctional peptides of the innate immunity. *J Leukoc Biol.* 2004; 75:39–48. [PubMed: 12960280]
43. Agerberth B, Charo J, Werr J, Olsson B, Idali F, Lindbom L, Kiessling R, Jornvall H, Wigzell H, Gudmundsson GH. The human antimicrobial and chemotactic peptides LL-37 and alpha - defensins are expressed by specific lymphocyte and monocyte populations. *Blood.* 2000; 96:3086–3093. [PubMed: 11049988]
44. Di Nardo A, Yamasaki K, Dorschner RA, Lai Y, Gallo RL. Mast Cell Cathelicidin Antimicrobial Peptide Prevents Invasive Group A Streptococcus Infection of the Skin. *J Immunol.* 2008; 180:7565–7573. [PubMed: 18490758]
45. Kumar A, Christopher Singh N, Inna Glybina V, Tamer Mahmoud H, Shin F, Yu X. Toll-like Receptor 2 Ligand-Induced Protection against Bacterial Endophthalmitis. *The Journal of Infectious Diseases.* 201:255–263. [PubMed: 19995266]
46. Gordon YJ, Huang LC, Romanowski EG, Yates KA, Proske RJ, McDermott AM. Human cathelicidin (LL-37), a multifunctional peptide, is expressed by ocular surface epithelia and has potent antibacterial and antiviral activity. *Curr Eye Res.* 2005; 30:385–394. [PubMed: 16020269]
47. Kumar A, Zhang J, Yu F-SX. Toll-like receptor 2-mediated expression of [beta]-defensin-2 in human corneal epithelial cells. *Microbes and Infection.* 2006; 8:380–389. [PubMed: 16242370]

48. Li Q, Kumar A, Gui J-F, Yu F-SX. Staphylococcus aureus lipoproteins trigger human corneal epithelial innate response through toll-like receptor-2. *Microbial Pathogenesis*. 2008; 44:426–434. [PubMed: 18191935]
49. Smeianov V, Scott K, Reid G. Activity of cecropin P1 and FA-LL-37 against urogenital microflora. *Microbes and Infection*. 2000; 2:773–777. [PubMed: 10955957]
50. Travis SM, Anderson NN, Forsyth WR, Espiritu C, Conway BD, Greenberg EP, McCray PB Jr, Lehrer RI, Welsh MJ, Tack BF. Bactericidal Activity of Mammalian Cathelicidin-Derived Peptides. *Infect Immun*. 2000; 68:2748–2755. [PubMed: 10768969]
51. Overhage J, Campisano A, Bains M, Torfs ECW, Rehm BHA, Hancock REW. Human Host Defense Peptide LL-37 Prevents Bacterial Biofilm Formation. *Infect Immun*. 2008; 76:4176–4182. [PubMed: 18591225]
52. Huang LC, Petkova TD, Reins RY, Proske RJ, McDermott AM. Multifunctional roles of human cathelicidin (LL-37) at the ocular surface. *Invest Ophthalmol Vis Sci*. 2006; 47:2369–2380. [PubMed: 16723446]
53. Wu M, McClellan SA, Barrett RP, Zhang Y, Hazlett LD. {beta}-Defensins 2 and 3 Together Promote Resistance to *Pseudomonas aeruginosa* Keratitis. *J Immunol*. 2009; 183:8054–8060. [PubMed: 19933858]
54. Ni M, Tam C, Verma A, Ramphal R, Hawgood S, Evans DJ, Fleiszig SMJ. Expression of surfactant protein D in human corneal epithelial cells is upregulated by *Pseudomonas aeruginosa*. *FEMS Immunology & Medical Microbiology*. 2008; 54:177–184. [PubMed: 18657106]
55. Molhoek EM, den Hertog AL, de Vries AM, Nazmi K, Veerman EC, Hartgers FC, Yazdanbakhsh M, Bikker FJ, van der Kleij D. Structure-function relationship of the human antimicrobial peptide LL-37 and LL-37 fragments in the modulation of TLR responses. *Biol Chem*. 2009; 390:295–303. [PubMed: 19166322]
56. Yin J, Yu FS. LL-37 Promotes High Glucose-Attenuated Epithelial Wound Healing via EGFR Transactivation in Organ Cultured Corneas. *Invest Ophthalmol Vis Sci*. 2009
57. Gewirtz AT. Flag in the crossroads: flagellin modulates innate and adaptive immunity. *Curr Opin Gastroenterol*. 2006; 22:8–12. [PubMed: 16319670]
58. Salazar-Gonzalez RM, McSorley SJ. Salmonella flagellin, a microbial target of the innate and adaptive immune system. *Immunology Letters*. 2005; 101:117–122. [PubMed: 15975666]
59. Hamrah P, Dana MR. Corneal antigen-presenting cells. *Chem Immunol Allergy*. 2007; 92:58–70. [PubMed: 17264483]
60. Munz C, Steinman RM, Fujii S. Dendritic cell maturation by innate lymphocytes: coordinated stimulation of innate and adaptive immunity. *J Exp Med*. 2005; 202:203–207. [PubMed: 16027234]
61. O'Brien RL, Taylor MA, Hartley J, Nuhsbaum T, Dugan S, Lahmers K, Aydinoglu MK, Wands JM, Roark CL, Born WK. Protective Role of {gamma}{delta} T Cells in Spontaneous Ocular Inflammation. *Invest Ophthalmol Vis Sci*. 2009; 50:3266–3274. [PubMed: 19151391]
62. Hazlett LD, McClellan SA, Barrett RP, Liu J, Zhang Y, Lighvani S. Spantide I decreases type I cytokines, enhances IL-10, and reduces corneal perforation in susceptible mice after *Pseudomonas aeruginosa* infection. *Invest Ophthalmol Vis Sci*. 2007; 48:797–807. [PubMed: 17251480]
63. Lighvani S, Huang X, Trivedi PP, Swanborg RH, Hazlett LD. Substance P regulates natural killer cell interferon-gamma production and resistance to *Pseudomonas aeruginosa* infection. *Eur J Immunol*. 2005; 35:1567–1575. [PubMed: 15832292]

Fig. 1. Topical application of flagellin is effective in inducing protection against *Pseudomonas* keratitis

B6 mice were pretreated with topical application of flagellin (500ng) or PBS, After 6 or 24h pretreatment the cornea was scarified and challenged topically with 1.0×10^4 CFU of PA.

(A) Clinical scores for the ocular disease response were assigned on 1, 3, and 5 dpi and represented as mean clinical score (B) microphotographs of infected corneas were taken at 5 dpi and (C) Histopathological analysis performed on representative infected corneas at 5 dpi. A nonparametric Mann-Whitney U test was performed to compare each flagellin pretreatment to the PBS group (* $p < 0.05$, ** $p < 0.01$).

Fig. 2. Flagellin pretreatment dampens cytokine production and PMN infiltration in B6 mouse cornea

B6 mouse corneas were pretreated with topically applied flagellin or PBS for 6 or 24h as described in figure 1, followed by PA inoculation. The eyes were enucleated at 5 dpi for IL-1β ELISA (A) and MPO determination (B). Experiments were repeated twice and indicated *p* values were generated using paired Student's *t* test.

Fig. 3. Flagellin pretreatment induces cytokine and CRAMP production in the wounded cornea
 B6 mouse corneas were either needle wounded and topically challenged with flagellin (Flag) or PBS. At the indicated time points, the corneas (n=3) were excised and subjected for ELISA measurement of MIP-2 or IL-1 β (A) or Western blot analysis of 17-kDa (CRAMP) (B). The band intensity of western blot was quantitated by densitometric analysis and normalized with ERK, used as a protein loading control (B, low panel). Indicated *p* values were generated using paired Student's *t* test.

Fig. 4. Flagellin pretreatment induces the recruitment of PMNs to the cornea

B6 mouse corneas, normal or needle injured, were challenged topically with flagellin or PBS, either OCT snap frozen, followed by sectioning and immunostaining with antibody NIMP-R14 to detect PMNs (A), or homogenized for MPO assay to assess PMN infiltration (B). * $P < 0.05$ (student's t -test), significant increase in PMN infiltration was detected in flagellin treated compared to both normal and injured corneas.

Fig. 5. Flagellin pretreatment does not protect PMN depleted mice

Control (IgG isotype) and neutropenic B6 mouse corneas were pretreated with topical flagellin for 6 h and infected with PA as described in the legend of figure 1. (A) Slit lamp examination of infected corneas performed at 1dpi and (B) bacterial load was determined at 2 dpi and indicated *p* values were generated using paired Student's *t* test.

Fig. 6. Topical application of flagellin is ineffective in inducing protection against PA infection in Camp^{-/-} mice cornea

Camp^{-/-} or WT (B6) mice were pretreated with topical application of flagellin (500ng) or PBS. After 6 pretreatment the cornea was scarified and challenged topically with 1.0×10^4 CFU of PA (A) clinical scores for the ocular disease response were assigned on 1, 3, and 5 dpi and represented as mean clinical score, (B) Slit lamp examination of infected corneas performed at 1, 3 and 5 dpi and (C) Bacterial load, (D) cytokine ELISA and (E) MPO assay was performed on representative infected corneas at 5 dpi. * $P < 0.01$, ** $P < 0.05$ (student's *t*-test), significant increase in bacterial load, cytokines and PMN infiltration was detected in Camp^{-/-} compared to WT mice with or without flagellin treatment.