Studies of a human T lymphocyte antigen recognized by a monoclonal antibody

(hybridoma/leukemia)

E. G. ENGLEMAN*, R. WARNKE*, R. I. FOX†, J. DILLEY†, C. J. BENIKE*, AND R. LEVY†‡

†The Howard Hughes Medical Institute Laboratory and the Departments of †Medicine and *Pathology, Stanford University School of Medicine, Stanford, California 94305

Communicated by Henry S. Kaplan, October 3, 1980

ABSTRACT A monoclonal antibody (designated L17F12) detects an antigen present on 95–100% of human peripheral T lymphocytes, the majority of thymocytes, and acute lymphocytic leukemia T cells but not B cells, B-cell lines, or monocytes. Examination of frozen tissue sections by the immunoperoxidase method revealed that the cells expressing this antigen were found predominantly in the medulla of thymus and in T-cell zones of lymph node and spleen. The antigen recognized by L17F12 was associated with a cell-surface glycoprotein of 67,000 daltons. L17F12 was used to isolate this molecule from human thymocytes, normal peripheral T cells, leukemic T cells, and T-cell lines. Expression of this antigen on normal T cells was not diminished by prolonged exposure *in vitro* to various T-cell stimuli. In the absence of complement, L17F12 bound to T cells without altering proliferative functions, thus enabling rapid purification of functionally intact T cells. In the presence of complement, L17F12 was cytolytic for T cells, providing the basis for depletion of T cells from heterogeneous populations. These data suggest that the monoclonal antibody L17F12 recognizes a specific T-cell differentiation protein. This antibody will be useful in studies of the human immune system.

Human T cells have traditionally been defined on the basis of their spontaneous binding to sheep erythrocytes (1, 2). Heterologous anti-T cell antisera have been developed (3–6) but, in general, these reagents have not been widely used because of limitations in titer and quantity. Recently, we and others have utilized the technique of somatic cell hybridization (7) to produce monoclonal antibodies to human T lymphocytes (2–13). These reagents have the advantages of purity, homogeneity, and unlimited supply. In this report we describe the production of a monoclonal antibody (L17F12) to an antigen expressed by all human peripheral T cells. Serological, functional, and biochemical studies of the T-cell antigen are presented.

MATERIALS AND METHODS

Production of Monoclonal Antibodies. BALB/c mice were immunized with the blast cells from a patient with acute lymphocytic leukemia of T-cell type (T-ALL). The immunization scheme, fusion of the immune spleen cells with the myeloma cell line NS1-Ag4, cloning, and initial screening of hybridomas have been described in detail (11). One of the monoclonal antibodies produced in this fusion, designated 12E7, has been shown (11) to react predominantly with thymocytes and acute leukemia cells. A second antibody, designated L17F12, from the same fusion is the subject of the current study. Monoclonal antibodies to human Ia antigen (L243) and to human β_2 -microglobulin (L368) were produced in a separate immunization with cells from the human B-cell line RPMI 8866. An analysis

of these antibodies has appeared (14). All of the monoclonal antibodies in the present studies were used in the form of tissue culture supernatants of cloned cell lines.

Isolation of Human Rosetting and Nonrosetting Cells from Peripheral Blood. Peripheral blood mononuclear leukocytes (PBML) were obtained from normal volunteers by Ficoll/Hypaque density gradient centrifugation of fresh defibrinated blood (15). Populations enriched for T or non-T cells were prepared with a sheep erythrocyte rosetting technique as described (12). Cells recovered from the rosetted pellet by treatment with 0.155 M NH_4^+ were 96–100% rosette positive and contained <3% surface immunoglobulin-positive cells on the basis of a direct fluoresceinated antibody binding assay. In addition, <3% of this T-cell-enriched fraction were monocytes, based on staining with α -naphthyl acetate (16). Nonrosetting (non-T) cells consisted of 48–64% surface Ig-positive cells and 24–46% monocytes.

"Panning" with L17F12 Antibody. In some experiments the technique of Mage *et al.* (17) as modified by Wysocki and Sato (18) was used to fractionate PBML into L17F12-positive and L17F12-negative cells.

Microcytotoxicity Assays. Cytotoxicity assay were performed at room temperature in microtiter wells as described (19).

Analysis and Cell Separation with the Fluorescence-Activated Cell Sorter (FACS). The binding specificity of monoclonal antibodies was studied by indirect immunofluorescence using a fluorescein-conjugated rabbit anti-mouse Ig (R/M-FITC) that had been preabsorbed to remove reactivity with human immunoglobulin. Cells were stained and analyzed with a FACS II (Becton-Dickinson Electronic Laboratories, Mt. View, CA) (20). Normal mouse serum or culture fluid from the IgG1 (K)-secreting myeloma MOPC-21 served as a negative control. To obtain cells for *in vitro* culture, labeling and sorting were carried out aseptically.

Immunoperoxidase Analysis of Solid Tissue with Monoclonal Antibody. Lymph node and thymus were obtained at the time of surgical removal and 1-mm cross sections were placed in embedding medium (O.C.T. Compound, Miles, Naperville, IL) in air-tight plastic capsules (no. 130 Beem, size 00, Palco, Tustin, CA) and frozen in isopentane/dry ice. Cryostat sections (4 μm) were fixed by a brief immersion in acetone, washed with phosphate-buffered saline, and incubated with monoclonal antibodies (21), with biotinylated goat anti-mouse antibody, and then with avidin conjugated to horseradish peroxidase (E. Y. Laboratories, San Mateo, CA). Color was developed by the addition of 0.5% 3,3-diaminobenzidine/0.01% H_2O_2 in phosphate-buffered saline.

The publication costs of this article were defrayed in part by page charge payment. This article must therefore be hereby marked "advertisement" in accordance with 18 U. S. C. §1734 solely to indicate this fact.

Abbreviations: FACS, fluorescence-activated cell sorter; MLR, mixed leukocyte reaction; PBML, peripheral blood mononuclear leukocytes; PHA, phytohemagglutinin; T-ALL, T-cell acute lymphocytic leukemia.

Proliferative Responses to Mitogens and Antigens. Stimulation with concanavalin A (Pharmacia) or phytohemagglutinin (PHA-P, Burroughs Wellcome Research, Beckingham, Kent, England) was carried out as described (12). Assays of [^3H]thymidine uptake were performed in triplicate after 4 days of culture. Stimulation of cells with tetanus toxoid [0.5 limes flocculation (LF) unit/ml] or diphtheria toxoid (0.5 limes flocculation unit/ml) (Massachusetts State Department of Public Health) was dependent on the presence of irradiated [1500 rad (15 grays)] autologous non-T cells. These cultures were assayed after 7 days. Allogeneic mixed leukocyte reactions (MLRs) were carried out between 5×10^4 T responder cells and 5×10^4 irradiated (3000 rad) allogeneic non-T cells and harvested as above at day 7. Autologous MLRs were carried out between 5×10^4 T cells and 1×10^5 irradiated autologous non-T cells in a manner otherwise identical to the allogeneic MLR.

Immunoprecipitation and NaDodSO₄/Polyacrylamide Gel Electrophoresis. Cells were radiolabeled either biosynthetically with [^{35}S]methionine or on their surface with [^{125}I]iodine and lactoperoxidase (22). Nonidet P-40 solubilized cellular proteins were subjected to indirect immunoprecipitation with monoclonal antibodies and an excess of either goat anti-mouse Fab or formalin-fixed *Staphylococcus aureus* Cowan strain I (23). Washed immunoprecipitates were solubilized by boiling in gel sample buffer (10% glycerol/5% 2-mercaptoethanol/3% NaDodSO₄/62.5 mM Tris) and electrophoresed in polyacrylamide gels (24). Fixed, dried gels were exposed to Kodak RP X-Omat film in the presence of intensifying screens (Dupont Lightning Plus).

RESULTS

Hybridomas with T-Cell Specificity. Hybridomas were cloned and screened for the production of antibody by an indirect cell binding assay using [^{125}I]labeled goat anti-mouse Fab (11). A series of clones was identified as producing antibodies that could distinguish between the immunizing T-ALL cells and an isologous Epstein-Barr virus-transformed B lymphoblastoid cell line. A second screening identified a subgroup of these clones whose antibody products bound equally to normal peripheral T cells and to the immunizing T-ALL cells. The binding specificity of one such monoclonal antibody, L17F12, is shown in Fig. 1. High levels of binding occurred with T-ALL cell line RPMI 8402 but no binding occurred with the isologous B-cell line RPMI 8392. In other studies, L17F12 bound the leukemia cells from eight of eight patients with T-ALL and from four of four T-ALL-derived cell lines.

This antibody bound to 60–75% of normal PBML but failed to bind to erythrocytes, platelets, or granulocytes. The reactivity of L17F12 was restricted to the erythrocyte-rosetting (T) cells, 94–98% of which bound this antibody. No binding was

FIG. 1. Binding specificity of L17F12. Target cells were incubated with varying dilutions of the hybridoma culture supernatant, washed, and then incubated with [^{125}I]labeled goat anti-mouse Fab. ●, RPMI 8402, a T-ALL-derived cell line; ○, RPMI 8392, a B-cell line from the same individual.

seen against rosette-negative (non-T) cells (data not shown).

L17F12 is a complement-fixing IgG2a. In the presence of rabbit complement, 60–80% of unseparated PBML were lysed by this antibody. Lympholysis with this antibody was confined to the rosetting cells, of which >95% routinely were killed. Thus, L17F12 can be used with complement to remove T cells from heterogeneous populations.

Nonlytic Separation of T and Non-T Populations with L17F12. Incubation of PBML with L17F12, and even subsequently with fluorescein-conjugated rabbit anti-mouse Ig, did not interfere with the capacity of the cells to form erythrocyte rosettes. Thus, rosetting could be used as an independent marker of T cells purified by the use of L17F12. PBML were separated by FACS into cells positively or negatively stained with L17F12. The positively stained cells were 95% rosette positive. The cells unstained by L17F12 consisted of 4% rosetting cells, 51% surface Ig-positive cells, and 30% monocytes.

An alternative method for the isolation or depletion of T cells with L17F12 was the use of coated plastic dishes for "panning." Plastic petri dishes were coated with affinity-purified goat anti-mouse Ig. T cells preincubated with L17F12 were found to bind specifically to these dishes. The purified T cells could be retrieved by vigorous pipetting. Typically, ≥95% of the cells that bound to the dish were rosette positive whereas the majority

Table 1. Influence of antibody L17F12 on T-cell function

Responder cells*	Response to various stimuli, cpm†						
	None	Autologous non-T cells	Allogeneic PBML	PHA	Con A	Tetanus toxoid	T-GAL
A	204 ± 49	14,912 ± 1,313	37,419 ± 2,554	96,423 ± 7,505	42,314 ± 3,780	34,315 ± 2,866	22,987 ± 1,872
B	381 ± 55	16,411 ± 1,418	38,823 ± 3,198	102,871 ± 5,292	38,594 ± 3,451	31,499 ± 3,080	28,272 ± 2,795
C	415 ± 68	16,140 ± 1,875	41,322 ± 3,467	100,712 ± 6,886	45,330 ± 4,668	37,210 ± 2,541	24,269 ± 3,601
D	353 ± 42	15,567 ± 1,224	42,540 ± 3,671	103,252 ± 7,243	44,812 ± 3,762	32,954 ± 3,800	27,170 ± 2,613

* Responder cells consisted of: A, erythrocyte rosette-positive cells; B, rosette-positive cells incubated with L17F12 plus fluorescein-conjugated rabbit anti-mouse Ig and then washed; C, L17F12-positive cells obtained after FACS fractionation of PBML; D, rosette-positive cells in the presence of saturating amounts of L17F12 added at the beginning of culture.

† Results of triplicate cultures are expressed as mean ± SEM. Stimulation with PHA, concanavalin A (Con A), tetanus toxoid, or T-GAL [poly(tyrosine, glutamic acid)-polyDL-alanine-polylysine] was performed in the presence of 5×10^3 irradiated (6000 rad) autologous non-T cells.

FIG. 2. Frozen sections of a human lymph node stained with monoclonal antibodies L17F12 (*Upper*) and L243 (*Lower*) by the immunoperoxidase technique. (*Upper*) L17F12. The majority of the cells in the paracortex are stained (arrow). Only occasional cells in the follicle are stained. (*Lower*) L243 (anti-Ia). The cells in the germinal center and mantle of the secondary follicle are stained (arrow), as well as occasional cells in the adjacent paracortex.

of unbound cells were rosette-negative Ig-positive lymphocytes (data not shown).

Functional Studies. Erythrocyte rosette-positive cells were coated with the antibody and tested for their capacity to respond to specific and nonspecific proliferative stimuli. No interference with T-cell proliferation in response to mitogens, soluble protein antigens, or alloantigens was seen with saturating concentrations of L17F12 (Table 1). This was true whether free antibody was removed prior to stimulation or in the continuous presence of saturating quantities of antibody. Thus, L17F12 could be used to isolate T cells without loss of their function. Moreover, the expression of L17F12 antigen was maintained on T cells during activation by mitogens, soluble antigens, and alloantigens (data not shown). In general, culture in PHA or concanavalin A resulted in more brightly stained cells whereas soluble antigens and alloantigens did not change relative fluorescence despite vigorous blast transformation as measured by incorporation of [^3H]thymidine and enumeration of blast cells microscopically.

Tissue Localization of L17F12-Reactive Cells. To determine the distribution of L17F12-positive cells in solid tissues, frozen sections were stained by an immunoperoxidase method (21). Fig. 2 shows normal lymph node and compares the staining by antibody L17F12 with that by antibody L243 (anti-Ia). The anti-Ia antibody detected cells predominantly in the germinal center and mantle of the follicle, whereas L17F12 detected cells in the

FIG. 3. Sections of human thymus stained with L17F12 (*Upper*) and 12E7 (*Lower*). (*Upper*) L17F12. The intensely stained cells reside in the medulla which contains a Hassall corpuscle (arrow). (*Lower*) 12E7. The stained cells are restricted to the cortex (arrow).

T-cell zone and only occasional cells within the follicle (Fig. 2). Similar results (not shown) were obtained on sections of normal spleen. With normal human thymus, cells reacting the strongest with L17F12 were concentrated in the thymic medulla whereas those detected by 12E7 [an antibody shown to react predominantly with cortical thymocytes (11)] were restricted to the cortex (Fig. 3).

Molecular Characterization of the Target Antigen of L17F12. For structural studies of L17F12 antigen, T cells or cell lines were radiolabeled either externally with ^{125}I or internally with [^{35}S]methionine. The radiolabeled proteins were extracted with Nonidet P-40, immunoprecipitated with L17F12 and goat anti-mouse Ig, and analyzed by electrophoresis in NaDodSO₄/polyacrylamide slab gels. The pattern of immunoprecipitable material obtained from surface- ^{125}I -labeled T lymphoma cells is shown in Fig. 4. The reduced and unreduced samples were electrophoresed in parallel on the same gel. A predominant band at 67,000 daltons was apparent under reducing conditions. The unreduced material showed a similarly predominant but slightly smaller band. Additional minor bands were seen in the range 60,000–65,000 daltons. These were of variable intensity depending on the T-cell source and they were always accompanied by the major 67,000-dalton band.

Fig. 5 shows immunoprecipitates from [^{35}S]methionine-labeled T cells of various sources. In each example, a single band of 67,000 daltons is seen. All of the material immunoprecipitable by L17F12 could be removed by passage over a lentil lectin affinity column (data not shown). From these experiments, we conclude that the molecule bearing the determinant recognized

FIG. 4. NaDodSO₄/10% polyacrylamide gel electrophoresis of L17F12 immunoprecipitable material from ¹²⁵I-labeled T lymphoma cells analyzed before (*Right*) and after (*Left*) reduction with 2-mercaptoethanol. In each case an IgG was electrophoresed in parallel with the immunoprecipitates to provide size markers.

by L17F12 is a glycosylated polypeptide with an apparent size of approximately 67,000 daltons.

Reactivity of L17F12 for PBML from Other Species. PBML were obtained from various animal species and assayed for binding of L17F12 as detected by ¹²⁵I-labeled goat anti-mouse Ig. Cells from mice, rats, and dogs and from various lower primate species including rhesus, owl, and squirrel monkeys failed to react. However, cells from higher primates, including gibbon apes and chimpanzees, were reactive. In these latter primate species, the reactive cells were restricted to the erythrocyte-rosetting fraction (R. H. Newbauer, Frederick Cancer Research Center, personal communication).

FIG. 5. NaDodSO₄/polyacrylamide gel electrophoresis of [³⁵S]methionine-labeled cell proteins immunoprecipitated with L17F12. Material from three different cell sources was analyzed: T lymphoma cells from a patient with the Sézary syndrome, cells from a patient with chronic lymphocytic leukemia of T cell type (T-CLL), and normal thymocytes. For comparison, the HLA and β_2 -microglobulin chains immunoprecipitated by monoclonal antibody L368 (anti- β_2 -microglobulin) from Sézary cells are shown.

DISCUSSION

These studies demonstrate that monoclonal antibody L17F12 recognizes a glycoprotein expressed on the surface of peripheral blood T cells, thymocytes, and T-ALL cells but not on B cells, transformed B-cell lines, monocytes, or other elements of normal peripheral blood. NaDodSO₄/polyacrylamide gel electrophoresis of the L17F12 target antigen revealed it to be a 67,000-dalton glycoprotein. The relatively intense expression of L17F12 antigen on medullary thymocytes and peripheral T cells would suggest that the antigen is a marker of a late stage of T-cell differentiation. Its absence from various animal species, including Old World monkeys, and its presence in the gibbon ape and chimpanzee suggests that it is a recent marker in the phylogeny of T cells.

The available information suggests that this antigen is distinct from some other reported human pan-T-cell markers. For example, L17F12 antigen apparently differs from the sheep erythrocyte receptor because lymphocytes bound by L17F12 are unimpeded from forming erythrocyte rosettes, and rosetted cells bind L17F12 to the same extent as do unrosetted T cells. On the basis of marked differences in size, the L17F12 antigen can also be distinguished from another human antigen, thought to be the analog of murine Thy-1 (25). Furthermore, unlike the L17F12 antigen, this antigen is not expressed on normal peripheral T cells in man. The OKT₃ monoclonal antibody described by Kung *et al.* (8, 26) recognizes all peripheral T cells but, unlike L17F12, it binds to only 15% of thymocytes and fails to bind to most leukemia T cells (27). On the other hand, the OKT₁ (26), SK5 (28), T101 (29), and 10-2 (30) monoclonal pan-T antibodies share a number of similarities with L17F12.

The current study shows that the L17F12 antigen continues to be expressed under various culture conditions; that is, it is a stable T-cell marker. The failure of L17F12 to alter T-cell responses to such stimuli as lectin mitogens, soluble antigens, and alloantigens suggests that the L17F12 antigen does not mediate any of the functions tested. However, these studies do not rule out a functional role for the molecule bearing the L17F12 antigen. For example, certain monoclonal antibodies to human Ia antigens have only slight effect on stimulation in MLRs or presentation of antigen by accessory cells but clearly precipitate the same molecules as do antibodies that block these functions (31).

Regardless of the function of its target antigen, L17F12 promises to be a useful reagent. The current study demonstrates that the antibody may be useful in studies of lymphoid organ architecture. Also, the antibody can be used to separate T cells from heterogeneous lymphocyte populations without affecting T proliferative functions. In this regard, the antibody has been used to confirm that cells expressing Ia antigen after activation in an MLR are T cells (32). The observation that L17F12 antigen is expressed on T-ALL cells but not on non-T-ALL cells suggests that this antibody might be useful as a standard reagent for the classification of leukemias (33). Alternatively, in the presence of complement, L17F12 is lytic for T cells and can be used to remove "contaminating" T cells from non-T preparations for purposes of HLA-DR typing. Finally, it has now been shown that myeloid stem cells in human bone marrow are unaffected by treatment with L17F12 and complement under conditions in which all T cells could be killed [ref. 34; R. Parkman (Sydney Farber Cancer Center), H. Kaiser (Johns Hopkins University), personal communications]. Conceivably, the same approach might be used on a larger scale for bone marrow transplantation to deplete donor bone marrow of T cells prior to transplantation in order to reduce the risk of graft-vs.-host reaction.

We are grateful to Hidetoshi Kaneoka and Meme Pederson for technical assistance. We also thank Drs. Peter Greenberg, Herbert Kaiser,

Robertson Parkman, and Russell Neubauer for allowing us to refer to their unpublished results. This work was supported by Grants CA 21223-03 and CA 24607-02 from the National Institutes of Health and a grant from the Veterans Administration.

1. Jondall, M., Golm, G. & Wigzell, H. J. (1972) *J. Exp. Med.* **136**, 217–215.
2. Wybran, J., Carr, M. C. & Fudenberg, (1972) *J. Clin. Invest.* **51**, 2537–2543.
3. Aiuti, F. & Wigzell, H. (1973) *Clin. Exp. Immunol.* **13**, 171–181.
4. Smith, R. W., Terry, W. D., Buell, D. M. & Sell, K. W. (1973) *J. Immunol.* **110**, 884–887.
5. Williams, R. C., Jr., DeBoard, J. R., Mellbye, O. J., Messner, R. P. & Lundstrom, F. D. (1973) *J. Clin. Invest.* **52**, 283–295.
6. Evans, R. L., Beard, J. M., Lazarus, A., Schlossman, S. F. & Chess, L. (1977) *J. Exp. Med.* **145**, 221–233.
7. Köhler, G. & Milstein, C. (1975) *Nature (London)* **256**, 495–497.
8. Kung, P. C., Goldstein, G., Reinherz, E. L. & Schlossman, S. F. (1979) *Science* **206**, 347–349.
9. McMichael, A. J., Pilch, J. F., Galfre, G., Mason, D. Y., Fabre, J. W. & Milstein, C. (1979) *Eur. J. Immunol.* **9**, 205–210.
10. Haynes, G. V., Eisenbarth, G. S. & Fauci, A. S. (1979) *Proc. Natl. Acad. Sci. USA* **76**, 5829–5833.
11. Levy, R., Dilley, J., Fox, R. I. & Warnke, R. (1979) *Proc. Natl. Acad. Sci. USA* **76**, 6552–6556.
12. Engleman, E. G., Benike, C., Osborne, B. & Goldsby, R. (1980) *Proc. Natl. Acad. Sci. USA* **77**, 1607–1611.
13. Levy, R., Dilley, J. & Lampson, L. A. (1978) *Curr. Top. Microbiol. Immunol.* **81**, 164–169.
14. Lampson, L. A. & Levy, R. (1980) *J. Immunol.* **125**, 293–299.
15. Boyum, A. (1968) *Scand. J. Clin. Lab. Invest. Suppl.* **21** **97**, 77–89.
16. Yam, L. P., Li, L. Y. & Crosby, W. K. (1971) *Am. J. Clin. Pathol.* **65**, 283–290.
17. Mage, M. G., McHugh, L. & Rothstein, T. (1977) *J. Immunol. Methods* **15**, 47–56.
18. Wysocki, L. J. & Sato, V. L. (1978) *Proc. Natl. Acad. Sci. USA* **75**, 2844–2848.
19. Ray, J. G. (ed.) (1979) National Institutes of Health publication no. 80-545 (U.S. Department of Health, Education, and Welfare), p. 39.
20. Julius, M. D., Sweet, R. G., Fathman, C. G. & Herzenberg, L. A. (1974) in *Mammalian Cells: Probes and Problems*. Proceedings of the First Los Alamos Life Science Symposium AEC Series (CONS:73-1007), eds. Richmond, C. R., Petersen, D. F., Mullaney, P. F. & Anderson, E. C. (Technical Information Center, Office of Public Affairs, U.S. Energy Research and Development Administration, Washington, DC), p. 107.
21. Warnke, R. & Levy, R. (1980) *J. Histochem. Cytochem.* **28**, 771–776.
22. Hausteine, D. (1975) *J. Immunol. Methods* **7**, 25–38.
23. Kessler, S. W. (1975) *J. Immunol.* **115**, 1617–1624.
24. Laemmli, U. K. (1970) *Nature (London)* **227**, 680–685.
25. Ades, E. W., Zwerner, R. K., Acton, R. T. & Balch, C. M. (1980) *J. Exp. Med.* **151**, 400–406.
26. Reinherz, E. L., Kung, P. C., Goldstein, G. & Schlossman, S. F. (1979) *J. Immunol.* **123**, 1312–1317.
27. Reinherz, E. L., Kung, P. C., Goldstein, G., Levey, R. H. & Schlossman, S. F. (1980) *Proc. Natl. Acad. Sci. USA* **77**, 1588–1592.
28. Chang, Y. W., Good, R. A., Ammirati, P., Dymfort, G. & Evans, R. L. (1980) *J. Exp. Med.* **151**, 1539–1544.
29. Royston, I., Majda, J. A., Baird, S. M., Meserve, B. L. & Griffiths, J. C. (1980) *J. Immunol.* **125**, 725–731.
30. Martin, P. J., Hansen, J., Nowinski, R. C. & Brown, M. A. (1980) *Immunogenetics* **11**, 429–439.
31. Engleman, E. G., Charron, D., Benike, C. J. & Stewart, G. J. (1980) *J. Exp. Med.* **152**, 99–113s.
32. Charron, D., Engleman, E. G., Benike, C. J. & McDevitt, H. O. (1980) *J. Exp. Med.* **152**, 127s–136s.
33. Zipf, T., Fox, R. I. & Levy, R. *Cancer Res.*, in press.
34. Greenberg, P., Grossman, M., Charron, O. & Levy, R. (1981) *Exp. Hematol.*, in press.