Published in final edited form as:

Tetrahedron Lett. 2011 April 27; 52(17): 2162–2164. doi:10.1016/j.tetlet.2010.11.134.

STUDIES DIRECTED TOWARD THE SYNTHESIS OF NAKADOMARIN A

Thomas Haimowitz[♦], Mark E. Fitzgerald[♦], and Jeffrey D. Winkler

Department of Chemistry, University of Pennsylvania, Philadelphia, PA 19104

Abstract

The application of a Pummerer-initiated tandem reaction cascade leads to the highly stereoselective formation of the tetracyclic core of Nakadomarin A.

Keywords

alkaloid; Pummerer; tandem reactions; nakadomarin; manzamine

Nakadomarin A **1** was first isolated from the Okinawan marine sponge *Amphimedon sp.* by Kobayashi in 1997.ⁱ The structure of **1** was elucidated spectroscopically and consists of an unprecedented hexacyclic ring system (8/5/5/5/15/6). Kobayashi has proposed an interesting biogenetic transformation for the conversion of ircinal A **2**, the precursor to manzamine A **3**,ⁱⁱ to nakadomarin A **1**, underscoring the striking structural similarities among this compound class.ⁱⁱⁱ

Nakadomarin A **1** demonstrates a range of promising biological activities including cytotoxic activity against murine lymphoma L1210 cells (IC₅₀ 1.3 µg/mL), inhibitory activity against cyclin dependent kinase 4 (IC₅₀ 9.9 µg/mL), anti-microbial activity against a fungus (*Trichophyton mentagrophytes*, MIC 23 µg/mL) and a Gram-positive bacterium (*Cornebacterium xerosis*, MIC 11 µg/mL). However, its limited availability from natural sources (1.8 × 10⁻³% of sponge wet weight) has prevented a complete study of its biological activity. The structural complexity of nakadomarin A coupled with its intriguing and as yet not fully explored biological activity led to considerable effort directed toward the synthesis

© 2010 Elsevier Ltd. All rights reserved.

Correspondence to: Jeffrey D. Winkler.

[♦]Current Address: TetraLogic Pharmaceuticals, 343 Phoenixville Pike, Malvern, PA 19355

[♦]Current Address: Chemical Biology Platform, The Broad Institute of MIT and Harvard, 7 Cambridge Center, Cambridge, MA 02142

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

of **1**, culminating in a series of elegant total syntheses.^{iv} We describe herein a conceptually novel approach to the synthesis of the manzamine core based on a Pummerer-initiated tandem reaction cascade.^v

We envisioned that the ABCDE pentacyclic core **4** of nakadomarin A **1** could be obtained by global reduction of bislactam sulfide **5**, which we envisioned as the product of a tandem Pummerer-initiated tandem reaction cascade of **7** via the intermediacy of thionium ion **6**. We note that the elegant studies of Martin,^{2b} Mukai,^{4f} and Zhai^{vi} suggest that the C-14 stereocenter would lead to the desired diastereoselectivity in the cascade cyclization reaction via approach of the thionium ion **6** from the α -face of the pyrroline D-ring to give **5**. We anticipated that the requisite cyclization substrate **7** could be prepared via coupling of **8** and **9**.

To examine the feasibility of the key cascade sequence, we examined the reaction in the simpler model system **16**,^{vii} lacking the C-14 stereocenter, as outlined in Scheme 2. Metalation of **10**,^{viii} based on the work of Das, and reaction of the derived anion with ethylene oxide delivered alcohol **11**, which on Mitsunobu reaction with phthalimide afforded **12**. Deprotection of **12** with hydrazine and reductive alkylation with p-bromobenzaldehyde gave secondary amine **14**, which was coupled with acid chloride **15**^{2b} to give the cascade cyclization substrate **16**. Exposure of **16** to dimethyl(methylthio)sulfonium tetrafluoroborate (DMTSF)^{ix} led to the exclusive formation of the desired tetracycle **17**, the structure of which was unambiguously confirmed via X-ray crystallographic analysis of the hydrochloride salt of the derived tertiary amine **18**, as shown in Scheme 2.^x

The extension of these results to a substrate containing the C-14 stereocenter is outlined in Scheme 3. Reaction of the known acid **19**^{2b} with the p-methoxybenzyl amine **20** (obtained from **13** by reductive alkylation) gave amide **21**. While reaction of **21** with DMTSF resulted in extensive decomposition of the starting material without formation of the desired product, we were delighted to find that exposure of the corresponding benzoate **22** to DMTSF led to the formation of the desired cyclization product **23** in 50% yield, which was fully characterized as the corresponding Cbz analog **24**.^{xi} The stereoselective formation of **23** is consistent with the addition of the thionium ion from the α -face of the pyrroline ring, as noted in related systems by Zhai⁶ and Martin.^{2b} Further support for this stereochemical assignment, albeit indirect, was obtained by cyclization of **24** to the cyclic urethane **25**, which revealed no nOe between the two methines (C-6 and C-14 using the numbering of the nakadomarin ring system) of the pyrrolidine ring of **25**.^{xii}

We have developed an alternative strategy to the elegant work of Zhai⁶ on the use of the proline-derived stereochemistry to control the establishment of the key stereochemical relationships of the core structure of nakadomarin A. In both Zhai's work and the current study, a single stereocenter leads to the establishment of all of the key stereochemical relationships of the nakadomarin tetracyclic core. Further studies on the application of this strategy to the synthesis of biologically relevant systems are underway in our laboratories and our results will be reported in due course.

Acknowledgments

This paper is dedicated with deep gratitude to Professor Harry Wasserman, a scientist who has taught us so much about the art of organic synthesis. We would like to thank the NIH (CA134983) for their generous support of this work.

LITERATURE CITED

- i. Kobayashi J, Watanabe D, Kawasaki N, Tsuda M. *J Org Chem*. 1997; 62:9236–9239.
- ii. For the total synthesis of manzamine A, see a) Winkler JD, Axten JM. *J Am Chem Soc*. 1998; 120:6425–6426. b) Martin SF, Humphrey JM, Ali A, Hillier MC. *J Am Chem Soc*. 1999; 121:866–867. c) Toma T, Kita Y, Fukuyama TJ. *Am Chem Soc*. 2010; 132:10233–10235.
- iii. Kobayashi J, Tsuda M, Ishibashi M. *Pure Appl Chem*. 1999; 71:1123–1126.
- iv. For a recent review, see a) Martin D, Vanderwal C. *Ang Chem Int Ed*. 2010; 49:2830–2832. b) Nagata T, Nakagawa M, Nishida A. *J Am Chem Soc*. 2003; 125:7484–7485. [PubMed: 12812466] c) Ono K, Nakagawa M, Nishida A. *Ang Chem Int Ed*. 2004; 43:2020–2023. d) Young I, Kerr M. *J Am Chem Soc*. 2007; 129:1465–1469. [PubMed: 17263433] e) Jakubec P, Cockfield D, Dixon D. *J Am Chem Soc*. 2009; 131:16632–16633. [PubMed: 19883080] f) Milson M, Funk R. *Org Lett*. 2010; 12:4912–4915. [PubMed: 20932022] For a formal synthesis of nakadomarin A, see g) Inagaki F, Kinebuchi M, Miyakoshi N, Mukai C. *Org Lett*. 2010; 12:1800–1803. [PubMed: 20232853]
- v. For recent reviews, see Padwa A. *J Org Chem*. 2009; 74:6421–6441. [PubMed: 19663447] Bur S, Padwa A. *Adv Het Chem*. 2007; 94:1–105. Bur S, Padwa A. *Chem Rev*. 2004; 104:2401–243. [PubMed: 15137795]
- vi. Deng H, Yang X, Tong Z, Li Z, Zhai H. *Org Lett*. 2008; 10:1791–1793. [PubMed: 18393518]
- vii. All new compounds were fully characterized by ^1H and ^{13}C NMR, IR, and high resolution mass spectrometry.
- viii. Das B, Ramyu R, Reddy M, Mahender G. *Synthesis*. 2005:250–254.
- ix. Trost B, Murayama E. *J Am Chem Soc*. 1981; 103:6529–6530.
- x. CCDC 796390 Cambridge Crystallographic Data Centre.
- xi. Characterization data for **24**: $[\alpha]_{\text{D}}^{24.1} = +83.36$, $c = 1.00$ (CDCl_3); ^1H NMR (500 MHz, CDCl_3 , 335 °K) δ 7.87 (d, $J = 7.6$ Hz, 2H), 7.57 (d, $J = 8.4$ Hz, 2H), 7.47 (br s, 1H), 7.46–7.28 (m, 5H), 7.16 (d, $J = 8.4$ Hz, 2H), 6.72 (d, $J = 8.2$ Hz, 2H), 6.32 (d, $J = 1.6$ Hz, 1H), 5.76 (s, 1H), 5.70 (d, $J = 5.0$ Hz, 1H), 5.28 (d, $J = 2.4$ Hz, 2H), 4.91 (d, $J = 13.5$ Hz, 1H), 4.73 (dd, $J = 4.7$, 10.5 Hz, 1H), 4.63 (t, $J = 9.4$ Hz, 1H), 4.35 (br, 1H), 4.33 (d, $J = 14.3$ Hz, 1H), 4.03 (d, $J = 17.3$ Hz, 1H), 3.74 (dd, $J = 6.2$, 17.4 Hz, 1H), 3.69 (s, 3H), 2.44 (d, $J = 13.7$ Hz, 1H), 2.21 (dd, $J = 8.6$, 13.7 Hz, 1H); ^{13}C NMR (125 MHz, CDCl_3 , 335 °K) δ 171.0, 165.3, 159.4, 148.9, 136.5, 131.7, 131.3, 129.5, 129.2, 128.4, 128.2, 127.9, 124.2, 114.4, 109.0, 105.0, 67.6, 64.5, 62.0, 58.3, 55.2, 50.0, 47.5, 38.1, 29.7; IR (neat) 2917, 2850, 2253, 1714, 1698, 1652, 1590, 1514, 1486, 1404, 1354, 1272, 1248, 1175, 1118, 1030 cm^{-1} ; HRMS (ESI) m/z calc. for $\text{C}_{36}\text{H}_{31}\text{BrN}_2\text{NaO}_7$ $[\text{MNa}^+]$ 705.1212, obsd. 705.1236.
- xii. Characterization data for **25**: $[\alpha]_{\text{D}}^{24.2} = +122.85$, $c = 1.00$ (CDCl_3); ^1H NMR (500 MHz, CDCl_3) δ 7.49 - 7.45 (m, 1H), 7.18 (d, $J = 8.5$ Hz, 2H), 6.88 - 6.81 (m, 2H), 6.35 (d, $J = 1.9$ Hz, 1H), 5.94 (s, 1H), 5.72 (dd, $J = 2.0$, 6.5 Hz, 1H), 4.61 (d, $J = 14.4$ Hz, 1H), 4.54 (d, $J = 14.4$ Hz, 1H), 4.48 (dd, $J = 7.4$, 8.9 Hz, 1H), 4.17 (dd, $J = 2.3$, 9.1 Hz, 1H), 4.01 - 3.93 (m, 2H), 3.79 (s, 3H), 3.75 (dd, $J = 6.5$, 17.3 Hz, 1H), 2.36 (dd, $J = 7.1$, 12.8 Hz, 1H), 2.00 (dd, $J = 9.1$, 12.8 Hz, 1H); ^{13}C NMR (125 MHz, CDCl_3) δ 170.5, 159.4, 159.2, 159.1, 149.8, 136.5, 129.6, 128.9, 127.0, 114.2, 109.8, 105.5, 67.7, 66.9, 62.9, 58.0, 55.3, 49.4, 47.8, 44.0; IR (neat) 2969, 2925, 2856, 1750, 1646, 1611, 1510, 1398, 1375, 1248, 1222, 1178, 1029 cm^{-1} ; HRMS (ESI) m/z calc. for $\text{C}_{22}\text{H}_{21}\text{N}_2\text{O}_5$ $[\text{MH}^+]$ 392.1372, obsd. 392.1348.

Figure 1.

Scheme 1.

Scheme 2.

Scheme 3.