POSTER PRESENTATION

Open Access

Granuloma-in-follicles in pediatric Crohn's disease

Carl El Janssen^{1*}, Isabelle Cleynen², Gert De Hertogh¹, Valeer J Desmet¹, Tania Roskams¹, Carine H Wouters³

From 18th Pediatric Rheumatology European Society (PReS) Congress
Bruges, Belgium. 14-18 September 2011

Background

SNPs in NOD2 are a major susceptibility factor for Crohn's disease (CD). (Sub)mucosal granulomas are a diagnostic feature of CD.

Aim

To better describe morphological and immunohistochemical features of pediatric CD (pCD) granulomas.

Methods

17 pCD patients with granulomas were genotyped for CD-associated mutations or polymorphisms in NOD2 and autophagy-related genes ATG1 and ATG16L1. Granulomas were found in intestinal biopsies of 22 out of 83 endoscopic and 9 out of 12 surgical procedures. Of each procedure we selected one biopsy per organ to study the cellular composition and cytokine profile of granulomas. We stained 43 paraffin-embedded, formalin-fixed pCD biopsies with hematoxylin&eosin and monoclonal antibodies targeting leukocyte markers (HLA-DR, CD68, CD4, CD8, CD20, IL23R) and cytokines (TNF α , IFN γ , IL6, IL10, IL17, TGF β) (Figure 1)

Morphology and immunohistochemistry were scored semi-quantitatively.

Results

In addition to small isolated (sub)mucosal granulomas, we found granulomas in the follicle(GIF) centre of mucosa-associated lymphoid tissue, visualized by CD20 staining (fig). GIF were found in half the number of procedures (more often in surgical than in endoscopic) in only 7 out of 17 pCD patients. Exceptional lymphocyte emperipolesis in multinucleated giant cells (MGCs) and IL-17 expression were observed, only in granulomas of pCD with GIF (fig). The granulomas of patients with GIF often had a lymphocytic corona, but polycyclic granuloma architecture was rare. No clear relation was seen between the presence of GIF and the CD-associated NOD2, ATG1 and ATG16L1 genotypes.

Conclusion

We defined GIF as a morphological characteristic in few selected pCD patients, and demonstrate association with

Figure 1

* Correspondence: Carljanssen39@hotmail.com

¹Departments of Pathology, University Hospital Leuven, Belgium

Full list of author information is available at the end of the article

emperipolesis of lymphocytes in MGCs and weak IL17 expression.

Author details

¹Departments of Pathology, University Hospital Leuven, Belgium. ²Gastroenterology, University Hospital Leuven, Belgium. ³Pediatric Rheumatology, University Hospital Leuven, Belgium.

Published: 14 September 2011

doi:10.1186/1546-0096-9-S1-P294

Cite this article as: Janssen *et al.*: Granuloma-in-follicles in pediatric Crohn's disease. *Pediatric Rheumatology* 2011 **9**(Suppl 1):P294.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit