Published in final edited form as:

European J Org Chem. 2011 August ; 2011(23): 4439–4447. doi:10.1002/ejoc.201100382.

Rapid and Efficient Radiosyntheses of Meta-substituted [¹⁸F]Fluoroarenes from [¹⁸F]Fluoride Ion and Diaryliodonium Tosylates within a Microreactor

Joong-Hyun Chun^a, Shuiyu Lu^a, and Victor W. Pike^{a,*}

^aMolecular Imaging Branch, National Institute of Mental Health, National Institutes of Health, 10 Center Drive, Building 10, Room B3 C346A, Bethesda, MD 20892 USA

Abstract

Effective methods for the introduction of the short-lived positron-emitter fluorine-18 ($t_{1/2} = 109.7$ min) at high specific radioactivity into fluoroarenes are valuable for the development of radiotracers for molecular imaging with positron emission tomography. Here we have explored the scope of the radiofluorination of diaryliodonium salts with no-carrier-added [¹⁸F]fluoride ion for the preparation of otherwise difficult to access meta-substituted [¹⁸F]fluoroarenes. A microfluidic reaction platform was used to establish optimal radiochemical yields. Rapid, high yielding and selective radiofluorinations were achieved in unsymmetrical diaryliodonium tosylates ($\text{ArI}^+\text{Ar}'\text{TsO}^-$) in which Ar carried either a meta electron-withdrawing (CN, NO₂, CF₃) or a meta electron-donating (Me or MeO) group, and in which the partner aryl group (Ar') was relatively electron-rich, such as Ph, 3-Me-C₆H₄, 4-MeO-C₆H₄, 2-thienyl or 5-Me-2-thienyl. The radiofluorination of appropriate diaryliodonium tosylates is therefore a generally useful method for the preparation of simple [¹⁸F]*m*-fluoroarenes ([¹⁸F]ArF).

Keywords

Isotopic labeling; Fluorine-18; Hypervalent compounds; Microreactors; Iodonium salts

Introduction

The short-lived positron-emitter, fluorine-18 ($t_{1/2} = 109.7$ min), has gained great importance as a radiolabel for probes used with positron emission tomography (PET)^[1] - a molecular imaging technique that is widely applied both in clinical research^[2] and in drug development.^[3–8] Expansion of the utility of PET depends on the development and availability of radioactive probes that are specific to particular biochemical targets or pathways. Radioligands bind to specific protein targets such as enzymes, channels, receptors, or transporters to enable their concentrations to be determined in vivo.^[9] Yet other radiotracers may act as enzyme substrates and so permit other types of measurement, such as [¹⁸F]2-fluoro-2-deoxy-D-glucose ([¹⁸F]FDG) for measuring glucose metabolism.^[10] Fluorine-18 is attractive as a radiolabel for several reasons. These include i) that fluorine may to some extent mimic a hydrogen atom or hydroxyl group in an organic molecule, ii) the half-life of fluorine-18 is well suited to monitoring kinetics in vivo over an acceptably short scanning session, and iii) fluorine-18 can be produced with a moderate energy

Fax: +1-301-480-5112, pikev@mail.nih.gov.

Supporting information for this article is available on the WWW under <http://www.eurjoc.org/> or from the author.

Supporting Information Copies of ¹H, ¹³C and ¹⁹F NMR spectroscopic data and sample radiochromatograms.

cyclotron in very high activities and specific activities through the $^{18}\text{O}(\text{p},\text{n})^{18}\text{F}$ reaction on ^{18}O -enriched water.^[11,12] The ease of production of high amounts of fluorine-18 coupled with its almost 2-h half-life also allows this isotope to be distributed over distances that may be covered in a few hours to sites where a production cyclotron is unavailable. Hence, some useful ^{18}F -labeled radiotracers, such as [^{18}F]FDG, have become commercially available for clinical application through a network of regional distribution centers.

The $^{18}\text{O}(\text{p},\text{n})^{18}\text{F}$ reaction on ^{18}O -enriched water provides the fluorine-18 as aqueous fluoride ion. Therefore, all radiochemistry with fluorine-18 from this method of production must first involve a reaction of fluoride ion. In general, this usually implies an aliphatic or aromatic nucleophilic substitution reaction.^[13] Labeling at aliphatic carbon atoms can be accomplished efficiently. However, fluorine-18 bonded to aliphatic carbon may be prone to defluorination in vivo, giving rise to [^{18}F]fluoride ion which then binds avidly to bone. Defluorination can be problematic for PET scanning. For example, the uptake of [^{18}F]fluoride ion by skull may compromise PET measurements with parent radiotracer in nearby brain. For this reason, attachment of fluorine-18 to an aryl carbon is often more attractive, since tendency for radiodefluorination is usually greatly reduced.

Several methods for preparing ^{18}F -labeled aryl fluorides have been evaluated and applied. These include the Balz-Schiemann reaction,^[14] the Wallach reaction,^[15] and classical aromatic nucleophilic substitution^[13,16,17] (Scheme 1). The Balz-Schiemann reaction suffers from low maximal theoretical radiochemical yield (25%) which in practice may be very much lower on complex substrates, and also dilution of specific radioactivity due to the co-production of non-radioactive fluoro product from the usual tetrafluoroborate anion. The latter problem may be circumvented by use of non-fluorine containing anions, but the process is difficult and radiochemical yields remain low.^[18] The Wallach reaction may be performed at high specific radioactivity but again radiochemical yields tend to be very low on complex substrates. Over recent decades, classical aromatic nucleophilic substitution has become the most established and popular method for preparing [^{18}F]fluoroarenes at high specific radioactivity.^[13,16,17] High radiochemical yields can be obtained, especially when deploying nitro or trimethylamino as the leaving group. Nevertheless, this method also has some limitations. In particular, an electron-withdrawing group, generally in ortho or para orientation to the leaving group, is required on the ring to be fluorinated.

A more recently explored method for the preparation of [^{18}F]fluoroarenes is the reaction of [^{18}F]fluoride ion with diaryliodonium salts (Scheme 2).^[19–23] This method permits the introduction of [^{18}F]fluoride ion not only into electron-deficient rings but also into electron-rich rings, and into ortho,^[24] meta or para orientation to substituents. A mechanism that is wholly distinct from classical aromatic nucleophilic substitution probably accounts for these useful features.^[24,25] Despite the great potential of this labeling method, its scope has not been studied systematically. Here, we take advantage of the benefits of a microfluidic apparatus^[26–29] for performing sequential small-volume radiofluorination reactions under well-controlled conditions as we described previously,^[30,31] in order to study the reactions of diaryliodonium salts with [^{18}F]fluoride ion to prepare meta-substituted [^{18}F]fluoroarenes. We show that these [^{18}F]fluoroarenes can be obtained rapidly and in high radiochemical yield regardless of the electron-donating or withdrawing nature of the meta substituent, provided that the partner ring in the diaryliodonium salt is relatively electron-rich.

Results and Discussion

Our aim was to study the reactivity of no-carrier-added [^{18}F]fluoride ion towards diaryliodonium salts ($\text{ArI}^+\text{Ar}'\text{X}^-$) bearing one of five meta substituents (OMe, Me, CN, CF_3 or NO_2) in one ring (Ar), where this ring is partnered in the salt with another relatively

electron-rich ring (Ar') selected from Ph, 3-MeC₆H₄, 4-MeOC₆H₄, 2-thienyl or 5-Me-2-thienyl. It was necessary to prepare each of these salts, with poorly nucleophilic tosylate as the preferred counter anion (X⁻ = TsO⁻).

Synthesis of diaryliodonium salts

Essentially three methods were used to prepare the diaryliodonium salts required for this study. These methods are based on the use of an isolated or in-situ generated meta-substituted [hydroxy(tosyloxy)iodo]arene (HTIA) with i) an electron-rich arene, ii) an arylboronic acid^[32] or, iii) a *tri-n*-butylarylstannane^[33] (Scheme 3).

The meta-substituted (diacetoxyiodo)arenes (**3–7**), required to prepare HTIAs as reactive intermediates, were obtained by oxidation of the corresponding iodoarenes with peracetic acid in acetic acid in generally moderate to high yields (16–75%). The required meta-substituted HTIAs were then prepared by treating the appropriate substituted (diacetoxyiodo)arene with tosic acid monohydrate in acetonitrile. The HTIAs bearing a meta Me (**8**), CN (**9**) or CF₃ (**10**) substituent were isolated in high yields (72–91%), and in some examples these were also prepared and used in situ. HTIAs with a meta NO₂ or MeO group were prepared and used in situ only. Regiospecific control was required for the synthesis of unsymmetrically substituted diaryliodonium salts, in which each aromatic ring is differently substituted. The direct treatment of reactive electron-rich arenes, such as anisole, thiophene and 2-methylthiophene, with HTIAs was found to be regiospecific and produced the 4-methoxyphenyl, 2-thienyl and 5-methyl-2-thienyl arylidonium tosylates, respectively. In this manner, we produced the diaryliodonium tosylates **16–21** from isolated HTIAs in moderate to high yields (36–86%) and **22–28** from in-situ generated HTIAs in high yields (81–98%). In the syntheses of other diaryliodonium salt, regioselectivity was conferred by treatment of a substituted arylboronic acid or a substituted aryl *tri-n*-butylarylstannane with the appropriate HTIA.^[32,33] Thus, 3-methylphenyl(phenyl)iodonium tosylate (**11**) was obtained from 3-tolylboronic acid in low but useful yield (20%). *tri-n*-Butylarylstannanes gave the iodonium tosylates **12–15** in low to moderate yields (25–59%). The requisite 3-cyano-*tri-n*-butylstannylbenzene (**1**) and 3-methoxy-*tri-n*-butylstannylbenzene (**2**) were readily obtained from the corresponding iodoarenes with hexabutylditin in moderately high yields (73%). All prepared diaryliodonium salts were stable to long-term storage at 4–5 °C under argon.

Radiochemistry

All radiochemistry was performed in a microfluidic apparatus. We have recently described the construction and operation of this apparatus.^[24] In essence, the apparatus has a coiled silica glass capillary tube micro-reactor with an internal volume of 31.4 μL. This reactor is housed in a heater which can be thermostatted to any set temperature up to 200 °C. The reactor can be infused simultaneously with two solutions, a solution of diaryliodonium salt and a solution of the [¹⁸F]fluoride ion reagent, each from a separate reservoir at a set flow rate. The reaction time is taken to equate to the residence time of reagents in the micro-reactor, which is the internal volume of the reactor divided by its effluent flow rate. Decay-corrected radiochemical yields (RCYs) were estimated from the reverse phase HPLC analysis of reactor effluent based on the percentage of radioactivity in the radiochromatogram represented by radioactive product. All injected radioactivity was found to elute from the HPLC column. The HPLC measurement was subsequently corrected for the proportion of used radioactivity not recovered from the micro-reactor apparatus which was assumed to be adsorbed [¹⁸F]fluoride ion. Radioactivity recovery was moderate and variable (48 ± 27%, *n* = 19). Adsorption is a common phenomenon in reactions of dry no-carrier-added [¹⁸F]fluoride ion, regardless of reactor material (e.g., glass, glassy carbon or platinum).^[34]

The microfluidic apparatus allows radiofluorination reactions to be performed in rapid sequence under very controlled conditions of reagent concentrations, residence time and temperature. Therefore, we tested each reaction under different conditions, in a search for the optimal radiochemical yield of [^{18}F]*m*-fluoroarenes. Usually, between 10 and 14 different conditions were tried, involving several different reaction temperatures. Decay-corrected radiochemical yields are reported for conditions found to give the highest incorporation of [^{18}F]fluoride ion into [^{18}F]fluoroarenes. The diaryliodonium tosylates were adequately soluble in acetonitrile and this was chosen as a suitable polar aprotic reaction medium. Acetonitrile is widely used for reactions with [^{18}F]fluoride ion, particularly aliphatic nucleophilic substitution reactions, since the [^{18}F]fluoride ion is readily solubilized in the presence of K^+ -K 2.2.2 as counter anion. In general, acetonitrile solutions of diaryliodonium salts were infused into the reactor at a concentration of 10 mM and at an infusion rate equal to that of the acetonitrile solution of no-carrier-added [^{18}F]fluoride ion K^+ -K 2.2.2 complex. Thus, the concentration of salt within the micro-reactor was generally 5 mM. These concentrations are similar to those used in our preceding study of the radiofluorination of ortho-substituted diaryliodonium salts.^[24] Reaction times were generally less than 5 minutes.

Micro-reactor effluents were quenched by dilution in acetonitrile-water (1:1 v/v) at room temperature and then analyzed by reverse phase HPLC with radioactivity detection. Radiochromatograms were in all cases simple, showing only separated [^{18}F]fluoride ion at or near the solvent front and either one or two later-eluted [^{18}F]fluoroarenes. The [^{18}F]*m*-fluoroarenes were identified by their comobility in HPLC with reference fluoroarenes.

We first investigated whether the radiofluorination of diaryliodonium salts was a viable route to [^{18}F]fluoroarenes bearing an electron-withdrawing meta substituent (Table 1). The reactions of 12 salts were studied (Scheme 4). These carried a meta CN, NO_2 or CF_3 group in one ring. The other ring was relatively electron-rich phenyl, 3-toluyyl, 4-anisyl, 2-thienyl or 5-Me-2-thienyl, or in one case 3-CN- C_6H_4 in a symmetrical salt.

Radiofluorination of the symmetrical salt (**15**) produced [^{18}F]*m*-fluorobenzonitrile as the only radioactive product in 55% RCY in just over 3 min at 160 °C (Table 1, entry 1). Use of the unsymmetrical salt in which one ring was phenyl (**12**) reduced the RCY of [^{18}F]*m*-fluorobenzonitrile to 25% at 130 °C (Table 1, entry 2). The selectivity for this product was however very high (> 25). When the *m*-cyanophenyl ring was partnered with a more electron-rich aryl ring, substantially improved RCYs (58–82%) of [^{18}F]*m*-fluorobenzonitrile were produced with generally high selectivities (Table 1, entries 3–6). The radiofluorination of diaryliodonium salts having an electron-rich aryl ring (4-MeOC $_6\text{H}_4$, 2-thienyl or 5-Me-2-thienyl) and a phenyl ring bearing a meta NO_2 (**22–24**) or CF_3 group (**19**, **20**, **25**) also produced meta-substituted [^{18}F]fluoroarenes in moderate to quite high RCYs (31–66%) and with high selectivities (Table 1, entries 7–12). RCYs varied somewhat with the nature of the electron-rich ring, especially for the *m*-nitro salts. Overall, these results showed that diaryliodonium tosylates can be effective precursors for the rapid (< 5 min) preparation of [^{18}F]fluoroarenes bearing meta electron-withdrawing substituents. Hence, this method is a useful adjunct to classical aromatic nucleophilic substitution with [^{18}F]fluoride ion in arenes carrying a leaving group in meta orientation to a nitro group.^[35–37]

We next studied the radiofluorination of diaryliodonium tosylates bearing meta electron-donating methyl or methoxy substituents in one ring, and in which the partner ring was electron-rich Ph, 2-thienyl or 4-methoxyphenyl (Table 2). The reactions of 6 salts were studied (Scheme 4). RCYs of [^{18}F]*m*-fluoroarenes and product selectivities greatly depended on the nature of the non-meta substituted electron-rich ring. Where this ring was phenyl as in **11**, the RCY of [^{18}F]*m*-fluorotoluene was low (12%) and less than that of

[¹⁸F]fluorobenzene (15%) (Table 2, entry 1). By contrast, the RCY of [¹⁸F]*m*-fluoroanisole from the phenyl compound **13** was exceptionally high (87%) and co-production of [¹⁸F]fluorobenzene was low (6% RCY) (Table 2, entry 5). Use of 2-thienyl as the electron-rich ring partner gave low RCYs of the target meta-substituted [¹⁸F]fluoroarenes but with acceptable selectivity (Table 2, entries 3 and 6). Use of 4-methoxyphenyl as electron-rich ring partner gave [¹⁸F]*m*-fluorotoluene from **21** in moderately high RCY (47%) and good selectivity (Table 1, entry 2), and gave [¹⁸F]*m*-fluoroanisole from **28** in moderately high RCY (36%) with very high selectivity (Table 2, entry 7). These results show that the treatment of diaryliodonium salts with [¹⁸F]fluoride ion can rapidly produce simple [¹⁸F]*m*-fluoroarenes with electron-donating substituents in impressively high yields under selected conditions. Diaryliodonium tosylates that have 4-methoxyphenyl as one of the aryl rings gave generally moderate to high RCYs with high product selectivity. Thus, the reactions of appropriately designed diaryliodonium salts with [¹⁸F]fluoride ion is a uniquely effective method for the rapid preparation of NCA [¹⁸F]fluoroarenes bearing meta electron-donating groups, such as Me or OMe.

In this study, we did not systematically study the influence of solvent on RCY; the vast majority of reactions were conducted in acetonitrile, which in the microfluidic apparatus could be used well above its boiling point. The RCY of [¹⁸F]*m*-fluoroanisole from the 2-thienyl salt **26** was dramatically improved to 82% when the reaction was conducted in DMF instead of acetonitrile (Table 2, entry 4). Thus, reaction outcomes might well be influenced by solvent (compare entries 4 and 3 in Table 2), and this may warrant a further detailed study. In our previous study on the radiofluorination of ortho-substituted diaryliodonium salts, we mainly used DMF (with trace water) as a solvent with impressive results.^[24]

In a previous study, the free radical scavenger TEMPO, was reported to be beneficial for improved yield and reproducibility in the reactions of fluoride ion with diaryliodonium salts.^[23] As in our previous study of the radiofluorination of ortho substituted diaryliodonium salts with NCA [¹⁸F]fluoride ion in the same microfluidic apparatus,^[24] here we observed high and consistent RCYs in the absence of TEMPO, possibly because photoinitiated decomposition of the salt is avoided.

Conclusions

This study demonstrates that the radiofluorination of unsymmetrical diaryliodonium tosylates (ArI⁺Ar'TsO⁻), in which Ar' is 3-Me-C₆H₄, 4-MeO-C₆H₄, 2-thienyl or 5-Me-2-thienyl, is a rapid and effective method for producing simple NCA [¹⁸F]*m*-fluoroarenes with either electron-withdrawing or electron-donating meta substituents from [¹⁸F]fluoride ion.

Experimental Section

General Methods

¹H NMR spectra were recorded at 400 MHz, ¹³C-NMR at 100 MHz and ¹⁹F-NMR at 376 MHz. No-carrier-added (NCA) [¹⁸F]fluoride ion was obtained through the ¹⁸O(p,n)¹⁸F nuclear reaction by irradiating [¹⁸O]water (95 atom %) for 90–120 min with a proton beam (17 MeV; 20 μA) produced by a PETrace cyclotron. MP-1 or QMA anion exchange cartridges for [¹⁸F]fluoride ion trapping and release were supplied by ORTG (Oakdale, TN).

Precaution—Peracetic acid (32 wt%) is a strong oxidant and may cause explosion in presence of reducing agent and organic materials.

Syntheses of Arylstannanes

3-(Tri-*n*-butylstannyl)benzonitrile (1)—3-Iodobenzonitrile (2.2 mmol, 0.50 g) was mixed with hexabutyltin (2.5 mmol, 1.45 g) and a catalytic amount of tetrakis(triphenylphosphine)palladium (0.5 mol%, 15 mg) in anhydrous toluene (20 mL) under argon. The resulting mixture was refluxed overnight (*ca.* 14 h). Palladium catalyst was filtered off and toluene was removed on a rotary evaporator. The crude oily product was purified by column chromatography on silica gel (10% EtOAc/hexane, R_f = 0.42) to give **1** as a pale yellow oil (0.63 g, 73%); ^1H NMR (CDCl_3): δ 7.72 (t, J = 0.8 Hz, 1H), 7.68 (dt, J = 0.8, 4.4 Hz, 1H), 7.57 (dt, J = 1.6, 5.6 Hz, 1H), 7.40 (t, J = 7.8 Hz, 1H), 1.48–1.54 (m, 6H), 1.27–1.36 (m, 6H), 1.01–1.05 (m, 6H), 0.88 (t, J = 7.2 Hz, 9H); ^{13}C NMR (CDCl_3): δ 144.2, 140.5, 139.6, 131.4, 128.1, 119.5, 112.1, 28.9, 27.3, 13.6, 9.7.

3-(Tri-*n*-butylstannyl)methoxybenzene (2)—3-Iodoanisole (3 mmol, 0.7 g) was mixed with hexabutyltin (3.5 mmol, 2.1 g) and tetrakis(triphenylphosphine)-palladium (0.5 mol %, 17 mg) in anhydrous toluene (20 mL) under argon. The reaction mixture was refluxed overnight (*ca.* 12 h). Palladium catalyst was filtered off and toluene was removed on a rotary evaporator. Crude product was separated by column chromatography on silica gel (10% EtOAc/hexane, R_f = 0.46) to give **2** as a colorless oil (0.87 g, 73%); ^1H NMR (CDCl_3): δ 7.29–7.22 (m, 1H), 7.01–6.96 (m, 2H), 6.85–6.82 (m, 1H), 3.80 (s, 3H), 1.57–1.52 (m, 6H), 1.38–1.28 (m, 6H), 1.07–1.03 (m, 6H), 0.89 (t, J = 7.2 Hz, 9H); ^{13}C NMR (CDCl_3): δ 158.9, 143.4, 128.8, 128.7, 122.1, 112.9, 55.0, 29.1, 27.4, 13.7, 9.6.

Syntheses of (Diacetoxyiodo)arenes

3-(Diacetoxyiodo)toluene (3)—Peracetic acid (33 mmol; 32 wt%, diluted in AcOH, 7 mL) was added dropwise to 3-iodotoluene (8.0 mmol, 1.74 g) at 0 °C. The reaction mixture was then slowly warmed to rt and stirred *ca.* 4 h. H_2O (*ca.* 1 mL) was then added and the white precipitate was filtered off, washed with Et_2O , dried in air to give **3** as a white solid (1.25 g, 46%); mp = 156–159 °C; Lit. mp 150–152 °C^[38]; ^1H NMR (CDCl_3): δ 7.92–7.89 (m, 2H), 7.40–7.38 (m, 2H), 2.43 (s, 3H), 2.01 (s, 6H); ^{13}C NMR (CDCl_3): δ 176.6, 141.6, 135.6, 132.9, 132.3, 130.9, 121.7, 21.6, 20.6.

3-(Diacetoxyiodo)cyanobenzene (4)—Peracetic acid (9.5 mmol, 32 wt.%, diluted in AcOH, 2 mL) was added dropwise to 3-iodobenzonitrile (3.0 mmol, 0.68 g) while cooled in an ice-bath. The reaction mixture was then slowly warmed to rt and stirred overnight (*ca.* 14 h). The resulting white precipitate was filtered off, washed with Et_2O , and dried in air to give **4** as a white solid (0.78 g, 75%); mp = 188–190 °C; Lit. mp 188–189 °C^[39]; ^1H NMR (CDCl_3): δ 8.37 (t, J = 1.4 Hz, 1H), 8.31 (dt, J = 1.2, 6.4 Hz, 1H), 7.89 (dt, J = 1.0, 8.8 Hz, 1H), 7.66 (t, J = 7.8 Hz, 1H), 2.04 (s, 6H); ^{13}C NMR (CDCl_3): δ 176.8, 138.9, 138.1, 137.1, 131.3, 120.9, 116.7, 115.1, 20.4.

3-(Diacetoxyiodo)nitrobenzene (5)—Peracetic acid (33 mmol; 32 wt%, diluted in AcOH, 7 mL) was added dropwise with vigorous stirring to 1-iodo-3-nitrobenzene (7.0 mmol, 1.74 g) cooled in an ice-bath. The reaction mixture was stirred below 5 °C for 1 h and then gradually warmed to rt. Reaction progress was monitored by TLC. After 6 h the white precipitate was filtered off, washed with H_2O (20 mL) followed by Et_2O (20 mL), and dried in air to give **5** as a white solid (1.5 g, 59%); mp = 155–158 °C; Lit. mp 148–150 °C^[39]; ^1H NMR (CDCl_3): δ 8.95 (t, J = 2 Hz, 1H), 8.44 (m, 1H), 8.40 (m, 1H), 7.23 (t, J = 8 Hz, 1H), 2.04 (s, 6H); ^{13}C NMR (CDCl_3): δ 176.8, 148.7, 140.4, 131.5, 130.1, 126.3, 120.6, 20.3.

3-(Diacetoxyiodo)trifluoromethylbenzene (6)—The procedure described for **5** was applied. 1-iodo-3-(trifluoromethyl)benzene (10 mmol, 2.72 g) with peracetic acid (23.7 mmol; 32 wt%, diluted in AcOH, 5 mL) gave **6** as a white solid (1.69 g, 43%); mp = 138–

139 °C; Lit. mp 146–147 °C^[40]; ¹H NMR (CDCl₃): δ 8.33 (s, br, 1H), 8.28 (d, *J* = 8.4 Hz, 1H), 7.85 (dd, *J* = 0.4, 8 Hz, 1H), 7.65 (t, *J* = 8 Hz, 1H), 2.03 (s, 6H); ¹³C NMR (CDCl₃): δ 176.6, 138.1, 133.0 (q, *J* = 34 Hz), 131.8 (q, *J* = 4 Hz), 131.2, 131.0, 128.5 (q, *J* = 4 Hz), 124.2, 121.4, 121.0, 20.3; ¹⁹F NMR (CDCl₃): δ –62.8.

3-(Diacetoxyiodo)methoxybenzene (7)—Peracetic acid (33 mmol; 32 wt %, diluted in AcOH, 7 mL) was added dropwise to 3-iodoanisole (15 mmol, 3.51 g) in an ice-salt bath (*ca.* –10 °C). The reaction mixture was gradually warmed to rt and became a strong yellow solution over 1 h. H₂O (10 mL) was added to quench the reaction and the aqueous mixture was extracted with CH₂Cl₂ (10 mL × 3). The organic layer was dried over MgSO₄ and concentrated *in vacuo*. The resulting yellow oil was triturated with Et₂O, and the generated white solid was filtered off, washed with Et₂O, and dried in air to give **7** as a pale yellow solid (0.84 g, 16%); mp = 128–130 °C; Lit. mp 129–131 °C^[38]; ¹H NMR (CDCl₃): δ 7.68–7.64 (m, 2H), 7.42 (t, *J* = 8.4 Hz, 1H), 7.11 (dq, *J* = 0.8, 2.8 Hz, 1H), 3.87 (s, 3H), 2.02 (s, 6H); ¹³C NMR (CDCl₃): δ 176.5, 160.5, 131.6, 127.1, 121.4, 120.5, 118.0, 55.8, 20.4.

Synthesis of [Hydroxy(tosyloxy)iodo]arenes

3-[Hydroxy(tosyloxy)iodo]toluene (8) was prepared using a published method^[41] (0.74 g, 72%); mp = 126–129 °C; ¹H NMR (DMSO-*d*₆): δ 9.75 (s, br., 1H), 8.06–8.00 (m, 2H), 7.51–7.47 (m, 4H), 7.12 (dd, *J* = 0.4, 8.4 Hz, 2H), 2.38 (s, 3H), 2.29 (s, 3H); ¹³C NMR (DMSO-*d*₆): δ 145.4, 141.1, 137.8, 134.7, 133.1, 131.7, 130.8, 128.1, 125.5, 123.4, 20.8, 20.7.

3-[Hydroxy(tosyloxy)iodo]cyanobenzene (9)—*p*-TsOH·H₂O (1.1 mmol, 0.2 g) in MeCN (10 mL) was added portion-wise to a solution of **4** (1.0 mmol, 0.35 g) in hot MeCN (10 mL) to give a yellow solution. A white solid precipitated out when the solution was cooled to rt. The precipitate was filtered off, washed with Et₂O (20 mL), and dried to give **4** as a slightly yellow solid (0.328 g, 79%); mp = 142–144 °C; ¹H NMR (DMSO-*d*₆): δ 9.98 (s, br., 1H), 8.72 (s, 1H), 8.50 (d, *J* = 8 Hz, 1H), 8.16 (d, *J* = 7.6 Hz, 1H), 7.80 (t, *J* = 7.6 Hz, 1H), 7.47 (d, *J* = 7.6 Hz, 2H), 7.12 (d, *J* = 7.6 Hz, 2H), 2.29 (s, 3H); ¹³C NMR (DMSO-*d*₆): δ 145.3, 138.7, 137.9, 137.5, 135.9, 131.9, 128.2, 125.5, 123.4, 117.2, 113.5, 20.8; HRMS [M–OTs]⁺ calcd for C₇H₅INO: 245.9416, found: 245.9411.

3-[Hydroxy(tosyloxy)iodo]trifluoromethylbenzene (10) was prepared using a published method^[42] (1.67 g, 91%); mp = 154–156 °C; Lit. mp 158–162 °C^[42]; ¹H NMR (DMSO-*d*₆): δ 9.94 (s, br., 1H), 8.60 (s, 1H), 8.49 (d, *J* = 8 Hz, 1H), 8.06–8.04 (m, 1H), 7.83 (t, *J* = 7.6 Hz, 1H), 7.49 (dd, *J* = 1.6, 6.4 Hz, 2H), 7.13 (d, *J* = 8 Hz, 2H), 2.29 (s, 3H); ¹³C NMR (DMSO-*d*₆): δ 145.0, 138.1, 132.1, 131.1, 130.7 (q, *J* = 3 Hz), 128.9 (q, *J* = 3 Hz), 128.2, 125.5, 124.5, 123.7, 121.8, 20.8; ¹⁹F NMR (DMSO-*d*₆): δ –61.2. HRMS [M–OTs]⁺ calcd for C₇H₅F₃IO: 288.9337, found: 288.9345.

Synthesis of Diaryliodonium Salts

By reaction of [Hydroxy(tosyloxy)iodo]arenes with an Arylboronic Acid^[32]

3-Methylphenyl(phenyl)iodonium tosylate (11): 3-Tolylboronic acid (4 mmol, 0.54 g) was added portion-wise to a suspension of Koser's reagent (4.0 mmol, 1.57 g) in CH₂Cl₂ (20 mL) at 0 °C. The resulting mixture became a yellow solution over 30 min at rt. CH₂Cl₂ was removed *in vacuo* and the resulting brown oil was treated with Et₂O (20 mL). The resulting solid was filtered off, washed with Et₂O, air-dried and recrystallized from MeOH/Et₂O to give **11** as a white solid (0.36 g, 20%); mp = 182–184 °C; Lit. mp 169–170 °C^[43]; ¹H NMR (CDCl₃): δ 7.94 (d, *J* = 7.6 Hz, 2H), 7.77 (s, 1H), 7.72 (d, *J* = 8 Hz, 1H), 7.51–7.48 (m, 3H), 7.51–7.27 (m, 3H), 7.21 (t, *J* = 7.6 Hz, 1H), 7.02 (d, *J* = 7.6 Hz, 2H), 2.30 (s, 3H), 2.28 (s, 3H); ¹³C NMR (CDCl₃): δ 142.2, 139.3, 135.5, 135.1, 132.5, 132.2, 131.6, 131.5, 131.3,

128.4, 126.0, 115.3, 115.1, 21.3, 21.2; HRMS $[M-OTs]^+$: calcd for $C_{13}H_{12}I$: 294.9984, found: 294.9986.

By Reaction of [Hydroxy(tosyloxy)i]darenes with Aryltri-n-butylstannanes

3-Cyanophenyl(phenyl)iodonium tosylate (12): The 3-cyano derivative of Koser's reagent (**9**; 1.0 mmol, 0.39 g) was added portion-wise to a solution of **1** (1.0 mmol, 0.39 g) in CH_2Cl_2 (20 mL) at rt. The mixture was refluxed for 4 h. Solvent was removed *in vacuo*, and the resulting yellow oil was triturated with Et_2O (10 mL). The solid was filtered off, washed with Et_2O , dried in air and recrystallized from MeOH/ Et_2O to give **12** as a white solid (0.15 g, 32%); mp = 154–155 °C; 1H NMR (DMSO- d_6): δ 8.86 (s, 1H), 8.58 (d, J = 8.4 Hz, 1H), 8.29 (d, J = 7.6 Hz, 2H), 8.15 (d, J = 8 Hz, 1H), 7.75–7.68 (m, 2H), 7.56 (t, J = 8 Hz, 2H), 7.47 (d, J = 8 Hz, 2H), 7.15 (d, J = 8 Hz, 2H), 2.29 (s, 3H); ^{13}C NMR (DMSO- d_6): δ 139.7, 138.4, 137.9, 137.3, 135.7, 135.3, 132.4, 132.3, 131.9, 128.0, 125.5, 117.0, 115.8, 113.9, 20.8; HRMS $[M-OTs]^+$: calcd for $C_{13}H_9IN$: 305.9780, found: 305.9775.

3-Methoxyphenyl(phenyl)iodonium tosylate (13): The procedure described for **12** was applied. **2** (2.2 mmol, 0.87 g) with Koser's reagent (2.5 mmol, 0.98 g) gave **13** as a white solid (0.63 g, 59%); mp = 136–139 °C; Lit. mp 147–152 °C^[33]; 1H NMR ($CDCl_3$): δ 7.96 (dd, J = 1.2, 8.4 Hz, 2H), 7.59 (t, J = 2 Hz, 1H), 7.50–7.47 (m, 3H), 7.41 (dd, J = 0.4, 7.2 Hz, 1H), 7.33 (t, J = 7.6 Hz, 2H), 7.21 (t, J = 8.4 Hz, 1H), 7.02–6.97 (m, 3H), 3.73 (s, 3H), 2.30 (s, 3H); ^{13}C NMR ($CDCl_3$): δ 141.6, 138.3, 134.2, 130.9, 130.5, 127.4, 125.9, 124.9, 119.1, 117.7, 114.4, 114.3, 54.8, 20.3; HRMS $[M-OTs]^+$: calcd for $C_{13}H_{12}IO$: 310.9933, found: 310.9926.

By Reaction of Isolated HTIAs with Aryltri-n-butylstannanes

3-Cyanophenyl(3'-methylphenyl)iodonium tosylate (14): Isolated HTIA **8** (1.7 mmol, 0.69 g) was added portion-wise to a solution of **1** (1.7 mmol, 0.70 g) in CH_2Cl_2 (20 mL). The reaction mixture was refluxed for 4 h and solvent was removed *in vacuo*. The resulting crude oil was triturated with Et_2O and the generated solid washed with Et_2O and dried in air to give **14** as a white solid (0.21 g, 25%); mp = 185–188 °C; 1H NMR (DMSO- d_6): δ 8.84 (t, J = 1.6 Hz, 1H), 8.57 (dq, J = 0.8, 8.4 Hz, 1H), 8.15–8.13 (m, 2H), 8.09 (dd, J = 0.4, 8 Hz, 2H), 7.72 (t, J = 8 Hz, 1H), 7.51–7.42 (m, 4H), 7.11 (d, J = 7.6 Hz, 2H), 2.35 (s, 3H), 2.29 (s, 3H); ^{13}C NMR (DMSO- d_6): δ 145.8, 141.9, 139.6, 138.4, 137.5, 135.6, 135.4, 133.0, 132.4, 132.3, 131.6, 128.0, 125.5, 116.9, 116.7, 116.5, 113.8, 20.8; HRMS $[M-OTs]^+$: calcd for $C_{14}H_{11}IN$: 319.9936, found: 319.9931.

Bis(3-cyanophenyl)iodonium tosylate (15): The procedure described for **14** was applied to **1** (1.0 mmol, 0.39 g) with **9** (1.0 mmol, 0.42 g) to give **15** as a white solid (0.63 g, 59%); mp = 219–220 °C; 1H NMR (DMSO- d_6): δ 8.86 (t, J = 1.6 Hz, 2H), 8.60 (dq, J = 0.8, 5.6 Hz, 2H), 8.17 (dd, J = 1.2, 6 Hz, 2H), 7.75 (t, J = 8 Hz, 2H), 7.46 (d, J = 8 Hz, 2H), 7.11 (d, J = 8 Hz, 2H), 2.29 (s, 3H); ^{13}C NMR (DMSO- d_6): δ 139.8, 138.6, 137.6, 135.9, 132.5, 128.0, 125.5, 117.0, 116.9, 114.0, 20.8; HRMS $[M-OTs]^+$: calcd for $C_{14}H_8N_2I$: 330.9732, found: 330.9734.

By Reaction of Isolated HTIAs with Electron-rich Arenes

3-Cyanophenyl(4'-methoxyphenyl)iodonium tosylate (16): A mixture of anisole (0.5 mL, 4.6 mmol) and **9** (0.25 mmol, 0.11 g) in chloroform (10 mL) was refluxed for 4 h. After concentration of solvent under vacuum, the resulting oil was treated with Et_2O . The generated solid was filtered off, washed with Et_2O and dried in air to give **16** as a white solid (69 mg, 56%); mp = 172–175 °C; 1H NMR (DMSO- d_6): δ 8.80 (s, 1H), 8.52 (d, J = 8 Hz, 1H), 8.20 (d, J = 8.8 Hz, 2H), 8.12 (d, J = 7.6 Hz, 1H), 7.71 (t, J = 8 Hz, 1H), 7.47 (d, J = 8 Hz, 2H), 7.10 (t, J = 6.8 Hz, 4H), 3.80 (s, 3H), 2.28 (s, 3H); ^{13}C NMR (DMSO- d_6): δ

162.1, 145.7, 139.4, 138.1, 137.6, 137.4, 135.5, 132.3, 128.0, 125.5, 117.6, 117.1, 117.0, 113.7, 105.8, 55.7, 20.8; HRMS [M-OTs]⁺: calcd for C₁₄H₁₁INO: 335.9885, found: 335.9894.

3-Cyanophenyl(2'-thienyl)iodonium tosylate (17): The procedure described for **16** was applied. Thiophene (50 mg, 6 mmol) with **9** (0.50 mmol, 0.21 g) gave **17** as a white solid (0.19 g, 78%); mp = 151–152 °C; ¹H NMR (CDCl₃): δ 8.29 (d, *J* = 8.4 Hz, 1H), 8.12 (t, *J* = 1.2 Hz, 1H), 7.87 (dd, *J* = 1.2, 4 Hz, 1H), 7.67 (d, *J* = 1.2 Hz, 1H), 7.56 (dd, *J* = 1.2, 4 Hz, 1H), 7.42–7.35 (m, 3H), 7.05–6.99 (m, 3H); ¹³C NMR (CDCl₃): δ 141.0, 140.1, 138.7, 137.2, 136.2, 134.5, 131.6, 129.6, 128.7, 125.8, 118.9, 116.5, 114.9, 100.0, 21.3; HRMS [M-OTs]⁺: calcd for C₁₁H₇NSI: 311.9344, found: 311.9352.

3-Cyanophenyl(5'-methyl-2'-thienyl)iodonium tosylate (18): The procedure described for **16** was applied. 2-Methylthiophene (0.7 mmol, 68 mg) with **9** (0.50 mmol, 0.21 g) gave **18** as a white solid (90 mg, 36%); mp = 146–148 °C; ¹H NMR (DMSO-*d*₆): δ 8.23 (s, 1H), 8.57 (d, *J* = 7.6 Hz, 1H), 8.12 (d, *J* = 6.8 Hz, 1H), 7.95 (s, 1H), 7.72 (t, *J* = 7.2 Hz, 1H), 7.47 (d, *J* = 7.2 Hz, 2H), 7.12 (d, *J* = 6.8 Hz, 2H), 6.92 (s, 1H), 2.56 (s, 3H), 2.29 (s, 3H); ¹³C NMR (DMSO-*d*₆): δ 151.7, 145.5, 141.3, 139.1, 137.8, 135.6, 132.3, 128.3, 128.1, 125.5, 119.7, 117.0, 113.6, 97.4, 20.8, 15.0; HRMS [M-OTs]⁺: calcd for C₁₂H₉NSI: 325.9500, found: 325.9503.

3-Trifluoromethylphenyl(4'-methoxyphenyl)iodonium tosylate (19): The procedure described for **16** was applied. Anisole (2.7 mmol, 0.3 mL) in chloroform (2 mL) with **10** (1.5 mmol, 0.69 g) gave **19** as a white solid (0.70 g, 85%); mp = 174–175 °C; ¹H NMR (CDCl₃): δ 8.17 (d, *J* = 8 Hz, 1H), 8.03 (s, 1H), 7.84 (dd, *J* = 2, 6.8 Hz, 1H), 7.59 (d, *J* = 8 Hz, 1H), 7.36–7.30 (m, 3H), 6.91 (d, *J* = 8 Hz, 1H), 6.74 (dd, *J* = 2, 7.2 Hz, 2H), 3.71 (s, 3H), 2.22 (s, 3H); ¹³C NMR (CDCl₃): δ 162.4, 142.0, 139.6, 138.3, 137.6, 133.0 (q, *J* = 34 Hz), 131.6, 131.1 (q, *J* = 4 Hz), 128.5, 127.9 (q, *J* = 4 Hz), 125.8, 121.4, 117.4, 116.3, 104.6, 55.5, 21.2; ¹⁹F NMR (CDCl₃): δ -62.7; HRMS [M-OTs]⁺: calcd for C₁₄H₁₁F₃IO: 378.9807, found: 378.9804.

3-Trifluoromethylphenyl(2'-thienyl)iodonium tosylate (20): The procedure described for **16** was applied. Thiophene (6.0 mmol, 0.5 mL) with **10** (1.5 mmol, 0.69 g) at rt (*ca.* 14 h) gave **20** as a white solid (0.68 g, 86%); mp = 151–152 °C; ¹H NMR (CDCl₃): δ 8.24 (d, *J* = 8.4 Hz, 1H), 8.15 (s, 1H), 7.85 (dd, *J* = 1.2, 3.6 Hz, 1H), 7.66 (d, *J* = 8 Hz, 1H), 7.53 (dd, *J* = 1.2, 5.2 Hz, 1H), 7.42 (t, *J* = 7.6 Hz, 1H), 7.32 (d, *J* = 8.4 Hz, 2H), 7.00–6.97 (m, 3H), 2.29 (s, 3H); ¹³C NMR (CDCl₃): δ 141.7, 140.8, 139.8, 138.0, 136.0, 133.0 (q, *J* = 31 Hz), 131.6, 130.9 (q, *J* = 3 Hz), 129.5, 128.5, 128.0 (q, *J* = 4 Hz), 125.7, 121.2, 118.8, 100.0, 21.2; ¹⁹F NMR (CDCl₃): δ -62.8; HRMS [M-OTs]⁺: calcd for C₁₁H₇F₃IS: 354.9265, found: 354.9254.

3-Methylphenyl(4'-methoxyphenyl)iodonium tosylate (21): The procedure described for **16** was applied. Anisole (2.7 mmol, 0.3 mL) with **8** (0.7 mmol, 0.28 g) gave **21** as a white solid (0.70 g, 85%); mp = 122–126 °C; ¹H NMR (CDCl₃): δ 7.88 (d, *J* = 9.2 Hz, 2H), 7.75 (s, 1H), 7.70 (d, *J* = 8 Hz, 1H), 7.50 (d, *J* = 8 Hz, 2H), 7.27–7.16 (m, 2H), 7.02 (d, *J* = 8 Hz, 2H), 6.81 (d, *J* = 8.4 Hz, 2H), 2.30 (s, 3H), 2.26 (s, 3H); ¹³C NMR (CDCl₃): δ 162.2, 142.4, 142.0, 139.4, 137.3, 135.0, 132.3, 131.6, 131.2, 128.4, 126.0, 117.3, 115.5, 103.6, 55.5, 21.2; HRMS [M-OTs]⁺: calcd for C₁₄H₁₄IO: 325.0089, found: 325.0078.

By Reaction of In-situ Generated HTIAs with Electron-rich Arenes

3-Nitrophenyl(4'-methoxyphenyl)iodonium tosylate (22): *p*-TsOH·H₂O (0.50 mmol, 95 mg) was added to a suspension of **5** (0.50 mmol, 0.18 g) in MeCN (5 mL). A yellow

solution developed instantly, and then a precipitate. Chloroform (15 mL) was added, followed by anisole (2.7 mmol, 0.3 mL). The mixture was refluxed for 3 h. MeCN-chloroform was then removed *in vacuo* and the resulting brown oil was triturated with Et₂O (10 mL). The generated solid was filtered off, washed with Et₂O (20 mL), and dried in air to give **22** as a white solid (0.23 g, 87%); mp = 167–169 °C; Lit. mp 175–180 °C^[43]; ¹H NMR (DMSO-*d*₆): δ 9.13 (s, 1H), 8.61 (d, *J* = 8 Hz, 1H), 8.44 (d, *J* = 8 Hz, 1H), 8.27 (d, *J* = 9.2 Hz, 2H), 7.79 (t, *J* = 8 Hz, 1H), 7.47 (d, *J* = 8 Hz, 2H), 7.12–7.09 (m, 4H), 3.80 (s, 3H), 2.29 (s, 3H); ¹³C NMR (DMSO-*d*₆): δ 162.2, 148.2, 145.6, 140.9, 137.7, 132.7, 129.5, 128.1, 126.5, 125.5, 117.6, 116.9, 105.9, 55.7, 20.8; HRMS [M–OTs]⁺: calcd for C₁₃H₁₁INO₃: 355.9784, found: 355.9778.

3-Nitrophenyl(2'-thienyl)iodonium tosylate (23): The procedure described for **22** was applied. **5** (0.50 mmol, 0.18 g) with thiophene (1.2 mmol, 0.1 mL) in the presence of *p*-TsOH·H₂O (0.50 mmol, 95 mg) gave **23** as a white solid (0.21 g, 82%); mp = 143–146 °C; ¹H NMR (DMSO-*d*₆): δ 9.18 (s, 1H), 8.66 (d, *J* = 7.6 Hz, 1H), 8.44 (d, *J* = 8 Hz, 1H), 8.17 (d, *J* = 3.2 Hz, 1H), 8.00 (d, *J* = 4.8 Hz, 1H), 7.81 (t, *J* = 8.4 Hz, 1H), 7.46 (d, *J* = 7.6 Hz, 2H), 7.21 (t, *J* = 3.6 Hz, 1H), 7.11 (d, *J* = 7.6 Hz, 2H), 2.29 (s, 3H); ¹³C NMR (DMSO-*d*₆): δ 148.2, 145.6, 141.1, 140.6, 137.9, 137.7, 132.7, 129.8, 129.3, 128.1, 126.7, 125.5, 119.2, 101.3, 20.8; HRMS [M–OTs]⁺: calcd for C₁₀H₇INO₂S: 331.9242, found: 331.9242.

3-Nitrophenyl(5'-methyl-2'-thienyl)iodonium tosylate (24): The procedure described for **22** was applied. **5** (0.50 mmol, 0.18 g) with 2-methylthiophene (0.7 mmol, 68 mg) in the presence of *p*-TsOH·H₂O (0.50 mmol, 95 mg) to give **24** as a white solid (0.23 g, 90%); mp = 142–144 °C; ¹H NMR (DMSO-*d*₆): δ 9.15 (s, 1H), 8.64 (d, *J* = 7.6 Hz, 1H), 8.43 (d, *J* = 8 Hz, 1H), 7.99 (d, *J* = 3.6 Hz, 1H), 7.80 (t, *J* = 8 Hz, 1H), 7.46 (d, *J* = 7.6 Hz, 2H), 7.11 (d, *J* = 7.6 Hz, 2H), 6.93 (d, *J* = 2.4 Hz, 1H), 2.56 (s, 3H), 2.29 (s, 3H); ¹³C NMR (DMSO-*d*₆): δ 151.8, 148.2, 145.5, 141.5, 140.5, 137.7, 132.7, 129.2, 128.3, 128.1, 126.7, 125.5, 119.4, 20.8, 15.0; HRMS [M–OTs]⁺: calcd for C₁₁H₉NO₂SI: 345.9399, found: 345.9409.

3-Trifluoromethylphenyl(5'-methyl-2'-thienyl)iodonium tosylate (25): The procedure described for **22** was applied. **6** (2.0 mmol, 0.78 g) with 2-methylthiophene (2.5 mmol, 0.24 g) in the presence of *p*-TsOH·H₂O (2.2 mmol, 0.42 g) gave **25** as a white solid (1.03 g, 98%); mp = 140–142 °C; ¹H NMR (CDCl₃): δ 8.22 (d, *J* = 8.4 Hz, 1H), 8.10 (s, 1H), 7.68–7.63 (m, 2H), 7.75–7.27 (m, 3H), 7.01 (d, *J* = 4.4 Hz, 2H), 6.65 (dd, *J* = 1.2, 3.6 Hz, 1H), 2.53 (s, 3H), 2.31 (s, 3H); ¹³C NMR (CDCl₃): δ 152.0, 141.8, 141.4, 139.7, 137.6, 132.9 (q, *J* = 34 Hz), 131.5, 130.6 (q, *J* = 3 Hz), 128.5, 128.0, 125.8, 123.9, 121.2, 118.9, 21.2, 15.4; ¹⁹F NMR (CDCl₃): δ –62.8; HRMS [M–OTs]⁺: calcd for C₁₂H₉F₃IS: 368.9422, found: 368.9412.

3-Methylphenyl(2'-thienyl)iodonium tosylate (26): The procedure described for **22** was applied. **3** (0.5 mmol, 0.17 g) with thiophene (1.2 mmol, 0.1 mL) in the presence of *p*-TsOH·H₂O (0.5 mmol, 95 mg) gave **26** as a white solid (0.21 g, 90%); mp = 146–147 °C; Lit. mp 151–154 °C^[44]; ¹H NMR (CDCl₃): δ 7.77–7.76 (m, 2H), 7.73 (d, *J* = 8 Hz, 1H), 7.56 (dd, *J* = 1.2, 5.6 Hz, 1H), 7.51 (d, *J* = 8 Hz, 2H), 7.28 (s, 1H), 7.21 (t, *J* = 8 Hz, 1H), 7.06–7.01 (m, 3H), 2.32 (s, 3H), 2.29 (s, 3H); ¹³C NMR (CDCl₃): δ 142.3, 142.2, 140.3, 139.5, 135.9, 134.4, 132.4, 131.2, 131.1, 129.6, 128.5, 126.0, 118.9, 21.4, 21.3; HRMS [M–OTs]⁺: calcd for C₁₁H₁₀IS: 300.9548, found: 300.9539.

3-Methoxyphenyl(2'-thienyl)iodonium tosylate (27): The procedure described for **22** was applied. **7** (1 mmol, 0.35 g) with thiophene (6.0 mmol, 0.5 mL) in the presence of *p*-TsOH·H₂O (1 mmol, 0.19 g) gave **27** as a white solid (0.42 g, 86 %); mp = 155–157 °C; Lit. mp = 161 °C^[45]; ¹H NMR (DMSO-*d*₆): δ 8.08 (dd, *J* = 1.6, 4 Hz, 1H), 7.97 (dd, *J* = 1.6, 7.6 Hz, 1H), 7.94–7.93 (m, 1H), 7.79 (dt, *J* = 0.4, 2 Hz, 1H), 7.79–7.42 (m, 3H), 7.21–7.16 (m,

2H), 7.13 (d, $J = 7.60$ Hz, 2H), 3.79 (s, 3H), 2.29 (s, 3H); ^{13}C NMR (DMSO- d_6): δ 160.3, 145.6, 140.4, 137.7, 132.4, 129.6, 128.1, 126.6, 125.5, 120.1, 119.3, 117.8, 100.9, 55.9, 20.8; HRMS $[\text{M}-\text{OTs}]^+$: calcd for $\text{C}_{11}\text{H}_{10}\text{IOS}$: 316.9497, found: 316.9492.

3-Methoxyphenyl(4'-methoxyphenyl)iodonium tosylate (28): The procedure described for **22** was applied. **7** (0.3 mmol, 0.13 g) with anisole (4.6 mmol, 0.5 mL) in the presence of *p*-TsOH·H₂O (0.4 mmol, 76 mg) gave **28** as a white solid (0.15 g, 81%); mp = 131–133 °C; ^1H NMR (CDCl₃): δ 7.89 (dd, $J = 2$, 6.8 Hz, 2H), 7.58 (t, $J = 2$ Hz, 1H), 7.54 (dd, $J = 1.6$, 6.4 Hz, 2H), 7.41 (dt, $J = 0.8$, 7.2 Hz, 1H), 7.20 (t, $J = 8$ Hz, 1H), 7.21 (d, $J = 8$ Hz, 2H), 6.96 (dd, $J = 0.8$, 2.4 Hz, 1H), 6.83 (dd, $J = 2$, 7.2 Hz, 2H), 3.79 (s, 3H), 3.74 (s, 3H), 2.31 (s, 3H); ^{13}C NMR (CDCl₃): δ 162.4, 161.0, 142.7, 139.3, 137.4, 131.9, 128.5, 126.2, 126.0, 119.5, 118.2, 117.4, 115.7, 103.6, 55.9, 55.6, 21.3; HRMS $[\text{M}-\text{OTs}]^+$: calcd for $\text{C}_{14}\text{H}_{14}\text{IO}_2$: 341.0039, found: 341.0045.

Microfluidic NCA Radiofluorination of Diaryliodonium Salts

The detailed configuration and operation of the microfluidic device are described in our previous publication.^[24,30]

Cyclotron-produced NCA [^{18}F]fluoride ion (100–200 mCi) in [^{18}O]water (225–350 μL) was first adsorbed onto an anion exchange resin (*i.e.*, MP-1 or QMA) cartridge within the CE module of a NanoTek apparatus (Advion), and then released with a solution of K₂CO₃ (0.8 mg; 5 μmol) plus K 2.2.2 (4.5 mg; 11 μmol) in MeCN-H₂O (9: 1 v/v; 150 μL) into a 5-mL V-vial. The solution was dried by two cycles of azeotropic evaporation with MeCN (0.45 mL) at 100 °C. Dry $^{18}\text{F}^-$ -K 2.2.2-K⁺ complex (70–150 mCi) was dissolved in MeCN or DMF (350 μL). A 10 mM-solution of diaryliodonium salt was prepared in the same solvent. Each of the two solutions (255 μL) was loaded into a separate storage loop of the microfluidic apparatus. For radiofluorination reactions, each solution (10–15 μL) was infused simultaneously into 4 m-length coiled silica glass tube microreactor of the apparatus at a set flow rate in the range 5–10 $\mu\text{L}/\text{min}$ and at different temperatures. The reaction product exiting the microreactor was quenched with MeCN-H₂O (1:1 v/v; 1 mL). Amounts of precursor, temperature and flow rates were varied to optimize RCYs. RCYs of [^{18}F]m-fluoroarenes were measured by reverse phase radio-HPLC on a Luna C18 column (250 $\times$ 4.6 mm i.d., 10 μm) eluted at 1.75 mL/min with a gradient of MeCN-H₂O with the percentage of MeCN increased linearly from 60 to 90% over 7 min. Retention times were 4.5, 5.6, 3.3, 4.0, 6.2, 5.4, 4.3, 5.0 and 5.1 min for [^{18}F]fluorobenzene, [^{18}F]3-fluorotoluene, [^{18}F]3-fluorobenzonitrile, [^{18}F]3-fluoronitrobenzene, [^{18}F]1-fluoro-3-(trifluoromethyl)benzene, [^{18}F]3-fluoroanisole, [^{18}F]4-fluoroanisole, [^{18}F]2-fluorothiophene, [^{18}F]2-fluoro-5-methylthiophene, respectively. When the gradient was altered to increase MeCN from 20 to 90% over 11 min, [^{18}F]fluorobenzene and [^{18}F]3-fluoroanisole eluted with retention times of 10.9 and 12.2 min, respectively, and when altered to increase MeCN from 55 to 90% over 7 min, [^{18}F]3-fluoroanisole and [^{18}F]4-fluoroanisole eluted at 5.4 and 5.0 min, respectively. These products were identified by their comobility with the respective reference fluoro compounds.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This research was supported by the Intramural Research Program of the National Institutes of Health (NIMH). The authors are grateful to the NIH Clinical Center PET Department (Chief, Dr. Peter Herscovitch) for the cyclotron production of fluorine-18.

References

1. Phelps ME. Proc. Natl. Acad. Sci., USA. 2000; 97:9226–9233.
2. Wong DF, Gründer G, Brašić JR. Int. Rev. Psychiatry. 2007; 19:541–558. [PubMed: 17896234]
3. Gibson RE, Burns HD, Hamill TG, Eng WS, Francis BE, Ryan C. Curr. Radiopharm. Design. 2000; 6:973–989.
4. Lee CM, Farde L. TIPs. 2006; 27:310–316. [PubMed: 16678917]
5. Burns HD, Hamill TG, Eng W, Francis B, Fioravanti C, Gibson RE. Curr. Opin. Chem. Biol. 1999; 3:388–394.
6. Miller JM, Kumar D, Mann JJ, Parsey RV. Curr. Radiopharm. 2008; 1:12–16.
7. Halldin C, Gulyás B, Langer O, Farde L. Q. J. Nucl. Med. 2001; 45:139–152. [PubMed: 11476163]
8. Ametamey SM, Honer M, Schubiger PA. Chem. Rev. 2008; 108:1501–1516. [PubMed: 18426240]
9. Guilloteau D, Chalon S. Curr. Pharm. Design. 2005; 11:3237–3245.
10. Fowler JS., Ido, T. Handbook of Radiopharmaceuticals, Radiochemistry and Applications. Welch, MJ.; Redvanly, CS., editors. Vol. Ch. 9. Chichester: Wiley; 2003. p. 307-321.
11. Ruth TJ, Wolf AP. Radiochim. Acta. 1979; 26:21–24.
12. Guillaume M, Luxen A, Nebeling B, Argentini M, Clark JC, Pike VW. Appl. Radiat. Isot. 1991; 42:749–762.
13. Cai LS, Lu SY, Pike VW. Eur. J. Org. Chem. 2008; 17:2853–2873.
14. Balz G, Schiemann G. Chem. Ber. 1927; 60:1186–1190.
15. Wallach O. Annalen. 1886; 235:233–255.
16. Coenen, HH. PET Chemistry – The Driving Force in Molecular Imaging. Schubiger, PA.; Lehman, L.; Friebe, M., editors. Vol. Ch. 1. Berlin: Springer-Verlag; 2007. p. 15-50. Ernst Schering Research Foundation Workshop 62.
17. Wuest, F. PET Chemistry – The Driving Force in Molecular Imaging. Schubiger, PA.; Lehman, L.; Friebe, M., editors. Vol. Ch. 2. Berlin: Springer-Verlag; 2007. p. 51-78. Ernst Schering Research Foundation Workshop 62.
18. Knochel A, Zwernemann O. Appl. Radiat. Isot. 1991; 42:1077–1080.
19. Pike VW, Aigbirhio FI. J. Chem. Soc., Chem. Commun. 1995; 21:2215–2216.
20. Shah A, Pike VW, Widdowson DA. J. Chem. Soc., Perkin Trans. 1. 1998:2043–2046.
21. Zhang MR, Kumata K, Suzuki K. Tetrahedron Lett. 2007; 48:8632–8635.
22. Ross L, Ermert J, Hocke C, Coenen HH. J. Am. Chem. Soc. 2007; 129:8018–8025. [PubMed: 17536798]
23. Carroll MA, Nairne J, Smith G, Widdowson DA. J. Fluorine Chem. 2007; 128:127–132.
24. Chun J-H, Lu S, Lee Y-S, Pike VW. J. Org. Chem. 2010; 75:3332–3338. [PubMed: 20361793]
25. Lee Y-S, Hodošček M, Chun J-H, Pike VW. Chem. Eur. J. 2010; 16:10418–10423.
26. Lu SY, Watts P, Chin FT, Hong J, Musachio JL, Briard E, Pike VW. Lab Chip. 2004; 4:523–525. [PubMed: 15570360]
27. Lee CC, Sui GD, Elizarov A, Shu CYJ, Shin YS, Dooley AN, Huang J, Daridon A, Wyatt P, Stout D, Kolb HC, Witte ON, Satyamurthy N, Heath JR, Phelps ME, Quake SR, Tseng HR. Science. 2005; 310:1793–1796. [PubMed: 16357255]
28. Elizarov AM. Lab Chip. 2009; 9:1326–1333. [PubMed: 19417895]
29. Miller PW, deMello AJ, Gee AD. Curr. Radiopharm. 2010; 3:254–262.
30. Lu S, Giamis AM, Pike VW. Curr. Radiopharm. 2009; 2:49–55.
31. Lu S, Pike VW. J. Fluorine Chem. 2010; 131:1032–1038.
32. Carroll MA, Pike VW, Widdowson DA. Tetrahedron Lett. 2000; 41:5393–5396.
33. Pike VW, Butt F, Shah A, Widdowson DA. J. Chem. Soc., Perkin Trans. 1999; 1:245–248.
34. Brodack JW, Kilbourn MR, Welch MJ, Katzenellenbogen JA. Appl. Radiat. Isot. 1986; 37:217–221.
35. Constantinou M, Shah A, Pike VW. Radiofluorinations of *m*-substituted nitrobenzenes. J. Label. Compd. Radiopharm. 2001; 44 Suppl. 1:S889–S891.

36. Lazarova N, Siméon FG, Musachio JL, Lu SY, Pike VW. *J. Label. Compd. Radiopharm.* 2007; 50:463–465.
37. Hamill TG, Krause S, Ryan C, Bonnefous C, Govek S, Seiders TJ, Cosford NDP, Roppe J, Kamenecka T, Patel S, Gibson RE, Sanabria S, Riffel K, Eng WS, King C, Yang X, Green MD, O'Malley SS, Hargreaves R, Burns HD. *Synapse.* 2005; 56:205–216. [PubMed: 15803497]
38. Zielinska A, Skulski L. *Molecules.* 2002; 7:806–809.
39. Kaźmierczak P, Skulski L. *Synthesis.* 1998:1721–1723.
40. Hossain MD, Kitamura T. *Synthesis.* 2005:1932–1934.
41. Carman CS, Koser GF. *J. Org. Chem.* 1983; 48:2534–2539.
42. Nabana T, Togo H. *J. Org. Chem.* 2002; 67:4362–4365. [PubMed: 12054975]
43. Caserio MC, Glusker DL, Roberts JD. *J. Am. Chem. Soc.* 1959; 81:336–342.
44. Margida AJ, Koser GF. *J. Org. Chem.* 1984; 49:3643–3646.
45. Ross TL, Ermert J, Hocke C, Coenen HH. *J. Am. Chem. Soc.* 2007; 129:8018–8025. [PubMed: 17536798]

Scheme 1.

Well-explored approaches to the radiosyntheses of $[^{18}\text{F}]$ fluoroarenes from $[^{18}\text{F}]$ fluoride ion.

Scheme 2.

Synthesis of [^{18}F]fluoroarenes from the radiofluorination of diaryliodonium salts.

Reagents, conditions and yields: i) peracetic acid, AcOH, rt; 4 h for **3** (46%); 14 h for **4** (75%); 6 h for **5** (59%); 6 h for **6** (43%); 1 h for **7** (16%); ii) TsOH·H₂O, MeCN; iii) CHCl₃, reflux, 3 h for **22** (87%), **23** (82%), **24** (90%), **25** (98%), **26** (90%), **27** (86%) and **28** (81%); iv) TsOH·H₂O, MeCN for **8** (72%), **9** (79%) and **10** (91%); v) Sn₂Bu₆, Pd(PPh₃)₄, toluene, reflux, 14 h for **1** (73%) and **2** (73%); vi) CH₂Cl₂, reflux for 4 h for **14** (25%) and **15** (59%); vii) CHCl₃, reflux, 4 h, for **16** (54%), **17** (69%), **18** (36%), **19** (85%), **20** (86%) and **21** (85%); viii) Koser's reagent, CH₂Cl₂; 30 min, rt for **11** (20%); 4 h, reflux for **12** (32%) and **13** (59%).

Scheme 3.

Preparation of meta-substituted diaryliodonium tosylates **11–28**.

Scheme 4.

Preparation of [^{18}F]m-fluoroarenes from NCA radiofluorination of diaryliodonium salts within a micro-reactor.

RCYs of [¹⁸F]*m*-fluoroarenes (Ar¹⁸F) from the NCA radiofluorination of diaryliodonium tosylates bearing a *meta* electron-withdrawing substituent.

Table 1

$$\text{Ar}-\text{I}^+-\text{Ar}' \xrightarrow[\text{MeCN}]{^{18}\text{F}^-} \text{Ar}^{18}\text{F} + \text{Ar}'^{18}\text{F}$$

Entry	Diaryliodonium salt (ArI ⁺ Ar'TsO ⁻)		Temp (°C)	Time ^[a] (s)	RCY of [¹⁸ F]fluoroarene (% ^[b])		Selectivity for Ar ¹⁸ F
	#	Ar	Ar'		Total	Ar ¹⁸ F	Ar' ¹⁸ F
1	15	3-CN-C ₆ H ₄	3-CN-C ₆ H ₄	160	189	55	55
2	12	3-CN-C ₆ H ₄	Ph	130	146	26	25
3	16	3-CN-C ₆ H ₄	4-MeO-C ₆ H ₄	130	146	93	82
4	14	3-CN-C ₆ H ₄	3-Me-C ₆ H ₄	160	189	81	76
5	17	3-CN-C ₆ H ₄	2-thienyl	160	189	60	58
6	18	3-CN-C ₆ H ₄	5-Me-2-thienyl	180	189	79	78
7	22	3-NO ₂ -C ₆ H ₄	4-MeO-C ₆ H ₄	130	191	65	58
8	23	3-NO ₂ -C ₆ H ₄	2-thienyl	130	140	35	31
9	24	3-NO ₂ -C ₆ H ₄	5-Me-2-thienyl	160	189	39	37
10	19	3-CF ₃ -C ₆ H ₄	4-MeO-C ₆ H ₄	190	189	54	53
11	20	3-CF ₃ -C ₆ H ₄	2-thienyl	180	236	66	66
12	25	3-CF ₃ -C ₆ H ₄	5-Me-2-thienyl	190	189	57	57

^[a] Residence time in micro-reactor.

^[b] Decay-corrected, optimized yield chosen from 10–14 different runs.

NA = not applicable.

RCYs of [¹⁸F]*m*-fluoroarenes (Ar¹⁸F) from the NCA radiofluorination of diaryliodonium tosylates bearing a *meta* electron-donating substituent.

Table 2

Entry	Diaryliodonium salt (Ar ⁺ Ar ⁺ TsO ⁻)			Temp (°C)	Time ^[a] (s)	RCY of [¹⁸ F]fluoroarene (%)/[b]		Selectivity for Ar- ¹⁸ F	
	#	Ar	Ar ⁺			Total	Ar- ¹⁸ F		
1	11	3-Me-C ₆ H ₄	Ph	160	236	27	12	15	1
2	21	3-Me-C ₆ H ₄	4-MeO-C ₆ H ₄	180	236	52	47	5	9
3	26	3-Me-C ₆ H ₄	2-thienyl	190	236	27	26	<1	>26
4 ^[c]	26	3-Me-C ₆ H ₄	2-thienyl	200	236	82	82	0	>82
5	13	3-MeO-C ₆ H ₄	Ph	150	236	93	87	6	15
6	27	3-MeO-C ₆ H ₄	2-thienyl	200	314	10	9	1	9
7	28	3-MeO-C ₆ H ₄	4-MeO-C ₆ H ₄	180	236	37	36	1	36

^[a] Residence time in micro-reactor.

^[b] Decay-corrected, optimized yield chosen from 10–14 different runs.

^[c] Reaction performed in DMF